

This is a repository copy of *Elucidation of the genetic basis of variation for stem strength characteristics in bread wheat by Associative Transcriptomics*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/100158/>

Version: Accepted Version

---

**Article:**

Miller, Charlotte N, Harper, Andrea Louise [orcid.org/0000-0003-3859-1152](https://orcid.org/0000-0003-3859-1152), Trick, Martin et al. (3 more authors) (2016) Elucidation of the genetic basis of variation for stem strength characteristics in bread wheat by Associative Transcriptomics. BMC Genomics. ISSN 1471-2164

<https://doi.org/10.1186/s12864-016-2775-2>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

This is an author produced version of *Elucidation of the genetic basis of variation for stem strength characteristics in bread wheat by Associative Transcriptomics*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/100158/>

---

**Article:**

Bancroft, Ian (orcid.org/0000-0001-7707-1171) (2016) Elucidation of the genetic basis of variation for stem strength characteristics in bread wheat by Associative Transcriptomics. BMC Genomics. ISSN 1471-2164 (In Press)

---

[Click here to view linked References](#)

1 **Title: Elucidation of the genetic basis of variation for stem strength characteristics in**  
2 **bread wheat by Associative Transcriptomics**

3 **Running title: Genetic control of stem strength in wheat**

4 Charlotte N. Miller<sup>1\*</sup>, Andrea L. Harper<sup>1†\*</sup>, Martin Trick<sup>1</sup>, Peter Werner<sup>2</sup>, Keith Waldron<sup>3</sup>, Ian  
5 Bancroft<sup>1†</sup>

6  
7 <sup>1</sup>John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK

8 <sup>2</sup>KWS UK Ltd., 56 Church Street, Thriplow, Hertfordshire SG8 7RE, UK

9 <sup>3</sup>Institute of Food Research, Norwich Research Park, Norwich, NR4 7UH, UK

10 † Present Address: Department of Biology, University of York, York, YO10 5DD, UK

11 \*Authors contributed to the manuscript equally

12  
13 **Corresponding Author Address:**

14 Prof. Ian Bancroft  
15 Department of Biology  
16 University of York  
17 Heslington  
18 York  
19 YO10 5DD  
20 Email: [ian.bancroft@york.ac.uk](mailto:ian.bancroft@york.ac.uk)  
21 Tel: +44 (0) 1904 328778

22  
23 **Authors**

24 Charlotte N. Miller [charlotte.miller@jic.ac.uk](mailto:charlotte.miller@jic.ac.uk)  
25 Andrea L. Harper [andrea.harper@york.ac.uk](mailto:andrea.harper@york.ac.uk)  
26 Martin Trick [martin.trick@jic.ac.uk](mailto:martin.trick@jic.ac.uk)  
27 Peter Werner [Peter.Werner@kws.com](mailto:Peter.Werner@kws.com)  
28 Keith Waldron [keith.waldron@ifr.ac.uk](mailto:keith.waldron@ifr.ac.uk)  
29 Ian Bancroft [ian.bancroft@york.ac.uk](mailto:ian.bancroft@york.ac.uk)

31 **Abstract**

32 **Background**

33 The current approach to reducing the tendency for wheat grown under high fertilizer  
34 conditions to collapse (lodge) under the weight of its grain is based on reducing stem height  
35 via the introduction of *Rht* genes. However, these reduce the yield of straw (itself an  
36 important commodity) and introduce other undesirable characteristics. Identification of  
37 alternative height-control loci is therefore of key interest. In addition, the improvement of  
38 stem mechanical strength provides a further way through which lodging can be reduced.

39 **Results**

40 To investigate the prospects for genetic alternatives to *Rht*, we assessed variation for plant  
41 height and stem strength properties in a training genetic diversity panel of 100 wheat  
42 accessions fixed for *Rht*. Using mRNAseq data derived from RNA purified from leaves,  
43 functional genotypes were developed for the panel comprising 42,066 Single Nucleotide  
44 Polymorphism (SNP) markers and 94,060 Gene Expression Markers (GEMs). In the first  
45 application in wheat of the recently-developed method of Associative Transcriptomics, we  
46 identified associations between trait variation and both SNPs and GEMs. Analysis of  
47 marker-trait associations revealed candidates for the causative genes underlying the trait  
48 variation, implicating xylan acetylation and the COP9 signalosome as contributing to stem  
49 strength and auxin in the control of the observed variation for plant height. Predictive  
50 capabilities of key markers for stem strength were validated using a test genetic diversity  
51 panel of 30 further wheat accessions.

52 **Conclusions**

53 This work illustrates the power of Associative Transcriptomics for the exploration of complex  
54 traits of high agronomic importance in wheat. The careful selection of genotypes included in  
55 the analysis, allowed for high resolution mapping of novel trait-controlling loci in this staple  
56 crop. The use of Gene Expression markers coupled with the more traditional sequence-  
57 based markers, provides the power required to understand the biological context of the  
58 marker-trait associations observed. This not only adds to the wealth of knowledge that we  
59 strive to accumulate regarding gene function and plant adaptation, but also provides  
60 breeders with the information required to make more informed decisions regarding the  
61 potential consequences of incorporating the use of particular markers into future breeding  
62 programmes.

63

64 **Keywords:** Modulus of Rupture - lodging - Associative Transcriptomics - xylan acetylation -  
65 COP9 signalosome - auxin

66

## 67 **Background**

68 Lodging is defined as the permanent displacement of a crop from its usually vertical growth  
69 habit. This phenomenon may be divided into two main categories: lodging caused by  
70 anchorage failure, or root lodging; and lodging caused by stem mechanical failure, also  
71 known as brackling or stem lodging. Lodging is a complex trait, influenced by environmental,  
72 agronomic and genetic factors and continues to be a widespread problem in wheat grown  
73 worldwide. In years where lodging is particularly severe, yield losses as great as 80% can be  
74 expected [1].

75 Previous efforts to reduce the occurrence of lodging in wheat have centred on reducing the  
76 height of plants through incorporation of semi-dwarfing alleles into accessions and the use of  
77 plant growth regulators (PGR). The most common semi-dwarfing genes found in modern  
78 wheat accessions are the GA-insensitive *Rht-B1* and *Rht-D1*, which markedly increased the  
79 yield potential of wheat following their introduction [2]. However, these genes may not be  
80 beneficial under some environmental conditions, and efforts to identify other semi-dwarfing  
81 genes with different physiological functions are ongoing. Another potential strategy is to  
82 breed accessions with increased mechanical strength in the plant stems. While stem  
83 mechanical strength is considered an important agronomic trait, few studies have focused on  
84 the identification of genetic markers for this trait which may be utilised in marker-assisted  
85 breeding. Furthermore, the few mapping studies that have been conducted with this aim  
86 have been limited by low marker density and mapping resolution through the utilisation of the  
87 traditional bi-parental cross, QTL analysis approach [3, 4].

88 In recent years we have seen the successful application of GWAS in a number of different  
89 plant species[5-7]. This method makes use of historical recombination events which, when  
90 coupled with high marker density, provides increased mapping resolution. Furthermore,  
91 recent advances have expanded this powerful mapping approach to combine the exploration  
92 of marker variation at both the sequence and gene expression level in a method termed,  
93 Associative Transcriptomics (AT) [8].The ordered transcriptome resource necessary for  
94 implementation of AT in hexaploid wheat has been established [9] .

95 Our aim was to explore the variation available in European wheat breeding material for both  
96 height and stem mechanical strength, and in the first application of AT in wheat, to identify  
97 molecular markers associated with this variation. This will provide breeders with both

98 insights into the bases of variation for these traits and molecular markers to underpin  
99 marker-assisted breeding of wheat accessions with improved lodging resistance.

100

## 101 **Methods**

### 102 ***Plant material and phenotyping***

103 A panel of 100 European accessions of hexaploid bread wheat, *Triticum aestivum*  
104 (Supplementary Data. 1), was grown at a single site (KWS, Thriplow, UK) across two years  
105 (2011 and 2012). In 2012, prior to harvesting, a subset of accessions was screened for stem  
106 lodging risk. Using a pulley system attached to the base of the ear of each plant tested (Fig.  
107 1c), stems were pulled through a reproducible arc path (to ground level), a similar motion to  
108 that which would be induced by wind or heavy rainfall. Following this, any stem mechanical  
109 failure induced by the bending of the stem was recorded. Stems found to suffer stem  
110 breakage were scored with a “1” and those for which no mechanical failure was observed  
111 were scored with a “0”. This experiment was performed for 6 plants per accession and a  
112 mean “breakage score” determined.

113 Harvesting of material was carried out by hand, cutting the stems as close to the soil as  
114 possible using secateurs. Ten plants were harvested for each accession in 2011 and five  
115 plants per accession in 2012. Prior to further processing, all plants were dried thoroughly at  
116 room temperature. Any plants showing signs of stem tissue damage were excluded from the  
117 study. To allow for an in-depth analysis of the relationship between plant morphology, stem  
118 structure and stem mechanical strength, the following measurements were determined for  
119 each plant harvested: Plant height; main stem (determined as the tallest) weight and  
120 threshed weight; length of the second internode (from plant base) and stem width (measured  
121 using digital callipers at the midway point of the second internode). A 5cm section was then  
122 removed from the 2<sup>nd</sup> internode of the main stem using a scalpel. The basal end of this  
123 section was marked using a permanent marker pen. To obtain stem cross-sectional  
124 measurements (required for the later calculation of stem second moment of area ( $I$ )), the  
125 transverse of the marked stem end was photographed. All images were later analysed using  
126 the digital analysis software Sigma Scan (Stystat Software Inc., San Jose, USA), allowing  
127 the following cross-sectional measurements to be determined: whole stem area (used in the  
128 later calculation of  $D2$ ); stem hollow area (used in the later calculation of  $d1a$ ); the area of  
129 stem parenchyma and the thickness of the stem outer cortex. Following these initial  
130 measurements, all samples were stored at 55% relative humidity at 23<sup>o</sup>C for a minimum of 2  
131 days in a silica chamber to ensure equilibration of moisture content between samples.

132 Mechanical testing of the material was carried out using a Texture Analyser (TA) (Analyser  
133 (TA-XT2®- Stable Microsystems, Godalming, UK) with a three-point bend test setup (Institute  
134 of Food Research, Norwich, UK) (Fig. 1a). These methods were adapted from Kern et al  
135 (2005) [10]. The TA was fitted with a load cell with maximum loading capacity of 5kg. The  
136 support stands were set at 2.5cm apart (across which the 5cm stem sample was placed) and  
137 the testing probe was set to move at a constant speed of 2mm/sec. The TA, connected to a  
138 computer, produces a real-time graphical output, representing the mechanical profile of the  
139 stem sample being tested. From this graph,  $F_{max}$ , the absolute resistance of the stem  
140 sample to break under-load, and  $F/V$ , the resistance of the stem sample to bend elastically,  
141 were obtained (Fig. 1b). These are 'absolute strength measures', being the result of a  
142 combination of both strength due to structure and material strength. These absolute  
143 measures of strength, together with the stem sample second moment of area ( $I$ ) (Eq. (1)),  
144 were used in calculating the material strength of the stem samples: the Modulus of Rupture  
145 ( $MOR$ ), describing the resistance of the stem material to break under-load (Eq. (3)) and the  
146 Modulus of Elasticity ( $MOE$ ) describing the resistance of the stem material to bend elastically  
147 (Eq. (2)).

## 2.2 Equations

Equation 1

$$I = \pi (D2^4 - D1a^4) / 64$$

Where:

$D2$  = diameter of whole stem calculated from stem cross-sectional area

$D1a$  = diameter of stem hollow calculated from stem hollow area

Equation 2

$$MOR = (F_{max} * a * D2) / I$$

Equation 3

$$MOE = (F/V) * (a^2/12) * (3L-4a) / I$$

161 Where:

162  $L$  = the length of the stem sample between the two supports

163  $a = L/2$

164

### 165 **Statistical analysis of data**

166 Following the assessment of year by year interactions, traits were assessed for significant  
167 genotypic variation and REML-predicted means calculated for use in the consequent  
168 correlation (Genstat 15<sup>th</sup> edition) and Associative Transcriptomics analyses. These statistical  
169 analyses were carried out using Genstat 15<sup>th</sup> edition (VSN International, Hemel Hempstead,  
170 UK).

171

### 172 **mRNA-seq and marker scoring**

173 For the mRNA-seq, second true leaves from each of four plant replicates per accession were  
174 harvested approximately 14 days after pricking out (21 d after sowing) as close to the  
175 midpoint of the light period as possible, pooled and immediately frozen in liquid nitrogen.  
176 Samples were extracted using the Omega Biotek EZNA Plant RNA Kit according to  
177 manufacturer's instructions.

178 Transcriptome sequence data was then obtained for each of the 100 wheat accessions  
179 included in the training panel. This was achieved using Illumina transcriptome sequencing  
180 (mRNA-seq). Illumina sequencing, quality checking and processing were conducted as  
181 described previously [11] except that, for SNP calling and transcript quantification, 100 base  
182 reads obtained from the HiSeq platform were trimmed in order to retain comparability with 80  
183 base reads generated on GAllx instruments, and capped at 35 million reads to maintain  
184 comparable read depth. Maq was used for mapping with default parameters, meaning that  
185 reads with no more than two mismatches with summed  $Q \geq 70$  were mapped.

186 The alignment of these reads for SNP detection was facilitated by the development of a  
187 reference sequence, as described previously [9]. Briefly, the reference sequence was  
188 generated based on *de novo* transcriptome assemblies of *Triticum urartu*, *Aegilops*  
189 *speltoides* and *Aegilops tauschii* (representing the A, B and D genomes, respectively)  
190 generated using the Trinity assembly package [12]. The B genome was further improved by  
191 "curing" [13] using sequence information from the tetraploid *T. turgidum* ssp. *dicoccoides*,  
192 which more closely represents the B genome in hexaploid wheat. This resulted in a



193 reference transcriptome sequence comprising 105,069, 132,363 and 85,296 transcript  
194 assemblies for the A, B and D genomes respectively. Based on linkage map information and  
195 conserved synteny between wheat and *Brachypodium distachyon*, these assemblies were  
196 arranged into their hypothetical gene order, providing a set of pseudomolecules [14]. Based  
197 on sequence similarity to *Brachypodium*, rice, sorghum and Maize, these pseudomolecules  
198 were annotated with probable gene functions. The A, B and D reference assemblies were  
199 sufficiently distinct to enable reads to be aligned in a genome-specific manner.

200 The alignment of the diversity panel mRNA-seq reads to this reference sequence enabled  
201 the detection of 42,066 SNP markers. SNP-calling was conducted essentially as described  
202 previously for *Brassica napus*, with read mapping and SNP calls made for each accession  
203 using Maq and Maq.pl commands, before integrating calls across the panel using the Perl  
204 script combiner.pl [11]. Simple SNPs were called by the meta-analysis of alignments against  
205 the Trinity unigene reference from mRNA-Seq reads obtained from each of 100 bread wheat  
206 lines. SNP positions were excluded from further analysis if more than two alleles were  
207 detected across the accessions, and a noise threshold of 0.15 was employed to reduce false  
208 SNP calls due to sequencing errors.

209  
210 In addition, quantification of transcript abundance (as reads per kb per million aligned reads;  
211 RPKM) provided a measure of expression for each transcript assembly. This provided the  
212 information required to explore any relationships between gene expression and the trait of  
213 interest in what has been termed a Gene Expression Marker (GEM) analysis [8].

### 214 215 **SNP marker analysis**

216 SNP-based association analysis was performed using 12,456 SNPs (following the removal  
217 of SNPs present at minor allele frequency <5%). The results were assessed visually by  
218 plotting the obtained  $P$  values (as  $-\log_{10}P$ ) in pseudomolecule order.

219 The filtered SNP dataset was used to construct a kinship matrix using the software TASSEL  
220 V3.0[15]. In addition, broad-scale population structure was assessed using the Bayesian  
221 clustering method, STRUCTURE[16]. Based on the SNP data, this method was used to  
222 identify ancestral groups to which the different accessions could be apportioned. The SNP  
223 data was processed using an admixture model with independent allele frequencies. To allow  
224 for likelihood estimates of a range of ancestral populations to be made, the model was set to  
225 run with hypothetical population (K) estimates of 1 to 5. The SNP data was processed for

226 each value of K three times with a burn-in length of 100,000. This was followed by 100,000  
227 iterations of the Monte Carlo Markov Chain algorithm. To allow for a more accurate estimate  
228 of K, the results obtained from STRUCTURE were further analysed using the methods  
229 described by Evanno et al (2005) [17] allowing for the assessment of variance between  
230 iterations. The output of the STRUCTURE analysis was used as a Q matrix (Supplementary  
231 Data. 2) for the subsequent Associative Transcriptomics analyses. The trait data, SNP data,  
232 Q matrix and kinship matrix were incorporated into a Mixed Linear Model (MLM) algorithm  
233 performed using TASSEL.

### 235 ***Gene Expression Marker (GEM) analysis***

236 Following the quantification and normalisation of the transcript levels as reads per kb per  
237 million aligned reads (RPKM), and the filtering of transcripts with an RPKM of less than 0.4  
238 across accessions, linear regression was performed. This analysis made use of 94,060 GEM  
239 markers. By fitting the RPKM values for each unigene as the dependant variable and the  
240 trait data as the independent variable, it was possible to assess the relationship between this  
241 measure of gene expression and the traits. The results were explored visually by plotting the  
242 obtained  $P$  values (as  $-\log_{10}P$ ) in pseudomolecule order. The scripts used in this analysis  
243 were developed in R (cran.r-project.org).

### 245 ***Validation of markers***

246 Leaf material was collected from a test panel of 96 hexaploid wheat accessions grown as  
247 part of the WAGTAIL panel (a diversity panel developed for the BBSRC LINK project “Wheat  
248 Association Genetics for Trait Analysis and Improved Lineages” (BB/J002607/1)) at KWS,  
249 Thriplow in 2013. DNA was extracted according to an adapted method of that described by  
250 Pallotta et al (2003) [18]. Genome specific primers were designed for each of the marker loci  
251 analysed. All marker assays were first tested on wheat accessions of known genotype (a  
252 subset of the Associative Transcriptomics panel). Following confirmation that the marker  
253 assays were able to effectively screen for the target variation, they were further used to  
254 genotype the 96 WAGTAIL accessions. All genotyping was performed using AMPLITAQ  
255 Gold polymerase (250 u – Life Technologies Ltd (Invitrogen Division, Paisley, UK)). Prior to  
256 sequencing, PCR reactions were purified using the ExoSAP protocol [19]. Following this,  
257 sequencing reactions were set up in 0.2 ml tubes according to a revised protocol from  
258 BigDye V3.1 terminator cycle sequencing kit [20]. All PCR and sequencing reactions were

259 performed using a G-Storm GS1 thermal cycler (Somerton, UK). Capillary sequencing was  
1 performed by GATC Biotech AG, Germany and all sequencing trace files obtained were  
2 performed by GATC Biotech AG, Germany and all sequencing trace files obtained were  
3 analysed using Contig Express (Vector NTI advance® 11.5.2, Paisley, UK).  
4

5  
6 262 Following genotyping, a subset of 30 wheat accessions (Supplementary Data. 3) showing  
7 representative variation at the chosen marker loci, were selected for mechanical testing.  
8  
9 264 These accessions were mechanically tested as described previously. Using a T-test  
10 (Genstat 15<sup>th</sup> edition) the trait data and genotype data obtained were assessed for any  
11 265 significant marker-trait segregation patterns.  
12 266  
13  
14

15 267

16  
17 268

18  
19  
20 269

21  
22 270

## 23 24 25 271 **Results**

### 26 27 272 ***Variation for stem structural and material strength***

28  
29  
30 273 The diversity panel of 100 wheat accessions was analysed for a range of traits indicative of  
31 stem structural and material strength. With the exception of second moment of area,  
32 274 significant variation was present for all traits included in the analysis ( $P<0.05$ )  
33 275 (Supplementary Data. 1). The absolute strength traits Fmax and F/V showed respective trait  
34 276 ranges of 7.45-38.55 and 29.82-80.44 N/sec. The wheat accession displaying highest stem  
35 277 absolute strength (for both Fmax and F/V) was Orlando. The lowest trait values were seen in  
36 278 Battalion and Escorial for F/V and Fmax respectively. For the material strength traits, MOR  
37 279 and MOE, respective trait ranges of 0.70-8.05 and 121.6-1490.3 Nmm<sup>-2</sup> were recorded. Of  
38 280 the wheat accessions screened, Gatsby exhibited the lowest trait values for both MOE and  
39 281 MOR. Accessions displaying the highest material strength were Alba (for MOR) and Cordiale  
40 282 (for MOE). A wide range of variation was also observed for the various stem structural traits  
41 283 assessed. For example, mean stem hollow area ranged from 1.16 mm<sup>2</sup> (for Capelle-  
42 284 Desprez) and 6.51mm<sup>2</sup> (for Starke2). For outer cortex thickness, trait means ranging  
43 285 between 0.24mm (as seen for Hyperion) and 0.46 mm (as seen for Alba) were recorded. For  
44 286 plant height, despite a lack of segregation at the *Rht* loci, a trait range of 42.8-98.4cm was  
45 287 recorded. The tallest accession included within the panel was Steadfast whereas the  
46 288 shortest stem measurements were recorded for Equinox.  
47 289  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

290 A correlation analysis was performed to analyse the relationships between the absolute  
291 strength and the structural and morphological traits to assess which may be good breeding  
292 targets (Table 1). Several highly significant ( $P \leq 0.001$ ) relationships were detected between  
293 the absolute strength measures (Fmax and F/V) and the structural traits, however, despite  
294 such high statistical significance, in the majority of cases, the amount of variation in stem  
295 absolute strength explained by stem structure was found to be modest. Stem parenchyma  
296 area ( $R^2 = 0.27$  and  $0.17$  for Fmax and F/V respectively) and outer cortex thickness ( $R^2 =$   
297  $0.19$  and  $0.13$  for Fmax and F/V respectively) show the closest positive relationships with  
298 absolute strength. These traits may therefore be the most promising targets for the  
299 improvement of stem structural strength in wheat. In contrast to the modest contributions  
300 made by stem geometry, a much closer correlation is seen between the absolute strength  
301 measures and stem weight ( $R^2 = 0.42$  and  $0.47$  for Fmax and F/V respectively). These  
302 correlations may represent a combined effect of several different stem structural  
303 components (each contributing to weight) or may more specifically relate to the density of  
304 the materials that make up the plant stem. Plant height also correlates positively with stem  
305 absolute strength ( $R^2 = 0.21$  and  $0.25$  for Fmax and F/V respectively).

306 The lack of strong correlations observed between stem structure and absolute strength may  
307 suggest that stem material properties are of high value for the improvement of stem  
308 mechanical strength in wheat. Consistent with this, the relationship between the field-based  
309 measure of stem lodging risk (utilising the pulley system illustrated in Figure 1c) and the  
310 absolute and material strength traits, showed a stronger correlation for the material strength  
311 trait Modulus of Rupture (MOR;  $R^2$  of  $0.41$ ,  $P < 0.001$ ) in comparison to absolute strength  
312 traits such as Fmax ( $R^2$  of  $0.27$ ,  $P < 0.001$ ) (Supplementary Data. 4).

### 314 ***The development of functional genotypes for Associative Transcriptomics***

315 Illumina mRNAseq data were produced from leaf RNA from the diversity panel of 100 wheat  
316 accessions. These sequences were mapped to the ordered transcriptome reference  
317 reported recently (Harper et al, 2015), with an average number of input reads across the full  
318 panel of 29.5 million, providing an average read coverage of 5.87. The panel was scored for  
319 SNPs and transcripts were quantified as RPKM. In total, 42,066 SNPs were scored, of  
320 which 12456 were present at  $MAF > 0.05$ , so were considered suitable for use in AT.  
321 Abundance was measured as  $>0.4$  RPKM across the population for 94,060 transcripts,  
322 which were considered suitable for use in AT. Full association plots for the following traits  
323 can be found in Supplementary Figures 1-9.

### 324 ***Associative Transcriptomics for Plant height***

1  
2 325 In order to identify loci controlling plant height, AT was conducted using the functional  
3  
4 326 genotypes scored and the plant height trait data obtained. Supplementary Data 5  
5  
6 327 summarises the results obtained. Two major association peaks were identified: one on  
7  
8 328 chromosome 6A and the other on 5B, each exhibiting SNP and GEM associations (Figure  
9  
10 329 2). To identify candidates for the causative genes for control of the trait underlying the  
11  
12 330 association peaks, the sequence similarities of unigenes to gene models in Brachypodium,  
13  
14 331 Sorghum, rice and Arabidopsis were used as a guide to gene function. This revealed that the  
15  
16 332 gene corresponding to the highest significance GEM on 6A is an orthologue of a rice Auxin  
17  
18 333 Response Factor (*OsARF16*, Os02g41800; Panel a). The peak found on chromosome 5B  
19  
20 334 coincided with a cluster of *SMALL AUXIN UP RNA* (SAUR) genes, with high significance  
21  
22 335 GEMs occurring in three of the unigenes with BLAST identity to SAUR genes (Panel b).  
23  
24 336 Although these loci have not been implicated previously in the control of plant height in  
25  
26 337 wheat, the genes identified are excellent candidates for controlling this trait: ARFs are  
27  
28 338 transcription factors that bind specifically to auxin response elements (*AuxREs*) found in the  
29  
30 339 promoters of early auxin response genes such as the large family of SAUR genes, and  
31  
32 340 mediate their response to auxin [21]. In wheat, we found that the GEM for the ARF on 6A  
33  
34 341 had a positive correlation with stem height. These results suggest that this Auxin Response  
35  
36 342 Factor may have a developmental role in wheat. Although the actual function of the SAURs  
37  
38 343 is not known, it has been reported that some have an important role in control of cell  
39  
40 344 expansion and patterning [22]. On closer inspection of their sequence similarities, the SAUR  
41  
42 345 genes in the region of 5B are putative orthologues of some of the members of a cluster of 17  
43  
44 346 SAURs found on rice chromosome 9 (*OsSAUR39-55*) and an orthologous cluster can also  
45  
46 347 be found on Arabidopsis chromosome 1 (*AtSAUR61-68*) [23]. In rice, *OsSAUR39* has been  
47  
48 348 found to negatively regulate auxin synthesis and transport, leading to reduced growth  
49  
50 349 phenotypes when over-expressed [24]. Our observation that all of the highly associated  
51  
52 350 SAURs in this cluster exhibited gene expression that was negatively correlated with height is  
53  
54 351 concordant with this.

55 352

### 56 353 ***Associative Transcriptomics for Modulus of Rupture***

57 354 AT for MOR identified three SNP association peaks. On chromosome 2D, two association  
58  
59 355 peaks were found. The first of these (marked with an arrow in Figure 3a) was found to be in  
60  
61 356 close proximity to a gene orthologous to a rice acetyl xylan esterase (*AxeA*; Os04g01980).  
62  
63 357 *AxeA*, is thought to have hydrolase activity, specifically acting on ester bonds in the  
64  
65 358 deacetylation of xylans in the plant cell wall [25]. The second association peak found on

359 chromosome 2D for MOR exhibited both SNP and GEM associations (shown within the grey  
1 360 dotted lines on Figure 3a and 3b). Several genes in this region show a consistent, positive,  
2 361 relationship of their expression with variation in MOR, which may be indicative of a large-  
3 362 scale rearrangement such as a deletion.  
4  
5  
6

7 363 A final SNP association peak was seen on chromosome 1B (Figure 3d). On closer  
8 364 inspection, it was revealed that the locus with the most highly associated marker has high  
9 365 sequence similarity to an Arabidopsis GDSL-like Lipase/Acylhydrolase superfamily gene  
10 366 (At1g54790). GDSL-like lipases are thought to be involved in the hydrolysis of ester bonds in  
11 367 cell wall xylans and have been found to have xylan acetylase activity [26]. This is a very  
12 368 similar function to that previously described for the candidate detected on chromosome 2D.  
13 369 Previous work in Arabidopsis has shown that xylan acetylation is an important contributor to  
14 370 stem strength. For example, the *eskimo-1* mutant, which displays reduced xylan acetylation,  
15 371 exhibits reduced cell wall thickening and significantly weaker stems in comparison to wild-  
16 372 type plants [27].  
17  
18  
19  
20  
21  
22  
23  
24

25 373 In addition to the GEM association peak seen on chromosome 2D, several individual GEMs  
26 374 were also found to show significant association with material strength. An example of this  
27 375 can be seen in Figure 3b (GEM marked at the foot of orange line). Transcript abundance for  
28 376 this GEM correlates positively with MOR. This marker corresponds to an orthologue of  
29 377 Arabidopsis *SERINE CARBOXYPEPTIDASE-LIKE 49* (At3g10410). The Tobacco  
30 378 orthologue of this gene, *NtSCP1*, is known to be important for cell elongation and it has been  
31 379 proposed that this gene may target proteins involved in cell wall remodelling [28], making  
32 380 this a very plausible candidate gene for stem material strength. Another example was found  
33 381 on chromosome 7B with a GEM corresponding to an orthologue of Arabidopsis  
34 382 *QUASIMODO 1* (At3g25140). Mutants defective in this gene exhibit a number of defects  
35 383 including reduced homogalacturonan (a cell wall pectin) content in the cell wall and reduced  
36 384 cell adhesion [29]. Previous studies have shown that variation in pectin can have a dramatic  
37 385 effect of stem mechanical strength in plants [30]. As a final example, on chromosome 6B, a  
38 386 marker located within a gene orthologous to that which, in rice, has been described as a  
39 387 translation initiation factor, *EIF-2B* epsilon subunit (Os02g56740), shows a high association  
40 388 with MOR. In rice, this gene is thought to have a role in the recruitment of mRNAs and the  
41 389 machinery required for translation. A related protein however, *EIF-5A*, has been found to be  
42 390 involved in a signalling pathway contributing to cell wall integrity and formation [31]. It is  
43 391 therefore possible that *EIF-2B* also has a similar, additional function.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

58 392 To further analyse the individual GEM associations detected, their respective transcript  
59 393 abundances (measured as RPKM) were mapped as traits against the SNP data of the wheat  
60  
61  
62  
63  
64  
65

394 accessions. Interestingly, for each of the described GEMs, a strong SNP peak was detected  
395 on chromosome 2D, the same region previously described for MOR in both the SNP and  
396 GEM analyses. An example of this can be seen in Figures 3c for the previously mentioned  
397 single GEM detected on chromosome 2D (Figure 3b). Figure 3c shows a clear SNP  
398 association on chromosome 2D following the mapping of the transcript abundance values for  
399 this GEM as a trait against the SNP data. All additional GEMs found to show this relationship  
400 with the 2D locus can be seen marked with a red asterisk in Supplementary Data 5. This  
401 finding could be indicative of an interaction between those genes detected as single marker  
402 associations and one or more genes located within the 2D region. Due to the many genes  
403 showing associations in the 2D region detected in the GEM analysis for MOR, it is difficult to  
404 propose a candidate gene. However, one gene, which corresponds to one of the most highly  
405 associating GEM markers within this peak, may be considered a very plausible target. This  
406 gene corresponds to an orthologue in rice described as a *COP9 SIGNALOSOME SUBUNIT*  
407 *5B (CSN5B)* (Os04g56070). The COP9 signalosome is a multi-protein complex which is  
408 known to be involved in protein degradation and has, in plants, been implicated in a number  
409 of developmental processes including photomorphogenesis (light-mediated growth), cell  
410 cycle progression and gene expression [32]. Interestingly, in Fungi, the COP9 signalosome  
411 has been implicated in cell wall remodelling. Work conducted by Nahlik *et al.* (2012), found  
412 that in the absence of a functional COP9 complex, *Aspergillus nidulans* exhibits altered  
413 expression of genes involved in cell wall remodelling [33]. Furthermore, one of the single  
414 GEM associations detected for material strength, corresponds to a eukaryotic translation  
415 initiation factor (*EIF-2B*) gene. Previous studies have shown evidence of interactions  
416 between EIF-related genes and the COP9 complex [34]. Given this, it is plausible that the  
417 genes associated with material strength are interacting with the COP9 (or more specifically,  
418 *CSN5B*) complex by means of a pathway analogous to that seen in *Aspergillus nidulans*,  
419 contributing to cell wall remodelling.

### 421 **Marker validation**

422 To test the power of Associative Transcriptomics for the identification of predictive markers,  
423 a marker validation study was carried out using a panel of 96 additional wheat accessions  
424 and focusing on three SNP associations previously described for MOR. This analysis  
425 involved the screening of a completely independent panel of wheat accessions (taken from  
426 the WAGTAIL panel) for variation at the three marker loci. These accessions were then  
427 phenotyped using the three-point bend test as before and any marker-trait segregation  
428 patterns assessed statistically.

429 Although this analysis would ideally focus on segregating variation of the most significant  
1 430 SNP within the association peak, the development of genome-specific marker assays for two  
2 431 of the targeted loci (B\_comp6657\_c0\_seq1:3733 and For D\_comp970\_c0\_seq1:1030)  
3 432 proved problematic (due to mixed traces in sequencing reads). However, genome-specific  
4 433 marker assays were successfully developed for alternative, highly associating SNPs within  
5 434 the corresponding peaks. Supplementary Table 1 provides an overview of the marker  
6 435 assays used for successful amplification of the targeted loci. Although variation was seen for  
7 436 two of the targeted marker loci, the WAGTAIL panel was monomorphic for  
8 437 D\_comp1058\_c0\_seq1:1573, so it was not used.

15 438 Based on the marker variation uncovered from the remaining two marker assays, 30  
16 439 accessions were chosen for mechanical testing. These accessions were chosen based on  
17 440 genotype alone to ensure non-biased trait prediction and to ensure that all possible marker  
18 441 allele combinations were represented in downstream analyses. Following mechanical  
19 442 testing, a student T-test was used to assess whether, on average, a higher trait value is  
20 443 observed in accessions carrying the increasing alleles of the markers uncovered through AT,  
21 444 thus proving that the markers identified have trait predictive capability. **Supplementary Figure**  
22 445 **10** summarises the results for each marker locus. As predicted, significantly increased trait  
23 446 values are seen in segregation with increasing alleles at both loci (with segregation patterns  
24 447 being assigned  $P$  values of  $\leq 0.01$  and  $\leq 0.001$  for D\_comp19374\_c0\_seq1:702 and  
25 448 B\_comp2391\_c0\_seq1:284 respectively), proving that these markers have robust trait  
26 449 prediction capability. It is also promising to note, that the WAGTAIL accessions showing  
27 450 particularly high mean MOR (between 25.9 – 34.9 N mm<sup>-2</sup>), are among those carrying  
28 451 increasing alleles at both marker loci (**Supplementary Figure 11**).

## 40 452 Discussion

41 453 Despite great efforts, lodging continues to be one of the key factors threatening wheat yield  
42 454 worldwide. The selection of elite accessions with alternative semi-dwarfing alleles or high  
43 455 stem mechanical strength may be a powerful approach to reducing this problem.

44 456 As previously mentioned, the selection of dwarfing alleles is a commonly employed method  
45 457 for lodging control in wheat. The lack of segregation of these loci (*Rht-B1* and *Rht-D1*) in our  
46 458 training panel has enabled the identification of additional candidate genes that may  
47 459 contribute to controlling height in this species, implicating auxin-related genes as key  
48 460 regulators. Importantly, the loci implicated in plant height control are completely independent  
49 461 to those seen for stem strength and may therefore be used to further maximise lodging  
50 462 resistance in future elite wheat accessions, or to develop taller lodging resistant accessions.



1 463 Such accessions would also improve the achievable profit margin by increasing the amount  
2 464 of straw that can be harvested for use as animal bedding or biorefining feedstocks.  
3

4 465 In recent years we have seen increased interest in the possible exploitation of agricultural  
5 466 residues (such as waste straw of the wheat crop) as a feedstock for lignocellulosic ethanol  
6 467 production. However, at present, high costs related to the breakdown of lignocellulosic  
7 468 biomass is hindering this fuel source becoming a feasible future alternative. One way  
8 469 through which processibility may be improved, is through altering the composition of the  
9 470 lignocellulosic matrix. In this study, we have shown evidence for the importance of xylan  
10 471 acetylation in contributing to stem material strength in wheat, but xylan acetylation is also  
11 472 known to impede the enzymatic breakdown of lignocellulosic biomass and therefore reduced  
12 473 xylan acetylation is a desirable target for this industry [35]. The results presented here  
13 474 suggest that alterations in xylan acetylation may affect stem mechanical strength, so given  
14 475 this, it is essential that any effects of altering cell wall xylan acetylation on agronomic  
15 476 performance are assessed.  
16  
17  
18  
19  
20  
21  
22  
23  
24

25 477 In addition to the potential role of xylan acetylation, this study has uncovered a possible role  
26 478 of the COP9 signalosome in contributing to stem mechanical strength in wheat. The  
27 479 detection of several interactions between single GEMs and the *CSN5* locus is very  
28 480 interesting. One of the associating GEMs showing a relationship with the 2D locus,  
29 481 corresponds to an orthologue of *EIF-2B*. Previous studies have shown evidence of  
30 482 interactions between the COP9 complex and EIF-related genes. This suggests that the  
31 483 utilisation of the GEM data for the assessment of gene-gene interactions through mapping is  
32 484 effective. Several of the GEMs found to interact with the 2D locus are expected to have a  
33 485 role in cell wall remodelling/biosynthesis. Previous work has shown that, in fungi, the COP9  
34 486 complex has a role in cell wall remodelling, an important aspect of growth [33]. It is possible,  
35 487 based on the results presented here, that there is a pathway, analogous to that described in  
36 488 fungi, where the COP9 complex (or at the very least subunit *CSN5*) regulates the expression  
37 489 of genes involved in cell wall remodelling, and that this is an important contributor to stem  
38 490 material strength in wheat. To our knowledge, this is the first instance of reporting such a  
39 491 role of the COP9 complex *in planta*.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

## 51 492 **Conclusions**

52  
53

54 493 This work illustrates the power of Associative Transcriptomics for the exploration of even the  
55 494 most complex, environmentally sensitive traits. With careful selection of the genotypes  
56 495 included, we have shown that even a relatively small diversity set can, when coupled with  
57 496 high marker density and low linkage disequilibrium, provide the power required for the  
58 497 discovery of novel and agronomically valuable genetic variation. In this study, we  
59  
60  
61  
62  
63  
64  
65

498 successfully identified and validated markers for two loci that provide increased Modulus of  
1 499 Rupture, an important measure of the resistance of the plant material to breakage. We have  
2  
3 500 also shown that this method has the potential to uncover novel targets for breeding of  
4  
5 501 important morphological traits, such as plant height. Furthermore, the coupling of SNP  
6  
7 502 variation with variation at the gene expression level has provided the power required to gain  
8  
9 503 a deeper understanding of the biological context of variation underlying important agronomic  
10 504 traits. This not only adds to the wealth of knowledge that we strive to accumulate regarding  
11 505 gene function and plant adaptation, but also provides breeders with the information required  
12 506 to make more informed decisions regarding the potential consequences of incorporating the  
13 507 use of particular markers into future breeding programmes.  
14  
15  
16

## 17 508 **Declarations**

### 18 19 20 509 **Availability of data and material**

21  
22 510 Illumina sequence reads are available from SRA (accession number ERA283619) and  
23 511 transcript assemblies from:

24  
25 512 [http://opendata.tgac.ac.uk/associative\\_transcriptomics/wheat/v1/Trinity\\_ABD\\_cured.fasta.gz](http://opendata.tgac.ac.uk/associative_transcriptomics/wheat/v1/Trinity_ABD_cured.fasta.gz).  
26  
27

### 28 513 **Competing interests**

29  
30  
31 514 The authors declare that they have no competing interests.  
32

### 33 515 **Funding**

34  
35  
36 516 This work was supported by the UK Biotechnology and Biological Sciences Research  
37 517 Council (BBSRC BB/H004351/1 (IBTI Club), BB/L002124/1, BB/L027844/1).  
38  
39

### 40 518 **Authors' contributions**

41  
42  
43 519 IB, KW, ALH and CNM conceived and planned the project. CNM and ALH performed the  
44 520 experiments. MT performed data analysis. PW planned and supervised field trials. CNM,  
45 521 ALH and IB wrote the manuscript and all authors reviewed it.  
46  
47

### 48 522 **Acknowledgements**

49  
50  
51 523 We thank Klaus Wellner at the Institute of Food Research for technical support. Next-  
52 524 generation sequencing and library construction was delivered via the BBSRC National  
53 525 Capability in Genomics (BB/J010375/1) at The Genome Analysis Centre by members of the  
54 526 Platforms and Pipelines Group.  
55  
56  
57  
58

59 527  
60  
61  
62  
63  
64  
65

- 1  
2 529 1. Foulkes MJ, Slafer GA, Davies WJ, Berry PM, Sylvester-Bradley R, Martre P,  
3 530 Calderini DF, Griffiths S, Reynolds MP: **Raising yield potential of wheat. III.**  
4 531 **Optimizing partitioning to grain while maintaining lodging resistance.** Journal of  
5 532 Experimental Botany 2011, **62**(2):469-486.
- 6 533 2. Russell GE: **Progress in Plant Breeding—1:** Elsevier Science; 2013.
- 7 534 3. Verma V, Worland AJ, Savers EJ, Fish L, Caligari PDS, Snape JW: **Identification**  
8 535 **and characterization of quantitative trait loci related to lodging resistance and**  
9 536 **associated traits in bread wheat.** Plant Breeding 2005, **124**(3):234-241.
- 10 537 4. Keller M, Karutz C, Schmid JE, Stamp P, Winzeler M, Keller B, Messmer MM:  
11 538 **Quantitative trait loci for lodging resistance in a segregating wheat×spelt**  
12 539 **population.** Theoret Appl Genetics 1999, **98**(6-7):1171-1182.
- 13 540 5. Atwell S, Huang YS, Vilhjalmsson BJ, Willems G, Horton M, Li Y, Meng D, Platt A,  
14 541 Tarone AM, Hu TT et al: **Genome-wide association study of 107 phenotypes in**  
15 542 **Arabidopsis thaliana inbred lines.** Nature 2010, **465**(7298):627-631.
- 16 543 6. Pasam R, Sharma R, Malosetti M, van Eeuwijk F, Haseneyer G, Kilian B, Graner A:  
17 544 **Genome-wide association studies for agronomical traits in a world wide spring**  
18 545 **barley collection.** BMC Plant Biology 2012, **12**(1):16.
- 19 546 7. Hwang E-Y, Song Q, Jia G, Specht J, Hyten D, Costa J, Cregan P: **A genome-wide**  
20 547 **association study of seed protein and oil content in soybean.** BMC Genomics 2014,  
21 548 **15**(1):1.
- 22 549 8. Andrea LH, Martin T, Janet H, Fiona F, Leah C, Rachel W, Chie H, Peter W, Ian B:  
23 550 **Associative transcriptomics of traits in the polyploid crop species Brassica napus.**  
24 551 Nature Biotechnology 2012, **30**(8):798-802.
- 25 552 9. Harper AL, Trick M, He Z, Clissold L, Fellgett A, Griffiths S, Bancroft I: **Genome**  
26 553 **distribution of differential homoeologue contributions to leaf gene expression in**  
27 554 **bread wheat.** Plant Biotechnology Journal 2016, **14**(5):1207-1214.
- 28 555 10. Kern KA, Ewers FW, Telewski FW, Koehler L: **Mechanical perturbation affects**  
29 556 **conductivity, mechanical properties and aboveground biomass of hybrid**  
30 557 **poplars.** Tree Physiology 2005, **25**(10):1243-1251.
- 31 558 11. Bancroft I, Morgan C, Fraser F, Higgins J, Wells R, Clissold L, Baker D, Long Y,  
32 559 Meng J, Wang X et al: **Dissecting the genome of the polyploid crop oilseed rape by**  
33 560 **transcriptome sequencing.** Nat Biotechnol 2011, **29**(8):762-766.
- 34 561 12. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X,  
35 562 Fan L, Raychowdhury R, Zeng Q et al: **Full-length transcriptome assembly from**  
36 563 **RNA-Seq data without a reference genome.** Nat Biotechnol 2011, **29**(7):644-652.
- 37 564 13. Higgins J, Magusin A, Trick M, Fraser F, Bancroft I: **Use of mRNA-seq to**  
38 565 **discriminate contributions to the transcriptome from the constituent genomes of**  
39 566 **the polyploid crop species Brassica napus.** BMC Genomics 2012, **13**(1):247.
- 40 567 14. Harper AL, Trick M, He Z, Clissold L, Fellgett A, Griffiths S, Bancroft I: **Genome**  
41 568 **distribution of differential homoeologue contributions to leaf gene expression in**  
42 569 **bread wheat.** Plant Biotechnology Journal 2015:n/a-n/a.
- 43 570 15. Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES:  
44 571 **TASSEL: software for association mapping of complex traits in diverse samples.**  
45 572 Bioinformatics 2007, **23**(19):2633-2635.
- 46 573 16. Pritchard JK, Stephens M, Donnelly P: **Inference of population structure using**  
47 574 **multilocus genotype data.** Genetics 2000, **155**(2):945-959.
- 48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

575 17. Evanno G, Regnaut S, Goudet J: **Detecting the number of clusters of individuals**  
1 576 **using the software structure: a simulation study.** *Molecular Ecology* 2005,  
2 577 **14(8):2611-2620.**

3 578 18. Pallotta MA, Warner P, Fox RL, Kuchel H, Jefferies SJ, Langridge P: **Marker-**  
4 579 **assisted wheat breeding in the southern region of Australia.** *Proceedings of the*  
5 580 *Tenth International Wheat Genetics Symposium* 2003:789-791.

6 581 19. Etchevers H: **DNA sequencing and quick clean-up.** 2007.

7 582 20. Applied.Biosystems: **BigDye™ Terminator v3.1 Ready Reaction Cycle**  
8 583 **Sequencing Kit Protocol.** . 2002.

9 584 21. Guilfoyle TJ, Ulmasov T, Hagen G: **The ARF family of transcription factors and**  
10 585 **their role in plant hormone-responsive transcription.** *Cellular and molecular life*  
11 586 *sciences : CMLS* 1998, **54(7):619-627.**

12 587 22. Chae K, Isaacs CG, Reeves PH, Maloney GS, Muday GK, Nagpal P, Reed JW: **Arabidopsis**  
13 588 **SMALL AUXIN UP RNA63 promotes hypocotyl and stamen**  
14 589 **filament elongation.** *The Plant Journal* 2012, **71(4):684-697.**

15 590 23. Jain M, Tyagi AK, Khurana JP: **Genome-wide analysis, evolutionary expansion,**  
16 591 **and expression of early auxin-responsive SAUR gene family in rice (Oryza**  
17 592 **sativa).** *Genomics* 2006, **88(3):360-371.**

18 593 24. Kant S, Bi Y-M, Zhu T, Rothstein SJ: **SAUR39, a Small Auxin-Up RNA Gene, Acts**  
19 594 **as a Negative Regulator of Auxin Synthesis and Transport in Rice.** *Plant*  
20 595 *Physiology* 2009, **151(2):691-701.**

21 596 25. Alalouf O, Balazs Y, Volkinshtein M, Grimpel Y, Shoham G, Shoham Y: **A New**  
22 597 **Family of Carbohydrate Esterases Is Represented by a GDSL**  
23 598 **Hydrolase/Acetylxyylan Esterase from Geobacillus stearothermophilus.** *Journal of*  
24 599 *Biological Chemistry* 2011, **286(49):41993-42001.**

25 600 26. Akoh CC, Lee G-C, Liaw Y-C, Huang T-H, Shaw J-F: **GDSL family of serine**  
26 601 **esterases/lipases.** *Progress in Lipid Research* 2004, **43(6):534-552.**

27 602 27. Yuan Y, Teng Q, Zhong R, Ye Z-H: **The Arabidopsis DUF231 Domain-Containing**  
28 603 **Protein ESK1 Mediates 2-O- and 3-O-Acetylation of Xylosyl Residues in Xylan.**  
29 604 *Plant and Cell Physiology* 2013, **54(7):1186-1199.**

30 605 28. Bienert MD, Delannoy M, Navarre C, Boutry M: **NtSCP1 from Tobacco Is an**  
31 606 **Extracellular Serine Carboxypeptidase III That Has an Impact on Cell**  
32 607 **Elongation.** *Plant Physiology* 2012, **158(3):1220-1229.**

33 608 29. Bouton S, Leboeuf E, Mouille G, Leydecker M-T, Talbotec J, Granier F, Lahaye M,  
34 609 Höfte H, Truong H-N: **QUASIMODO1 Encodes a Putative Membrane-Bound**  
35 610 **Glycosyltransferase Required for Normal Pectin Synthesis and Cell Adhesion in**  
36 611 **Arabidopsis.** *The Plant Cell Online* 2002, **14(10):2577-2590.**

37 612 30. Hongo S, Sato K, Yokoyama R, Nishitani K: **Demethylesterification of the Primary**  
38 613 **Wall by PECTIN METHYLESTERASE35 Provides Mechanical Support to the**  
39 614 **Arabidopsis Stem.** *The Plant cell* 2012, **24(6):2624-2634.**

40 615 31. Valentini SR, Casolari JM, Oliveira CC, Silver PA, McBride AE: **Genetic**  
41 616 **Interactions of Yeast Eukaryotic Translation Initiation Factor 5A (eIF5A)**  
42 617 **Reveal Connections to Poly(A)-Binding Protein and Protein Kinase C Signaling.**  
43 618 *Genetics* 2002, **160(2):393-405.**

44 619 32. Nezames CD, Deng XW: **The COP9 Signalosome: Its Regulation of Cullin-Based**  
45 620 **E3 Ubiquitin Ligases and Role in Photomorphogenesis.** *Plant Physiology* 2012,  
46 621 **160(1):38-46.**

47 622 33. Nahlik K, Dumkow M, Bayram Ö, Helmstaedt K, Busch S, Valerius O, Gerke J,  
48 623 Hoppert M, Schwier E, Opitz L et al: **The COP9 signalosome mediates**  
49 624 **transcriptional and metabolic response to hormones, oxidative stress protection**

625  
1 626  
2 627  
3 628  
4 629  
5 630  
6 631  
7 631  
8  
9 632  
10  
11 633  
12  
13  
14 634  
15  
16 635  
17  
18  
19 636  
20  
21 637  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**and cell wall rearrangement during fungal development.** *Molecular Microbiology* 2010, **78**(4):964-979.

34. Kim T-H, Hofmann K, von Arnim AG, Chamovitz DA: **PCI complexes: pretty complex interactions in diverse signaling pathways.** *Trends in Plant Science* 2001, **6**(8):379-386.

35. Pawar PM-A, Koutaniemi S, Tenkanen M, Mellerowicz EJ: **Acetylation of woody lignocellulose: significance and regulation.** *Frontiers in Plant Science* 2013, **4**.

638


**Table 1.** Pearson's correlation coefficient (tested against zero) for traits measured across wheat panel

1											
2											
3											
4	Fmax (N/sec)										
5											
6	F/V (N/sec)	***0.85									
7											
8	Stem width (mm)	0.01	0								
9											
10	Stem hollow area (mm <sup>2</sup> )	***0.16	***0.12	***0.27							
11											
12	Second moment of area (N/mm <sup>4</sup> )	**0.07	*0.06	***0.33	***0.16						
13											
14	Parenchyma area (mm <sup>2</sup> )	***0.27	***0.17	***0.11	***0.11	**0.09					
15											
16	Outer cortex thickness (mm)	***0.19	***0.13	0.032	0	0.02	0				
17											
18	Length of 2nd internode (cm)	0.014	0.037	0	**0.09	*0.06	*0.06	0			
19											
20	Height minus ear (cm)	***0.21	***0.25	0.011	0.01	**0.09	0	**0.08	***0.38		
21											
22	Threshed stem weight (g)	***0.49	***0.51	***0.13	0.01	***0.22	***0.15	**0.09	**0.1	***0.55	
23											
24											
25											
26											
27											
28											
29											
30											
31		Fmax (N/sec)	F/V (N/sec)	Stem width (mm)	Stem hollow area (mm <sup>2</sup> )	Second moment of area (N/mm <sup>4</sup> )	Parenchyma area (mm <sup>2</sup> )	Outer cortex thickness (mm)	Length of 2nd internode (cm)	Height minus ear (cm)	Threshed stem weight (g)
32											
33											
34											
35											
36											
37											
38	639										
39											
40											
41	640										
42											
43	641										
44											
45											
46	642										
47											
48											
49	643										
50											
51	644										
52											
53											
54											
55											
56											
57											
58											
59		*** indicates significance at $P \leq 0.001$ and ** indicates significance at $P \leq 0.01$ and * indicates									
60		significance at $P \leq 0.05$ .									
61											
62											
63											
64											
65											

645 **Figure 1. Apparatus used for assessment of stem mechanical strength in wheat.** A lab-  
646 based three-point bend test setup (a) allowed for the absolute strength traits,  $F_{max}$  (the  
647 resistance of the stem sample to break under load) and  $F/V$  (the resistance of the stem  
648 sample to bend elastically), to be obtained (b). A field-based stem lodging risk measure was  
649 obtained using a pulley device (c).


650  
651 **Figure 2. SNP and GEM marker associations detected for plant height.** Marker  
652 associations are illustrated, for both sequence-based (SNP) and gene expression-based  
653 (GEM) markers, with significance of association (as  $-\log_{10}P$  values) plotted against position  
654 within specific chromosomes. The inferred order of unigenes is illustrated below the scans  
655 with colour coding by sequence similarity to chromosomes of *B. distachyon* (blue=Bd1;  
656 yellow=Bd2; purple=Bd3; red=Bd4 and green=Bd5). Two associating loci for height are  
657 shown, one on chromosome 6A (a, b) and one on chromosome 5B (c, d). Both loci show  
658 associating SNP and GEM marker variation. The positions of candidate genes are indicated  
659 by arrows.


660  
661 **Figure 3. Variation at both the sequence (SNP) and gene expression (GEM) level show**  
662 **high association with MOR.** Two SNP association peaks for MOR were seen on  
663 chromosome 2D (a). The peak to the right of panel a was also identified in the GEM analysis  
664 (b). Several single GEM associations were also detected for MOR (see single GEM at the  
665 foot of the orange line in panel b as an example). Mapping transcript abundance (as RPKM)  
666 as a trait against the SNP data revealed the same 2D SNP peak for several single GEMs  
667 (see panel c for an example). A further SNP association for MOR was detected on  
668 chromosome 1B. The positions of candidate genes are indicated by arrows.  $-\log_{10}P$  values  
669 are plotted in wheat pseudomolecule order. Unigene order is colour-coded according to  
670 sequence similarity to *B. distachyon* chromosomes (blue=Bd1; yellow=Bd2; purple=Bd3;  
671 red=Bd4 and green=Bd5). Position of candidate genes are indicated by arrows.




**Figure. 1**



**Figure. 2**

**Figure. 3**




Click here to access/download  
**Supplementary Material**  
Supplementary Figures\_REVISED.pdf





Click here to access/download  
**Supplementary Material**  
Supplementary Table 1.docx



Click here to access/download

**Supplementary Material**

Supplementary dataset

1\_Association\_panel\_trait\_data\_adjusted\_REVISED.xlsx





Click here to access/download

**Supplementary Material**

Supplementary dataset

[2\\_Population\\_structure\\_Q\\_matrix\\_REVISED.xlsx](#)





Click here to access/download

**Supplementary Material**

Supplementary dataset

3\_trait\_data\_WAGTAIL\_genotypes\_adjusted\_REVISED.

xlsx



Click here to access/download

**Supplementary Material**

Supplementary dataset 4\_Field-  
based\_stem\_lodging\_risk\_data\_adjusted\_REVISED.xlsx





Click here to access/download

**Supplementary Material**

Supplementary dataset

5\_SNP\_and\_GEM\_association\_summary\_REVISED.xlsx