



UNIVERSITY OF LEEDS

This is a repository copy of *Cardiovascular Outcomes Following Rotational Atherectomy: A UK Multicentre Experience*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/100858/>

Version: Accepted Version

Article:

Eftychiou, C, Barmby, DS, Wilson, SJ et al. (7 more authors) (2016) Cardiovascular Outcomes Following Rotational Atherectomy: A UK Multicentre Experience. *Catheterization and Cardiovascular Interventions*, 88 (4). pp. 546-553. ISSN 1522-1946

<https://doi.org/10.1002/ccd.26587>

© 2016 Wiley Periodicals, Inc. This is the peer reviewed version of the following article: Eftychiou, C., Barmby, D. S., Wilson, S. J., Ubaid, S., Markwick, A. J., Makri, L., Blaxill, J. M., Spratt, J. C., Gunning, M. and Greenwood, J. P. (2016), Cardiovascular Outcomes Following Rotational Atherectomy: A UK Multicentre Experience. *Cathet. Cardiovasc. Intervent.*, 88: 546–553. doi: 10.1002/ccd.26587, which has been published in final form at <https://doi.org/10.1002/ccd.26587>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Table 4: Outcomes by SYNTAX category comparing ≤ 22 vs. ≥ 23 and ≤ 32 vs. ≥ 33

	SYNTAX ≤ 22	SYNTAX ≥ 23	p-value	SYNTAX ≤ 32	SYNTAX ≥ 33	p-value
MACE	44 (13.2%)	47 (26.0%)	<0.001	68 (15.3%)	23 (31.9%)	0.001
• Cardiac Death	15 (4.5%)	21 (11.6%)	0.004	24 (5.4%)	12 (16.7%)	0.002
• MI	26 (7.8%)	34 (18.8%)	<0.001	43 (9.7%)	17 (23.6%)	0.002
• TVR	22 (6.6%)	17 (9.4%)	0.295	33 (7.4%)	6 (8.3%)	0.810
Death (all cause)	33 (9.9%)	37 (20.4%)	0.001	47 (10.6%)	23 (31.9%)	<0.001
Definite Stent Thrombosis	1 (0.3%)	6 (3.3%)	0.009	5 (1.1%)	2 (2.8%)	0.255
Stroke	8 (2.5%)	7 (4.2%)	0.284	11 (2.6%)	4 (5.9%)	0.139
Peri-procedural Complications	16 (4.8%)	16 (8.8%)	0.085	22 (5.0%)	10 (13.9%)	0.008
Vascular Access Complications	6 (1.8%)	4 (2.2%)	0.747	9 (2.0%)	1 (1.4%)	1.000

Data shown as N (%),

MACE=Major Adverse Cardiac Events, MI= Myocardial Infarction, TVR= Target Vessel

Revascularization.