

This is a repository copy of Balancing Anti-Tumor Efficacy with Local Inflammatory Toxicity for the Treatment of Diffuse Intrinsic Pontine Glioma and Other Brain Tumors.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/101793/

## **Proceedings Paper:**

Schuelke, M, Evgin, L, Kottke, T et al. (8 more authors) (2016) Balancing Anti-Tumor Efficacy with Local Inflammatory Toxicity for the Treatment of Diffuse Intrinsic Pontine Glioma and Other Brain Tumors. In: Molecular Therapy. 19th Annual Meeting of the American-Society-of-Gene-and-Cell-Therapy (ASGCT), 04-07 May 2016, Washington DC, USA. Nature Publishing Group, S77-S77.

© 2016 Author(s). This is an author produced version of a paper published in Molecular Therapy. Uploaded in accordance with the publisher's self-archiving policy.

## Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

## Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/ Balancing Anti-Tumor Efficacy with Local Inflammatory Toxicity for the Treatment of Diffuse Intrinsic Pontine Glioma and Other Brain Tumors.

Matthew Schuelke, Laura Evgin, Timothy Kottke, Jill Thompson, Christopher Driscoll, Elizabeth Ilett, Julia Cockle, Amulya NageswaraRao, Richard Bram, Alan Melcher, Richard Vile.

We have shown previously that multiple rounds of a systemic treatment of GM-CSF, followed by intravenous reovirus, leads to effective treatment of subcutaneous melanomas (Ilett et al., Mol. Ther. 2015). We show here that a similar regimen is also effective at treating both melanoma (B16) and glioma (GL261) tumors growing intra-cranially. As a result of these pre-clinical studies, we initiated a Phase I clinical trial in paediatric patients with gliomas of GM-CSF and reovirus therapy. To date, three patients have been treated. In two of three of these patients there were possible indications of pseudoprogression and intra-cranial inflammation, both of which were clinically resolved upon treatment with dexamethasone.

Whilst this trial continues to recruit, we have investigated additional treatments for paediatric brain tumors, especially those in which direct virus injection may allow greater local access to tumor. In this respect, there is a constant tension between the pro-inflammatory, anti-tumor nature of oncolytic viroimmunotherapy and the need to reduce potentially toxic local inflammatory reactions within the brain. Our overall hypothesis is that it will be possible to balance the anti tumor effects of viral oncolysis, in part caused by inflammatory reactions to the virus, with the associated toxicity. Nowhere is this more relevant than for the treatment of Diffuse Pontine Gliomas (DIPG), which grow in the brain stem. Our experiments show that it is indeed possible to treat tumors growing in the pons/and or medulla, with inflammatory, immune based therapies (both T cell mediated and locally cytotoxic) without increasing toxicity over that caused by tumor growth alone. Therefore, we have screened multiple oncolytic virus types for their ability to kill DIPG cell lines both in vitro and in vivo. In particular, we have compared viruses with different speeds of oncolvsis and different inflammatory properties to investigate how the efficacy can be balanced with local toxicity, using standard of care anti inflammatory treatments (with dexamethasone) as well as with novel agents and viruses designed to suppress local inflammatory reactions.