

This is a repository copy of *Cysteine peptidase B regulates Leishmania mexicana virulence through the modulation of GP63 expression*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/102642/>

Version: Accepted Version

Article:

Mottram, Jeremy Charles orcid.org/0000-0001-5574-3766, Pierre-Andre, Casgrain, McMaster, W. Robert et al. (3 more authors) (2016) Cysteine peptidase B regulates *Leishmania mexicana* virulence through the modulation of GP63 expression. PLOS PATHOGENS. e1005658. ISSN 1553-7366

<https://doi.org/10.1371/journal.ppat.1005658>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Cysteine Peptidase B Regulates *Leishmania mexicana* Virulence**
2 **through the Modulation of GP63 Expression**

3
4
5 Pierre-André Casgrain¹, Caroline Martel², W. Robert McMaster³, Jeremy C.
6 Mottram⁴, Martin Olivier², and Albert Descoteaux^{1*}

7
8
9 ¹INRS- Institut Armand-Frappier and the Center for Host-Parasite Interactions, Laval, Canada

10 ²The Research Institute of the McGill University Health Centre, Montréal, Canada.

11 ³Immunity and Infection Research Centre, Vancouver Coastal Health Research Institute,
12 Department of Medical Genetics, University of British Columbia, Vancouver, Canada

13 ⁴Wellcome Trust Centre for Molecular Parasitology, Institute of Infection, Immunity and
14 Inflammation, College of Medical, Veterinary and Life Sciences, University of Glasgow,
15 Glasgow, United Kingdom

16
17
18
19
20
21
22
23 *Corresponding author:

24 E-mail: : albert.descoteaux@iaf.inrs.ca

ABSTRACT

26
27
28 Cysteine peptidases play a central role in the biology of *Leishmania*. In this work, we sought to
29 further elucidate the mechanism(s) by which the cysteine peptidase CPB contributes to *L.*
30 *mexicana* virulence and whether CPB participates in the formation of large communal
31 parasitophorous vacuoles induced by these parasites. We initially examined the impact of *L.*
32 *mexicana* infection on the trafficking of VAMP3 and VAMP8, two endocytic SNARE proteins
33 associated with phagolysosome biogenesis and function. Using a CPB-deficient mutant, we
34 found that both VAMP3 and VAMP8 were down-modulated in a CPB-dependent manner. We
35 also discovered that expression of the virulence-associated GPI-anchored metalloprotease GP63
36 was inhibited in the absence of CPB. Expression of GP63 in the CPB-deficient mutant was
37 sufficient to down-modulate VAMP3 and VAMP8. Similarly, episomal expression of GP63
38 enabled the CPB-deficient mutant to establish infection in macrophages, induce the formation of
39 large communal parasitophorous vacuoles, and cause lesions in mice. These findings implicate
40 CPB in the regulation of GP63 expression and provide evidence that both GP63 and CPB are key
41 virulence factors in *L. mexicana*.

42

43

AUTHOR SUMMARY

44
45
46 The parasite *Leishmania mexicana* expresses several cysteine peptidases of the papain family that
47 are involved in processes such as virulence and evasion of host immune responses. The cysteine
48 peptidase CPB is required for survival within macrophages and for lesion formation in
49 susceptible mice. Upon their internalization by macrophages, parasites of the *L. mexicana*
50 complex induce the formation of large communal parasitophorous vacuoles in which they
51 replicate, and expansion of those large vacuoles correlates with the ability of the parasites to
52 survive inside macrophages. Here, we found that CPB contributes to *L. mexicana* virulence
53 (macrophage survival, formation and expansion of communal parasitophorous vacuoles, lesion
54 formation in mice) through the regulation of the virulence factor GP63, a *Leishmania* zinc-
55 metalloprotease that acts by cleaving key host cell proteins. This work thus elucidates a novel
56 *Leishmania* virulence regulatory mechanism whereby CPB controls the expression of GP63.

57

58

59

INTRODUCTION

60
61
62 The protozoan *Leishmania* parasitizes macrophages and causes a spectrum of human
63 diseases ranging from self-healing cutaneous lesions to a progressive visceral infection that can
64 be fatal if left untreated. Infection is initiated when promastigote forms of the parasite are
65 inoculated into the mammalian host by infected sand flies and are internalized by phagocytes.
66 Inside macrophages, promastigotes differentiate into amastigotes to replicate within
67 phagolysosomal compartments also known as parasitophorous vacuoles (PVs). Upon their
68 internalization, *L. donovani* and *L. major* promastigotes arrest phagolysosomal biogenesis and
69 create an intracellular niche favorable to the establishment of infection and to the evasion of the
70 immune system [1, 2]. Disruption of the macrophage membrane fusion machinery through the
71 action of virulence factors plays an critical role in this PV remodeling. Hence, insertion of the
72 promastigote surface glycolipid lipophosphoglycan (LPG) into the PV membrane destabilizes
73 lipid microdomains and causes exclusion of the membrane fusion regulator synaptotagmin V
74 from the PV [2-4]. Similarly, the parasite GPI-anchored metalloprotease GP63 [5, 6]
75 redistributes within the infected cells and cleaves key Soluble NSF Attachment Protein Receptors
76 (SNAREs) and synaptotagmins to impair phagosome functions [1, 7].

77
78 Whereas *L. major* and *L. donovani* multiply in tight individual PVs, parasites of the *L.*
79 *mexicana* complex (*L. mexicana*, *L. amazonensis*) replicate within large communal PVs.
80 Relatively little is known about the host and parasite factors involved in the biogenesis and
81 expansion of those communal PVs. Studies with *L. amazonensis* revealed that phagosomes
82 containing promastigotes fuse extensively with late endosomes/lysosomes within 30 minutes
83 post-infection [8]. At that stage, parasites are located within small individual compartments and

84 by 18 to 24 hours large PVs containing several parasites are observed. The rapid increase in the
85 size of those PVs requires extensive fusion with secondary lysosomes and correlates with the
86 depletion of those organelles in infected cells [9-11]. Homotypic fusion between *L.*
87 *amazonensis*-containing PVs also occurs, but its contribution to PV enlargement remains to be
88 further investigated [12]. These studies highlighted the contribution of the host cell membrane
89 fusion machinery in the biogenesis and expansion of large communal PVs and are consistent with
90 a role for endocytic SNAREs in this process [13]. Interestingly, communal PVs interact with the
91 host cell's endoplasmic reticulum (ER) and disruption of the fusion machinery associated with
92 the ER and Golgi inhibits parasite replication and PV enlargement [14-16].

93
94 The *Leishmania*-derived molecules involved in the expansion of the communal PVs
95 remains to be identified. LPG and other phosphoglycans do not play a significant role in *L.*
96 *mexicana* promastigote virulence and PV formation [17], in contrast to *L. major* and *L. donovani*
97 [2]. Cysteine peptidases (CP) are a large family of papain-like enzymes that play important roles
98 in the biology of *Leishmania* [18]. Three members of these papain-like proteases are expressed
99 by *L. mexicana* and the generation of CP-deficient mutants revealed that CPB contributes to the
100 ability to infect macrophages and to induce lesions in BALB/c mice [19-21]. The precise
101 mechanism(s) by which CPB participates in the virulence of *L. mexicana* is poorly understood.
102 Previous studies revealed that CPB traffics within and outside infected macrophages [18]. In
103 infected macrophages, CPB alters signal transduction and gene expression through the activation
104 of the protein tyrosine phosphatase PTP-1B and the cleavage of transcription factors responsible
105 for the expression of genes involved in host defense and immunity [20, 22]. The observation that
106 CPs interfere with the host immune response through the degradation of MHC class II molecules

107 and invariant chains present in PVs housing *L. amazonensis* [23], raises the possibility that CPB
108 participates in the modulation of PV composition and function.

109

110 In this study, we sought to gain insight into the mechanism by which CPB contributes to
111 *L. mexicana* virulence, with a focus on the PV. We provide evidence that CPB participates in PV
112 biogenesis and virulence through the regulation of GP63 expression.

113

RESULTS

114
115
116 ***CPB enables L. mexicana to down-modulate VAMP3 and VAMP8.*** Formation and expansion
117 of communal PVs hosting *L. mexicana* involve fusion between PVs and endocytic organelles, as
118 well as homotypic fusion among PVs [10-12]. To identify the host and parasite factors involved
119 in this process, we embarked on a study to elucidate the fate of endosomal SNAREs during
120 infection of macrophages with *L. mexicana*. Given the requirement of CPB for *L. mexicana* to
121 replicate normally inside macrophages [19], we included a *L. mexicana* CPB-deficient mutant
122 (Δcpb) in our investigation. We infected BMM with either WT or Δcpb *L. mexicana*
123 promastigotes for 2 h and we assessed the distribution of the endosomal SNAREs VAMP3 and
124 VAMP8 by confocal immunofluorescence microscopy. As previously observed during infection
125 with *L. major* promastigotes [1], we found a notable reduction in the staining intensity for both
126 VAMP3 (Figure 1A) and VAMP8 (Figure 1B) in BMM infected with WT *L. mexicana*, but this
127 was not observed with Δcpb . This reduction in staining intensity correlated with a down-
128 modulation of VAMP3 and VAMP8 proteins in BMM infected with WT *L. mexicana*, compared
129 to cells infected with Δcpb (Figure 1C). These results suggested that *L. mexicana* causes the
130 reduction of VAMP3 and VAMP8 levels in infected BMM through the action of CPB. However,
131 we considered the possibility that CPB acted indirectly on VAMP3 and VAMP8 because we
132 previously found that GP63 targets those SNAREs in *L. major*-infected BMM [1]. We therefore
133 ensured that similar levels of GP63 were present in lysates of BMM infected with WT and Δcpb
134 *L. mexicana* promastigotes. As shown in Figure 2, GP63 was detected in lysates of BMM
135 infected with WT *L. mexicana* up to 72 h post-infection, when the parasites replicate as
136 amastigotes. The important reduction in GP63 levels at this time point is consistent with
137 previously published data showing a 90% reduction in the amount of GP63 detected in

138 amastigotes with respect to promastigotes [24, 25]. Surprisingly, we found that GP63 was barely
139 detectable in BMM infected with Δcpb at all time points tested. This observation raised the
140 possibility that the lack of VAMP3 and VAMP8 down-regulation in Δcpb -infected BMM was
141 due to defective expression of GP63.

142
143 **Figure 1. Down-modulation of VAMP3 and VAMP8 by *L. mexicana*.** BMM were infected
144 with serum-opsonized stationary phase *L. mexicana* (WT and Δcpb) promastigotes for 2 h.
145 VAMP3 (A) and VAMP8 (B) levels (green) were then visualized by confocal microscopy.
146 Macrophage and parasite nuclei are shown in blue (DAPI). Internalized parasites are denoted by
147 white arrowheads. In (C), VAMP3 and VAMP8 levels in total cell extracts were assessed by
148 Western blot analysis. Each immunofluorescence assay was done on 300 phagosomes on
149 triplicate coverslips in two independent experiments and Western blot analyses were performed
150 twice in three independent experiments. VAMP3 and VAMP8/ β -actin ratios were determined by
151 densitometry. Original magnification X63.

152
153 **Figure 2. GP63 is down-modulated in the *L. mexicana* Δcpb mutant.** BMM were infected
154 with serum-opsonized stationary phase *L. mexicana* (WT and Δcpb) promastigotes for 2 h, 24 h,
155 48 h and 72 h. Total cell extracts were assayed for GP63 levels by Western blot analysis. GP63/
156 β -actin ratios were determined by densitometry. Similar results were obtained in three
157 independent experiments.

158
159 ***CPB is required for GP63 expression.*** To address the issue of GP63 down-regulation in *L.*
160 *mexicana* Δcpb , we first determined whether complementation of Δcpb with the *CPB* gene array

161 ($\Delta cpb+CPB$) restores wild type GP63 levels. As shown in Figure 3A, GP63 levels and activity
162 are down-modulated in the Δcpb mutant, and complementation with the CPB array restored
163 GP63 levels and activity similar to those observed in WT parasites. It was previously reported
164 that expression of the cell surface glycolipid LPG and of GP63 may share common biosynthetic
165 steps [26-29]. We therefore evaluated the levels of LPG in lysates of WT, Δcpb , $\Delta cpb+CPB$, and
166 $\Delta cpb+GP63$ parasites by Western blot analysis. Strikingly, similar to GP63, LPG levels were
167 also down-modulated in the Δcpb mutant and complementation with either the CPB array or
168 $GP63$ restored wild type LPG levels. To further investigate the possible role of CPB in the
169 regulation of GP63 expression, we determined the levels of $GP63$ mRNA in WT, Δcpb ,
170 $\Delta cpb+CPB$, and $\Delta cpb+GP63$ parasites by RT-PCR. As shown in Figure 3B, $GP63$ mRNA levels
171 were highly down-regulated in Δcpb and complementation with the CPB array restored wild type
172 levels of GP63 mRNA. Interestingly, complementation of Δcpb with *L. major* $GP63$ did not
173 increase endogenous $GP63$ mRNAs. RT-PCR using *L. major* $GP63$ -specific primers showed that
174 this gene is expressed only in $\Delta cpb+GP63$. Together, these results suggest that CPB controls
175 GP63 mRNA levels at the post-transcriptional level. Clearly, additional studies will be required
176 to elucidate the underlying mechanism(s). Our results also raised the possibility that down-
177 modulation of GP63 in the Δcpb mutant may have accounted for the inability of Δcpb to down-
178 regulate VAMP3 and VAMP8. The finding that expression of GP63 in Δcpb restored LPG levels
179 was unexpected and suggested a role for GP63 in the expression of LPG in *L. mexicana*. As it is
180 estimated that at least 25 genes are required for the synthesis, assembly, and transport of the
181 various components of LPG [30], it may be difficult to determine whether GP63 acts on the
182 expression of a LPG biosynthetic gene or on a biosynthetic step. Assessment of $LPG2$ gene
183 expression revealed that it was equally expressed WT, Δcpb , $\Delta cpb+CPB$, and $\Delta cpb+GP63$
184 parasites. Further studies will be necessary to understand how GP63 expression restores LPG

185 synthesis in Δcpb . Since LPG does not play a major role in the virulence of *L. mexicana* [17], the
186 Δcpb mutant expressing exogenous GP63 provides a unique opportunity to address the impact of
187 GP63 on SNARE cleavage, as well as on the *in vitro* and *in vivo* virulence of *L. mexicana*.

188
189 **Figure 3. Expression of GP63 and LPG is impaired in the absence of CPB.** (A) Stationary
190 phase promastigotes were lysed and total cell extracts were analysed by Western blotting and
191 zymography for GP63 levels and activity and for LPG levels. Aldolase was used as a loading
192 control. GP63 and LPG/aldolase ratios were determined by densitometry. (B) Promastigote total
193 RNA was extracted and reverse transcription followed by PCR was performed to assess mRNA
194 levels for *L. mexicana* GP63-C1, LPG2, and α -tubulin, and *L. major* GP63-1. Similar results
195 were obtained in three independent experiments.

196
197 **GP63 is responsible for the cleavage of VAMP3 and VAMP8 by *L. mexicana*.** We next
198 assessed the impact of GP63 on VAMP3 and VAMP8 during *L. mexicana* infection. To this end,
199 we infected BMM with either WT, Δcpb , $\Delta cpb+CPB$, or $\Delta cpb+GP63$ *L. mexicana* promastigotes
200 for various time points, and we assessed VAMP3 and VAMP8 levels and intracellular
201 distribution. Figure 4A shows that GP63 is present at high levels in lysates of BMM infected for
202 2 h with WT, $\Delta cpb+CPB$, and $\Delta cpb+GP63$ promastigotes (compared to lysates of BMM infected
203 with Δcpb). At 72 h post-infection, GP63 levels are strongly reduced in BMM infected with WT
204 and $\Delta cpb+CPB$, whereas they remain elevated in BMM infected with the $\Delta cpb+GP63$ (Figure
205 4A) [25]. The high levels of GP63 present in BMM infected with $\Delta cpb+GP63$ for 72 h may be
206 related to the fact that expression of the *L. major* GP63 gene from the pLEXNeo episomal vector
207 [31] is not under the control of endogenous GP63 3' untranslated regions. Western blot analyses

208 revealed that down-regulation of VAMP3 and VAMP8 correlated with GP63 levels expressed by
209 the parasites. Consistently, the staining intensity of VAMP3 and VAMP8 was reduced in BMM
210 infected with GP63-expressing parasites, as assessed by confocal immunofluorescence
211 microscopy (Figure 4D and E). These results suggest that GP63 is responsible for the down-
212 modulation of the endosomal SNAREs VAMP3 and VAMP8 in *L. mexicana*-infected BMM.
213 Interestingly, we observed recruitment of VAMP3 to PVs containing *L. mexicana* parasites at
214 later time points, when promastigotes have differentiated into amastigotes, with the exception of
215 $\Delta cpb+GP63$ *L. mexicana* promastigotes (Figure 4B). In contrast, we found that VAMP8 is
216 excluded from *L. mexicana*-containing PVs both at early and later time points post-infection,
217 independently of GP63 levels, suggesting that additional mechanisms regulate VAMP8
218 recruitment to *L. mexicana* PVs.

219
220 **Figure 4. GP63 is responsible for the down-modulation of VAMP3 and VAMP8 in *L.***
221 ***mexicana*-infected macrophages.** BMM were infected with serum-opsonized stationary phase
222 *L. mexicana* (WT, Δcpb , $\Delta cpb+CPB$ and $\Delta cpb+GP63$) promastigotes for 2 h and 72 h. Total cell
223 extracts were analysed by Western blot (A). Similar results were obtained in three independent
224 experiments. VAMP3 and VAMP8 recruitment to the phagosome was visualized by
225 immunofluorescence microscopy and quantified for 300 phagosomes on triplicate coverslips (B
226 and C). Representative pictures from each condition are shown (D and E). Immunofluorescence
227 assays were performed on 300 phagosomes on triplicate coverslips for three independent
228 experiments. * $p < 0.0001$. Original magnification X63.

229
230 ***GP63* expression restores virulence of Δcpb .** Since GP63 was shown to contribute to *L. major*
231 virulence [32], we next sought to determine whether expression of GP63 is sufficient to restore

232 the ability of Δcpb to replicate inside macrophages and to cause lesions in mice [19]. To this
233 end, we first infected BMM with either WT, Δcpb , $\Delta cpb+CPB$, or $\Delta cpb+GP63$ stationary phase
234 promastigotes and we assessed parasite burden and PV surface area at various time points post-
235 infection. We found that Δcpb had an impaired capacity to replicate inside macrophages and to
236 induce the formation of large communal PVs compared to WT and $\Delta cpb+CPB$ parasites (Figure
237 5A, B, C). Strikingly, expression of GP63 in Δcpb restored its ability to replicate in macrophages
238 and to induce large communal PVs up to 72 h post-infection. These results underline the role of
239 GP63 in the ability of *L. mexicana* to infect and replicate in macrophages, even in the absence of
240 CPB. Following inoculation inside the mammalian host, promastigotes are exposed to
241 complement and both GP63 and LPG confer resistance to complement-mediated lysis [32, 33].
242 *L. mexicana* promastigotes were therefore analyzed for their sensitivity to complement-mediated
243 lysis in the presence of fresh human serum. As shown in Figure 6A, over 40% of Δcpb was
244 killed after 30 min in the presence of 20% serum, whereas $\Delta cpb+CPB$, and $\Delta cpb+GP63$ were
245 more resistant to serum lysis at 14% and 10%, respectively. Absence of both GP63 and LPG
246 may be responsible for the serum sensitivity of Δcpb . Finally, to assess the impact of GP63 on
247 the ability of Δcpb to cause lesions, we used a mouse model of cutaneous leishmaniasis.
248 Susceptible BALB/c mice were infected in the hind footpad with either WT, Δcpb , $\Delta cpb+CPB$,
249 or $\Delta cpb+GP63$ promastigotes and disease progression was monitored for 9 weeks. Consistent
250 with its reduced capacity to replicate inside macrophages, Δcpb failed to cause significant lesions
251 compared to WT parasites [19] and Δcpb complemented with *CPB* (Figure 6B). Remarkably,
252 expression of GP63 in Δcpb restored its capacity to cause lesions, albeit to a lower level than
253 Δcpb complemented with *CPB*. Lesion size correlated with parasite burden, as measured at 9
254 weeks post-infection (Figure 6C). Collectively, these results indicate that expression of GP63 is
255 sufficient to restore virulence of Δcpb .

256
257 **Figure 5. GP63 enables *L. mexicana* Δcpb to infect macrophages and induce large PVs.**
258 BMM were infected with stationary phase serum-opsonized *L. mexicana* (WT, Δcpb , $\Delta cpb+cpb$
259 and $\Delta cpb+GP63$) promastigotes for 2 h, 24 h, 48 h and 72 h. Macrophages were stained with the
260 HEMA 3TM kit. Representative pictures from each condition are shown (A) Parasites were
261 counted in 300 macrophages on triplicate coverslips (B). Macrophages were stained with the
262 LAMP1 antibody and vacuole sizes were measured with the ZEN 2012 software (C). Parasitemia
263 and vacuole size was determined on 300 phagosomes in triplicate in three independent
264 experiments. *p<0.0001.

265
266 **Figure 6. GP63 confers virulence to *L. mexicana* Δcpb .** Stationary phase *L. mexicana* (WT,
267 Δcpb , $\Delta cpb+cpb$ and $\Delta cpb+GP63$) promastigotes were incubated in the presence of 20% human
268 serum for 30 min, stained with a fixable viability dye, and then subjected to flow cytometry (A).
269 Mice were challenged with 5×10^6 late-stationary phase *L. mexicana* (WT, Δcpb , $\Delta cpb+cpb$ and
270 $\Delta cpb+GP63$) promastigotes that were injected subcutaneously into the hind footpad. Disease
271 progression was monitored at weekly intervals, by measuring the thickness of the infected
272 footpad and the contralateral uninfected footpad. (B). Parasite burden was obtained by limiting
273 dilution assay from infected homogenized footpads 9 weeks after inoculation (C). Human serum
274 lyses were performed in two independent experiments and six mice per group were used for the
275 determination of lesion formation and parasite burden. Each data point represents the average \pm
276 SEM of 6 mice per group, and statistical significance was denoted by * p \leq 0.01, and ***
277 p \leq 0.0001.

DISCUSSION

278
279
280 This study aimed at investigating the mechanism(s) by which CPB contributes to *L.*
281 *mexicana* virulence. To this end, we initially examined PV biogenesis by assessing the impact of
282 *L. mexicana* infection on the trafficking of VAMP3 and VAMP8, two endocytic SNAREs
283 associated with phagosome biogenesis and function [1, 34]. We found that both SNAREs were
284 down-modulated in a CPB-dependent manner, which hampered VAMP3 recruitment to PVs. We
285 also discovered that expression of GP63, which we previously showed to be responsible for
286 cleaving SNAREs in *L. major*-infected macrophages [1], was down-modulated in the *L.*
287 *mexicana* Δcpb . Strikingly, restoration of GP63 expression in Δcpb bypassed the need for CPB
288 for SNARE cleavage. Similarly, episomal expression of GP63 enabled the Δcpb mutant to
289 establish infection in macrophages, induce larger PVs and cause lesions in mice. These findings
290 imply that CPB contributes to *L. mexicana* virulence in part through the regulation of GP63
291 expression, and provide evidence that GP63 is a key virulence factor for *L. mexicana*.

292
293 The observation that CPB regulates GP63 expression at the mRNA levels was both
294 unexpected and intriguing. Insight into the possible mechanism(s) may be deduced from a recent
295 study on the role of cathepsin B in *L. donovani*, which is homologous to the *L. mexicana* CPC
296 [35]. Similar to *L. mexicana* Δcpb , *L. donovani* $\Delta catB$ displays reduced virulence in
297 macrophages. To investigate the role of cathepsin B in virulence, the authors performed
298 quantitative proteome profiling of WT and $\Delta catB$ parasites and identified 83 proteins whose
299 expression is altered in the absence of cathepsin B, with the majority being down-modulated [35].
300 Among those were a group of proteins involved in post-transcriptional regulation of gene
301 expression (RNA stability, processing, translation) [35]. Whether this is the case in Δcpb

302 deserves further investigation. Clearly, a detailed analysis of wild-type and Δcpb parasites may
303 provide the information required to understand the extent of the impact of CBP on the expression
304 and synthesis of virulence factors and the exact role of CPB in *L. mexicana* virulence. The
305 observation that episomal expression of GP63 in Δcpb restored LPG synthesis is an intriguing
306 issue, as it suggests that GP63 acts on a LPG biosynthetic step. This role for GP63 is likely
307 redundant, since *L. major* $\Delta gp63$ promastigotes express LPG levels similar to that of wild type
308 parasites (S1 Fig).

309
310 It has been proposed that expansion of the PVs hosting parasites of the *L. mexicana*
311 complex leads to the dilution of the microbicidal effectors to which the parasites are exposed,
312 thereby contributing to parasite survival [36]. Both host and parasite factors may be involved in
313 the control of PV enlargement. On the host side, it has been previously reported that *L.*
314 *amazonensis* cannot survive in cells overexpressing *LYST*, a host gene that restricts *Leishmania*
315 growth by counteracting PV expansion [37]. Similarly, disrupting the fusion between PVs
316 housing *L. amazonensis* and the endoplasmic reticulum resulted in limited PV expansion and
317 inhibition of parasite replication [15, 16]. On the parasite side, virulence of *L. amazonensis*
318 isolates was shown to correlate with the ability to induce larger PVs [38]. Our results indicate
319 that the inability of Δcpb to multiply inside macrophages is associated with smaller PV size, and
320 that expression of GP63 is sufficient to restore the capacity of Δcpb to survive within
321 macrophages and to induce PV expansion. How does GP63 modulate *L. mexicana* virulence and
322 PV expansion? In addition to the numerous macrophage proteins known to be targeted by GP63,
323 it is possible that SNARE cleavage is one of the factors associated with *L. mexicana* virulence
324 and PV expansion. For instance, we previously reported that VAMP8 is required for phagosomal
325 oxidative activity [1]. One may envision that its degradation by GP63 is part of a strategy used

326 by *L. mexicana* to establish infection in an environment devoid of oxidants, thereby contributing
327 to parasite survival. The LYST protein is a regulator of lysosome size and its absence leads to
328 further PV expansion and enhanced *L. amazonensis* replication [37]. It is interesting to note that
329 LYST was proposed to function as an adaptor protein that juxtaposes proteins such as SNAREs
330 that mediate intracellular membrane fusion reactions [39]. In this context, cleavage of SNAREs
331 that interact with LYST may interfere with its function and promote PV expansion. Further
332 studies will be necessary to clarify these issues, including the potential role of VAMP3 and
333 VAMP8 in PV biogenesis and expansion.

334
335 Previous studies using Δcpb parasites led to the conclusion that CPB enables *L. mexicana*
336 to alter host cell signaling and gene expression through the cleavage of various host proteins [20,
337 22]. Hence, CPB-dependent cleavage of PTP-1B, NF- κ B, STAT1, and AP1 in *L. mexicana*-
338 infected macrophages was associated with the inhibition of IL-12 expression and generation of
339 nitric oxide, both of which are important for initiation of a host immune response and parasite
340 killing, respectively. Our finding that GP63 expression is down-modulated in the Δcpb mutant
341 raises the possibility that cleavage of those transcription factors may actually be mediated by
342 GP63. Indeed, GP63 cleaves numerous host macrophage effectors, including PTP-1B, NF- κ B,
343 STAT1, and AP1 [40]. Revisiting the role of CPB in the context of GP63 expression will be
344 necessary to elucidate whether, and to which extent, CPB is acting directly on the host cell
345 proteome.

346
347 In sum, we discovered that CPB contributes to *L. mexicana* virulence in part through the
348 regulation of GP63 expression. Complementation of Δcpb revealed the importance of GP63 for

349 the virulence of *L. mexicana*, as it participates in the survival of intracellular parasites, PV
350 expansion, and the formation of cutaneous lesions.

351

MATERIALS AND METHODS

352

353

354 **Ethics statement**

355 Experiments involving mice were done as prescribed by protocol 1406-02, which was approved
356 by the *Comité Institutionnel de Protection des Animaux* of the INRS-Institut Armand-Frappier. *In*
357 *vivo* infections were performed as per Animal Use Protocol #4859, which was approved by the
358 Institutional Animal Care and Use Committees at McGill University. These protocols respect
359 procedures on good animal practice provided by the Canadian Council on Animal Care (CCAC).

360

361 **Antibodies**

362 The mouse anti-GP63 monoclonal antibody was previously described [41]. The mouse anti-
363 phosphoglycans CA7AE monoclonal antibody [42] was from Cedarlane and the rabbit polyclonal
364 anti-aldolase was a gift from Dr. A. Jardim (McGill University). Rabbit polyclonal antibodies for
365 VAMP3 and VAMP8 were obtained from Synaptic Systems.

366

367 **Cell culture**

368 Bone marrow-derived macrophages (BMM) were differentiated from the bone marrow of 6- to 8-
369 week-old female 129XB6 mice (Charles River Laboratories) as previously described [43]. Cells
370 were cultured for 7 days in complete medium (DMEM [Life Technologies] supplemented with L-
371 glutamine [Life Technologies], 10% heat-inactivated FBS [PAA Laboratories], 10 mM HEPES at
372 pH 7.4, and antibiotics) containing 15% v/v L929 cell-conditioned medium as a source of M-
373 CSF. Macrophages were kept at 37°C in a humidified incubator with 5% CO₂. To render BMM
374 quiescent prior to experiments, cells were transferred to 6- or 24-well tissue culture microplates
375 (TrueLine) and kept for 16 h in complete DMEM without L929 cell-conditioned medium.

376 Promastigotes of *L. mexicana* wild-type strain (MNYC/BZ/62/M379) and of *L. major* NIH S
377 (MHOM/SN/74/Seidman) clone A2 were grown at 26°C in *Leishmania* medium (Medium 199
378 supplemented with 10% heat-inactivated FBS, 40 mM HEPES pH 7.4, 100 µM hypoxanthine, 5
379 µM hemin, 3 µM biopterin, 1 µM biotin, and antibiotics). The isogenic *L. mexicana* CPB-
380 deficient mutant Δcpb^{pac} (thereafter referred to as Δcpb) and its complemented counterpart
381 Δcpb^{pac} [pGL263] (thereafter referred to as $\Delta cpb+CPB$) were described previously [21]. *L.*
382 *mexicana* Δcpb promastigotes were electroporated as described [44] with the pLEXNeoGP63.1
383 plasmid [32] to generate $\Delta cpb+GP63$ parasites. *L. mexicana* $\Delta cpb+CPB$ and $\Delta cpb+GP63$
384 promastigotes were grown in the presence of 50 µg/ml hygromycin or 50 µg/ml G418,
385 respectively. The *L. major* NIH clone A2 isogenic $\Delta gp63$ mutant and its complemented
386 counterpart $\Delta gp63+gp63$ have been previously described [32]. Cultures of $\Delta gp63+gp63$
387 promastigotes were supplemented with 50 µg/ml G418.

388

389 **Synchronized phagocytosis assays**

390 Prior to macrophage infections, promastigotes in late stationary phase were opsonized with
391 DBA/2 mouse serum. For synchronized phagocytosis using parasites, macrophages and parasites
392 were incubated at 4°C for 10 min and spun at 167 g for 1 min, and internalization was triggered
393 by transferring cells to 34°C. Macrophages were washed twice after 2h with complete DMEM to
394 remove the non-internalized parasites and were further incubated at 34°C for the required times.
395 Cells were then washed with PBS and stained using the Hema 3™ Fixative and Solutions kit
396 (Fisher HealthCare), or prepared for confocal immunofluorescence microscopy.

397

398

399 **Confocal immunofluorescence microscopy**

400 Macrophages on coverslips were fixed with 2% paraformaldehyde (Canemco and Mirvac) for 40
401 min and blocked/permeabilized for 17 min with a solution of 0.05% saponin, 1% BSA, 6% skim
402 milk, 2% goat serum, and 50% FBS. This was followed by a 2 h incubation with primary
403 antibodies diluted in PBS. Macrophages were then incubated with a suitable combination of
404 secondary antibodies (anti-rabbit Alexa Fluor 488 and anti-rat 568; Molecular Probes) and DAPI
405 (Life technologies). Coverslips were washed three times with PBS after every step. After the final
406 washes, Fluoromount-G (Southern Biotechnology Associates) was used to mount coverslips on
407 glass slides, and coverslips were sealed with nail polish (Sally Hansen). Macrophages were
408 imaged with the 63X objective of an LSM780 microscope (Carl Zeiss Microimaging), and
409 images were taken in sequential scanning mode. Image analysis and vacuole size measurements
410 were performed with the ZEN 2012 software.

411

412 **Electrophoresis, Western blotting, and zymography**

413 Prior to lysis, macrophages were placed on ice and washed with PBS containing 1 mM sodium
414 orthovanadate and 5 mM 1,10-phenanthroline (Roche). Cells were scraped in the presence of
415 lysis buffer containing 1% Nonidet P-40 (Caledon), 50 mM Tris-HCl (pH 7.5) (Bioshop), 150
416 mM NaCl, 1 mM EDTA (pH 8), 5 mM 1,10-phenanthroline, and phosphatase and protease
417 inhibitors (Roche). Parasites were washed twice with PBS and lysed in the presence of lysis
418 buffer containing 0.5% Nonidet P-40 (Caledon), 100mM Tris-HCl (Bioshop) and 150 mM NaCl
419 at -70° C. Lysates were thawed on ice and centrifuged for 10 min to remove insoluble matter.
420 After protein quantification, protein samples were boiled (100°C) for 6 min in SDS sample buffer
421 and migrated in 12% or 15% SDS-PAGE gels. Three micrograms and 15 µg of protein were
422 loaded for parasite and infected macrophage lysates, respectively. Proteins were transferred onto

423 Hybond-ECL membranes (Amersham Biosciences), blocked for 1 h in TBS 1X-0.1% Tween
424 containing 5% skim milk, incubated with primary antibodies (diluted in TBS 1X-0.1% Tween
425 containing 5% BSA) overnight at 4°C, and thence with appropriate HRP-conjugated secondary
426 antibodies for 1 h at room temperature. Then, membranes were incubated in ECL (GE
427 Healthcare), and immunodetection was achieved via chemiluminescence. Membranes were
428 washed 3 times between each step. For zymography, 2 µg of lysate were incubated at RT for 6
429 min in sample buffer without DTT and then migrated in 12% SDS-PAGE gels with 0.2% gelatin
430 (Sigma). Gels were incubated for 1 h in the presence of 50 mM Tris pH 7.4, 2,5% Triton X-100,
431 5 mM CaCl₂ and 1 µM ZnCl₂ and incubated overnight in the presence of 50 mM Tris pH 7.4, 5
432 mM CaCl₂, 1µM ZnCl₂ and 0,01% NaN₃ at 37°C [45].

433

434 **FACS analysis**

435 Late stationary phase promastigotes were incubated for 30 min in complete DMEM medium with
436 20% human serum from healthy donors. Parasites were then incubated in LIVE/DEAD® Fixable
437 Violet Dead Cell Stain Kit (Life technologies) and fixed in 2% paraformaldehyde. Flow
438 cytometric analysis was carried out using the LSRFortessa cytometer (Special Order Research
439 Product; BD Biosciences), and the BD FACSDiva Software (version 6.2) was used for data
440 acquisition and analysis.

441

442 **Mouse infections**

443 Male BALB/c mice (6 to 8 weeks old) were purchased from Charles River Laboratories and
444 infected in the right hind footpad with 5x10⁶ stationary phase *L. mexicana* promastigotes as
445 described [46]. Disease progression was monitored by measuring footpad swelling weekly with a

446 metric caliper, for up to 9 weeks post-infection. Footpads were then processed to calculate
447 parasite burden using the limiting dilution assay.

448

449 **Limiting dilution assay**

450 After 9 weeks of infection, mice were euthanized under CO₂ asphyxiation and subsequently by
451 cervical dislocation. The infected footpads were surface-sterilized with a chlorine dioxide based
452 disinfectant followed by ethanol 70% for 5 min. Footpads were washed in PBS, lightly sliced,
453 transferred to a glass tissue homogenizer containing 6 ml of PBS, and manually homogenized.
454 The last step was repeated two to three times, until complete tissue disruption was achieved.
455 Final homogenate was then centrifuged at 3,000 x g for 5 min and resuspended in the appropriate
456 volume of PBS. 100 µl of homogenate were added in duplicates to 96-well plates containing 100
457 µl of complete Schneider's medium in each well (twenty-four 2-fold dilutions for each duplicate).
458 Plates were incubated at 28°C. After 7-10 days, the number of viable parasites was determined
459 from the highest dilutions at which promastigotes were observed using an inverted microscope
460 [47].

461

462 **Reverse transcription-PCR (RT-PCR)**

463 Total RNA was extracted from promastigotes using the TRIzol reagent (Invitrogen Life
464 Technology, Carlsbad, CA) and reverse transcribed. One-tenth of the resulting cDNA was
465 amplified by PCR on a DNA thermal cycler, version 2.3 (Perkin-Elmer Corporation, Norwalk,
466 CT), with the following primer pairs: for the *L. mexicana* GP63 C-1 5'-
467 ACCGTCTGAGAGTCGGA ACT-3' (forward), 5'-GTAGTCCAGGAATGGCGAGT-3'
468 (reverse); the *L. major* GP63-1 5'-TCTGAGGCACATGCTTCGTT-3' (forward), 5'-
469 GTCAGTTGCCTTCGGTCTGA-3' (reverse), the *L. mexicana* LPG2

470 5'CATTGGTATCCTGGTGCTG-3' (forward), 5'-GAGGAAGCCACTGTTAGCC-3' (reverse),
471 and the *L. mexicana* α -tubulin 5'-CTATCTGCATCCACATTGGC-3' (forward), 5'-
472 ACTTGTCAGAGGGCATGGA-3' (reverse). The PCR products were analyzed by
473 electrophoresis on a 3% (wt/vol) agarose gel, and the pictures were taken using AlphaImager
474 3400 imaging software (Alpha Innotech Corporation, San Leandro, CA).

475

476 **Statistical analyses**

477 Statistically significant differences were analyzed by ANOVA followed by the Tukey post-hoc
478 test using the Graphpad Prism program (version 5.0). For the limiting dilution assay, the non-
479 parametric Mann-Whitney or Kruskal-Wallis test was used. Values starting at $P < 0.05$ were
480 considered statistically significant. All data are presented as mean \pm SEM.

481

482

483 **Acknowledgments**

484 We are grateful to Dr. Armando Jardim (McGill University) for providing the anti-aldolase
485 antiserum, Jessy Tremblay for assistance in immunofluorescence experiments, and Guillermo
486 Arango Duque and Dr. Simona Stäger for critical comments.

487

488

REFERENCES

489

490 1. Matheoud D, Moradin N, Bellemare-Pelletier A, Shio MT, Hong WJ, Olivier M, et al.
491 *Leishmania* evades host immunity by inhibiting antigen cross-presentation through direct
492 cleavage of the SNARE VAMP8. *Cell Host Microbe*. 2013;14: 15-25.

493 2. Moradin N, Descoteaux A. *Leishmania* promastigotes: building a safe niche within
494 macrophages. *Front Cell Infect Microbiol*. 2012;2: 121.

495 3. Desjardins M, Descoteaux A. Inhibition of phagolysosomal biogenesis by the *Leishmania*
496 lipophosphoglycan. *J Exp Med*. 1997;185: 2061-2068.

497 4. Vinet AF, Fukuda M, Turco SJ, Descoteaux A. The *Leishmania donovani*
498 lipophosphoglycan excludes the vesicular proton-ATPase from phagosomes by impairing the
499 recruitment of synaptotagmin V. *PLoS Pathog*. 2009;5: e1000628.

500 5. Olivier M, Atayde VD, Isnard A, Hassani K, Shio MT. *Leishmania* virulence factors:
501 focus on the metalloprotease GP63. *Microbes Infect*. 2012;14: 1377-1389.

502 6. Arango Duque G, Descoteaux A. *Leishmania* survival in the macrophage: where the ends
503 justify the means. *Curr Opin Microbiol*. 2015;26: 32-40.

504 7. Arango Duque G, Fukuda M, Turco SJ, Stäger S, Descoteaux A. *Leishmania*
505 promastigotes induce cytokine secretion in macrophages through the degradation of
506 synaptotagmin XI. *J Immunol*. 2014;193: 2363-2372.

507 8. Courret N, Frehel C, Gouhier N, Pouchelet M, Prina E, Roux P, et al. Biogenesis of
508 *Leishmania*-harbouring parasitophorous vacuoles following phagocytosis of the metacyclic
509 promastigote or amastigote stages of the parasites. *J Cell Sci*. 2002;115: 2303-2316.

- 510 9. Alexander J, Vickerman K. Fusion of host cell secondary lysosomes with the
511 parasitophorous vacuoles of *Leishmania mexicana*-infected macrophages. J Protozool. 1975;22:
512 502-508.
- 513 10. Barbieri CL, Brown K, Rabinovitch M. Depletion of secondary lysosomes in mouse
514 macrophages infected with *Leishmania mexicana amazonensis*: a cytochemical study. Z
515 Parasitenkd. 1985;71: 159-168.
- 516 11. Real F, Pouchelet M, Rabinovitch M. *Leishmania (L.) amazonensis*: fusion between
517 parasitophorous vacuoles in infected bone-marrow derived mouse macrophages. Exp Parasitol.
518 2008;119: 15-23.
- 519 12. Real F, Mortara RA. The diverse and dynamic nature of *Leishmania* parasitophorous
520 vacuoles studied by multidimensional imaging. PLoS Negl Trop Dis. 2012;6: e1518.
- 521 13. Stow JL, Manderson AP, Murray RZ. SNAREing immunity: the role of SNAREs in the
522 immune system. Nat Rev Immunol. 2006;6: 919-929.
- 523 14. Ndjamen B, Kang BH, Hatsuzawa K, Kima PE. *Leishmania* parasitophorous vacuoles
524 interact continuously with the host cell's endoplasmic reticulum; parasitophorous vacuoles are
525 hybrid compartments. Cell Microbiol. 2010;12: 1480-1494.
- 526 15. Canton J, Ndjamen B, Hatsuzawa K, Kima PE. Disruption of the fusion of *Leishmania*
527 parasitophorous vacuoles with ER vesicles results in the control of the infection. Cell Microbiol.
528 2012.
- 529 16. Canton J, Kima PE. Targeting host syntaxin-5 preferentially blocks *Leishmania*
530 parasitophorous vacuole development in infected cells and limits experimental *Leishmania*
531 infections. Am J Pathol. 2012;181: 1348-1355.
- 532 17. Ilg T. Lipophosphoglycan is not required for infection of macrophages or mice by
533 *Leishmania mexicana*. EMBO J. 2000;19: 1953-1962.

- 534 18. Mottram JC, Coombs GH, Alexander J. Cysteine peptidases as virulence factors of
535 *Leishmania*. *Curr Opin Microbiol*. 2004;7: 375-381.
- 536 19. Mottram JC, Souza AE, Hutchison JE, Carter R, Frame MJ, Coombs GH. Evidence from
537 disruption of the *lmcpb* gene array of *Leishmania mexicana* that cysteine proteinases are
538 virulence factors. *Proc Natl Acad Sci USA*. 1996;93: 6008-6013.
- 539 20. Cameron P, McGachy A, Anderson M, Paul A, Coombs GH, Mottram JC, et al. Inhibition
540 of lipopolysaccharide-induced macrophage IL-12 production by *Leishmania mexicana*
541 amastigotes: the role of cysteine peptidases and the NF-kappaB signaling pathway. *J Immunol*.
542 2004;173: 3297-3304.
- 543 21. Denise H, McNeil K, Brooks DR, Alexander J, Coombs GH, Mottram JC. Expression of
544 multiple CPB genes encoding cysteine proteases is required for *Leishmania mexicana* virulence
545 in vivo. *Infect Immun*. 2003;71: 3190-3195.
- 546 22. Abu-Dayyeh I, Hassani K, Westra ER, Mottram JC, Olivier M. Comparative study of the
547 ability of *Leishmania mexicana* promastigotes and amastigotes to alter macrophage signaling and
548 functions. *Infect Immun*. 2010;78: 2438-2445.
- 549 23. De Souza Leao S, Lang T, Prina E, Hellio R, Antoine JC. Intracellular *Leishmania*
550 *amazonensis* amastigotes internalize and degrade MHC class II molecules of their host cells. *J*
551 *Cell Sci*. 1995;108: 3219-3231.
- 552 24. Bahr V, Stierhof YD, Ilg T, Demar M, Quinten M, Overath P. Expression of
553 lipophosphoglycan, high-molecular weight phosphoglycan and glycoprotein 63 in promastigotes
554 and amastigotes of *Leishmania mexicana*. *Mol Biochem Parasitol*. 1993;58: 107-121.
- 555 25. Medina-Acosta E, Karess RE, Schwartz H, Russell DG. The promastigote surface
556 protease (gp63) of *Leishmania* is expressed but differentially processed and localized in the
557 amastigote stage. *Mol Biochem Parasitol*. 1989;37: 263-273.

- 558 26. Garami A, Ilg T. Disruption of mannose activation in *Leishmania mexicana*: GDP-
559 mannose pyrophosphorylase is required for virulence, but not for viability. EMBO J. 2001;20:
560 3657-3666.
- 561 27. Garami A, Ilg T. The role of phosphomannose isomerase in *Leishmania mexicana*
562 glycoconjugate synthesis and virulence. J Biol Chem. 2001;276: 6566-6575.
- 563 28. Garami A, Mehlert A, Ilg T. Glycosylation defects and virulence phenotypes of
564 *Leishmania mexicana* phosphomannomutase and dolicholphosphate-mannose synthase gene
565 deletion mutants. Mol Cell Biol. 2001;21: 8168-8183.
- 566 29. Descoteaux A, Avila HA, Zhang K, Turco SJ, Beverley SM. *Leishmania* LPG3 encodes a
567 GRP94 homolog required for phosphoglycan synthesis implicated in parasite virulence but not
568 viability. EMBO J. 2002;21: 4458-4469.
- 569 30. Beverley SM, Turco SJ. Lipophosphoglycan (LPG) and the identification of virulence
570 genes in the protozoan parasite *Leishmania*. Trends Microbiol. 1998;6: 35-40.
- 571 31. Joshi PB, Webb JR, Davies JE, McMaster WR. The gene encoding streptothricin
572 acetyltransferase (sat) as a selectable marker for *Leishmania* expression vectors. Gene. 1995;156:
573 145-149.
- 574 32. Joshi PB, Kelly BL, Kamhawi S, Sacks DL, McMaster WR. Targeted gene deletion in
575 *Leishmania major* identifies leishmanolysin (GP63) as a virulence factor. Mol Biochem Parasitol.
576 2002;120: 33-40.
- 577 33. Spath GF, Garraway LA, Turco SJ, Beverley SM. The role(s) of lipophosphoglycan
578 (LPG) in the establishment of *Leishmania major* infections in mammalian hosts. Proc Natl Acad
579 Sci USA. 2003;100: 9536-9541.
- 580 34. Murray RZ, Kay JG, Sangermani DG, Stow JL. A role for the phagosome in cytokine
581 secretion. Science. 2005;310: 1492-1495.



- 582 35. Gerbaba TK, Gedamu L. Cathepsin B gene disruption induced *Leishmania donovani*
583 proteome remodeling implies cathepsin B role in secretome regulation. PLoS One. 2013;8:
584 e79951.
- 585 36. Sacks D, Sher A. Evasion of innate immunity by parasitic protozoa. Nat Immunol.
586 2002;3: 1041-1047.
- 587 37. Wilson J, Huynh C, Kennedy KA, Ward DM, Kaplan J, Aderem A, et al. Control of
588 parasitophorous vacuole expansion by LYST/Beige restricts the intracellular growth of
589 *Leishmania amazonensis*. PLoS Pathog. 2008;4: e1000179.
- 590 38. Franca-Costa J, Wanderley JL, Deolindo P, Zarattini JB, Costa J, Soong L, et al. Exposure
591 of phosphatidylserine on *Leishmania amazonensis* isolates is associated with diffuse cutaneous
592 leishmaniasis and parasite infectivity. PLoS One. 2012;7: e36595.
- 593 39. Tchernev VT, Mansfield TA, Giot L, Kumar AM, Nandabalan K, Li Y, et al. The
594 Chediak-Higashi protein interacts with SNARE complex and signal transduction proteins. Mol
595 Med. 2002;8: 56-64.
- 596 40. Isnard A, Shio MT, Olivier M. Impact of *Leishmania* metalloprotease GP63 on
597 macrophage signaling. Front Cell Infect Microbiol. 2012;2: 72.
- 598 41. Button LL, Wilson G, Astell CR, McMaster WR. Recombinant *Leishmania* surface
599 glycoprotein GP63 is secreted in the baculovirus expression system as a latent metalloproteinase.
600 Gene. 1993;134: 75-81.
- 601 42. Tolson DL, Turco SJ, Beecroft RP, Pearson TW. The immunochemical structure and
602 surface arrangement of *Leishmania donovani* lipophosphoglycan determined using monoclonal
603 antibodies. Mol Biochem Parasitol. 1989;35: 109-118.
- 604 43. Descoteaux A, Matlashewski G. *c-fos* and tumor necrosis factor gene expression in
605 *Leishmania donovani*- infected macrophages. Mol Cell Biol. 1989;9: 5223-5227.


- 606 44. Descoteaux A, Garraway LA, Ryan KA, Garrity LK, Turco SJ, Beverley SM.
607 Identification of genes by functional complementation in protozoan parasite *Leishmania*. In:
608 Adolph KW, editor. Methods in Molecular Genetics (Molecular Microbiology Techniques).
609 Molecular Biology Techniques. 3. San Diego: Academic Press; 1994. p. 22-48.
- 610 45. Hassani K, Shio MT, Martel C, Faubert D, Olivier M. Absence of metalloprotease GP63
611 alters the protein content of *Leishmania* exosomes. PLoS One. 2014;9: e95007.
- 612 46. Gomez MA, Contreras I, Halle M, Tremblay ML, McMaster RW, Olivier M. *Leishmania*
613 GP63 alters host signaling through cleavage-activated protein tyrosine phosphatases. Sci Signal.
614 2009;2: ra58.
- 615 47. Lima HC, Bleyenbergh JA, Titus RG. A simple method for quantifying *Leishmania* in
616 tissues of infected animals. Parasitol Today. 1997;13: 80-82.
- 617







618 **Supporting Information**

619

620 **S1 Fig. *L. major* Δ gp63 promastigotes express normal levels of LPG.** Stationary phase
621 promastigotes were lysed and total cell extracts were analysed by Western blotting for LPG
622 levels. Similar results were obtained in two separate experiments.







L. mexicana

