

This is a repository copy of *How do Zimbabweans value health states?*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/1038/>

Article:

Jelsma, J., Hansen, K., de Weerd, W. et al. (2 more authors) (2003) How do Zimbabweans value health states? Population Health Metrics. ISSN 1478-7954

<https://doi.org/10.1186/1478-7954-1-11>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Research

Open Access

How do Zimbabweans value health states?

Jennifer Jelsma*¹, Kristian Hansen², Willy de Weerd³, Paul de Cock⁴ and Paul Kind⁵

Address: ¹Division of Physiotherapy, University of Cape Town, Anzio Road, Observatory, South Africa, ²Department of Health Services Research, University of Copenhagen, ³Faculteit Lichamelijke Opvoeding en Kinesitherapie, Katholieke Universiteit Leuven, Belgium, ⁴Centrum voor Ontwikkelingsstoornissen, Faculteit Geneeskunde, Katholieke Universiteit, Leuven, Belgium and ⁵University of York; Department of Preventive Medicine, University of Wisconsin

Email: Jennifer Jelsma* - jjelsma@uctgsh1.uct.ac.za; Kristian Hansen - K.S.Hansen@pubhealth.ku.dk; Willy de Weerd - willy.deweerd@flok.kuleuven.ac.be; Paul de Cock - Paul.DeCock@uz.kuleuven.ac.be; Paul Kind - pk1@york.ac.uk

* Corresponding author

Published: 16 December 2003

Received: 03 July 2003

Population Health Metrics 2003, 1:11

Accepted: 16 December 2003

This article is available from: <http://www.pophealthmetrics.com/content/1/1/11>

© 2003 Jelsma et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Quality of life weights based on valuations of health states are often used in cost utility analysis and population health measures. This paper reports on an attempt to develop quality of life weights within the Zimbabwe context.

Methods: 2,384 residents in randomly selected small residential plots of land in a high-density suburb of Harare valued descriptors of 38 health states based on different combinations of the five domains of the EQ-5D (mobility, self-care, usual activities, pain or discomfort and anxiety or depression). The English version of the EQ-5D was used. The time trade-off method was used to determine the values, and 19,020 individual preferences for health states were analysed. A residual maximum likelihood linear mixed model was used to estimate a function for predicting the values of all possible combinations of levels on the five domains. The model was fit to a random subset of two-thirds of the observations, with the remaining observations reserved for analysis of predictive validity. The results were compared to a similar study undertaken in the United Kingdom.

Results: A credible model was developed to predict the values of states that were not valued directly. In the subset of observations reserved for validation, the mean absolute difference between predicted and observed values was 0.045. All domains of the EQ-5D were found to contribute significantly to the model, both at the moderate and severe levels. Severe pain was found to have the largest negative coefficient, followed by the inability to wash and dress oneself.

Conclusion: Despite a generally lower education level than their European counterparts, urban Zimbabweans appear to value health states in a consistent manner, and the determination of a global method of establishing quality of life weights may be feasible and valid. However, as the relative weightings of the different domains, although correlated, differed from the standard set of weights recommended by the EuroQol Group, the locally determined coefficients should be used within the Zimbabwean context.

Background

The resources available to health care are obviously finite, and prioritisation or rationing of public health provision is on government agendas across the world [1]. Cost-utility analysis is one method of investigating the relationship between the costs and benefits of health care that allows for comparison of different interventions across different health states. The quality-adjusted life year (QALY) forms the basic unit of measure in such evaluation and is the most widely used method for measuring health outcomes [2]. The QALY is the arithmetic product of data on quantity of life and quality of life. Whilst the former is typically measured in life years, the latter is measured in terms of utility weights. There is little consensus as to how these weights should be developed, but the measure should have at least interval properties and should represent the preferences of society [3].

There are a plethora of instruments for describing health-related quality of life, most of which demonstrate acceptable psychometric properties [4]. Some of these measures, such as the SF-36 [5], are primarily profile measures that provide descriptors of health states. Others, such as the Health Utilities Index (HUI) [6] and the EQ-5D [7], are linked directly to utility estimates, derived from population studies using some method of eliciting population preferences, such as the standard gamble.

The EQ-5D describes health-related quality of life in terms of five dimensions: mobility (MO), self-care (SC), usual activities (UA) (work, study, housework, family or leisure), pain/discomfort (PD) and anxiety/depression (AD). Each dimension is subdivided into three levels indicating no problem, a moderate problem or an extreme problem [7]. Different health states can be described by a five-digit code number relating to the relevant level of each dimension, with the dimensions always listed in the order given above. Thus a health state of 11223 means:

Dimension MO: No problems in walking about (= 1)

Dimension SC: No problems with self care (= 1)

Dimension UA: Some problems with performing usual activities (= 2)

Dimension PD: Moderate pain or discomfort (= 2)

Dimension AD: Extremely anxious or depressed (= 3) [8]

The validity and reliability of the EQ-5D have been found acceptable in Europe among different populations and patient groups [9-11]. Despite the limited number of dimensions and levels, the instrument has been found to be sensitive to improvements in health-related quality of

life [12]. A test-retest study was undertaken in Zimbabwe to determine the reliability of the English language version of the EQ-5D. Forty-four randomly selected subjects who had a minimum of seven years of education and whose health status had remained static over the previous seven days completed the instrument twice, one week apart. In all domains except SC, approximately half of the respondents reported some or severe problems. The kappa statistics were 0.695 (fair to good agreement) for SC, 0.878 for MO, 0.884 for UA, 0.892 for PD and 0.893 for AD (all excellent agreement beyond chance [13]). A similar reliability study on the version of the EQ-5D in Shona, the local Zimbabwean language, reported that the kappa statistics between the two sets of scores were high and ranged from 0.78 to 1.00 for different domains [14]. Although the Shona version was not used in the current exercise, multiple translators examined the cross-cultural equivalence of meaning of the EQ-5D during the process of forward and back translation. One of the conclusions of the translators of the instrument was that "although it is likely that the Shona respondents will identify it as a foreign instrument, Shona is able to capture the EQ-5D concepts. The respondents will be able to recognise the concepts and respond appropriately..." [15]. It was concluded that, despite the different cultural understanding of determinants of ill health, the English version of the EQ-5D could be used with confidence in an educated urban Zimbabwe population.

Several methods of valuation of health states have been developed, including rating scales or visual analogue scales, magnitude estimation, standard gamble, time trade-off and person trade-off methods [3]. The standard gamble has been extensively used to develop utility weights, and is regarded by some as being the most theoretically sound method of determining utility weights [16]. However, it is conceptually difficult and requires an ability to discriminate between probabilities close to one [3]. Nord [17] proposes that time trade-off techniques are likely to be the most valid technique for establishing preference weights for life years both in the clinical situation and in program evaluation.

The Measurement and Valuation of Health Group (MVH), headed by Williams, used time trade-off exercises to elicit preferences from 3,235 respondents in the United Kingdom for a range of different EQ-5D descriptor states [8]. Regression analysis was used to develop a set of values for each individual component of the five dimensions that can be used to calculate the value of health states not observed directly [8]. The test-retest reliability of health state valuations collected with the EQ-5D questionnaire is reported to be stable over time [18].

It is unlikely that preferences for health states are universal, although some health states might be given similar valuations across cultures [19]. The greater the divergences of the local culture from the Western worldview, the less likely that health state valuation will be the same [20]. Barker and Green [21] state that health state values should be developed locally based on the judgments and priorities of local communities, in the service of these communities.

Much work has been done in developed countries on the valuation of health states [7,8,22-24], but there is a need to develop locally applicable measures of health that may be used to monitor the impact of interventions in developing countries. The WHOQOL is one of the few attempts to develop a genuinely international quality of life assessment [25], but so far it has no direct link with a utility index. The primary objective of this study was the generation of a set of weights for the different health states as described by the EQ-5D that would represent the values of urban high-density dwellers in Zimbabwe. Urban dwellers were chosen, as they were more likely to have the numeracy and literacy skills necessary to participate in the exercise. Where appropriate the results were compared with the results of the MVH study [8].

Methods

Subjects

In March 2000, 2,488 residents of randomly selected small residential plots of land in Glenview, a high-density suburb of Harare, were interviewed in their homes. The entrance criteria included completion of primary school education and a minimum age of 15 years. The oldest person in each household who met the criteria was interviewed.

Instruments

English descriptors of 38 different health states based on the different combinations of the five EQ-5D domains used in the original MVH study [8] were compiled on flash cards (See Appendices I and II). (Thirty-eight, rather than the original 42 health states were used, as unconscious and death were not valued, and two other states were excluded due to an administrative error). Each respondent was asked to complete a self-assessment using the EQ-5D and to value his or her own health condition on the EQ-5D visual analogue scale (VAS), which ranges from the "Worst possible health state" at 0 to the "Best possible health state" at 100. Respondents then each valued a different set of seven randomly selected health states (which included one or two very mild, mild, moderate and severe states). All respondents also valued an eighth state, the 33333 state. Valuation of states was undertaken using the time trade-off (TTO) approach. For states better than death subjects choose between a length of time in

perfect health (11111), x , which was equivalent to spending ten years in the target state. In this case, a larger x indicates a better health state. For states worse than death, the choice was between dying immediately and spending a length of time $(10 - x_1)$ in the target state followed by x_1 years in full health. A visual aid was used to clarify this choice. The greater the number of years in full health perceived to compensate for the time spent in the target state, the worse the health state. States worse than death were thus given negative values in analysis.

Procedure

The full procedure is described in Appendix II and III – see additional file 1. Nine interviewers, all of whom had higher degrees or diplomas of some kind, participated in a training workshop over three days, which included a pilot study. All residential plot numbers in Glenview were identified from a municipal map of the area, and a random sample of 2,500 were chosen. In the event of no one being present at the identified residential plot, the interviewers returned once more. The research assistants were instructed to conduct interviews in the evenings and weekends to the extent possible, but this was difficult because of the political unrest and weekend rallies at the time, and many interviews took place during work hours. Before each interview, the research assistant shuffled the 38 health states (excluding the 33333 health state) and randomly chose seven states for the respondent to value. Check visits were conducted by a supervisor to ensure that the randomly chosen residential plot had been visited.

Data analysis

Statistical analyses were undertaken using GenStat version 4.2 [26] and SPSS for Windows, Release 10 [13]. Descriptive statistics and χ^2 and 95% confidence intervals (CI's) were used to delineate the demographic characteristics of the subjects and to compare them with population demographics of high-density dwellers in Harare derived from census findings. The health characteristics of the respondents in terms of the five EQ-5D domains were described. The sample of respondents was randomly divided into three, and analysis was performed on two-thirds of the sample, the *internal sample*. The results were then used to estimate the values of the remaining one-third, the *external sample*. The dependent variable was the TTO score divided by ten. A residual maximum likelihood (REML) linear mixed model was fitted. Residual maximum likelihood estimation is a method of estimating variance components in the context of unbalanced incomplete block designs. It takes account of the loss of degrees of freedom in estimating the mean and produces unbiased estimating equations for the variance parameters [27]. Interviewer effect and subject nested within interviewer were fitted as the random effects. The three levels of the five domains were entered as the fixed effects and a weighted least

Table 1: Demographic details of study sample compared to population figures.

	Sample Frequency	Sample proportion % (95% CI)	Population proportion % (95% CI)
<i>Gender*</i>			
Males	904	38.3 (36.3–40.3)	52.8 (52.4–53.1)
Females	1,457	61.7 (59.7–63.7)	47.3 (46.9–47.6)
Total	2,361		
Missing	23		
<i>Age category*</i>			
15–24 years	1,097	46.2 (44.2–48.2)	38.9 (38.6–39.3)
25–34 years	781	32.9 (31.0–34.8)	30.3 (30.0–30.7)
35–44 years	268	11.3 (11.0–12.6)	15.5 (15.2–15.8)
45–54 years	162	6.8 (5.8–7.9)	8.0 (7.7–8.2)
55–64 years	49	2.1 (1.5–2.7)	4.8 (4.6–5.0)
65+ years	17	0.7 (0.4–1.1)	2.4 (2.3–2.6)
Total	2,374		
Missing	10		
<i>Education Levels†</i>			
10 or fewer years	458	19.2 (17.7–20.9)	48.9 (48.8–49.1)
11–13 years	1,600	67.1 (65.2–69.0)	48.4 (48.3–48.5)
Post school	18	0.8 (0.49–1.2)	2.6 (2.6–2.7)
Total	2,076		
Missing	308		
<i>Occupation†</i>			
Scholar/studying further	361	16.8 (15.2–18.4)	11.6 (11.6–11.7)
Employed	718	33.3 (31.3–35.4)	54.2 (54.1–54.3)
Unemployed	1,075	49.9 (47.8–52.0)	34.2 (34.1–34.3)
Total	2,154		
Missing	230		
<i>Accommodation status</i>			
Owner	740	34.8 (36.9–34.8)	30
Renting	169	7.9 (6.8–9.2)	7
Lodging	961	45.2 (43.1–47.3)	48
Other	256	12.0 (10.7–13.5)	15
Total	2,126		
Missing	258		

*Compared to Highfield population, 1992 [28]. †Compared to Harare Province population, 1992 [28].

squares model was fitted. The fixed effects were entered in a forward and backward sequence and their effects assessed using Wald statistics. An ANOVA full factorial model with Type III least squares ($N = 15,671$) was used to establish the source of variance. The dependent variable was the TTO score, and random factors entered included research assistant, health state, occupational category and gender. Interactions between the main effects were also investigated.

Results

Subjects

Forty-eight respondents refused to answer the questionnaire. The data from 56 respondents were incomplete,

and the replies from 201 respondents demonstrated inconsistency but were included in the analysis (see discussion below). Inconsistent data included responses in which all states were given the same value, fewer than three states were valued, or there were more than three logical inconsistencies (e.g. if a 11112 state were valued as being more severe than a 11113 state). Ultimately, the responses of 2,384 subjects were analysed. The demographic details of the respondents were compared with the results for Highfield in the 1992 census Harare Profile [28] or, if not available, with results for Harare Province (Table 1). Males were underrepresented at 38.3%, a proportion which fell outside the 95% CI for the population (52.4 – 53.1%). There were more young adult

respondents (46.2%) than in the general population (38.9%, CI = 38.6 – 39.3), but the decline in numbers with increasing age was similar to the population of Harare.

Tenure status, which is an indicator of socio-economic status, indicated that, as in the population, the majority were lodgers (45.2%) or owners (34.8%). The sample was better educated than the population with 67.1% having attained 11 to 13 years of schooling compared to the population estimate of 48.3 – 48.5%. The percentage employed in the sample (33.3%) was smaller than the reference population (CI = 54.1 – 54.3). Approximately 72% of the sample interviewed by the female research assistants were female compared to 54.5% of the male sample ($\chi^2 = 552.0, p < 0.001$), and 57.3% were unemployed compared to the 38.6% of the male interviewers' samples ($\chi^2 = 1060.6, p < 0.001$).

Health status

Disability was reported by 104 (4.4%) of the sample, and 290 (12.2%) reported having a serious illness in the prior three months (total number with disability and/or illness was 14.7%). Approximately half (47.8%) reported no problem on any of the five dimensions. Information

regarding self-described health status appears in Table 2. With regard to the self-reported scores on the EQ-5D dimensions, nearly one-third of the respondents reported either some or severe problems in the dimensions of pain/discomfort and anxiety/depression. The mean score on the VAS was 79.8 (CI = 79.1 – 80.5).

Valuations

There were 19,020 values of 38 EQ-5D health states analysed, 12,663 in the internal sample and 6,357 in the external sample. Fifty two percent of the proportion of variance was due to the domain scores, 7% due to interviewer and health state interaction, 6% due to interviewer effects and 35% due to error. (Table 3).

The Wald statistic was highly significant for all main fixed effects, both when the effects were fitted in a forward and in a backward sequence ($p < 0.001$) (Table 4). Previous models included a variable (N3) which reflected that at least one domain was valued at the severe level. However, this led to only a small increase in R^2 and resulted in a model in which moderate problems in several domains were counter-intuitively valued as being worse than extreme problems. This model was subsequently discarded.

Table 2: Self-reported health status of subjects in urban Zimbabwe (N = 2,183) compared to the United Kingdom [9]. Frequency of levels on each dimension of EQ-5D are reported as percentages.

Dimension	Mobility		Self care		Usual Activities		Pain/discomfort		Anxiety/depression	
	Zim	UK	Zim	UK	Zim	UK	Zim	UK	Zim	UK
Level										
1.00	90.1	82	96.5	96	89.0	84	69.5	67	69.3	79
2.00	9.7	18	3.4	4	10.5	14	26.3	29	23.9	19
3.00	0.1	0.1	0.1	0.1	0.6	2	4.2	4	6.8	2

Table 3: Results of ANOVA full factorial model with Type III least squares (N = 12,663). The dependent variable was the TTO score, and random factors entered included research assistant, health state, occupational category and gender.

	Degrees of freedom	Mean Square	Degrees of freedom	Mean Square	F	p	% Variance
Error			Error				34.97
Health state	39	3.50	1,225	0.13	27.51	0.00	51.40
RA*health state	289	0.75	15,102	0.08	9.58	0.00	7.10
Research assistant (RA)	8	1.05	942	0.12	8.89	0.00	5.56
Occupation	3	0.59	74	0.10	5.77	0.00	0.72
RA*occupation	24	0.20	15,102	0.08	2.63	0.00	0.21
Gender*occupation	5	0.11	15,102	0.08	1.47	0.19	0.03
Occupation*health state	111	0.08	15,102	0.08	1.03	0.41	0.01
Gender	2	0.01	16	0.08	0.08	0.93	0.00
RA*gender	15	0.06	15,102	0.08	0.81	0.67	0.00
Gender*health state	74	0.05	15,102	0.08	0.59	1.00	0.00

Table 4: Wald tests for fixed effects (N = 12,663)

Fixed term	Wald statistic	d.f.	Wald/df.	χ^2 prob.
<i>Sequentially adding terms to fixed model</i>				
Mobility	9772.4	2	4886.2	<0.001
Self care	2070.1	2	1035.1	<0.001
Usual activities	426.1	2	213.0	<0.001
Pain/discomfort	2338.0	2	1169.0	<0.001
Anxiety/depression	544.7	2	272.3	<0.001
<i>Dropping individual terms from full fixed model</i>				
Mobility	544.7	2	272.4	<0.001
Self care	674.0	2	337.0	<0.001
Usual activities	253.2	2	126.6	<0.001
Pain/discomfort	1789.0	2	894.5	<0.001
Anxiety/depression	544.7	2	272.3	<0.001

The effects were largest for a level three on the dimensions of PD and SC, followed by level three MO and AD (Table 5).

There was some evidence of significant interaction effects. However, because not all combinations of health states are plausible and were not valued, numerical problems were experienced in the fitting of the model, and the estimates appeared unreliable.

Coefficients from the model were used to generate predicted values for each of the states included in the study. For example, for the health state 22331 (i.e., some problems in walking about, some problems with self care, unable to perform usual activities, severe pain and no anxiety/depression), the predicted value would equal $0.90 - 0.056 - 0.092 - 0.135 - 0.302 - 0.0 = 0.315$. The actual and predicted value and residuals for each health state for the internal and external samples and the observed results of the United Kingdom MVH study are reported in Table 6. The Pearson's correlation between mean values of health states was 0.914 ($p < 0.001$). The 33333 state was the only state that was valued as being worse than death in the current study (mean value = -0.24). There were three health states in the external sample for which the mean difference between the observed and predicted values was more than 0.1. In the internal sample, the mean absolute difference for all health states was 0.045.

Figure 1 depicts the predicted UK scores compared to the Zimbabwe internal sample scores for the 39 health states (the UK sample did not include the 11111 health state).

Whereas there is initial convergence in scores between the Zimbabwe and MVH sample at high health levels, values diverge as the health states become more severe and

domains at level three are included. Spearman's rank correlation between the values for the different states was 0.95 ($p < 0.001$).

Discussion

To the knowledge of the authors, this is the first paper to present the preferences for health states by urban Zimbabweans. The self-reported health-related quality of life of the Zimbabwe subjects was similar to that of UK counterparts. Kind et al. [9] found that 30% of a large UK sample reported some or severe pain/discomfort. However, the number reporting some or severe anxiety/depression was smaller in the UK sample (21%) than in the present study. The two samples were similar in finding very few people reporting problems in the area of self-care, or extreme problems with mobility or usual activities. The mean score on the Visual Analogue Scale (VAS) in the Zimbabwe sample was 79.8 (CI = 79.1 – 80.5), which was similar to the British sample (mean 82.5).

As the questionnaire was administered in English and numeracy was required, the methodology precluded gathering valuations from a truly representative sample. The educational inclusion criteria and the limitations imposed on the times for data gathering by female interviewers resulted in a sample in which females, younger people and those with a higher level of literacy were over-represented. In addition, the interviewer effect was considerable. As the interviewer and subject were entered in the computation as random effects, the REML linear mixed model allowed for the demographic deviations from census findings, the non-independence of the measures and the interviewer effect.

The interviewer effect appeared in spite of training sessions, piloting and standardisation of the format of the

Table 5: Main effects in residual maximum likelihood linear mixed model based on 12,663 observations.

	Level	Effect	Std. Err.
Constant		0.900	
Mobility	2	-0.056	0.0080
	3	-0.204	0.0089
Self care	2	-0.092	0.0074
	3	-0.231	0.0089
Usual activities	2	-0.043	0.0079
	3	-0.135	0.0088
Pain/discomfort	2	-0.067	0.0070
	3	-0.302	0.0076
Anxiety/depression	2	-0.046	0.0073
	3	-0.173	0.0076

interview. It is possible that the approach and amount of interpretation given by each interviewer differed. The effect of the gender of the interviewers was evident, and female interviewers apparently did not conduct interviews during the evenings or weekends to the same extent as their male counterparts. This imbalance might have compounded the interviewer effect, which suggests that the gender of interviewers should receive careful attention in community surveys, particularly in socially unstable conditions.

However, a credible model was ultimately developed in this study. The mean absolute difference between the actual and estimated means for the external sample (0.045) is comparable to that of the UK study (0.039) and a similar study in Japan (0.01) [24], although in each case different models were used. The inclusion of inconsistent responses is controversial, with some researchers excluding these data from analysis [29]. These responses were included in this analysis on the assumptions that inconsistencies do not necessarily indicate a lack of understanding of the task, that all those who participate have the right to have their data included, and that human beings are not always rational in their judgments regarding health states.

Significant interaction effects were found but, as noted above, appeared unreliable due to an incomplete data set. The inclusion of interaction effects resulted in a model that was difficult to interpret, which would likely limit the use of the model in practice. Similarly, the inclusion of the N3 term, which indicates severe problems on at least one domain, resulted in a more complicated and less intuitive model. It was therefore decided to adopt the simple main effects model.

The UK and Zimbabwe samples produced similar descriptions of their own health states and similar rank orderings of the hypothetical health states. (As a different model was used, the coefficients of the valuation function could not be compared directly between the UK and Zimbabwe samples.) The mean self reported VAS was 3% lower in the Zimbabwe sample compared to the UK sample. The Pearson's correlation for the predicted health state values (0.95) was high. Although previous studies based on the EQ-5D have reported similarities in valuations, with low sensitivity for socio-demographic variables across European countries [22], the results of this study were unexpected. A previous study on the rank ordering of health states had found no correlation between the international and locally determined Zimbabwean ranking [20]. It would appear that a deconstructed approach to valuation in which impairments or activity limitations (e.g. pain or problems in moving around) are valued [30,31] rather than disease conditions (e.g. rheumatoid arthritis) is more likely to tap into commonly understood constructs and yield universal preferences.

However, there were important differences between the samples that should be noted. Respondents in the UK study valued 16 health states as being worse than death, whereas in the Zimbabwe sample only the 33333 state was awarded a negative value. The inclusion of 16 negative values in the UK model resulted in generally lower values being assigned to health states in which an "extreme" problem was included. Consequently the predictions from the UK model for about two-thirds of the health states are lower than those from the Zimbabwe model. The reluctance to value states as worse than death in the Zimbabwe sample might reflect a fundamentally different attitude towards the sacrifice of years of life. There is, for example, no national debate on either euthanasia or abortion in Zimbabwe, and both are illegal and likely to remain so for the near future. The general state of health of the population might also contribute. The life expectancy is now dropping drastically because of the HIV/AIDS pandemic. The expected number of equivalent healthy years (Disability Adjusted Life Expectancy, DALE) is now estimated to be 32.9 (cf. UK 71.7), and Zimbabwe ranks 184 out of 191 nations [32]. (To calculate DALE, the years of ill-health are weighted according to severity and subtracted from the expected overall life expectancy to give the equivalent years of healthy life [32]). There may be a greater reluctance to sacrifice life years in a society in which each individual is likely to have had direct contact with death or illness. This conclusion is supported by the results of a Spanish study of preferences of 103 patients who were severely ill. The patients tended to rate the worst health states higher than proxies and rated no states as worse than death. The authors of that study concluded that within the EQ-5D descriptive system, there are no

Table 6: Mean observed and predicted time trade-off scores for 38 health states, for internal samples (N = 12,663 values) and external samples (N = 6,357) in Zimbabwe compared to observed values in the United Kingdom [8].

Health State	Internal sample		External sample		UK Sample
	Observed mean	Observed mean	Predicted mean*	Absolute difference	Observed
11111	0.94	0.93	0.90	0.03	
11112	0.87	0.90	0.85	0.05	0.834
11113	0.73	0.75	0.72	0.02	0.392
11121	0.85	0.83	0.83	0.00	0.850
11122	0.70	0.70	0.78	0.08	0.722
11131	0.56	0.59	0.60	0.00	0.200
11133	0.52	0.45	0.42	0.03	-0.049
11211	0.84	0.84	0.85	0.02	0.869
11312	0.67	0.69	0.72	0.03	0.552
12111	0.81	0.82	0.81	0.02	0.834
12121	0.69	0.67	0.74	0.06	0.742
12211	0.65	0.64	0.76	0.12	0.767
12222	0.58	0.59	0.65	0.06	0.552
12223	0.50	0.48	0.52	0.04	0.216
13212	0.63	0.58	0.58	0.00	0.389
13311	0.55	0.54	0.53	0.01	0.346
13332	0.28	0.28	0.18	0.10	-0.228
21222	0.62	0.60	0.69	0.09	0.553
21232	0.41	0.44	0.45	0.01	0.064
21312	0.61	0.62	0.66	0.04	0.536
22112	0.69	0.68	0.70	0.02	0.662
22121	0.65	0.69	0.68	0.01	0.645
22122	0.62	0.60	0.64	0.04	0.540
22222	0.56	0.59	0.59	0.01	0.500
22233	0.25	0.30	0.23	0.07	-0.142
22323	0.43	0.42	0.37	0.05	0.042
22331	0.33	0.33	0.32	0.02	-0.011
23232	0.28	0.27	0.22	0.05	-0.084
23321	0.37	0.39	0.41	0.02	0.147
32223	0.30	0.34	0.32	0.02	-0.174
32232	0.21	0.21	0.21	0.00	-0.223
32313	0.40	0.43	0.29	0.14	-0.152
32331	0.27	0.27	0.16	0.11	-0.276
33212	0.37	0.37	0.37	0.00	-0.022
33232	0.23	0.19	0.07	0.12	-0.332
33321	0.29	0.34	0.26	0.08	-0.120
33323	0.18	0.16	0.09	0.07	-0.386
33333	-0.24	-0.24	-0.15	0.09	-0.543
MAE†	0.049			0.045	0.039

*Based on estimates †MAE=Mean absolute difference between observed and predicted means

health states worse than death for seriously ill patients [33].

For Zimbabweans, the inability to wash and dress oneself is a major contributor to poor quality of life, and SC level 3 was ranked second. In contrast, SC level 3 was ranked fourth in the UK study. This difference may possibly be due to the importance that Zimbabweans attach to self-presentation. It is regarded as insulting to ask whether people are able to wash or dress themselves, if in any way it is implied that they have not done so [15,34]. In a

poorer country, self-presentation may also be regarded as indicative of socio-economic standing and hence valued more highly. The important differences between the results of the two studies illustrate the dangers of applying measures developed in one culture without adequate testing of items for cultural meaning and appropriateness.

Severe AD was ranked similarly in the UK (Rank 3) and Zimbabwe (Rank 4) samples. Of all the EQ-5D concepts, the idea of depression and anxiety is most difficult to capture in the sensibility of the Shona-speaking Zimba-




Figure 1
Observed TTO scores (divided by 10) of the full Zimbabwe sample (N = 19,020 values from 2,183 respondents), compared to observed scores from the Measurement and Valuation of Health study in the United Kingdom [8].

bwean. There is no specific word for depression; it is usually implied from symptoms rather than self-report. Anxiety and depression are not regarded simply as health states in Shona custom. They are understood as occasional psychological (social/alienation) or spiritual (religious) states. In addition, severe anxiety is seen to border on a psychiatric state known as "mhopu" [15]. It is therefore not surprising that extreme anxiety or depression should be regarded as being very serious.

The choice of the EQ-5D as the instrument to define the different domains of health-related quality of life needs justification. The measure is limited in that there are only five domains with three possible levels on each domain. The content validity may be questioned, as it may be that important areas that contribute to quality of life, such as cognitive function or energy, are excluded. However, even with this relatively crude measure, 243 hypothetical health states can be described. Researchers have to be cautious about transposing any measure across very different cultural contexts. The current study required a robust, relatively simple measure and, despite the shortcomings of the instrument, the EQ-5D appeared to be reliable and relatively insensitive to cultural context.

Conclusions

This study attempted to elicit cardinal values of health states from urban Zimbabweans. The limitation imposed by the educational criteria resulted in a sample that was

more educated than the general population of high-density areas, and the results should be generalized with care to other urban populations in the country. Despite this limitation, the values derived from the study are more likely to represent the values of urban Zimbabwe than values derived from valuation exercises performed in other countries. The parameter estimates for each level of the five domains generated by the TTO exercise are credible and are comparable to those generated by other studies. The ranking of observed preferences for health states by Zimbabweans and UK residents are remarkably similar, and if consensus could be reached on the valuation of states worse than death, it is possible that QALY weights based on EQ-5D descriptors might be developed which are valid globally.

However, the observed cardinal values for health states are much lower overall in the UK sample. It is therefore recommended that the parameter estimates developed in this study be used both to describe health-related quality of life and as an outcome measure of health interventions in the Zimbabwe urban population.

Authors' contributions

JJ carried out the data collection and analysis and wrote the paper. KH assisted in data collection and analysis of the data. WdW, PDC and PK assisted in conceptualising the study and developing the methodology. All authors have read and approved the manuscript.

Competing interests

None declared.

Additional material

Additional File 1

APPENDIX I: Example of cards used to describe the health states.
APPENDIX II: Detailed procedure. Instructions to interviewers. APPENDIX III: Information Sheet
Click here for file
[<http://www.biomedcentral.com/content/supplementary/1478-7954-1-11-S1.doc>]

Acknowledgements

Funding for this study was made available by the Zimbabwe Burden of Disease Steering Committee, which in turn is supported by DIFID and DANIDA. Thanks to Allan Williams, Aki Tsuchiya and Jan Busschbach for comments and insightful reviewing of the paper and to June Juritz and Jaqui Somerville for assistance with statistical analysis.

References

1. Bowling A: **Health care rationing: the public's debate.** *BMJ* 1996, **312**:670-674.
2. Bala M, Wood L, Zarkin G, Norton E, Gafni A, O'Brien B: **Valuing outcomes in health care: A comparison of willingness to pay**

- and quality adjusted life years. *Journal of Clinical Epidemiology* 1998, **51**:667-676.
3. Murray CJL: **Rethinking DALYs**. *The Global Burden of Disease Volume 1*. 1stth edition. Edited by: Murray CJL and Lopez AD. Cambridge, Harvard School of Public Health; 1996:1-89.
 4. Essink-Bot M, Krabbe P, Bonsel GJ, Aaronson NK: **An empirical comparison of four generic health status measures**. *Medical Care* 1997, **35**:522-537.
 5. Wagner AK, Gandek B, Aaronson N, Acquadro C, Alonso J, Apolone G, Bullinger M, Bjorner J, Fukuhara S, Kaasa S, Leplege A, Sullivan M, Wood-Dauphinee S, Ware JE: **Cross-cultural comparisons of the content of SF-36 translations across 10 countries: Results from the IQOLA project**. *Journal of Clinical Epidemiology* 1998, **51**:925-932.
 6. Holmes A: **A QALY based societal health statistic for Canada, 1985**. *Social Science and Medicine* 1995, **41**:1417-1427.
 7. Brooks R, Rabin R, De Charro F: **The measurement and valuation of health status using the EQ-5D: A European perspective**. Dordrecht, Kluwer; 2003.
 8. Dolan P: **Modeling valuations for EuroQol health states**. *Medical Care* 1997, **35**:1095-1108.
 9. Kind P, Dolan P, Gudex C, Williams A: **Variations in population health status: results from a United Kingdom national questionnaire survey**. *British Medical Journal* 1998, **316**:736-741.
 10. Brooks R, EuroQol Group: **EuroQol: the current state of play**. *Health Policy* 1996, **37**:53-72.
 11. Elvik R: **The validity of using health state indexes in measuring the consequences of traffic injury for public health**. *Social Science and Medicine* 1995, **40**:1385-1398.
 12. Macdonagh RP, Cliff AM, Speakman MJ, O'Boyle PJ, Ewings P, Gudex C: **The use of generic measures of health-related quality of life in the assessment of outcome from transurethral resection of the prostate**. *British Journal of Urology* 1997, **79**:401-408.
 13. SPSS Inc: **Statistical Package for the Social Sciences**. Release 10.1.3th edition. Chicago, SPSS Inc; 2001.
 14. Jelsma J, Mhundwa K, Weerd W De, Cock P De, Chimera J, Chivaura V: **The reliability of the Shona version of the EQ-5D**. *Central African Journal of Medicine* 2001, **47**:8-11.
 15. Jelsma J, Chivaura V, De Cock P, De Weerd W: **A bridge between cultures: A report on the process of translating the EQ-5D instrument into Shona**. *South African Journal of Physiotherapy* 2000, **56**:3-9.
 16. Torrance George W, Feeny David H: **Utilities and quality adjusted life years**. *International journal of technology assessment of health care* 1989, **5**:559-575.
 17. Nord E: **Methods for quality adjustment of life years**. *Soc Sci Med* 1992, **34**:559-569.
 18. van Agt HME, Essink-Bot M-L, Krabbe PFM, Bonsel GJ: **Test-retest reliability of health state valuations collected with the Euro-Qol questionnaire**. *Social Science and Medicine* 1994, **39**:1537-1544.
 19. Ustun TB, Saxena S, Rehm J, Bickenbach J: **Are disability weights universal?** *Lancet* 1999, **354**:1306.
 20. Jelsma J, Mhundwa K, Chivaura V, De Weerd W, De Cock P: **The global burden of disease disability weights**. *Lancet* 2000, **355**:2079.
 21. Barker C, Green A: **Opening the debate on DALYS**. *Health Policy and Planning* 1996, **11**:179-183.
 22. Krabbe PFM, Essink -Bot M-L, Bonsel GJ: **On the equivalence of collectively and individually collected responses: Standard gamble and Time-tradeoff judgements of health states**. *Medical Decision Making* 1996, **16**:120-132.
 23. Nord E, Richardson J, Macarounas-Kirchmann K: **Social evaluation of health care versus personal evaluation of health states**. *International Journal of Technology Assessment in Health Care* 1993, **9**:463-478.
 24. Tsuchiya A, Ikeda S, Ikegami N, Nishimura S, Sakai I, Fukuda T, Hamashima C, Hisashige A, Tamura M: **Estimating an EQ-5D population value set: the case of Japan**. *Health Econ* 2002, **11**:341-353.
 25. WHOQoL Group: **The World Health Organisation Quality of Life Assessment:(WHOQoL): Position paper from the World Health Organisation**. *Social Science and Medicine* 1995, **41**:1403-1409.
 26. Genstat: **Genstat Version 4.2**. Oxford, Numerical Algorithm Group; 2000.
 27. Smyth GK, A P Verbyla: **A conditional approach to residual maximum likelihood estimation in generalized linear models**. *Journal of the Royal Statistical Society B* 1996, **58**:565-572.
 28. Central Statistical Office: **Census 1992: Provincial Profile Harare**. CSO, Harare, Zimbabwe; 1994.
 29. Devlin NJ, Hansen P, Kind P, Williams A: **The health state preferences and logical inconsistencies of New Zealanders: A tale of two tariffs**. York, York Centre for Health Economics UK and University of Otago, New Zealand; 2000.
 30. World Health Organisation: **The International Classification of Functioning and Disability - Beta-2 Draft**. Geneva, WHO; 1999.
 31. World Health Organisation: **The International Classification of Functioning and Disability - Beta-2 Draft**. Geneva, WHO; 1999.
 32. World Health Organization: **The World Health Report 2000 Health Systems: Improving Performance**. Geneva; 2000.
 33. Badia X, Diaz-Prieto A, Rue, Patrick DL: **Measuring health and health state preferences among critically ill patients**. *Intensive Care Medicine* 1996, **22**:1379-1384.
 34. Jelsma J, Chivaura V: **Report on the process of translating the EQ-5D into Shona**. Harare, EuroQol Group, Rotterdam; 2000.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

