



This is a repository copy of *Struggling with Adversities of Life: The Role of Forgiveness in Patients Suffering from Fibromyalgia*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/104188/>

Version: Accepted Version

Article:

Offenbaecher, M., Dezutter, J., Kohls, N. et al. (9 more authors) (2017) Struggling with Adversities of Life: The Role of Forgiveness in Patients Suffering from Fibromyalgia. *Clinical Journal of Pain*, 33 (6). pp. 528-534. ISSN 0749-8047

<https://doi.org/10.1097/AJP.0000000000000432>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Please cite as:

Offenbaecher, M., Dezutter, J., Kohls, N., Sigl, C., Vallejo, M. A., Rivera, J., Bauerdorf, F., Schelling, J., Vincent, A., Hirsch, J., Sirois, F. M., & Toussaint, L. L. (accepted). Struggling with adversities of life: The role of forgiveness in patients suffering from fibromyalgia. *Clinical Journal of Pain*.

Title: Struggling with Adversities of Life: The Role of Forgiveness in Patients Suffering from Fibromyalgia

Authors: Offenbaecher M¹, Dezutter J², Kohls N³, Sigl C¹, Vallejo MA⁴, Rivera J⁵, Bauerdorf F¹, Schelling J¹, Vincent A⁶, Hirsch, JK⁷, Sirois, FM⁸, Webb, JR⁷, Toussaint LL⁹

Affiliations: ¹Institute of General Medicine, University Clinic, Munich, Germany
²Faculty of Psychology and Educational Sciences, KU Leuven University of Leuven, Belgium
³Division of Integrative Health Promotion, University of Applied Science and Arts, Coburg, Germany
⁴Psychology Faculty, UNED, Madrid, Spain
⁵Rheumatology Unit, Rehabilitation Provincial Institute, “Gregorio Marañón” General Hospital, Madrid, Spain
⁶Division of General Internal Medicine, Mayo Clinic, Rochester, MN, 55905, USA
⁷Department of Psychology, East Tennessee State University
⁸Department of Psychology, University of Sheffield, United Kingdom
⁹Department of Psychology, Luther College, Decorah, Iowa, USA

Corresponding author:

Loren Toussaint, Ph.D.
Department of Psychology
Luther College
700 College Dr.
Decorah, IA 52101
Phone: 563-387-1647
E-Mail: touslo01@luther.edu

Abstract

Objectives: We compared the magnitude and direction of associations between forgiveness and pain, mental and physical health, quality of life, and anger in a sample of fibromyalgia (FMS) patients and healthy controls. In addition, we compared FMS and controls on mean levels of these variables.

Methods: 173 FMS patients and 81 controls completed this study. Patients and controls were residents of Germany recruited with the support of the German Fibromyalgia Patient Association and several self-help groups. FMS patients and controls were about 53 years of age, mostly married (70%), Christians (81%), with levels of education ranging from 9 years to 13+. All participants completed assessments of forgiveness, pain, health, quality of life, and anger.

Results: Analyses revealed that FMS patients reported higher pain and anger and poorer health and quality of life. FMS patients also reported lower levels of both forgiveness of self and others. Size and direction of associations of forgiveness with pain, health, quality of life, and anger in were not significantly different between healthy individuals and patients with FMS.

Discussion: Forgiveness of self and others is beneficially associated with pain, health, quality of life, and anger in FMS patients at levels that are of similar size and direction as in healthy controls. However, FMS patients manifest lower levels of forgiveness of self and others. Therapeutic promotion of forgiveness as a psychosocial coping strategy may help patients with FMS to better manage psychological and physical symptoms, thereby enhancing well-being.

Key Words

Fibromyalgia, anger, forgiveness, chronic pain, coping, stress

Introduction

Fibromyalgia syndrome (FMS) is a common and chronic syndrome of unknown etiology characterized by diffuse pain, increased tenderness to palpation and additional symptoms (e.g. disturbed sleep, fatigue and psychological distress) [1] resulting in major effects on function, health, work status and quality of life [2-4] as well as high rates of health care utilization [5].

It has been proposed that FMS is a complex, multidimensional disorder with both biomedical and psychosocial-behavioral dimensions [6]. Past research has shown that these two dimensions are relatively independent in different chronic pain syndromes, suggesting the importance of considering heterogeneity of patients in each of the dimensions when planning treatment [7]. Turk et al [8] have suggested, for example, that psychological characteristics may be important factors in the identification of FMS patients who are more or less likely to benefit from an interdisciplinary intervention and in the customization of treatment plans for individual patients.

There is growing evidence that FMS is a stress-related syndrome. Studies in FMS have generally demonstrated altered functioning of the hypothalamic-pituitary-adrenal (HPA) axis and autonomic nervous system (ANS) [9, 10]. However, studies have revealed both hypo- and hyper-activity of the hypothalamic-pituitary-adrenal axis and the autonomic nervous system in FMS, and often the degree of abnormality is small or occurs in a subgroup of patients with substantial overlap between patients and controls. Some researchers have proposed that FMS can in fact be viewed as a failed attempt of our complex adaptive system to accommodate to a perceived hostile

external environment. FMS is, in this view, a disease in which distress is transformed into pain through sympathetic nervous system hyperactivity [10].

These results are corroborated by FMS patients' reports of symptom exacerbation when they are exposed to physical (e.g. acute infections, physical overload) or psychosocial stressors (e.g., bullying at work place, conflicts with partner). This notion has recently been endorsed by Fischer et al [11] who showed that higher stress at a given measurement time point (i.e. cortisol and alpha-amylase in saliva samples) was associated with higher reported pain levels at a subsequent time point. As well, it is well-established that negative emotions such as anger or hostility have detrimental effects on health [12].

Psychotherapeutic strategies for pain, including Acceptance and Commitment Therapy and various other cognitive behavioral therapies, aim to reduce negative emotions and stress resulting from pain catastrophizing, fear avoidance behaviors, and other maladaptive thoughts that contribute to disabling aspects of chronic pain [13-15]. Psychotherapy also aims to promote active coping strategies like pain acceptance, physical activity, and structured relaxation [16, 17]. Forgiveness might be a component of the larger part of the positive, health promoting thoughts, feelings, and behaviors encouraged by these types of therapies and could be considered an important target of therapy [18, 19]. Indeed, the process of forgiveness is consistent with and can be readily incorporated into acceptance-based treatment modalities, such as Acceptance and Commitment Therapy, Dialectical Behavior Therapy, and Mindfulness-Based Cognitive Therapy [20]. Moreover, some specific methods for facilitating forgiveness as a direct focus of therapy, including stand-alone forgiveness interventions have already been developed and have sound evidence of

effectiveness [21].

Forgiveness is a multidimensional construct [22] that can be directed toward oneself or others and is considered both a state and a trait. Trait forgiveness is often of greater interest in studies of health and well-being because it is thought to have its influence across time and situations and consequently has greater impact than a momentary state experience [23]. In healthy populations, forgiveness has been linked to a variety of indicators of psychological well-being including less anger, lower depression, and lower anxiety [24]. Forgiveness has been associated with decreased sympathetic arousal, increased parasympathetic tone, and improved physical health and longevity [25-27]. At present, only two known studies have examined forgiveness in pain patients. The first study examined a diverse sample of pain patients with various causes, locations, and durations of pain [28]. Results showed that pain patients' average levels of forgiveness of self and others did not differ from a healthy sample, and forgiveness was related to less time in pain, less pain intensity and interference, and better mental health. The second study examined forgiveness in a sample of chronic low back pain patients and showed that forgiveness was related to less pain, anger, and psychological distress [29]. The salutogenic associations with pain, health, and anger are likely due to the coping benefits of forgiveness [30] that impact the stress-disease connection [31]. Dealing with adversities of life through forgiveness may be especially useful for individuals with fibromyalgia who are interpersonally distressed [6, 32] as forgiveness applies well to inter-relational challenges in multiple life-domains that patients commonly struggle with [33].

Forgiveness and pain, mental and physical health, quality of life, and anger have not been previously examined simultaneously in fibromyalgia patients compared to

controls, and this was the goal of the current study. Based on the above review of the literature, we hypothesized that, compared to controls, FMS patients would show higher pain, anxiety, depression, and anger, lower levels of health and quality of life, and equal levels of forgiveness of self and others. Because FMS is a stressful condition and forgiveness has been proposed to be useful for managing the stresses of FMS [32, 34], it was hypothesized that the beneficial associations of forgiveness with pain, health, quality of life, and anger outcomes that have been demonstrated in other pain patients and healthy populations would extend to patients with FMS. Hence, beneficial associations of forgiveness with pain, health, quality of life, and anger should not be different for controls versus FMS patients.

Materials and Methods

Patients and controls

In total 320 questionnaire sets were sent out to several fibromyalgia self-help groups which were recruited with the support of the German Fibromyalgia Patient Association. 173 patients (response rate = 54%) filled out the set. The group leaders were contacted by the first author (MO) and were then sent the patient materials including the questionnaires, an information letter regarding the study objectives, and a consent form. Group leaders were asked to distribute the materials and to collect and return them when completed. Healthy controls (N = 81) were a German convenience sample of volunteers without fibromyalgia that completed the questionnaire set. Controls were recruited using similar snowball sampling methods to reach individuals of a variety of ages. Initial surveys were delivered to students who then delivered surveys to people they knew.

Measures

Forgiveness of Self and Others. The forgiveness of self and others scales, originally developed by Mauger et al [35], consist of 30 items. They are designed to assess general tendencies toward a forgiving disposition with regard to both oneself (e.g., 'I feel guilty because I don't do what I should for my loved ones') and others (e.g., 'It is hard for me to forgive those who hurt me'). Forgiveness of self is measured with 15 items, and forgiveness of others is measured with 15 items. Each item is responded to on a 1 (strongly disagree) to 5 (strongly agree) response scale. Both scale scores range from 15 to 60 and possess acceptable reliability and good construct validity [35]. Higher scores indicate higher forgiveness. Because there is no German version of the forgiveness scales, a back-forward translation from English according to scientific guidelines was performed [36]. Mauger's original psychometric analyses were replicated to evaluate the translated German instrument. Mauger reported internal consistency of English forgiveness of self and others scales to be .82 and .79, respectively, and the present study estimates from the German scales were .88 and .81, respectively. Mauger reported correlations of forgiveness of self and others with age, sex, and education that ranged between -.06 and -.21, and present study estimates similarly ranged between .09 and -.21. Translation to German appears to have not affected the integrity of the scales and psychometric estimates are similar or better than the original English version.

Pain. Duration of symptoms in years, and current and past (last three month) pain levels, were measured on a numerical rating scale from 0 (no pain) to 10 (extreme pain). Pain severity (range 0-120) and distribution (range 0-24) were assessed with the regional pain scale [37]. Higher scores indicate higher levels of pain.

Hospital Anxiety and Depression Scale. The Hospital Anxiety and Depression

Scale is a brief (14 item) self-report questionnaire measuring anxiety and depression [38]. It was developed for use in general medical out-patient clinics but is now widely used in clinical practice and research. The Hospital Anxiety and Depression Scale has good reliability and construct validity [39]. Responses are based on the relative frequency of symptoms over the past week, using a four point Likert scale ranging from 0 (not at all) to 3 (very often). The total score for each of the depression and anxiety scales can range from 0 to 21 and higher scores indicate higher depression and anxiety.

Quality of Life Scale. The Quality of Life Scale is a 16 item questionnaire designed specifically for use in chronic disease patients including patients with fibromyalgia [40]. Items assess various aspects of life such as physical and material well-being, relationships with other people, social, community, and civic activities, personal development and fulfillment, recreation, and independence. Satisfactory construct validity has been shown for the German version in a sample of fibromyalgia patients [41]. The items are scaled from 1 to 7 and aggregated into a sum score where a higher score indicates higher quality of life (possible sum score range 16-112).

Short Form 12. The Short Form 12 is a multidimensional general measurement instrument assessing health-related quality of life. It has become widely used in clinical trials and as a standard outcome assessment instrument because of its brevity and psychometric performance [42]. Both mental and physical health composite scores were used in this study ranging from 0 to 100 with higher scores indicating better mental and physical health.

State-Trait Anger Inventory-II. The State-Trait Anger Inventory-II is one of the most

widely used measures of anger [43]. It contains five subscales measuring: 1) state anger (raw score range 10-40), 2) trait anger (10-40), 3) anger-in (holding in or suppressing) (8-32), 4) anger-out (expressing anger toward other people or objects in environment) (8-32), and 5) anger-control (controlling retention or expression of anger) (8-32). The STAXI-2 contains 57 items responded to on a 1 (almost never) to 4 (almost always) response scale with higher scores indicating higher levels of anger.

Socio-demographics. Age, education, religion, sex and marital status were assessed. Two questions concerning the degree of religiosity (How religious are you?) and spirituality (How spiritual are you?) were included. The latter were measured with an 11-point rating scale (0=not religious/spiritual at all to 10=extremely religious/spiritual).

Statistical analysis

One way multivariate analyses of covariance were used to examine FMS patient versus control differences on all health and well-being outcome variables while holding constant the effects of age, marital status, education, religiosity, and spirituality. Subsequent univariate one way analyses of covariance were computed on each separate outcome. Wilk's lambda and F statistics are reported for the multivariate test and F is reported for the univariate tests. Partial eta squared (η_p^2) is reported as the effect size for both multivariate and univariate tests. Partial correlations (r_p) controlling socio-demographic and religious/spiritual variables, are reported for associations between forgiveness and health and well-being outcomes and to determine if the healthy associations expected to be present in control participants extended to FMS patients. Differences between FMS patients and controls in the magnitude of partial correlations were assessed using Cohen's Z-test

for differences between independent correlation coefficients [44]. Missing data was imputed using expectation maximization (EM) methods. Prior to imputation, missing data were determined to be missing at random. Missing values on any given variable were not related to missing values on any other variable (average $r = .11$). Results remain substantively unchanged in original and imputed datasets. Statistical significance was set at $p < .05$.

Results

Socio-demographic analyses

Patients and controls did not differ in terms of religious affiliation (i.e., mostly Christian) or sex (i.e., mostly female) (see Table 1). Patients were about 12 years older, less likely to be single and living alone, more likely to have been widowed, less educated, and more religious and spiritual.

Main analyses

As hypothesized, there was a large difference between FMS patients and controls on the overall multivariate composite of pain, anxiety, depression, anger, health, and quality of life ($\lambda = .19$, $F(14, 234) = 69.28$, $p < .001$, $\eta_p^2 = .81$). Not as hypothesized, there was a moderate difference between FMS patients and controls on the overall multivariate composite of forgiveness of oneself and others ($\lambda = .92$, $F(2, 246) = 10.90$, $p < .001$, $\eta_p^2 = .08$). Examining the univariate tests (see Table 2) revealed that the largest differences between FMS patients and controls were on pain and physical health followed by mental health and quality of life. Differences on anger and forgiveness were smaller but statistically significant. Differences on anger out and anger control were not statistically significant.

Partial correlations (r_p) between forgiveness and all outcomes are presented in Table 3. Associations between forgiveness of self and severity and number of regions of pain approached statistical significance ($ps < .10$) for both FMS patients and controls. Statistically significant correlations between forgiveness of self and mental health, quality of life, and anger ranged between $-.23$ to $-.56$ for FMS patients and between $-.45$ to $-.62$ for controls (see Table 3). Statistically significant correlations between forgiveness of others and mental health, quality of life, and anger ranged between $-.19$ to $-.48$ for FMS patients and between $-.21$ to $-.55$ for controls. Magnitude of associations of forgiveness of self and others with all outcomes did not differ significantly ($Zs < 1.96$, $ps > .05$) between FMS patients and controls.

Discussion

This is the first study, to our knowledge, to compare levels of forgiveness of oneself and others between FMS patients and controls. It is also the first time that the associations between forgiveness and pain, health, quality of life, and anger have been examined and directly compared between FMS patients and controls to examine the robustness of potentially beneficial associations between forgiveness and these outcomes.

Confirming prior research, FMS patients showed significantly higher levels of pain, anxiety, depression, and anger, and lower levels of health and quality of life, as compared to controls [1-5]. A new finding though was small to moderate and statistically significant differences between FMS patients and controls on forgiveness. FMS patients showed lower levels of forgiveness of oneself and others compared to controls. Given the number of interpersonal stressors and challenges that FMS patients face [32], it is important to consider that lower forgiveness may be an

important point of intervention for coping with these experiences. Most associations between forgiveness of self and others with anxiety, depression, anger, health, and quality of life were statistically significant, in the predicted direction, of moderate size, and consistent in direction and magnitude with much previous research in pain patients and healthy populations [28, 29, 45]. Associations between forgiveness of self and pain severity and number of pain regions approached statistical significance and remain clinically significant issues to address. There were no differences between FMS patients and controls in the magnitude of associations suggesting that potentially beneficial associations of forgiveness with these outcomes is equally available to patients with FMS as the healthy.

As compared to forgiveness of others, associations between forgiveness of self and pain, anxiety, depression, anger, health, and quality of life were stronger in both patients and controls. This is consistent with previous research [46, 47] in healthy populations and may reflect the especially difficult and chronic nature of not forgiving oneself. In essence, when an individual is considered one's own offender, it may be very difficult to escape lingering feelings of guilt, shame, and remorse and this may be associated with poor health and well-being [48]. Because one cannot escape oneself as an offender this might be a constant reminder of wrongdoing that stimulates continued guilt, shame, and remorse that may become a self-focused ruminative style. Self-forgiveness may, more so than forgiveness of others, offer the relief needed most from perfectionistic [49] and self-critical and uncompassionate tendencies [50, 51].

FMS patients exhibited significantly lower forgiveness of others and self, compared to controls. Recently the role of forgiveness in chronic pain and FMS [33] has been

comprehensively reviewed and several identified biographical incidents and life domains where (the lack of) forgiveness could potentially play a role in FMS patient groups has been discussed including, to name only a few, family, workplace, social situations, and medical interactions. This list, however, provides only a brief mention of some of the most obvious places where forgiveness might offer helpful relief for FMS patients, and other domains certainly exist (e.g., sociocultural stigma).

Knowing that forgiveness is lower in FMS patients, and yet, that its potential for benefiting pain, health, and quality of life is equally strong for these patients as for healthy controls, invites the question of how to promote forgiveness in FMS patients. Unfortunately, limited attention has been devoted to designing FMS-specific programs for enhancing forgiveness as part of a means of ensuring good mental health and quality of life in these patients. Two small scale studies [34, 52] suggest that forgiveness promotion can be systematically delivered and well-received by FMS patients and that these patients can enjoy and benefit (e.g. mental health) from the experience. More extensive forgiveness education research is required to fully address this issue in order to provide evidence-based programs to support the development of this coping skill in patients. Perhaps evidence-based forgiveness psycho-educational programs that have been repeatedly proven to be beneficial in healthy populations [53, 54] will be adopted and more regularly used in both clinical treatment and research with FMS patients. Similarly, focused clinical and research efforts incorporating the process of forgiveness into conceptually-related treatments (e.g., acceptance-based modalities) [20] will likely prove useful. These methods, whether stand-alone or conceptually-related interventions, offer effective techniques for facilitating empathy, compassion, altruism, and commitment, which are in and of themselves likely to be helpful to FMS patients, but all the more so to the extent that

these experiences promote forgiveness and its health benefits. Given the growing evidence that forgiveness is connected to health [15, 22, 31], promoting forgiveness in FMS patients may offer new avenues for encouraging healthy adjustment and health.

As with all observational studies, the present investigation has some limitations. First, our measurements are based on self-reports and response bias (i.e., faking good) is a possibility. Second, as an observational study, causal inferences cannot be made but can only be suggested or theorized. Third, this is a self-selected sample from only one region of the world. Last, forgiveness may reflect other positive coping strategies (e.g., problem-focused coping) and our analyses do not control these variables.

To conclude, forgiving oneself and others is one aspect of psychosocial care in FMS that could potentially be used to help promote enhanced mental health and quality of life and reduced anger. The present findings indicate that FMS patients report lower forgiveness of self and others than healthy controls, and yet, the beneficial associations of forgiveness with mental health, quality of life, and anger are equally strong in FMS patients as in healthy controls. Our findings may have clinical implications; bolstering forgiveness levels in FMS patients may result in improvements in other areas of healthy functioning for FMS patients. While future randomized controlled studies will have to bear out cause-and-effect relationships, this initial work provides the first known evidence that forgiveness is lower in FMS patients and that beneficial associations with pain, health, and quality of life are present and not diminished by FMS. Continued attention to building psychosocial resources for coping with FMS will bring greater attention to protective mechanisms

such as forgiveness that hold potential to ameliorate symptoms of pain and psychological distress and improve quality of life.

Conflicts of interest

The authors declare to have no conflicts of interest.

Acknowledgement

We would like to thank the German Fibromyalgia Patient Association (DFV) for their support.

References

1. Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, Tugwell P, Campbell SM, Abeles M, Clark P, et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990;33:160-72.
2. Henriksson C, Gundmark I, Bengtsson A, Ek AC. Living with fibromyalgia. Consequences for everyday life. *Clin J Pain* 1992;8:138-44.
3. Wolfe F, Anderson J, Harkness D, Bennett RM, Caro XJ, Goldenberg DL, Russell IJ, Yunus MB. Health status and disease severity in fibromyalgia: results of a six-center longitudinal study. *Arthritis Rheum* 1997;40:1571-9.
4. Salaffi F, Sarzi-Puttini P, Girolimetti R, Atzeni F, Gasparini S, Grassi W. Health-related quality of life in fibromyalgia patients: a comparison with rheumatoid arthritis patients and the general population using the SF-36 health survey. *Clin Exp Rheumatol* 2009;27:S67-74.
5. White KP, Speechley M, Harth M, Ostbye T. The London Fibromyalgia Epidemiology Study: direct health care costs of fibromyalgia syndrome in London, Canada. *J Rheumatol* 1999;26:885-9.
6. Turk DC, Rudy TE. Neglected factors in chronic pain treatment outcome studies--referral patterns, failure to enter treatment, and attrition. *Pain* 1990;43:7-25.
7. Turk DC, Rudy TE. The robustness of an empirically derived taxonomy of chronic pain patients. *Pain* 1990;43:27-35.
8. Turk DC, Okifuji A, Sinclair JD, Starz TW. Differential responses by psychosocial subgroups of fibromyalgia syndrome patients to an interdisciplinary treatment. *Arthritis Care Res* 1998;11:397-404.
9. Williams DA, Clauw DJ. Understanding fibromyalgia: lessons from the broader pain research community. *J Pain* 2009;10:777-91.

10. Martinez-Lavin M. Fibromyalgia: When Distress Becomes (Un)sympathetic Pain. *Pain Res Treat* 2012;2012:981565.
11. Fischer S, Doerr JM, Strahler J, Mewes R, Thieme K, Nater UM. Stress exacerbates pain in the everyday lives of women with fibromyalgia syndrome-The role of cortisol and alpha-amylase. *Psychoneuroendocrinology* 2015;63:68-77.
12. Chida Y, Steptoe A. The association of anger and hostility with future coronary heart disease: a meta-analytic review of prospective evidence. *J Am Coll Cardiol* 2009;53:936-46.
13. Sturgeon JA. Psychological therapies for the management of chronic pain. *Psychology Research and Behavior Management* 2014;7.
14. Cianfrini LR, Block C, Doleys DM. Psychological therapies. In: Deer T, Leong M, Ray A eds. *Treatment of chronic pain by integrative approaches: The American Academy of Pain Medicine textbook on patient management*. New York, NY: Springer Science + Business Media, 2015:83-100.
15. Borkum JM. Maladaptive cognitions and chronic pain: Epidemiology, neurobiology, and treatment. *Journal of Rational-Emotive & Cognitive-Behavior Therapy* 2010;28:4-24.
16. Dahl J, Lundgren T. Acceptance and Commitment Therapy (ACT) in the Treatment of Chronic Pain. In: Baer R ed. *Mindfulness-based treatment approaches: Clinician's guide to evidence base and applications*. San Diego, CA: Elsevier Academic Press, 2006:285-306.
17. Bernardy K, Klose P, Busch AJ, Choy EHS, Häuser W. Cognitive behavioural therapies for fibromyalgia. *Cochrane Database Syst Rev* 2013.
18. Legaree T-A, Turner J, Lollis S. Forgiveness and therapy: A critical review of conceptualizations, practices, and values found in the literature. *J Marital Fam Ther* 2007;33:192-213.

19. Zettle RD, Barner SL, Gird S. ACT with depression: The role of forgiving. *Acceptance and commitment therapy: Contemporary theory, research and practice*. Bowen Hills, QLD Australia: Australian Academic Press, 2009:151-173.
20. Webb JR, Hirsch JK, Toussaint L. Forgiveness as a positive psychotherapy for addiction and suicide: Theory, research, and practice. *Spiritual Clin Pract* 2015;2:48.
21. Wade NG, Hoyt WT, Kidwell JEM, Worthington Jr EL. Efficacy of psychotherapeutic interventions to promote forgiveness: A meta-analysis. *J Consult Clin Psychol* 2014;82:154-170.
22. Worthington Jr EL, Lavelock C, vanOyen Witvliet C, Rye MS, Tsang J-A, Toussaint L. Measures of Forgiveness: Self-Report, Physiological, Chemical, and Behavioral Indicators. In: Boyle GJ, Saklofske DH, Matthews G eds. *Measures of Personality and Social Psychological Constructs*. Oxford: Academic Press, 2014:474-502.
23. Toussaint L, Shields GS, Dorn G, Slavich GM. Effects of lifetime stress exposure on mental and physical health in young adulthood: How stress degrades and forgiveness protects health. *J Health Psychol* 2014:1-11.
24. Toussaint L, Webb JR. Theoretical and Empirical Connections Between Forgiveness and Mental Health and Well-Being. In: Worthington EL ed. *Handbook of Forgiveness*. New York, NY: Brunner-Routledge, 2005:349-362.
25. Toussaint LL, Owen AD, Cheadle A. Forgive to live: forgiveness, health, and longevity. *J Behav Med* 2012;35:375-86.
26. Whited MC, Wheat AL, Larkin KT. The influence of forgiveness and apology on cardiovascular reactivity and recovery in response to mental stress. *J Behav Med* 2010;33:293-304.

27. Witvliet CV, Ludwig TE, Vander Laan KL. Granting forgiveness or harboring grudges: implications for emotion, physiology, and health. *Psychol Sci* 2001;12:117-123.
28. Rippentrop AE, Altmaier EM, Chen JJ, Found EM, Keffala VJ. The relationship between religion/spirituality and physical health, mental health, and pain in a chronic pain population. *Pain* 2005;116:311-321.
29. Carson JW, Keefe FJ, Goli V, Fras AM, Lynch TR, Thorp SR, Buechler JL. Forgiveness and chronic low back pain: A preliminary study examining the relationship of forgiveness to pain, anger, and psychological distress. *J Pain* 2005;6:84-91.
30. Worthington EL, Scherer M. Forgiveness is an emotion-focused coping strategy that can reduce health risks and promote health resilience: Theory, review, and hypotheses. *Psychology & Health* 2004;19:385-405.
31. Worthington E. *Forgiveness and Reconciliation: Theory and Application*. New York: Routledge, 2006.
32. Toussaint L, Overvold-Ronningen M, Vincent A, Luedtke C, Whipple M, Schriever T, Luskin F. Implications of forgiveness enhancement in patients with fibromyalgia and chronic fatigue syndrome. *J Health Care Chaplain* 2010;16:123-39.
33. Offenbächer M, Dezutter J, Vallejo M, Toussaint L. The Role of Forgiveness in Chronic Pain and Fibromyalgia. In: Toussaint L, Worthington E, Williams D eds. *Forgiveness and health: Scientific evidence and theories relating forgiveness to better health*. Netherlands: Springer., 2016.
34. Toussaint LL, Vincent A, Whipple MO, McAllister SJ, Finnie DM, Hathaway JC, Oh TH, Fleming KC, Douglas KSV. Forgiveness education in fibromyalgia: A qualitative inquiry. *Pain Studies and Treatment* 2014;2014.

35. Mauger PA, Perry JE, Freeman T, Grove DC, McBride AG, McKinney K. The measurement of forgiveness: preliminary research. *J Psychol Christianity* 1992;11:170-180.
36. Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related quality of life measures: literature review and proposed guidelines. *J Clin Epidemiol* 1993;46:1417-32.
37. Wolfe F. Pain extent and diagnosis: development and validation of the regional pain scale in 12,799 patients with rheumatic disease. *J Rheumatol* 2003;30:369-78.
38. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-70.
39. Hinz A, Schwarz R. Hospital Anxiety and Depression Scale - Deutsche Version (HADS-D). *Diagnostica* 2002;48:112-113.
40. Burckhardt CS, Clark SR, Bennett RM. Fibromyalgia and quality of life: a comparative analysis. *J Rheumatol* 1993;20:475-9.
41. Offenbächer M, Sauer S, Kohls N, Waltz M, Schoeps P. Quality of life in patients with fibromyalgia: validation and psychometric properties of the German Quality of Life Scale (QOLS-G). *Rheumatol Int* 2012;32:3243-3252.
42. Gandek B, Ware JE, Aaronson NK, Apolone G, Bjorner JB, Brazier JE, Bullinger M, Kaasa S, Leplege A, Prieto L, Sullivan M. Cross-validation of item selection and scoring for the SF-12 Health Survey in nine countries: results from the IQOLA Project. International Quality of Life Assessment. *J Clin Epidemiol* 1998;51:1171-8.
43. Spielberger CD. *The State-Trait Anger Expression Inventory-2 (STAXI-2): Professional manual*. Odessa, FL: Psychological Assessment Resources (PAR), 1999.
44. Cohen J, Cohen P. *Applied multiple regression/correlation analysis for the behavioral sciences*. Hillsdale, NJ: Erlbaum, 1983.

45. Worthington EL, Witvliet CVO, Pietrini P, Miller AJ. Forgiveness, health, and well-being: A review of evidence for emotional versus decisional forgiveness, dispositional forgivingness, and reduced unforgiveness. *J Behav Med* 2007;30:291-302.
46. Svalina SS, Webb JR. Forgiveness and health among people in outpatient physical therapy. *Disabil Rehabil* 2012;34:383-92.
47. Toussaint LL, Marschall JC, Williams DR. Prospective associations between religiousness/spirituality and depression and mediating effects of forgiveness in a nationally representative sample of United States adults. *Depression Research and Treatment* 2012;2012.
48. Wilson T, Milosevic A, Carroll M, Hart K, Hibbard S. Physical health status in relation to self-forgiveness and other-forgiveness in healthy college students. *J Health Psychol* 2008;13:798-803.
49. Sirois FM, Molnar DS. *Perfectionism, health, and well-being*. Cham, Switzerland: Springer International Publishing, 2016.
50. Terry ML, Leary MR. Self-compassion, self-regulation, and health. *Self and Identity* 2011;10:352-362.
51. Sirois FM. A self-regulation resource model of self-compassion and health behavior intentions in emerging adults. *Prev Med Rep* 2015;2:218-222.
52. Lee YR, Enright RD. A forgiveness intervention for women with fibromyalgia who were abused in childhood: A pilot study. *Spiritual Clin Pract* 2014;1:203-217.
53. Enright RD, Fitzgibbons RP. Helping clients forgive: An empirical guide for resolving anger and restoring hope. In: Enright RD, Fitzgibbons RP eds. *The process model of forgiveness therapy*. Washington, DC: American Psychological Association, 2000: 65-88.
54. Worthington EL. *Five steps to forgiveness: The art and science of forgiving*. New York: Crown, 2001.

Table 1. Socio-Demographic Summary Statistics for Fibromyalgia Patients and Controls

	Patients	Controls	F/χ^2	p
Age in years ¹	58 (8.8)	47 (14.2)	57.76	< .001
Gender (female/male) ²	161/9	76/5	.09	.78
Religion ²			5.96	.11
Christian	138	65		
Muslim	1	4		
..Other	1	0		
No religion	30	12		
Marital status ²			14.26	< .01
Married/with partner	130	59		
Single	8	8		
Widowed/divorced	32	11		
Education (years) ²			68.76	< .001
9 or less	74	7		
10 or 11	65	23		
12 or more	16	44		
Advanced	12	5		
How religious are you? ¹	4.9 (2.7)	4.1 (2.7)	5.29	.02
How spiritual are you? ¹	3.8 (2.8)	3.0 (2.8)	4.72	.03

¹Mean (standard deviation) and F-test; ²N and Chi-square test

Table 2. Means and Standard Errors for Fibromyalgia Patients and Controls on Pain, Mental and Physical Health, Quality of Life, Anger, and Forgiveness

	Controls		FMS		F	***	η_p^2
	M	SE	M	SE			
Pain Today	.69	.20	6.10	.13	449.33	***	.65
Pain Severity (Last 3 mos)	.90	.20	6.78	.13	518.49	***	.68
Regional Pain Scale - Severity	2.33	1.21	31.84	.77	364.08	***	.60
Regional Pain Scale - Number of Regions	1.58	.44	14.18	.28	501.50	***	.67
Anxiety	5.05	.52	10.47	.34	65.36	***	.21
Depression	2.90	.45	9.31	.29	121.60	***	.33
Quality of Life	89.13	2.02	66.06	1.29	79.44	***	.24
SF-12 Physical Health	54.50	.80	30.14	.51	572.15	***	.70
SF-12 Mental Health	50.77	1.09	35.71	.70	116.00	***	.32
STAXI State Anger	11.74	.75	15.29	.48	13.71	***	.06
STAXI Trait Anger	19.50	.66	21.59	.42	6.06	*	.02
STAXI Anger In	15.42	.62	20.38	.40	38.88	***	.14
STAXI Anger Out	13.57	.50	13.41	.32	0.06		.00
STAXI Anger Control	23.08	.54	22.28	.34	1.38		.01
Forgiveness of Others	41.58	1.14	38.33	.73	5.00	*	.02
Forgiveness of Self	44.61	1.38	36.52	.88	21.13	***	.08

* $p < .05$, *** $p < .001$