



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/104514/>

Version: Accepted Version

Article:

Wielstra, B., Voeroes, J. and Arntzen, J.W. (2016) Is the Danube crested newt *Triturus dobrogicus* polytypic? A review and new nuclear DNA data. *Amphibia-Reptilia* , 37 (2). pp. 167-177. ISSN: 0173-5373

<https://doi.org/10.1163/15685381-00003041>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Is the Danube crested newt *Triturus dobrogicus* polytypic? A review and new nuclear DNA data

Short title: **Is the Danube crested newt *Triturus dobrogicus* polytypic?**

Ben Wielstra^{1,2,*}, Judit Vörös³, Jan W. Arntzen²

¹Department of Animal and Plant Sciences, University of Sheffield, S10 2TN Sheffield, UK.

²Naturalis Biodiversity Center, P. O. Box 9517, 2300 RA Leiden, The Netherlands. ³Department of Zoology, Hungarian Natural History Museum, 1088 Budapest, Hungary

* Corresponding author;

e-mail: b.wielstra@sheffield.ac.uk

Abstract

The Danube crested newt *Triturus dobrogicus* has been proposed to comprise two subspecies: *T. d. dobrogicus* and *T. d. macrosoma*. Uncertainty exists in the literature over their distribution and diagnosability. We conduct a multilocus phylogeographical survey and review published data to determine whether a two taxon treatment is warranted. Newly produced and published nuclear DNA data suggest intraspecific variation in the Pannonian Plain part of the range, but with extensive genetic admixture, whereas mitochondrial DNA data shows a lack of geographical structuring in *T. dobrogicus* altogether. None of the studied morphological characters suggest the presence of two geographical groups in *T. dobrogicus* unequivocally. Although Danube Delta newts do have relatively short bodies compared to the remainder of the range (the Pannonian and Lower Danube Plains and the Dnepr Delta), we argue that this finding can be explained by phenotypic plasticity – particularly in light of the incongruent evolutionary scenario suggested by genetic data. We conclude that the total body of evidence does not support the two subspecies hypothesis and recommend that *T. dobrogicus* is treated as a monotypic species.

Key words: Ion Torrent, Next-generation sequencing, Subspecies, Taxonomy, *Triturus cristatus* superspecies, *Triturus dobrogicus macrosoma*

Introduction

The taxonomy of crested newts (*Triturus cristatus* superspecies) has been regularly updated. Currently six species are recognized, with a seventh awaiting formal description (Wielstra et al., 2013b). All crested newt species are considered monotypic, except for *T. dobrogicus* (Kiritzescu, 1903), for which Litvinchuk and Borkin (2000) recognized two subspecies. They reinstated *T. d. macrosoma* (Boulenger, 1908) to reflect a perceived geographical differentiation within *T. dobrogicus*.

Litvinchuk and Borkin (2000) analyzed a suite of body measurements with principal component and discriminant analysis. In the formal description they mentioned two indices as most informative in separating the two subspecies, namely the ‘Wolterstorff-index’ (forelimb length divided by interlimb distance) and Ltc/L (head width divided by body length), that are both on average higher in the nominotypical subspecies. The authors noted that the two subspecies showed differences in the number of pre-sacral rib-bearing vertebrae (NRBV; with *T. d. dobrogicus* characterized by an NRBV count of 16, and *T. d. macrosoma* by an NRBV count of 17). Litvinchuk and Borkin (2000) also noted differences in coloration of the belly, which tends towards red in *T. d. dobrogicus* and orange or yellow in *T. d. macrosoma* and noted that the nominotypical subspecies has a more polished skin, more obvious costal grooves on the sides of the body and smaller and sparser rounded spots on the belly. Larvae of *T. d. dobrogicus* were said to be usually darker than those of *T. d. macrosoma*. Based on a crossing experiment the authors suggested a reduced fitness for offspring from crosses between representatives of the two subspecies.

According to Litvinchuk and Borkin (2000), the nominotypical subspecies is distributed in the Danube Delta “along the lower Danube, probably, eastward to Reni [Romania]” and ssp. *T. d. macrosoma* in the Pannonian and Lower Danube Plains (Fig. 1A). Other authors have suggested that the Iron Gate (the gorge where the Danube river makes its way through the Southern Carpathians) separates the ranges of the two subspecies (Raffaëlli, 2007; Thiesmeier et al., 2009). An allopatric crested newt population in the Dnepr Delta (Fig. 1A) was attributed to *T. dobrogicus* by Litvinchuk (2005), but not at the level of the subspecies.

The study by Litvinchuk and Borkin (2000) did not include samples from across the entire Danube crested newt range. Notably, material from the Lower Danube Plain and the Dnepr Delta (the latter not known to harbor *T. dobrogicus* at the time) was not included. This hampers determining the distribution of the two *T. dobrogicus* taxa. Furthermore, subsequent studies have casted doubt on the diagnosability of the two subspecies based on either genetic or morphological data (e.g. Vörös and Arntzen, 2010; Naumov and Biserkov, 2013) and the taxonomy of Litvinchuk and Borkin (2000) has not universally been adopted (e.g. Sparreboom, 2014).

In an attempt to shed light on the intraspecific taxonomy of *T. dobrogicus* we conduct a phylogeographic survey using an Ion Torrent next-generation sequencing protocol that provides large scale nuclear DNA data for crested newts (Wielstra et al., 2014a). We interpret the new results in the context of a literature review on intraspecific genetic and morphological variation in *T. dobrogicus* and use the total body of evidence to decide whether to support or contradict the two subspecies hypothesis. This raises the question on how to delineate a subspecies in the first place. For background on the difficulties of defining subspecies and applying the subspecies rank in taxonomy we refer to Mayr (1969). We here take a pragmatic approach. To accept the two subspecies hypothesis we would require to find geographically consistent intraspecific variation that cannot be explained by environmental plasticity.

Material and methods

Sampling and laboratory methods

Our sampling covered the entire range of *T. dobrogicus* (Fig. 1A; Table 1). We included 46 populations and 121 individuals (1-3 individuals with on average 2.6 individuals per population). Twenty-nine populations with 83 individuals were from the Pannonian Plain, 12 populations with 27 individuals were from the Lower Danube Plain, four populations with ten individuals were from the Danube Delta and one population with one individual was from the Dnepr Delta. We obtained data for 52 nuclear DNA markers following the protocol of Wielstra et al. (2014a). In brief, we amplified markers of c. 140 bp in length (excluding primers, see Table S1), positioned in 3' untranslated regions, in five multiplex PCRs. We pooled the multiplexes for each individual and ligated unique tags to be able to recognize the product belonging to each individual. We sequenced the amplicons on

the Ion Torrent next-generation sequencing platform and processed the output with a bioinformatics pipeline that filters out poor quality reads, identifies alleles and converts data to a format directly usable for population genetic analysis. Sequence data for a mtDNA marker (ND4, 658 bp) were taken from, or newly produced following the protocol in Wielstra et al. (2013a). For six individuals we did not manage to obtain mtDNA sequence data.

Nuclear DNA: testing for interspecific gene flow

Because crested newt species hybridize at their contact zones (Arntzen et al., 2014) we aimed to exclude potentially confounding effects of interspecific gene flow. Therefore we included data from Wielstra et al. (2014a) for the four species with which *T. dobrogicus* is in spatial contact, namely *T. carnifex* in the west, *T. cristatus* in the north, *T. ivanbureschi* in the southeast and *T. macedonicus* southwest. We analyzed the data with BAPS 6 (Corander et al., 2008; Cheng et al., 2013) and, considering that five crested species were involved in the comparison, enforced the number of distinct gene pools to five (i.e. $k = 5$). This should highlight any *T. dobrogicus* individuals showing introgression from other *Triturus* species. We conducted ten replicate runs and tested for admixture between gene pools. Newts ascribed to *T. dobrogicus* with a probability less than unity were excluded from further analyses.

Nuclear DNA: testing for intraspecific genetic structuring

We used FSTAT (Goudet, 1995) to determine gene diversity and the number of alleles per marker and the percentage of missing data. We conducted a spatially explicit Bayesian clustering analysis with BAPS 6 and TESS 2.3.1 (Chen et al., 2007; Durand et al., 2009). We determined the optimal number of gene pools k over a range of 1-44 (the upper limit is defined by the number of populations included), using ten replicates per k value. BAPS determines the optimum k value internally. In TESS the optimum k value was defined as the one where the average deviance information criterion value reached a plateau (as advocated in the TESS manual). In BAPS we incorporated the geographical origin of individuals ('spatial clustering') and tested for admixture between gene pools. In TESS we modelled admixture using the conditional autoregressive model, with 20,000 sweeps of which 5,000 were discarded as burn-in. The spatial interaction parameter was kept at the default with the option to update this parameter activated. The estimated admixture proportions for independent runs were averaged using CLUMPP 1.1.2 (Jakobsson and Rosenberg, 2007). All output was visualized with DISTRUCT (Rosenberg, 2004).

Nuclear DNA: population differentiation in *T. dobrogicus*

We constructed a population tree with the program POPTREE2 (Takezaki et al., 2010). To determine the position of the root we added *T. cristatus* as outgroup. We used the Neighbour Joining method based on uncorrected F_{ST} distance and the robustness of interpopulational relationships was tested with 1000 bootstrap replicates. Eight markers had to be excluded in this exercise because data were missing for one or more populations.

Mitochondrial DNA analyses

Mitochondrial DNA haplotypes were identified by comparison against the mtDNA haplotype database presented in Wielstra et al. (2013a). To determine to which *Triturus* species newly identified haplotypes belonged we constructed a Neighbour-Joining tree with 1000 bootstrap replicates in MEGA6 (Tamura et al., 2013). A Median Joining network (Bandelt et al., 1999) was created using Network 4.6.11 (www.fluxus-engineering.com).

Data accessibility

Sampling details are presented in Table S2 and GenBank Accession numbers for mtDNA are in Table S3. Raw Ion Torrent reads in FASTQ format; scripts associated with the bioinformatics pipeline; BWA alignments in SAM format; raw SNP reports in VCF and BCF format; the filtered SNP report used to construct consensus sequences; an overview of the number of reads in total and per marker and/or individual; nuclear DNA data in genotypic format; nuclear DNA sequence alignments; and input and output files for Network, FSTAT, BAPS, TESS and POPTREE; are available via Dryad Digital Repository entry [doi:10.5061/dryad.b9765](https://doi.org/10.5061/dryad.b9765).

Results

Results of the analysis testing for interspecific gene flow of nuclear DNA are summarized in Table 1 with details in Table S2. Five crested newt populations have *T. dobrogicus* in syntopy with another *Triturus* species (*T. carnifex* once, *T. macedonicus* once and *T. cristatus* three times) and seven populations have *T. dobrogicus* individuals showing genetic admixture with other *Triturus* species. Altogether we excluded 18 individuals from ten populations in further analyses, of which two populations in their entirety (Table S2).

The markers in the dataset used for the intraspecific analyses on average had 6.0 ± 3.1 (standard deviation) alleles and an average gene diversity of 0.35 (details in Table S1). The results for the analyses testing for intraspecific genetic clustering show the same overall pattern, with minor differences in details (Fig. 2, Table S2). BAPS yields $k = 3$ as the most probable number of gene pools present in *T. dobrogicus*. Although TESS also identifies $k = 3$ as the most likely solution, support for allocation of individuals to the third group is negligible. Hence we also provide results for TESS under the next lowest meaningful k value, i.e. $k = 2$ (Fig. 2; Table S2). The third TESS group under $k = 3$ is then subsumed in the first TESS group under $k = 2$ whereas the distinction between a first and a second group is similar under both k values (details in Table S2). From here on we only consider TESS results under $k = 2$.

Two of the three BAPS groups roughly correspond to one of the two groups in TESS. The overall pattern is one group occupying the southwest of the Pannonian Plain range (populations 12-21 and partially populations 7 and 10 in Fig. 1A) and another group occupying the remainder of the range (including the Pannonian and Lower Danube Plains and the Danube and Dnepr Deltas). The results differ in the amount of admixture inferred between these two groups (Fig. 2; details in Table S2). TESS generally suggests a more pronounced admixture between these groups than does BAPS. Conversely, BAPS suggests admixture in one individual from the Danube Delta (from population 42), which was not found by TESS.

The population tree based on nuclear DNA genetic distances constructed with POPTREE suggests that the four range sections, Pannonian and Lower Danube Plains and Danube and Dnepr Deltas, do not constitute reciprocally monophyletic groups (Fig. 1B). The populations comprising the southwest Pannonian Plain group identified by BAPS and TESS (populations 12-21 in Fig. 1A) cluster together, but bootstrap support for monophyly of this assemblage is low (Fig. 1B).

We observed one instance of a pure *T. dobrogicus* individual based on nuclear DNA possessing mtDNA originating from another species, namely *T. ivanbureschi* (Table S2). The haplotype network for the *T. dobrogicus* mtDNA indicates a lack of geographical structuring (Fig. 3) as e.g. illustrated by the central haplotype Tdob02, which is found all across the range of *T. dobrogicus*, in the Pannonian and Lower Danube Plains and in the Danube and Dnepr Deltas.

Discussion

Interpretation of published and newly produced genetic data

Mitochondrial DNA shows a shallow genetic divergence across the *T. dobrogicus* range, with identical haplotypes present in the Pannonian, Lower Danube and Dnepr range sections (Wallis and Arntzen, 1989; Wielstra et al., 2013a; this study). Allozyme data suggest a higher level of intraspecific genetic variation than mtDNA, but agree on a lack of geographical structure between the Pannonian and Lower Danube Plain populations (Vörös and Arntzen, 2010). Rather, populations from the southwest of the Pannonian Plain are relatively distinct from those in the remainder of the range. No material from the Danube Delta is included in Vörös and Arntzen (2010). However, the allozyme data presented in Litvinchuk et al. (1994) show a Nei's genetic distance of zero between a Danube Delta and Transcarpathian (northeast Pannonian Plain) population sample (Litvinchuk and Borkin, 2000; Arntzen, 2003; Naumov and Biserkov, 2013).

The newly produced nuclear DNA data improve upon previous studies in the number of markers analyzed and the geographical coverage of population samples. Although the results of the cluster-based analysis of the new data are in line with the presence of two genetic clusters within *T. dobrogicus*, the spatial arrangement differs distinctly from the one proposed by Litvinchuk and Borkin (2000). One group is distributed in the southwest of the Pannonian Plain range (mainly populations 12-21 in Fig. 1A) and another group occupies the remainder of the range, i.e. the rest of the Pannonian Plain, the Lower Danube Plain and the Danube and Dnepr Delta. These findings are similar to those based on allozyme data (Vörös and Arntzen, 2010). The new nuclear DNA data agree with all previous genetic studies that genetic admixture between intraspecific genetic groups in *T. dobrogicus* is extensive. Furthermore, we do not find significant bootstrap support for either group identified in the population tree based on the new data.

Crucially, our genetic results do not support long term limitations to gene flow between the Danube Delta and the remainder of the range (including the currently allopatric Dnepr Delta). This finding is not in line with the crossing experiment of Litvinchuk and Borkin (2000), which suggests

relatively low survival resulting from the cross of *T. d. dobrogicus* and *T. d. macrosoma* (it should be note that half of the offspring in the genus *Triturus* dies during embryonic development due to the peculiar 'chromosome 1 syndrome'; Macgregor and Horner, 1980; Ridley, 2004). However, it should be noted that sample sizes in this experiment are extremely low, with $n = 2$ for *T. d. dobrogicus* x *T. d. dobrogicus*, $n = 2$ for *T. d. dobrogicus* x *T. d. macrosoma* and $n = 0$ for *T. d. macrosoma* x *T. d. macrosoma*. The replicates for the *T. d. dobrogicus* x *T. d. macrosoma* differ widely in embryo and larval survival (1.9% versus 13.3% and 100% versus 51.1%). Embryo survival for *T. d. dobrogicus* x *T. d. macrosoma* is unrealistically low, even lower than for a cross between two distinct *Triturus* species (*T. carnifex* x *T. karelinii*; see Table 4 in Litvinchuk and Borkin, 2000). Because, in contrast to the two *T. dobrogicus* subspecies, the different *Triturus* species are characterized by distinct average genome sizes (Litvinchuk et al., 1999), genetic incompatibilities would *a priori* be expected to play a larger role in interspecific crosses. Considering the low sample size in the crossing experiment, alternative explanations for low survival (e.g. disease outbreak, a problem with temperature control, etc.) cannot be safely excluded. We consider the results of the crossing experiment unconvincing and put our trust in the molecular genetic data.

Interpretation of published morphological data

Litvinchuk and Borkin (2000) mention in their formal diagnosis of the two subspecies in *T. dobrogicus* that two indices, the 'Wolterstorff-index' (forelimb length divided by interlimb distance) and Ltc/L (head width divided by body length), are both on average higher in the nominotypical subspecies (the former for males only). These findings indicate that newts from the Danube Delta possess relatively short bodies. Naumov and Biserkov (2013) conducted a morphological survey for *T. dobrogicus*, adding material from the Lower Danube Plain that was not represented by Litvinchuk and Borkin (2000) and concluded that the new material is closer to *T. d. macrosoma* from the Pannonian Plain than it is to *T. d. dobrogicus* from the Danube Delta.

We summarized data for the Wolterstorff and Ltc/L indices for individual newts from Litvinchuk and Borkin (2009) and Naumov and Biserkov (2013) and Wolterstorff-index data only from Arntzen and Wallis (1999). We excluded populations located near the contact zones with other

Triturus species to minimize confounding effects of interspecific gene flow (see Table S4 for the total dataset). Data for males and females are plotted in Fig. 4 for the Pannonian Plain, the Lower Danube Plain, the Danube Delta and the Dnepr Delta separately. Student's *t*-tests confirm that for the Wolterstorff-index males ($t=5.93$, $P<0.01$) but not females ($t=1.24$, $P>0.05$) and for the Ltc/L both males ($t=4.80$, $P<0.001$) and females ($t=4.624$, $P<0.001$) from the Danube Delta on average show higher values – and therewith shorter bodies – compared to those from the Pannonian plus Lower Danube Plains.

One of us (JWA, pers. obs.) has noted a marked phenotypic plasticity in body shape in *T. dobrogicus* larvae, with particular stout bodies in water bodies with fish predators present, versus a more regular, elongated form typical for the genus as a whole. We suggest a similar phenotypic plasticity may apply to adults as well. This would explain why the genetic data suggest an incongruent evolutionary scenario, with the deepest intraspecific divergence found within the Pannonian Plain. In this light we doubt the taxonomical relevance of the observed body shape differentiation in *T. dobrogicus*.

The use of indices hampers interpretation of the diagnosticity of individual characters. Arntzen and Wallis (1994) find the number of rib-bearing vertebrae (NRBV) to better discriminate the different *Triturus* species. Litvinchuk and Borkin (2000) suggest that NRBV differs between *T. d. dobrogicus* and *T. d. macrosoma* on average, with *T. d. dobrogicus* more inclined to show an NRBV count of 16 and *T. d. macrosoma* an NRBV count of 17. However, their sample sizes are small, overlap in NRBV count is rampant (Table 3 in Litvinchuk and Borkin, 2000) and a G-test for independence does not suggest a significant difference between the two subspecies ($G=0.50$, $P>0.05$). A more comprehensive sampling shows that in the Pannonian Plain an NRBV count of 16 or 17 occurs at the same frequency (Arntzen et al., 2015). The limited sampling in the Lower Danube Plain shows two individuals with an NRBV count of 16 and two with 17.

Although NRBV is known to show intraspecific plasticity (Slijepčević et al., 2015), Litvinchuk and Borkin (2000) hypothesize that the frequency of an NRBV count of 16 in their material of *T. d. macrosoma* could be inflated due to genetic admixture with *T. cristatus* (and implicitly that this is not the case for *T. d. dobrogicus*). This hypothesis has some merit because

Arntzen et al. (2014) show that *T. dobrogicus* individuals from the contact zone with *T. cristatus* more often show an NRBV count of 16 and those away from contact zones an NRBV count of 17. However, we argue that without additional evidence that different mechanisms underlie an NRBV count of 16 in *T. d. macrosoma* and *T. d. dobrogicus*, we should stick to the simplest explanation of the data. We conclude that the currently available data do not support a frequency differentiation in NRBV count across the *T. dobrogicus* range.

Litvinchuk and Borkin (2000) consider the coloration of the belly, almost red in *T. d. dobrogicus* and orange or yellow in *T. d. macrosoma*, to distinguish the two subspecies (although raw data is not provided and it is not stated how color was quantified). Considering that the ‘redness’ of the belly in amphibians is influenced by diet (Matsui et al., 2003) and UVB exposure (Michaels et al., 2015), we question its applicability in crested newt taxonomy.

Although not mentioned in the taxonomic account as distinguishing the two subspecies, Litvinchuk and Borkin (2000) mention additional perceived differences in the main text. The frequency with which black spots merge on the belly and form a dorso-ventral black stripe is suggested to be lower in the Danube Delta than in the Pannonian Plain. As both the size and density of belly spots increase with age (Lantz, 1953; Arntzen and Teunis, 1993) we advise against using these characters in crested newt taxonomy. The obviousness of costal grooves seems to us to depend in the first place on the feeding condition of animals and we doubt its relevance for taxonomy. Without rigorous quantification it is impossible to interpret the relevance of differences in how polished the skin is. *Triturus dobrogicus* larvae stand out from the larvae of other *Triturus* species by a deep black coloration at large (> 50 mm) size (Arntzen, 2003), but we cannot confirm a geographical pattern within *T. dobrogicus*. The pigmentation of *Triturus* larva is most likely subject to phenotypic plasticity (Cvijanović et al., 2015) and we do not consider darkness of larvae taxonomically informative.

Taxonomy of T. dobrogicus

Based on the accumulated data from multiple molecular marker systems and morphological characters we can address the question whether the data so far support the two subspecies treatment of *T.*

dobrogicus. All molecular genetic data disagree with the treatment by Litvinchuk and Borkin (2000) of a distinct group inhabiting the Danube Delta. Furthermore, the molecular genetic data show that the Iron Gate, which has sometimes been interpreted as separating the ranges of two *T. dobrogicus* subspecies (Raffaëlli, 2007; Thiesmeier et al., 2009), does not pose a barrier to gene flow between the Pannonian and Lower Danube range sections (for further discussion see Arntzen et al., 1997; Gherghel and Papeş, 2015). Rather, for nuclear DNA the bulk of intraspecific genetic variability is to be found within the Pannonian range section of *T. dobrogicus*, with one distinct genetic cluster inhabiting the southwest of the Pannonian Plain and the other inhabiting the remainder of the range, including the Danube Delta. However, genetic admixture between these two genetic groups is rampant. Furthermore, we contest that the morphological data point towards a distinct taxon inhabiting the Danube Delta. We believe that other potential explanations for intraspecific variation within *T. dobrogicus* – related to age, environment and interspecific gene flow – have not been sufficiently excluded and doubt that some characters express geographical differentiation at all. Hence we conclude that there is insufficient support for the two subspecies treatment proposed by Litvinchuk and Borkin (2000) and recommend that *T. dobrogicus* is treated as monotypic.

Acknowledgements

BW conducted this work as a Newton International Fellow. JV was supported by the Hungarian Scientific Research Fund (OTKA K84071), and the Bolyai János Research Scholarship of the Hungarian Academy of Sciences. We are grateful to Peter Mikulíček for DNA samples from Slovakia and to Spartak Litvinchuk and Borislav Naumov for digital copies of their morphological data.

References

- Arntzen, J.W., Teunis, S.F.M. (1993): A six year study on the population dynamics of the crested newt (*Triturus cristatus*) following the colonization of a newly created pond. *Herpetol. J.* **3**: 99-110.
- Arntzen, J.W., Wallis, G.P. (1994): The 'Wolterstorff Index' and its value to the taxonomy of the crested newt superspecies. *Abhandlungen und Berichte für Naturkunde, Magdeburg* **17**: 57-66.

- Arntzen, J.W., Bugter, R.J.F., Cogalniceanu, D., Wallis, G.P. (1997): The distribution and conservation status of the Danube crested newt, *Triturus dobrogicus*. *Amphibia-Reptilia* **18**: 133-142.
- Arntzen, J.W., Wallis, G.P. (1999): Geographic variation and taxonomy of crested newts (*Triturus cristatus* superspecies): morphological and mitochondrial data. *Contrib. Zool.* **68**: 181-203.
- Arntzen, J.W. (2003): *Triturus cristatus* Superspecies - Kammolch-Artenkreis (*Triturus cristatus* (Laurenti, 1768) - Nördlicher Kammolch, *Triturus carnifex* (Laurenti, 1768) - Italienischer Kammolch, *Triturus dobrogicus* (Kiritzescu, 1903) - Donau-Kammolch, *Triturus karelinii* (Strauch, 1870) - Südlicher Kammolch). In: *Handbuch der Reptilien und Amphibien Europas. Schwanzlurche IIA*, p. 421-514. Grossenbacher, K., Thiesmeier, B., Eds., Wiebelsheim, Aula-Verlag.
- Arntzen, J.W., Wielstra, B., Wallis, G.P. (2014): The modality of nine *Triturus* newt hybrid zones, assessed with nuclear, mitochondrial and morphological data. *Biol. J. Linn. Soc.* **113**: 604-622.
- Arntzen, J.W., Beukema, W., Galis, F., Ivanović, A. (2015): Vertebral number is highly evolvable in salamanders and newts (family Salamandridae) and variably associated with climatic parameters. *Contrib. Zool.* **84**: 85-113.
- Bandelt, H.J., Forster, P., Röhl, A. (1999): Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* **16**: 37-48.
- Boulenger, G.A. (1908): Description of a new newt. *The Annals and Magazine of Natural History, Series 8* **2**: 32-33.
- Chen, C., Durand, E., Forbes, F., François, O. (2007): Bayesian clustering algorithms ascertaining spatial population structure: a new computer program and a comparison study. *Mol. Ecol. Notes* **7**: 747-756.
- Cheng, L., Connor, T.R., Sirén, J., Aanensen, D.M., Corander, J. (2013): Hierarchical and spatially explicit clustering of DNA sequences with BAPS software. *Mol. Biol. Evol.* **30**: 1224-1228.
- Corander, J., Sirén, J., Arjas, E. (2008): Bayesian spatial modeling of genetic population structure. *Computation. Stat.* **23**: 111-129.
- Cvijanović, M., Ivanović, A., Kalezić, M.L. (2015): Larval pigmentation patterns of closely related newt species (*Triturus cristatus* and *T. dobrogicus*) in laboratory conditions. *North-West. J. Zool.* **11**: 357-359.
- Durand, E., Jay, F., Gaggiotti, O.E., François, O. (2009): Spatial inference of admixture proportions and secondary contact zones. *Mol. Biol. Evol.* **26**: 1963-1973.
- Gherghel, I., Papeş, M. (2015): Landscape as a determinant of dispersal patterns and population connectivity in a newt species. *Ecol. Inform.* **28**: 1-6.
- Goudet, J. (1995): FSTAT (Version 1.2): a computer program to calculate F-statistics. *J. Hered.* **86**: 485-486.
- Jakobsson, M., Rosenberg, N.A. (2007): CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**: 1801-1806.
- Kiritzescu, C. (1903): Contribution à la faune des Batraciens de Roumanie. *Buletinul Societatii de Stiinte din Bucuresti* **12**: 243-265.
- Lantz, L.A. (1953): Adaptation to background in *Triturus cristatus* Laur. *J. Genet.* **51**: 502-514.
- Litvinchuk, S., Sokolova, T.M., Borkin, L.J. (1994): Biochemical differentiation of the crested newt (*Triturus cristatus* group) in the territory of the former USSR. *Abhandlungen und Berichte für Naturkunde, Magdeburg* **17**: 67-74.
- Litvinchuk, S.N., Borkin, L.J., Džukić, G., Kalezić, M.L., Khalturin, M.D., Rosanov, J.M. (1999): Taxonomic status of *Triturus karelinii* on the Balkans, with some comments about other crested newt taxa. *Russ. J. Herpetol.* **6**: 153-163.
- Litvinchuk, S.N., Borkin, L.J. (2000): Intraspecific taxonomy and nomenclature of the Danube crested newt, *Triturus dobrogicus*. *Amphibia-Reptilia* **21**: 419-430.
- Litvinchuk, S.N. (2005): A record of the Danube newt, *Triturus dobrogicus*, from the Dnepr River delta (Ukraine). *Russ. J. Herpetol.* **12**: 69-72.

- Litvinchuk, S.N., Borkin, L.J. (2009): Evolution, systematics and distribution of crested newts (*Triturus cristatus* complex) in Russia and adjacent countries St. Petersburg, Evropeisky Dom Press.
- Macgregor, H., Horner, H. (1980): Heteromorphism for chromosome 1, a requirement for normal development in crested newts. *Chromosoma* **76**: 111-122.
- Matsui, K., Mochida, K., Nakamura, M. (2003): Food habit of the juvenile of the Japanese newt *Cynops pyrrhogaster*. *Zoolog. Sci.* **20**: 855-859.
- Mayr, E. (1969): Principles of Systematic Zoology New York, McGraw-Hill, Inc.
- Michaels, C.J., Antwis, R.E., Preziosi, R.F. (2015): Impacts of UVB provision and dietary calcium content on serum vitamin D₃, growth rates, skeletal structure and coloration in captive oriental fire-bellied toads (*Bombina orientalis*). *J. Anim. Physiol. Anim. Nutr.* **99**: 391-403.
- Naumov, B., Biserkov, V. (2013): On the distribution and subspecies affiliation of *Triturus dobrogicus* (Amphibia: Salamandridae) in Bulgaria. *Acta Zool. Bulg.* **65**: 307-313.
- Raffaëlli, J. (2007): Les urodèles du monde Condé-sur-Noireau, Penclen édition.
- Ridley, M. (2004): Evolution. third Edition. Cambridge Massachusetts, Blackwell Scientific Publications.
- Rosenberg, N.A. (2004): DISTRUCT: a program for the graphical display of population structure. *Mol. Ecol. Notes* **4**: 137-138.
- Slijepčević, M., Galis, F., Arntzen, J.W., Ivanović, A. (2015): Homeotic transformations and number changes in the vertebral column of *Triturus* newts. *PeerJ* **3**: e1397.
- Sparreboom, M. (2014): Salamanders of the Old World Zeist, The Netherlands, KNNV Publishing.
- Takezaki, N., Nei, M., Tamura, K. (2010): POPTREE2: software for constructing population trees from allele frequency data and computing other population statistics with Windows interface. *Mol. Biol. Evol.* **27**: 747-752.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S. (2013): MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Mol. Biol. Evol.* **30**: 2725-2729.
- Thiesmeier, B., Kupfer, A., Jehle, R. (2009): Der Kammmolch Bielefeld, Laurenti-Verlag.
- Vörös, J., Arntzen, J.W. (2010): Weak population structuring in the Danube crested newt, *Triturus dobrogicus*, inferred from allozymes. *Amphibia-Reptilia* **31**: 339-346.
- Wallis, G.P., Arntzen, J.W. (1989): Mitochondrial-DNA variation in the crested newt superspecies: limited cytoplasmic gene flow among species. *Evolution* **43**: 88-104.
- Wielstra, B., Crnobrnja-Isailović, J., Litvinchuk, S.N., Reijnen, B.T., Skidmore, A.K., Sotiropoulis, K., Toxopeus, A.G., Tzankov, N., Vukov, T., Arntzen, J.W. (2013a): Tracing glacial refugia of *Triturus* newts based on mitochondrial DNA phylogeography and species distribution modeling. *Front. Zool.* **10**: 13.
- Wielstra, B., Litvinchuk, S.N., Naumov, B., Tzankov, N., Arntzen, J.W. (2013b): A revised taxonomy of crested newts in the *Triturus karelinii* group (Amphibia: Caudata: Salamandridae), with the description of a new species. *Zootaxa* **3682**: 441-453.
- Wielstra, B., Duijm, E., Lagler, P., Lammers, Y., Meilink, W.R.M., Ziermann, J.M., Arntzen, J.W. (2014a): Parallel tagged amplicon sequencing of transcriptome-based genetic markers for *Triturus* newts with the Ion Torrent next-generation sequencing platform. *Mol. Ecol. Resour.* **14**: 1080-1089.
- Wielstra, B., Sillero, N., Vörös, J., Arntzen, J.W. (2014b): The distribution of the crested and marbled newt species (Amphibia: Salamandridae: *Triturus*) – an addition to the New Atlas of Amphibians and Reptiles of Europe. *Amphibia-Reptilia* **35**: 376-381.

Tables

Table 1. Sampled *Triturus dobrogicus* populations. Population numbers correspond to Fig. 1 and more details can be found in Table S2.

Figure legends

Fig. 1. Sampling scheme and population tree for *T. dobrogicus*. (a) The distribution is outlined with a black line (based on Wielstra et al., 2014b). Populations are from four range sections: the Pannonian (yellow in the online version of this figure) and Lower Danube (green online) Plains and the Danube (blue online) and Dnepr (red online) Deltas. The Pannonian and Lower Danube Plain are separated by the Iron Gate and the Lower Danube Plain and the Danube Delta are bounded by populations 41 and 45 following Litvinchuk and Borkin (2000). Grey background shading reflects elevation above 150 meters above sea level. The Danube River is shown as a dark line (blue in the online version) running through the center of the range. Population numbers correspond to Table 1 and Table S2. (b) Genetic differentiation of populations based on nuclear DNA. In the online version populations are color coded according to the region of origin. Only bootstrap support values over 50 are shown. Populations with a thick line in (a) and in boldface in (b) are consistently identified as predominantly belonging to a relatively distinct genetic cluster in the intraspecific spatial Bayesian clustering analyses (see Fig. 2).

Fig. 2. Spatial Bayesian clustering results for *Triturus dobrogicus* individuals according to the programs BAPS and TESS. Numbers at the top are population numbers and in the online version colors reflect regions.

Fig. 3. Haplotype network for *Triturus dobrogicus* mtDNA. Only individuals identified as pure *T. dobrogicus* are included and one mtDNA haplotype introgressed from another crested newt species is excluded. Pies (or pie slices) are color coded according to the region of origin of haplotypes: the Pannonian (white in print and yellow in the online version) and Lower Danube (light grey in print, green online) Plains and the Danube (dark grey in print, blue online) and Dnepr (black in print, red online) Deltas. Numbers refer to haplotype code (with the suffix “Tdob” not shown) and correspond to Table S3.

Fig. 4. Body shape measurements for *Triturus dobrogicus*. Shown from left to right are the average, range and standard deviation for (a) the Wolterstorff-index (forelimb length divided by interlimb distance) and (b) Ltc/L (head width divided by body length) for females and males from the Pannonian (yellow in the online version) and Lower Danube (green online) Plains and the Danube (blue online) and Dnepr (red online) Deltas. Note that for females there are no data from the Dnepr Delta. Sample sizes are noted in italics below box plots and raw data are in Table S4. The thick grey lines show the optimal cut-off value as determined with weighted logistic regression, for the Wolterstorff-index – males ($1/(1+\exp(0.161*WI-8.375))$), for Ltc/L – females ($1/(1+\exp(0.811*HI-10.475))$) and Ltc/L – males ($1/(1+\exp(0.915*HI-12.669))$). For Wolterstorff-index – females no significant model was found.

Online appendices

Table S1. Marker details for intraspecific analysis.

Table S2. Sampling details for *Triturus dobrogicus*.

Table S3. GenBank Accession numbers for mtDNA.

Table S4. Morphological data for *Triturus dobrogicus*.