

This is a repository copy of *Hyperglycaemia and risk of adverse perinatal outcomes:A systematic review and meta-analysis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/104564/>

Version: Accepted Version

---

**Article:**

Farrar, Diane, Simmonds, Mark Crawford [orcid.org/0000-0002-1999-8515](https://orcid.org/0000-0002-1999-8515), Bryant, Maria [orcid.org/0000-0001-7690-4098](https://orcid.org/0000-0001-7690-4098) et al. (5 more authors) (2016) Hyperglycaemia and risk of adverse perinatal outcomes:A systematic review and meta-analysis. *BMJ*. i4694. pp. 1-11. ISSN 1756-1833

<https://doi.org/10.1136/bmj.i4694>

---


**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.


**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.


## Combining 75g and 100g OGTT tests – Fasting



## 50g OGCT



## Two-hour postload 75g OGTT



## Two-hour postload 100g OGTT

