

This is a repository copy of *Untangling complex organic mixture in prehistoric hearths*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/107554/>

Version: Accepted Version

Article:

Lucquin, Alexandre Jules Andre orcid.org/0000-0003-4892-6323 (2016) Untangling complex organic mixture in prehistoric hearths. *Proceedings of the National Academy of Sciences of the United States of America*. pp. 10456-10457. ISSN: 1091-6490

<https://doi.org/10.1073/pnas.1612429113>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Lucquin, A. (2016). Untangling complex organic mixture in prehistoric hearths. *Proceedings of the National Academy of Sciences of the United States of America*, 113(38), 10456–10457.

doi: 10.1073/pnas.1612429113

Untangling complex organic mixture in Prehistoric hearths

Alexandre Lucquin¹

¹ BioArCh, University of York, Environment Building, Wentworth Way, York YO10 5DD, UK.

In the last decades organic residue analysis of ceramic artefacts have advanced our understanding of culinary practices and economies in the past. Current approaches, deploying lipid residue analysis, have allowed a wide range of prehistoric commodities processed in pottery to be identified (1, 2). Recently, organic residues analysis has allowed us to gain a new perspective on the emergence and adoption of pottery by prehistoric hunter-gatherers. These data have shown a the strong reliance on the exploitation of aquatic resources (e.g. 3, 4). But this approach is limited by the fact that not all hunter-gatherer societies used pottery and that usage of pots seems specialised in comparison to the existing broader spectrum economies (4). Thus, looking at organic residues contained in fire structure remains would provide an extended testimony of diet and cooking practices complementary to the study of vegetal and faunal remains. Although organic residue analysis of prehistoric hearths began in the late eighties (5–7) they are still an underestimated source of information and only a few studies have been exploring their potential. In this issue of PNAS, Choy et al (8) uses isotopic and molecular analysis of organic residues from hearths to obtain new insights into salmonid exploitation and processing by Alaskan hunter-gatherer of the end of the Pleistocene.

Choy et al.'s paper (8) main objective was to resolve the complex mixture of organic matter (OM) found within hearth and to obtain quantifiable data of the contribution of the various sources considered (regrouped as salmon, freshwater and terrestrial animal) to allow a comparison of the content of the fire structures from the USR occupations. This is a relevant issue as OM preserved in hearths may unlock archaeological information, although complex to deconvolute. Indeed, preserved OM is a mixture resulting from the potential input of: sedimentary OM initially present or coming from post-depositional process (e.g. deposition and migration of sediment), plant or animal tissues used as fuel (e.g. wood, bones, dung) or processed during cooking activities (e.g. roasting or grilling of animal resources) as well as other domestic activities. Moreover, the OM have been subject to a natural oxidation and to heat, conducting to a partial or total removal of compounds and to molecular transformations.

Choy et al. have followed analytical procedures previously employed . OM study in hearths have focused on the molecular characterisation of lipids from archaeological or experimental replicate using gas chromatography mass spectrometry (GC-MS) (e.g. 6, 9–12), associated more recently with stable isotopic analysis of bulk organic matter (EA-IRMS) (13) or of individual molecular compound obtained by gas chromatography combined with isotope ratio mass spectrometry (GC-c-IRMS) (14, 15). By combining these biomolecular techniques, Choy et al. addressed the mixing issues .

The presence of anthropogenic OM is demonstrated by an higher $\delta^{15}\text{N}$ compared to the control samples. The $\delta^{15}\text{N}$ values also indicate a contribution of aquatic resources and the variability observed reflects the diversity of the composition of the OM. As the bulk $\delta^{13}\text{C}$ are possibly reflecting an input of wood fuel and as no specific biomarker have been recovered, further insight are obtained with the compound specific $\delta^{13}\text{C}$ value of individual fatty acids. Finally the composition of the mixture is resolved using SIAR, a Bayesian isotopic mixing model, allowing to calculate the relative contribution of the various sources (grouped as salmon, freshwater and terrestrial animal). Importantly the good bone preservation of the Component 3 of USR, allows Choy et al. to validate the result by obtaining a positive correlation between the calculated contribution of salmonids by SIAR and the frequency index of salmonids in the faunal assemblages. Thus Choy et al. can distinguish certain patterns of activity within Component 3 but also can infer the exploitation of anadromous salmon as early as in Component 2, showing that hunter-gatherers living in Alaska relied heavily for their diet on aquatic resources, and notably anadromous fishes, since the end of Pleistocene.

Control of fire is one of the exclusive and typical traits of humanity, as reflected in numerous myths where fire is the element that allows humanity to “get out” of the natural kingdom (16). But the fire in question is not any fire to heat or light, it is the fire that allows cooking (17), transforming a natural resource in cultural food. Once adopted, fire was never abandoned and as a consequence, there is no society that doesn't process by heat at least a part of its food (18, 19). Organic residues analysis of archaeological hearths provides a direct testimony on the evolution of diet and subsistence practices in human societies.

Reference

1. Regert M (2011) Analytical strategies for discriminating archeological fatty substances from animal origin. *Mass Spectrom Rev* 30(2):177–220.
2. Evershed RP (2008) ORGANIC RESIDUE ANALYSIS IN ARCHAEOLOGY: THE ARCHAEOLOGICAL BIOMARKER REVOLUTION*. *Archaeometry* 50(6):895–924.
3. Craig OE, et al. (2013) Earliest evidence for the use of pottery. *Nature* 496(7445):351–354.
4. Lucquin A, et al. (2016) Ancient lipids document continuity in the use of early hunter–gatherer pottery through 9,000 years of Japanese prehistory. *Proceedings of the National Academy of Sciences*. doi:10.1073/pnas.1522908113.
5. Rottländer RCA (1989) Chemische untersuchungen an sedimenten der Höhle

Geissenklösterle bei Blaubeuren. *Fundberichte Aus Baden-Württemberg* 14:23–32.

6. March RJ, Baldessari A, Gross EG (1989) Determinacion de compuestos orgánicos en estructuras de combustión arqueológicas. *Nature et Fonction Des Foyers Préhistoriques, Actes Du Colloque International de Nemours (1987)*, eds Olive M, Taborin Y (APRAIF, Nemours), pp 47–58.
7. Cliquet D, et al. (1989) Approches d'une étude comparative des matières organiques de foyers préhistoriques et de foyers expérimentaux : l'exemple du gisement moustérien de Saint-Germain-des-Vaux, Manche. *Nature et Fonction Des Foyers Préhistoriques*, Mémoires du Musée de préhistoire d'Ile de France., eds Olive M, Taborin Y (APRAIF, Nemours), pp 29–36.
8. Choy K, et al. Chemical profiling of ancient hearths reveals recurrent salmon use in Ice Age Beringia. *Proc Natl Acad Sci U S A*.
9. March RJ (1999) Chimie organique appliquée à l'étude des structures de combustion du site de Tunel I (Terre de Feu, Argentine). *Revue d'archéométrie* 23:127–156.
10. Kedrowski BL, et al. (2009) GC/MS ANALYSIS OF FATTY ACIDS FROM ANCIENT HEARTH RESIDUES AT THE SWAN POINT ARCHAEOLOGICAL SITE*. *Archaeometry* 51(1):110–122.
11. March RJ, Dumarçay G, Lucquin A, Joly D (2006) Les activités liées à l'utilisation du feu: Un dernier hiver à Pincevent : les Magdaléniens du niveau IV0 (Pincevent, La Grande Paroisse, Seine-et-Marne). *Gallia Préhistoire* 48(1):89–108.
12. Lejay M, Alexis M, Quénéa K, Sellami F, Bon F (2016/9) Organic signatures of fireplaces: Experimental references for archaeological interpretations. *Org Geochem* 99:67–77.
13. Heron C, Nilsen G, Stern B, Craig O, Nordby C (2010) Application of lipid biomarker analysis to evaluate the function of "slab-lined pits" in Arctic Norway. *J Archaeol Sci* 37(9):2188–2197.
14. Buonasera TY, Tremayne AH, Darwent CM, Eerkens JW, Mason OK (2015) Lipid biomarkers and compound specific $\delta^{13}\text{C}$ analysis indicate early development of a dual-economic system for the Arctic Small Tool tradition in northern Alaska. *J Archaeol Sci* 61:129–138.
15. March RJ (2013) Searching for the Functions of Fire Structures in Eynan (Mallaha) and their Formation Processes: a Geochemical Approach. *Natufian Foragers in the Levant: Terminal Pleistocene Social Changes in Western Asia*, Archaeological Series., eds Bar-Yosef O, Valla FR (International Monographs in Prehistory, Ann Arbor), pp 227–283.
16. Frazer JG (1967) *Mythes sur l'origine du feu* (Payot, Paris).
17. Barrau J (1983) *Les hommes et leurs aliments : esquisse d'une histoire écologique et ethnologique de l'alimentation humaine* (Messidor/temps actuels, Paris).
18. Lévi-Strauss C (1968) *L'origine des manières de table* (Plon, Paris).
19. Goody J (1984) *Cuisines, cuisine et classes* (Paris).