



UNIVERSITY OF LEEDS

This is a repository copy of *Structure and lipid binding properties of the kindlin-3 pleckstrin homology domain*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/109597/>

Version: Accepted Version

---

**Article:**

Ni, T, Kalli, AC [orcid.org/0000-0001-7156-9403](http://orcid.org/0000-0001-7156-9403), Naughton, FB et al. (6 more authors)  
(2017) Structure and lipid binding properties of the kindlin-3 pleckstrin homology domain.  
Biochemical Journal, 474 (4). pp. 539-556. ISSN 0264-6021

<https://doi.org/10.1042/BCJ20160791>

---

© 2016 The Author(s). This is an author produced version of a paper published in Biochemical Journal. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## **Structure and lipid binding properties of the kindlin-3 pleckstrin homology domain**

Tao Ni<sup>1,\*</sup>, Antreas C. Kalli<sup>2,5,\*</sup>, Fiona B. Naughton<sup>2</sup>, Luke A. Yates<sup>1,4</sup>, Omar Naneh<sup>3</sup>, Mirijam Kozorog<sup>3</sup>, Gregor Anderluh<sup>3</sup>, Mark S. P. Sansom<sup>2,†</sup> and Robert J. C. Gilbert<sup>1,†</sup>

<sup>1</sup>Division of Structural Biology, Wellcome Trust Centre for Human Genetics, University of Oxford, Roosevelt Drive, Oxford, OX3 7BN, UK.

<sup>2</sup>Department of Biochemistry, University of Oxford, South Parks Road, Oxford, OX1 3QU, UK.

<sup>3</sup> Department for Molecular Biology and Nanobiotechnology, National Institute of Chemistry, Hajdrihova 19, 1000 Ljubljana, Slovenia.

<sup>4</sup> Current address: Section of Structural Biology, Department of Medicine, Imperial College London, Sir Alexander Fleming Building, South Kensington, London, SW7 2AZ, UK

<sup>5</sup> Current address: Leeds Institute of Cancer and Pathology, School of Medicine, St James' University hospital, University of Leeds, Leeds, LS9 7TF, UK

\*These authors contributed equally to this work.

†For correspondence: [mark.sansom@bioch.ox.ac.uk](mailto:mark.sansom@bioch.ox.ac.uk) and [gilbert@strubi.ox.ac.uk](mailto:gilbert@strubi.ox.ac.uk)

## *Abstract*

Kindlins co-activate integrins alongside talin. They possess, like talin, a FERM domain comprising F0-F3 subdomains, but with a pleckstrin homology (PH) domain inserted in the F2 subdomain that enables membrane association. We present the crystal structure of murine kindlin-3 PH domain determined at 2.23Å resolution and characterise its lipid binding using biophysical and computational approaches. Molecular dynamics (MD) simulations suggest flexibility in the PH domain loops connecting  $\beta$ -strands forming the putative phosphatidylinositol phosphate (PtdInsP) binding site. Simulations with PtdInsP-containing bilayers reveal that the PH domain associates with PtdInsP molecules mainly via the positively charged surface presented by the  $\beta$ 1- $\beta$ 2 loop and that it binds with somewhat higher affinity to PtdIns(3,4,5)P<sub>3</sub> compared to PtdIns(4,5)P<sub>2</sub>. Surface plasmon resonance (SPR) with lipid headgroups immobilised and the PH domain as analyte indicate affinities of 300  $\mu$ M for PtdIns(3,4,5)P<sub>3</sub> and 1mM for PtdIns(4,5)P<sub>2</sub>. In contrast, SPR studies with immobilised PH domain and lipid nanodiscs as analyte show affinities of 0.40  $\mu$ M for PtdIns(3,4,5)P<sub>3</sub> and no affinity for PtdIns(4,5)P<sub>2</sub> when the inositol phosphate constitutes 5% of the total lipids (~5 molecules per nanodisc). Reducing the PtdIns(3,4,5)P<sub>3</sub> composition to 1% abolishes nanodisc binding to the PH domain, as does site-directed mutagenesis of two lysines within the  $\beta$ 1- $\beta$ 2 loop. Binding of PtdIns(3,4,5)P<sub>3</sub> by a canonical PH domain, Grp1, is not similarly influenced by SPR experimental design. These data suggest a role for PtdIns(3,4,5)P<sub>3</sub> clustering in the binding of some PH domains and not others, highlighting the importance lipid mobility and clustering for the biophysical assessment of protein-membrane interactions.

**Short title:** Kindlin-3 PH domain binding of clustered inositol phosphates

**Keywords:** PH domain; lipid clustering; X-ray crystallography; molecular dynamics; surface plasmon resonance

**Abbreviations:**

CG-MD: Coarse-grained molecular dynamics

EMT: epithelial-to-mesenchymal transition

FERM domain: 4.1-erythrin-radixin-moesin domain

LAD-III: Leukocyte adhesion deficiency type III

MD: molecular dynamics

PH: pleckstrin homology

PMF: Potential of mean force

PtdInsP/PIP: Phosphatidyl inositol phosphate

SPR: surface plasmon resonance

## Introduction

Integrin-mediated adhesion between cells and the extracellular matrix is essential for the development of multicellular organisms. Integrins are heterodimeric ( $\alpha$  and  $\beta$ ) transmembrane receptors comprising a large extracellular domain for ligand binding (such as fibronectin, collagen and vitronectin, *etc.*) and short cytoplasmic tails [1,2], and are responsible for bi-directional signal transduction across the cell membrane. The cytoplasmic tail of the  $\beta$  subunit, despite its short length, is critical for integrin activation and has been shown to interact with several regulatory proteins. Talin was the first essential intracellular activator of integrins to be identified, and it contains an amino-terminal FERM domain complemented with a long rod-like domain at its carboxy-terminal end [3,4]. The later identification of the kindlin family of proteins identified them as key co-activators of integrins, alongside talin. Talin and kindlin interact directly with the cytoplasmic tails of integrin  $\beta$  subunits, with kindlin binding to a membrane-distal NPxY motif and talin a membrane-proximal NPxY motif [5-8]. Biophysical data indicate that talin and kindlins are able to bind simultaneously to their respective NPxY sites [5] while kindlins may be earlier recruited to integrins during activation than talin is [9].

In humans, three isoforms of kindlin proteins, kindlins -1, -2 and -3, have been identified and shown to adopt tissue-specific patterns of expression. Kindlin-1 is expressed in the epithelial cells of the skin and gut, kindlin-2 is widely expressed but most notably in striated and smooth muscle, and kindlin-3 is expressed particularly in hematopoietic tissues but also in endothelial cells [10,11]. Abnormal expression of, or mutations in, kindlins can result in severe disease. For example, kindlin-1 mutations are associated with Kindler syndrome, a rare genetic dermatitis [12], while aberrant expression of kindlin-2 is associated with

oncogenesis, especially on relocation to the nucleus where it appears to have a role in the control of transcriptional activity leading to the loss of tumour-suppressor signals [13,14].

By contrast, a nuclear role has not been shown for kindlin-3 but it is over-expressed in chronic lymphocytic leukaemia (which led to its discovery) [15] and when over-expressed in breast cancer may play a key angiogenetic role [16]. The importance of kindlin-3 in blood clot formation is however clear – for example mice carrying a QW>AA substitution in the integrin-binding F3 subdomain have compromised clotting ability [17] because kindlin-3 activity is essential for platelet activation [18]. As detailed above, nuclear localisation of kindlin-2 is oncogenic [13,14,19] and another recent intriguing insight is that kindlin-2 is the preferred binding partner of  $\beta_1$  integrins over kindlin-3 [20]. This suggests one possible mechanism linking the two is that over-expression of kindlin-3 could dislodge kindlin-2 from an integrin-bound state, leading to its nuclear relocalisation.

In any case, membrane binding by kindlins is thus not only an important factor in integrin activation but may also determine their cell signalling impact, and even the cell's fate. The membrane binding capacity of kindlins has been proved to be essential for integrin activation and is dependent on their PH domains [4,21-23] (see Figure S1 for a sequence alignment) and on a long loop within the F1 FERM subdomain which is conserved and similar to an equivalent loop in talin [24]. Deletion of the kindlin-3 PH domain eliminates its ability to participate in the adhesion and migration of B cells mediated by the leukocyte integrin LFA-1 [25].

Our previous work on the kindlin PH domains focused on kindlin-1, where we solved its crystal structure to reveal an isoform-specific salt bridge occluding the canonical inositol phosphate binding site [23]. Molecular dynamics indicated that the salt bridge is dynamic and led us to make a mutant lacking it which had altered inositol phosphate binding properties. We showed however that in either case the affinity for inositol phosphate ligands is relatively weak ( $10^{-4}$  M), although with a specific preference for PtdIns(3,4,5)P<sub>3</sub>. We also showed that the apparent affinity was influenced by the buffer choice for an experiment: phosphate-based buffers seem to interfere with the apparent affinity measured, suggesting a lack of specificity in the PtdIns(3,4,5)P<sub>3</sub> interactions taking place [23].

Here we report the crystal structure of the kindlin-3 PH domain together with molecular dynamics-based and biophysical characterisation of its inositol phosphate lipid binding properties. We show that the kindlin-3 PH domain has a hydrogen bond-based occlusion to its (canonical) PtdInsP binding cleft, and that like the kindlin-1 PH domain it binds surface-immobilized inositol phosphates with rather low ( $10^{-4}$  M) affinity. Using lipid nanodiscs as a model membrane system and molecular dynamics simulations to study the interaction of the kindlin-3 PH domain with different combinations of lipid species including inositol phosphate lipids, we show that lipid clustering is likely to be a significant factor in its binding of PtdIns(3,4,5)P<sub>3</sub> in bilayer membranes. On this basis we propose that a subset of PH domains is able to bind to multiple inositol phosphates simultaneously and so via an avidity effect have its interaction with target membranes strengthened. This would explain how an apparently low affinity for inositol phosphates can be biologically relevant to the localisation of kindlins to the plasma membrane and their subsequent activation of integrins.

## Results

### *Overall structure of Kindlin-3 PH domain.*

The crystal structure of the kindlin-3 PH domain exhibits an archetypal pleckstrin homology (PH) superfamily fold at its core [26], with a partly open seven-stranded  $\beta$ -barrel capped at one end by a C-terminal  $\alpha$ -helix (Figure 1; see Table 1 for data collection and refinement statistics). Similar to that of kindlin-1 and 2 [22,23], the PH domain of kindlin-3 contains an additional C-terminal amphipathic helix extension, which is not present in other PH domains (Figure 1A). A salt bridge in kindlin-1 PH domain (Arg<sup>380</sup>-Glu<sup>416</sup>) has been shown to occlude its canonical phosphoinositol binding site [23], and in the kindlin-3 PH domain this salt bridge is replaced by a hydrogen bond (Arg<sup>360</sup>-Gln<sup>396</sup>) in the equivalent position (Figure 1A). As shown in Figure 1B, the kindlin-3 PH domain has a mixed surface charge distribution with a pronounced concentration of positive charge at the canonical inositol phosphate binding site.

The overall fold of the kindlin PH domain appears remarkably conserved. The kindlin-3 PH domain is closest in structure to the PH domain of kindlin-2 (RMSD = 0.69Å), with the major conformational difference being in the  $\beta$ 1- $\beta$ 2 loop. The close relationship between the kindlin PH domain structures is reflected in their position within a structural phylogenetic tree we constructed for a set of PH domains of different kinds of function and (where relevant) with different lipid binding properties (Figure S2). Of note, a previously-constructed structural phylogeny places the previously-determined kindlin-3 PH domain NMR structure in a different relationship to that of kindlins -1 and -2 to the crystal structure reported here [23]. This highlights the sensitivity of structural phylogenetic comparisons to the quality of the models used.



PH domains with canonical inositol phosphate binding sites typically exhibit a highly positively-charged core at the open end of the PH domain, whereby they interact with their ligand, as seen in PH domains of Grp1 [27] and PLC $\delta$ 1 [28]. Non-canonical PH domains, such as Slm1 [29], Tiam1 and ArhGAP9 [30], lack the same kind of positive charge distribution, and instead have their inositol phosphate binding site on the side of the barrel mainly between the  $\beta$ 1- $\beta$ 2 loop and  $\beta$ 5- $\beta$ 6 loop (Figure S3). The PH domains of kindlin-3 and kindlin-2 have similar surface charge distribution to the PH domains with canonical inositol phosphate binding sites (Figure 1B), indicating that inositol phosphate might interact in a canonical way through the  $\beta$ 1- $\beta$ 2 loop. The previously-reported crystal structure of the kindlin-1 PH domain is not as complete as either of the available crystallographic kindlin-2 or kindlin-3 models, lacking seven residues within the  $\beta$ 1- $\beta$ 2 loop; its surface charge characteristics however seem to be similar to those of the other kindlin PH domains.

Atomistic molecular dynamics simulation of the kindlin-3 PH crystal structure in solution (i.e. in the absence of any lipid bilayer) with two different force-fields showed that the overall fold of the PH domain is retained during the simulations. The seven-stranded  $\beta$ -barrel in the core of the PH domain is very stable, whereas the unstructured regions connecting the  $\beta$ -strands are somewhat more flexible (Figure 1C). In particular, the  $\beta$ 1- $\beta$ 2 loop is flexible with both simulation force-fields, in good agreement with the crystal structure (Figure 1A). At the end of the simulation the side chains of the positive residues on the  $\beta$ 1- $\beta$ 2 loop remain exposed towards the open end of the barrel, presumably providing a site for inositol phosphate binding. With the exception of one simulation in which the Arg<sup>360</sup>-Gln<sup>396</sup> hydrogen bond identified in the crystal structure breaks for an extended period of time, the Arg<sup>360</sup>-

Gln<sup>396</sup> hydrogen bond was generally preserved during the simulations although it breaks transiently before reforming (Figure S4).

*Molecular dynamics simulations reveal a membrane-bound state of kindlin-3 PH domain and its interaction with PIP lipids.*

Coarse-grained molecular dynamics (CG-MD) simulations can be used to identify the molecular mechanism of peripheral membrane proteins binding to model membranes incorporated with anionic e.g. PIP lipids [31,32]. In particular, MD simulations have been used to study the interaction of a number of PH domains e.g. DAPP1 PH [32] and Grp1 PH [33] with PIP lipids at the molecular level. A recent study has also demonstrated that MD simulations can be used to study the free energy of the interaction of Grp1 PH domain with PIP<sub>2</sub> and PIP<sub>3</sub> molecules [34].

To examine the association of the new crystal structure of the kindlin-3 PH domain with model membranes, the PH domain was displaced away from a preformed bilayer containing 4 PIP<sub>3</sub> molecules in each leaflet (concentration of ~1.5 % that mimics the *in vivo* concentration of PIP<sub>3</sub> lipids in the plasma membrane) and an ensemble of 20 CG-MD simulations of 1.5  $\mu$ s each was performed (Figure 2A). Calculation of the orientation of the PH domain relative to the bilayer (by calculating the R<sub>zz</sub> component of its rotational matrix) suggests that the PH domain has a preferred orientation relative to the bilayer (Figure 2B). In this orientation residues 360 to 372 (i.e. the positively charged loop between strands  $\beta$ 1/ $\beta$ 2) made the largest number of contacts with the PIP<sub>3</sub> molecules. Regions 385-386, 410-415 and 434-436 also face towards the bilayer (Figure 2) and interact with PIP<sub>3</sub> molecules. This orientation is rather

different from e.g. the canonical orientation of the Grp1 PH domain relative to a bilayer [27,35]. Indeed, the helix at the end of the Grp1 PH domain was shown to be in a parallel orientation relative to the bilayer. We note, however, that in our ensemble we also observe secondary binding modes in which the kindlin-3 PH domain adopts an orientation that is similar to that of the Grp1 PH domain (Figure S5E). Atomistic simulations starting from the preferred orientation of the PH domain on the PIP<sub>3</sub>-containing bilayer reveal that the PH domain retains this orientation relative to the bilayer.

To examine whether the concentration of PIP<sub>3</sub> in a membrane bilayer affects the association of the kindlin-3 PH domain to the membrane, MD simulations were then carried out with various concentrations of PIP<sub>3</sub> lipids in the bilayer (7% PIP<sub>3</sub> or 2 PIP<sub>3</sub> in each leaflet or 1 PIP<sub>3</sub> in each leaflet; see Table 2). Despite the fact that when we reduce the number of PIP<sub>3</sub> molecules in the bilayer the kindlin-3 PH domain associates with the bilayer less frequently, in all cases it adopts a preferred orientation which is similar to the preferred orientation described above (Figure S6). The PH domain adopted the same orientation when we ran simulations with 7% PIP<sub>2</sub> lipids or with both PIP<sub>3</sub> and PIP<sub>2</sub> lipids in the bilayer (Kin3PH-1 and Kin3PH-6 in Table 2, respectively). Interestingly, atomistic simulations starting from four different snapshots of the kindlin-3 PH domain bound to PIP<sub>2</sub>-containing bilayers at the preferred orientation (6 simulations overall; see Methods) resulted in three simulations in which the kindlin-3 PH domain retained its orientation relative to the bilayer and in three simulations in which the PH domain adopted the secondary binding mode at the end of the atomistic simulations (Figure S5) as in subsequent potential of mean force (PMF) calculations (see below).

Analysis of the contacts between the PIP lipids and the PH domain during the simulations suggested that residues R362, K363 and K367 and residues K363, K367 and R370 made the largest number of contacts with PIP<sub>2</sub> and PIP<sub>3</sub> lipids, respectively. Our analysis also revealed that the PH domain induces clustering of the PIP lipids when it binds to the bilayer (Figure 2D). The clustering of PIP<sub>2</sub> lipids is somewhat higher compared to PIP<sub>3</sub> lipids. Note that the clustering of the PIP lipids occurs only in the leaflet in which the protein is bound. In the simulations with 7% of PIP<sub>2</sub> or 7% PIP<sub>3</sub> lipids in the bilayer most of the time we observe four PIP<sub>3</sub> and five PIP<sub>2</sub> lipids respectively in an annulus of 1 nm from the protein surface. Given that recent studies have suggested an important role for anionic lipids and PIPs in integrin activation [36,37], the change in the local lipid environment by the binding of the PH domain may have some functional role.

*Free energy of binding of the kindlin-3 PH domain to PIP(4,5)P<sub>2</sub> and PIP(3,4,5)P<sub>3</sub>:*

Our analysis above revealed the molecular mechanism of the association of the kindlin-3 PH domain with the membrane, suggesting the occurrence of two different modes of interaction of the PH domain with the PIP-containing bilayer. To examine the strength of interaction of the PH domain with PIP<sub>2</sub> and PIP<sub>3</sub> molecules we performed PMF calculations using a protocol we developed recently (see Methods and reference [34]). PMF profiles for PIP<sub>3</sub> and PIP<sub>2</sub> lipids have a global minimum with a well depth of ~3 kcal/mol for PIP<sub>3</sub> and a well of ~2.5 kcal/mol for PIP<sub>2</sub> lipids. This suggests that the protein associates somewhat more strongly with PIP<sub>3</sub> lipids (Figure 3). The first well (corresponding to the preferred orientation of the PH domain) in the profile of the PIP<sub>3</sub> molecules was followed by a second shallower well (corresponding to a secondary PH orientation) at a distance of ~2.5 nm. Examination of the interactions of the kindlin-3 PH domain with the lipids showed that in the windows covering

~1.5 to ~2.3 nm protein-lipid separations, the protein is in a similar orientation to the preferred orientation (mode 1) described above in which the protein interacts with the bilayer mainly via the  $\beta 1/\beta 2$  unstructured loop. In these windows the PIP<sub>3</sub> molecule is located between the  $\beta 1/\beta 2$  and  $\beta 5/\beta 6$  loops (i.e in a non-canonical PIP binding site) for most of the time (Figure S7A). We note that we have also seen interactions of PIP molecules with this site in our encounter simulations as described above. Interestingly, in the second well the protein adopts an orientation that is similar to the secondary binding mode also observed in the simulations above, with the PIP<sub>3</sub> molecule now located between the  $\beta 1/\beta 2$  and  $\beta 3/\beta 4$  loops (i.e in the canonical PIP binding site). In contrast, the profiles for PIP<sub>2</sub> lipids and for the mutated form of kindlin-3 PH domain did not have a second well. In the simulations with a PIP<sub>2</sub> molecule, the PIP<sub>2</sub> is found both in the canonical (i.e. contacting the  $\beta 1/\beta 2$  and  $\beta 3/\beta 4$  loops) and in a non-canonical site (i.e. contacting the  $\beta 1/\beta 2$  and  $\beta 5/\beta 6$  loops; Figure S7B). We note that the depth of the well for the binding of the Grp1 PH domain to PIP lipids using the same method was -5.3 kcal/mol for PIP<sub>3</sub> and -3.8 kcal/mol for PIP<sub>2</sub> suggesting that the binding of kindlin-3 PH domain is weaker compared to Grp1 PH domain binding a single lipid [34]. Disruption of the PH/PIP interactions by mutating the K363 and K367 residues which were identified to form a large number of contacts with the PIP lipids resulted in a reduced binding to PIP<sub>3</sub> molecules by -1 kcal/mol.

*An SPR study confirms that the Kindlin-3 PH domain binds to multiple Phosphatidylinositol Phosphates*

To verify the potential interaction between clustered phosphatidylinositol phosphates and the kindlin-3 PH domain observed in our MD simulations, we conducted surface plasmon resonance (SPR) experiments. Firstly, a biotinylated inositol phosphate was immobilized onto

a streptavidin-derivatized sensor chip, as in previous studies [23]. The use of streptavidin-mediated immobilisation meant that the distance separating individual lipids would be larger than the size of the tetrameric streptavidin, and therefore than the PH domain itself meaning that one PH domain would be able to bind no more than one lipid head group simultaneously, imposing one-to-one binding in the interaction (Figure 4A). In agreement with the MD simulation results, the kindlin-3 PH domain displays a preference for PIP<sub>3</sub>, though with a dissociation constant ( $K_D$ ) of about 300  $\mu$ M, in agreement with previous measurements [23]. The apparent binding capacity for PIP<sub>2</sub> was much lower ( $K_D = 1100 \mu$ M). Introduction of two point mutations (K363A, K367A) effectively abolished the binding capacity of both inositol phosphate moieties, indicating that the interaction is specific even if it is low affinity (Figure 4B). We note that in the MD simulation with the mutated form of the kindlin-3 PH domain, the protein is able to bind to the bilayer in a similar orientation to the wild type however in many simulations it quickly dissociates from the bilayer (Figure S8). A naturally-occurred IPRR (isoleucine-proline-arginine-arginine) insertion in the  $\beta$ 1- $\beta$ 2 loop of kindlin-3 has been reported to disable integrin activation and thus lymphocyte adhesion and migration [21]. In agreement with results obtained by others using a pulldown assay with IP<sub>3</sub> and IP<sub>4</sub>-coated beads, our SPR study *in vitro* revealed that the IPRR insertion directly disrupted the interaction between the PH domain and phosphoinositol moieties (Figures 4B and 4C), presumably because of the steric effect of an elongated  $\beta$ 1- $\beta$ 2 loop.

The observation in MD simulation that multiple phosphoinositol molecules may bind to more than one site on a single PH domain (Figure 2 and Figure 3) led us to attempt further verification of this mode of interaction experimentally. Nanodiscs have been shown to be an effective model membrane system due to their stability and capacity to mimic the membrane bilayer [38,39], including the fluidity of lipids though within a limited bilayer area. To study

the effect of phosphatidylinositol lipid concentrations on their binding by PH domains, we performed the SPR experiments in an inverted setup. The biotinylated PH domain was immobilised onto the sensor chip and the nanodiscs were prepared as the analyte.

Nanodiscs incorporated with different components were stable in solution (Figures S9A and S8D), with an estimated molecular weight of 120 kDa; they behaved as single species with a sedimentation coefficient of 4 S as assessed by analytical ultracentrifugation (Figure S9B). It appears that the diameter of the nanodiscs correlates with the molar ratio of phosphatidylinositides incorporated in their membranes: nanodiscs with 5% PtdIns(3,4,5)P<sub>3</sub> (diameter,  $d = 5.87$  nm) were bigger than those with 3% PtdIns(3,4,5)P<sub>3</sub> ( $d = 4.64$  nm) (Figures S9C and S9E). Note that the diameters are small because they are average values; as shown in Figure S9D the nanodiscs are elliptical and of the expected size.

The SPR results showed that 5% PIP<sub>3</sub>-incorporated nanodiscs bind to kindlin-3 PH domain strongly, with a much-enhanced affinity ( $K_D = 0.4$   $\mu$ M) (Figure 5) compared to that shown by the PH domain towards immobilised inositol phosphates. In contrast, nanodiscs with 5% PIP<sub>2</sub> showed no binding to kindlin-3 PH domain, demonstrating that PIP<sub>3</sub> is its real ligand and suggesting that the apparent binding to immobilised PIP<sub>2</sub> is due to a non-specific interaction. We also performed the same experiments with nanodiscs without incorporated phosphatidylinositol moieties and observed no binding in SPR experiments.

There is evidence for PIP<sub>2</sub> clustering within lipid bilayers [40] while other data suggest protein-driven PIP<sub>2</sub> clustering in signalling complex formation [41]. Furthermore, while a

study of yeast proteins has revealed that the interaction between one set of PH domains and membranes can be enhanced by cooperative interaction between PIPs and other lipids (such as POPS) [42], the interaction between the kindlin-3 PH domain and PIP<sub>3</sub> seems to more rely on the concentration of PIP<sub>3</sub> in the membrane, i.e., cooperativity among copies of the same lipid.

#### *Grp1 PH domain binds inositol phosphate with 1:1 specificity*

We further asked whether our observations are specific to kindlin-3 PH domain (and similar domains) or could also be shown for PH domains which are already known to bind to inositol phosphate species in a classical, 1:1 interaction mode. If they could be shown for such well-characterised domains, then our results with the kindlin-3 PH domain would seem questionable. The crystal structure of the Grp1 PH domain in complex with Ins(1,3,4,5)P<sub>4</sub> (equivalent to the PtdIns(3,4,5)P<sub>3</sub> headgroup) reveals the ligand well embedded in a pocket formed by positively-charged residues [43]. In line with this, we showed by SPR that the analyte/ligand design of our experiment does not significantly affect the apparent affinity of Grp1 for PIP<sub>3</sub> compared to the effect it has on the kindlin-3 PIP<sub>3</sub> binding observed. Grp1 PH domain was shown to bind to immobilised PIP<sub>3</sub> headgroups with a  $K_D$  of  $2.8 \pm 0.4 \mu\text{M}$  (Figure 6A), and to immobilised PIP<sub>2</sub> headgroups with a  $K_D$  ten times greater (Figure S10), consistent with a previous study though with a lower absolute value due to methodological differences [27,44]. In contrast, the binding affinity to PIP<sub>3</sub> within nanodiscs (5% of constituent lipids) has a  $K_D$  of  $0.58 \mu\text{M}$ . So, in the case of Grp1, there is only a 5 times difference in affinity between immobilised lipid headgroups and whole lipids in a lipid nanodisc (rather than the ~100x enhancement observed with the kindlin-3 PH domain). This suggests that the Grp1-PH interaction with PIP<sub>3</sub> at high affinity does not rely on the binding



of multiple PIP<sub>3</sub>, though it does indicate an enhancement of lipid binding – most likely by a simple electrostatic attraction – in a membrane-like context [42].

## **Discussion**

The cytoplasmic surface of the plasma membrane serves as a platform for many critical biological processes. The recruitment of peripheral membrane proteins from the cytoplasm to the plasma membrane surface is spatiotemporally and accurately regulated. The PH domain is one of a selection of commonly-used folds directing protein-membrane association and has been shown to confer specificity to different inositol phosphates on protein-membrane interactions.

PH domains have been shown to bind phosphatidylinositol phosphates (PtdInsPs) with different degrees of specificity [45] and the PH domains of the kindlins have a limited affinity for them [23], although the membrane interaction mediated by the PH domain is essential for integrin activation [23]. Phosphatidylinositol lipids are present in membranes at different concentrations, which may lead to variable activity and roles in membrane structure and membrane recognition. PtdIns(4,5)P<sub>2</sub> (PIP<sub>2</sub>) has been reported to be the most abundant phosphatidylinositol lipid in cells [46], and can undergo phosphorylation to a PtdIns(3,4,5)P<sub>3</sub> (PIP<sub>3</sub>) state; both PIP<sub>2</sub> and PIP<sub>3</sub> are found at the plasma membrane. Monophosphorylated inositide lipids are, by contrast, associated with compartments such as the Golgi and endoplasmic reticulum [46].

In the present study, we firstly solved the crystal structure of kindlin-3 PH domain, and then used the crystal structure as the basis for a multiscale molecular dynamics simulation study.

MD simulation revealed that kindlin-3 PH domain interacts with the PIP lipids primarily via the loop connecting the  $\beta 1$  and  $\beta 2$  strands and that it has a somewhat higher affinity for binding PtdIns(3,4,5)P<sub>3</sub> than for PtdIns(4,5)P<sub>2</sub> in a membrane-like system. Moreover, one kindlin-3 PH domain appears to bind multiple inositol phosphate lipids, at both canonical and non-canonical sites, by means of which the affinity of interaction between the PH domain and the membrane is greatly enhanced by an avidity effect. In contrast, with a PH domain (from Grp1) previously shown to have canonical and specific one-to-one binding to inositol phosphate lipid, this one-to-more binding mode was not observed. These observations from MD simulation were supported by our surface plasmon resonance experiments in two complementary setups. The first one involves immobilisation of biotinylated inositol phosphates on the sensor chip and use of the PH domain (WT and mutant) as the analyte, to mimic an isolated one-to-one binding mode. In this case, each PH domain could only bind to no more than one phospholipid due to the distance restraints imposed by streptavidin-mediated immobilisation. The other setup attempted to mimic the bilayer plasma membrane, by immobilising the PH domain on the chip, and using soluble membranes (nanodiscs) as the analyte. The affinity of interaction measured between the PH domain and inositol phosphate lipids were remarkably affected by experimental design.

Compared with liposomes, nanodiscs have obvious advantages for protein-lipid interaction studies in a membrane-like environment: they are more stable and have a limited number of lipids per disc. Nanodiscs made of MSP1E3D1 are estimated to accommodate about 125 POPC lipids, with a diameter of about 13 nm [47]; this is in agreement with our cryo-EM map of the nanodisc (Figure S9D) which is fitted well (length ~14nm, width ~9nm) with the elongated and twisted conformation of the human apolipoprotein A1 [48]. In the SPR experiments here, the small and homogeneous size of the nanodisc membrane is critical for

1:1 model fitting and interpretation of the SPR results. Liposomes, with inherently larger diameters (e.g., 50 - 100 nm), would create multiple possible contact sites for PH domains, confounding binding affinity analysis. Nanodiscs however are able to bind no more than one immobilised PH domain due their limited size compared with the size and density of immobilised PH domains on the chip surface.

We find that PtsIns(3,4,5)P<sub>3</sub> clustering enhances the affinity of kindlin PH domains for target membranes. This is a charge-based interaction as shown by its loss when positively-charged lysine residues are mutated to alanine (Figure 4). Similarly, a number of other proteins have been shown to interact with clusters of PtdInsP<sub>2</sub> lipids on the basis of opposite charge attraction. For example, a series of basic residues in the MARCKS peptide from a protein kinase C substrate are known to bind to PtdIns(4,5)P<sub>2</sub> lipid clusters, as do syntaxin-1A, BAR domains and N-WASP. N-WASP is an activator of Arp2/3 which in turn nucleates actin filament formation at membranes [49,50].

There is increasing evidence showing that PH domains can interact with multiple lipid species simultaneously. A recent crystal structure of the ASAP1 PH domain suggested that there were two anionic lipid binding sites on the ASAP1 PH domain (a canonical site and an atypical site) [51], and recent comparative MD simulations have suggested both sites are present for a number of PH domains [33]. Moreover, Vonkova *et al.* studied 91 different PH domains and showed that lipid cooperativity is essential for their recruitment to the membrane [42]. Clustering of lipids by the kindlin-3 PH domain is expected to change the local lipid environment. Given that kindlins act in synergy with talin, which was also shown to interact with anionic lipids [52], this may provide a mechanism for how talin and kindlins

in turn associate with the integrin. Focal adhesion kinase has also been shown to bind to inositol phosphates, inducing clustering of a protein-mediated kind [2,23].

The orientation of the kindlin-3 PH domain on a target membrane indicated by this study, and the fact that kindlins have a rather long unstructured loop in the F1 domain, allow us to form a hypothesis for how the kindlin head domain may interact with the membrane and with the integrin receptor. In particular, the orientation of the PH domain that we find, in combination with the short linker regions plugging it into the F2 subdomain, suggest that it acts as a buffer or spacer between the kindlin and the membrane surface, and thereby allows the F3 subdomain to interact with the membrane distal NPxY integrin  $\beta$ -subunit tail motif rather than the membrane proximal NPxY bound by talin [53] (Figure 7). Thus in combination with the N-terminal F0 domain [54] and long F1 loop [24] the PH domain provides kindlins with a specific location and binding mode for integrin activation.

There are no high resolution structural data for the rest of kindlin-3, however it has been shown to have a similar structure to the talin head domain [5]. Therefore, further studies will focus on structure determination for kindlins and use structural and biophysical methods to test the hypothesis outlined above that binding of clustered phosphatidylinositol phosphate lipids is a mechanism employed by some proteins and specifically a subset of PH domains to enable the binding of membranes with an enhanced affinity.

## **Materials and Methods:**

### *Constructs Design, Cloning and Mutagenesis*

Mouse *FERMT3* (UniProtKB: Q8K1B8) encoding kindlin-3 PH domain (residues 344 - 479) and Grp1 (General receptor of phosphoinositides 1, UniProtKB: O43739) PH domain (residues 264 - 392) were cloned into a pOPINJ vector, which contains a tandem His<sub>6</sub> and GST (Glutathione S-transferase) tag followed by a Human Rhinovirus (HRV)-3C protease cleavage site in the N-terminus. The successful cloning was confirmed by DNA sequencing (Source Bioscience, UK). Mutagenesis and insertions were achieved by overlapping PCR following the standard protocol [55]. A DNA fragment encoding a 15-residue AviTag<sup>TM</sup> was inserted between the 3C protease cleavage site and PH domain by PCR to enable *in vivo* protein biotinylation [56] using a BirA+ *E. coli* strain for surface plasmon resonance experiments.

### *Expression of kindlin-3 PH domain and Grp1 PH domain*

The construct encoding recombinant kindlin-3 PH domain was transformed into *E. coli* B834 strain. A single colony was picked from fresh transformants on an LB-agar-carbenicillin plate and cultured in 100ml of LB supplemented with 50 µg/ml of carbenicillin overnight. The pre-culture was diluted 50 times into LB media with antibiotics and incubated at 37° with vigorous shaking at 200 rpm until OD<sub>600</sub> reached about 0.6. The expression of protein was induced by 1mM IPTG (Isopropyl β-D-1-thiogalactopyranoside) at 18°C overnight. Cells were harvested by centrifugation at 6000 g for 20 min and pellets were frozen in a -80°C freezer until further use. Other proteins used in this study were expressed using the same protocol.

### *Protein purification and crystallisation*

The frozen cell pellet of kindlin-3 PH domain (about 10 g of cell pellet from 2 l of overnight culture) was resuspended in lysis buffer (50 mM Tris pH 7.5, 500 mM NaCl and 0.1% Tween-20) and lysed with sonication and the debris were separated by centrifugation at 40,000g for 1hr. The resulting supernatant was incubated with glutathione sepharose equilibrated with lysis buffer for 2 hrs. The glutathione sepharose with protein bound was then washed with lysis buffer extensively. GST-3C protease was added to cleave the kindlin3-PH domain from sepharose overnight at 4°C with gentle shaking. Supernatant containing PH domain was concentrated and applied to size exclusion chromatography (Superdex 75 16/60, GE Healthcare Life Sciences) in 10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.7 mM TCEP (*tris*(2-carboxyethyl)phosphine) buffer to obtain the purified proteins. The purification of kindlin-3 PH domain mutants and Grp1 PH domain and biotinylated proteins was carried out in the same way. Successful *in vivo* biotinylation was verified by efficient binding to streptavidin magnetic beads (Sigma-49532).

Crystallisation screening was performed by means of sitting drop vapour diffusion methods. Briefly, 100 nl of proteins at 20 mg/ml in 10 mM Tris-HCl pH 7.5, 150 mM NaCl were mixed with 100 nl of reservoirs and equilibrated against 95 µl of crystallisation reservoir (Hampton Research) at 293K. Crystals appeared within 4 days and continued growing until 2 weeks in 4 M Sodium Nitrate, 0.1M Sodium Acetate, pH 4.6. Crystals were flash-cooled with 25% (v/v) glycerol in the mother liquor as cryoprotectant.

*Data collection, processing and structure determination*

Diffraction data were collected using beamline I24 at Diamond Light Source (Didcot, UK) and then indexed, integrated and scaled using the Xia2 pipeline [57]. The structure of kindlin-3 PH domain was solved by molecular replacement (MR) using PHASER [58] with a polyalanine model of kindlin-1 PH domain (PDB: 4BBK). The resulting model from MR was further built by *Buccanneer* [59] and corrected manually in *Coot* [60]. Refinement was accomplished using *PHENIX.refine* together with manual correction in *Coot*. Molecular graphics were prepared with *PyMOL*. Attempts at co-crystallising kindlin-3 PH domain with Ins(1,3,4,5)P<sub>4</sub> and at soaking the crystal with highly concentrated Ins(1,3,4,5)P<sub>4</sub> resulted in a PH domain structure without ligand bound, presumably due to the low affinity interaction between isolated Ins(1,3,4,5)P<sub>4</sub> and the kindlin-3 PH domain.

#### *Nanodisc preparation*

POPC (1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine) and POPE (1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphoethanolamine) were purchased from Avanti Polar Lipids and PtdIns(4,5)P<sub>2</sub> (P-4516), PtdIns(3,4,5)P<sub>3</sub> (P-3924) and Ins(1,3,4,5)P<sub>4</sub> (Q-1345) were purchased from Echelon Biosciences. Nanodiscs were prepared following standard protocols [38]. Briefly, to prepare nanodiscs containing 5% (mol/mol) PtdIns(3,4,5)P<sub>3</sub>, a mixture of 200 µl of 10 mg/ml POPC, 200 µl of 10 mg/ml POPE together with 670 µl of 0.5mg/ml PtdIns(3,4,5)P<sub>3</sub> in chloroform were dried on a clean pyrex tube under argon. Residual chloroform was then removed by desiccation in a desiccator attached on a VARIO-SP diaphragm pump (Vacuubrand). The lipid film was subsequently hydrated by adding of 1 ml solubilisation buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 17 mM Na-cholate) followed by vigorous vortexing. The hydrated lipid film was then sonicated for 10 min in a water bath sonicator before 250 µl of MSP1E1 (7 mg/ml) proteins were added to the lipid mixture. Nanodiscs

were self-assembled upon removal of Na-cholate by dialyzing against 20 mM HEPES, pH 7.5, 150 mM NaCl buffer overnight. The aggregate was removed by size-exclusion chromatography (Superdex 200 10/300, GE Healthcare Life Sciences) and the peak fractions corresponding to size of monomeric nanodiscs species were collected (Figure S9A) and used immediately for biophysical characterisation and SPR experiments.

### *Dynamic Light Scattering*

Dynamic light scattering measurements were conducted on a Protein Solutions DynaPro instrument at 20 °C. The nanodisc eluates from size exclusion chromatography were diluted to approximately 0.05 mg/ml in 10 mM HEPES, pH 7.5, 150 mM NaCl before measurement in a cuvette.

### *Cryo-electron microscopy*

After plunge-freezing in liquid ethane, nanodiscs on holey carbon grids were imaged using an FEI Tecnai F30 microscope operating at 200kV and at a nominal magnification of 59,000x. Images were picked using *Boxer* [61] and reconstructed *ab initio* using *IMAGIC* [62] followed by *SPIDER* [63] software. The resolution of the map was determined by Fourier shell correlation at FSC=0.5 and was 23.5Å.

### *Surface Plasmon Resonance*

SPR experiments were performed using a Biacore T200 machine (GE Healthcare Life Sciences) at 20°C in 10 mM HEPES, pH 7.5, 150 mM NaCl. PtdIns(4,5)P<sub>2</sub> (C-45B6a) and PtdIns(3,4,5)P<sub>3</sub> (C-39B6a) were purchased from Echelon Biosciences. The BIACore CM5



chip (GE Healthcare Life Sciences) was firstly covered with streptavidin following the manufacturer's instructions before the biotinylated PIPs or biotinylated proteins were immobilised. The analyte with 2-fold serial dilutions was applied at a flow-rate of 20  $\mu\text{l}/\text{min}$  for 180 s followed by 600 s of dissociation time. The biosensor chip was regenerated after each sample injection cycle with different buffer in the two experiment setups: in the case of immobilised PIPs on the sensor chip, 0.1 % SDS was used; when the PH domain was immobilised then the surface was regenerated by 2 M  $\text{MgCl}_2$ . The data were fit with a 1:1 Langmuir adsorption model ( $B = B_{\text{max}}C/(K_D + C)$ , where  $B$  is the response of bound analyte and  $C$  is the concentration of the analyte in the sample) to calculate the dissociation constant ( $K_D$ ) using BIAanalysis software. The molecular weight of the nanodiscs for SPR analysis was estimated to be 120 kDa.

### *Molecular Dynamics Simulations*

#### Coarse-grained Molecular Dynamics (CG-MD) Simulations:

The CG-MD simulations were performed using GROMACS 4.5.5 [64] with the Martini 2.1 force field [65,66]. The simulation set up is shown in Figure 2. A summary of all simulations performed is shown in Table 2. Prior to the production simulations all systems were energy minimized and then equilibrated for 500 ns with the protein backbone particles restrained. For each repeat simulation within an ensemble, the protein was rotated around the  $x$ ,  $y$ , and  $z$  axes to form a different initial configuration. An ensemble of 20 simulations of 1.5  $\mu\text{s}$  each were run with a time step of 20 fs. An elastic network model (ENM) was applied to all backbone particles with a cut-off distance of 0.7 nm [67]. The LINCS algorithm was used to constrain to equilibrium bond lengths [68]. Lennard-Jones interactions were shifted to zero between 0.9 nm and 1.2 nm. Coulombic interactions were shifted to zero between 0 and 1.2

nm. The pressure was 1 bar and the temperature was 323 K. A Berendsen's algorithm [69] was used to control the pressure and the temperature with a coupling time of 1 ps.

#### Atomistic Molecular Dynamics (AT-MD) Simulations:

Conversion of CG to atomistic systems was made using a fragment-based approach [70]. The snapshot that was converted to an atomistic representation corresponded to the preferred orientation of the kindlin-3 PH domain in the density landscapes shown in SI Figure S5. The GROMOS96 43a1 force field [71] was used with SPC water molecules. The temperature was 323 K. The velocity rescaling method [72] was used to control the temperature with a coupling time of 0.1 ps. The pressure was 1 bar and it was controlled with semi-isotropic pressure coupling using the Parrinello-Rahman barostat [73] with a coupling time of 1 ps. Bond lengths were constrained to equilibrium lengths using the LINCS method. The time step was set at 2 fs. The particle mesh Ewald (PME) method was used to model the electrostatic interactions. For the atomistic simulation with the kindlin-3 PH domain in solution both the GROMOS96 43a1 and the OPLS-AA forcefield within GROMACS were used. For these simulations isotropic pressure coupling was used. The temperature was 310 K.

#### Potential of Mean Force (PMF) simulations:

Initial structures of wild type or mutant kindlin-3 PH domain bound to PIP<sub>3</sub> or PIP<sub>2</sub> in a 80:20 POPC:POPS bilayer were generated by alignment with an existing Grp1-PH/PIP<sub>3</sub> complex (see [34]). The system was converted to a coarse-grained representation. The PH domain was pulled from the bound PIP lipid along the membrane normal ( $z$ -axis) by fixing the centre of mass to a reference point which was moved at a rate of 0.001 nm ps<sup>-1</sup> for a total distance of 2.6 nm, while the PIP phosphate atom was restrained to its initial position. Snapshots with

protein-lipid separations (measured from the PH domain centre-of-mass to the phosphate atom of the PIP) in 0.1 nm intervals for the first 2 nm and 0.2 nm thereafter were extracted and used as initial structures for umbrella sampling simulations. In each simulation, the phosphate atom was again restrained to its initial position and the  $z$ -distance of the PH centre-of-mass was restrained relative to the phosphate atom with a force constant of  $1000 \text{ kJ mol}^{-1} \text{ nm}^{-2}$ . To reduce the time required for adequate sampling, the PH domain centre-of-mass was also restrained in the  $x$  and  $y$  directions, by a force constant of  $100 \text{ kJ mol}^{-1} \text{ nm}^{-1}$ . Each window was simulated for  $1 \mu\text{s}$  (mutant),  $1.4 \mu\text{s}$  (WT/PIP<sub>3</sub>) or  $1.5 \mu\text{s}$  (WT/PIP<sub>2</sub>), with the distance between PH centre-of-mass and PIP atom bead along the  $z$ -axis recorded every 20 ps. PMF profiles were calculated using the GROMACS *g\_wham* tool [74]. The time taken to reach convergence was judged from profiles generated for sequential time intervals; data before this time was discarded as equilibration before a final profile was calculated. Errors were obtained from bootstrap analysis. All profiles were shifted so the value of the PMF in bulk was  $0 \text{ kcal mol}^{-1}$ .

### **Acknowledgements**

T.N. was a Chinese Scholarships Council-University of Oxford scholar within the Nuffield Department of Medicine. We gratefully acknowledge Diamond Light Source beamline staff, and the support of Dr David Staunton with use of the Oxford Molecular Biophysics Suite within the university's Department of Biochemistry; the Division of Structural Biology is a part of the Wellcome Trust Centre for Human Genetics, Wellcome Trust Core Grant Number 090532/Z/09/Z. Electron microscopy was performed using the Wellcome Trust-supported microscopy facility within the Division of Structural Biology established with a JIF award (060208/Z/00/Z) and further supported by a Trust equipment grant (093305/Z/10/Z). A.C.K. and M.S.P.S. were also funded by the Wellcome Trust.

## References:

- 1 Humphries, J. D., Byron, A. and Humphries, M. J. (2006) Integrin ligands at a glance. *J. Cell. Sci.* **119**, 3901–3903.
- 2 Harburger, D. S. and Calderwood, D. A. (2009) Integrin signalling at a glance. *J. Cell. Sci.* **122**, 159–163.
- 3 Critchley, D. R. (2009) Biochemical and Structural Properties of the Integrin-Associated Cytoskeletal Protein Talin. *Annual Reviews Biophysics* **38**, 235–254.
- 4 Calderwood, D. A., Campbell, I. D. and Critchley, D. R. (2013) Talins and kindlins: partners in integrin-mediated adhesion. *Nat. Rev. Mol. Cell Biol.* **14**, 503–517.
- 5 Yates, L. A., Füzéry, A. K., Bonet, R., Campbell, I. D. and Gilbert, R. J. C. (2012) Biophysical analysis of Kindlin-3 reveals an elongated conformation and maps integrin binding to the membrane-distal  $\beta$ -subunit NPXY motif. *J. Biol. Chem.* **287**, 37715–37731.
- 6 Brahme, N. N. and Calderwood, D. A. (2012) Cell Adhesion: A FERM Grasp of the Tail Sorts Out Integrins. *Curr. Biol.* **22**, R692–R694.
- 7 Moser, M., Legate, K. R., Zent, R. and Fässler, R. (2009) The Tail of Integrins, Talin, and Kindlins. *Science* **324**, 895–899.
- 8 Shattil, S. J., Kim, C. and Ginsberg, M. H. (2010) The final steps of integrin activation: the end game. *Nat. Rev. Mol. Cell Biol.* **11**, 288–300.
- 9 Bachir, A. I., Zareno, J., Moissoglu, K., Plow, E. F., Gratton, E. and Horwitz, A. R. (2014) Integrin-Associated Complexes Form Hierarchically with Variable Stoichiometry in Nascent Adhesions. *Curr. Biol.* **24**, 1845–1853.
- 10 Ussar, S., Wang, H.-V., Linder, S., Fässler, R. and Moser, M. (2006) The Kindlins: Subcellular localization and expression during murine development. *Exp. Cell Res.* **312**, 3142–3151.
- 11 Bialkowska, K., Ma, Y.-Q., Bledzka, K., Sossey-Alaoui, K., Izem, L., Zhang, X., Malinin, N., Qin, J., Byzova, T. and Plow, E. F. (2010) The integrin co-activator Kindlin-3 is expressed and functional in a non-hematopoietic cell, the endothelial cell. *J. Biol. Chem.* **285**, 18640–18649.
- 12 Siegel, D. H., Ashton, G. H. S., Penagos, H. G., Lee, J. V., Feiler, H. S., Wilhelmsen, K. C., South, A. P., Smith, F. J. D., Prescott, A. R., Wessagowit, V., et al. (2003) Loss of Kindlin-1, a Human Homolog of the *Caenorhabditis elegans* Actin–Extracellular-Matrix Linker Protein UNC-112, Causes Kindler Syndrome. *Am. J. Hum. Genet.* **73**, 174–187.
- 13 Zhao, T., Guan, L., Yu, Y., Pei, X., Zhan, J., Han, L., Tang, Y., Li, F., Fang, W. and Zhang, H. (2013) Kindlin-2 promotes genome instability in breast cancer cells. *Cancer*

*Lett.* **330**, 208–216.

- 14 Yu, Y., Wu, J., Wang, Y., Zhao, T., Ma, B., Liu, Y., Fang, W., Zhu, W. G. and Zhang, H. (2012) Kindlin 2 forms a transcriptional complex with  $\beta$ -catenin and TCF4 to enhance Wnt signalling. *EMBO Rep.* **13**, 750–758.
- 15 Boyd, R. S., Adam, P. J., Patel, S., Loader, J. A., Berry, J., Redpath, N. T., Poyser, H. R., Fletcher, G. C., Burgess, N. A., Stamps, A. C., et al. (2003) Proteomic analysis of the cell-surface membrane in chronic lymphocytic leukemia: identification of two novel proteins, BCNP1 and MIG2B. *Leukemia* **17**, 1605–1612.
- 16 Sossey-Alaoui, K., Pluskota, E., Davuluri, G., Bialkowska, K., Das, M., Szpak, D., Lindner, D. J., Downs-Kelly, E., Thompson, C. L. and Plow, E. F. (2014) Kindlin-3 enhances breast cancer progression and metastasis by activating Twist-mediated angiogenesis. *FASEB J.* **28**, 2260–2271.
- 17 Xu, Z., Chen, X., Zhi, H., Gao, J., Bialkowska, K., Byzova, T. V., Pluskota, E., White, G. C., Liu, J., Plow, E. F., et al. (2014) Direct interaction of kindlin-3 with integrin  $\alpha$ IIb $\beta$ 3 in platelets is required for supporting arterial thrombosis in mice. *Arterioscler. Thromb. Vasc. Biol.* **34**, 1961–1967.
- 18 Moser, M., Nieswandt, B., Ussar, S., Pozgajova, M. and Fässler, R. (2008) Kindlin-3 is essential for integrin activation and platelet aggregation. *Nat. Med.* **14**, 325–330.
- 19 Yu, Y., Wu, J., Guan, L., Qi, L., Tang, Y., Ma, B., Zhan, J., Wang, Y., Fang, W. and Zhang, H. (2013) Kindlin 2 promotes breast cancer invasion via epigenetic silencing of the microRNA200 gene family. *Inter. J. Cancer* **133**, 1368–1379.
- 20 Huet-Calderwood, C., Brahme, N. N., Kumar, N., Stiegler, A. L., Raghavan, S., Boggon, T. J. and Calderwood, D. A. (2014) Differences in binding to the ILK complex determines kindlin isoform adhesion localization and integrin activation. *J. Cell. Sci.* **127**, 4308–4321.
- 21 Hart, R., Stanley, P., Chakravarty, P. and Hogg, N. (2013) The Kindlin 3 Pleckstrin Homology Domain Has an Essential Role in Lymphocyte Function-associated Antigen 1 (LFA-1) Integrin-mediated B Cell Adhesion and Migration. *J. of Biol. Chem.* **288**, 14852–14862.
- 22 Liu, Y., Zhu, Y., Ye, S. and Zhang, R. (2012) Crystal structure of kindlin-2 PH domain reveals a conformational transition for its membrane anchoring and regulation of integrin activation. *Protein Cell* **3**, 434–440.
- 23 Yates, L. A., Lumb, C. N., Brahme, N. N., Zalyte, R., Bird, L. E., De Colibus, L., Owens, R. J., Calderwood, D. A., Sansom, M. S. P. and Gilbert, R. J. C. (2012) Structural and functional characterization of the kindlin-1 pleckstrin homology domain. *J. Biol. Chem.* **287**, 43246–43261.

- 24 Bouaouina, M., Goult, B. T., Huet-Calderwood, C., Bate, N., Brahme, N. N., Barsukov, I. L., Critchley, D. R. and Calderwood, D. A. (2012) A conserved lipid-binding loop in the kindlin FERM F1 domain is required for kindlin-mediated  $\alpha$ IIb $\beta$ 3 integrin coactivation. *J. Biol. Chem.* **287**, 6979–6990.
- 25 Mory, A., Feigelson, S. W., Yarali, N., Kilic, S. S., Bayhan, G. I., Gershoni-Baruch, R., Etzioni, A. and Alon, R. (2008) Kindlin-3: a new gene involved in the pathogenesis of LAD-III, *Blood* **112**, 2591–2591.
- 26 Lemmon, M. A., Ferguson, K. M. and Abrams, C. S. (2001) Pleckstrin homology domains and the cytoskeleton. *FEBS Lett.* **513**, 71–76.
- 27 Lai, C.-L., Srivastava, A., Pilling, C., Chase, A. R., Falke, J. J. and Voth, G. A. (2013) Molecular Mechanism of Membrane Binding of the GRP1 PH Domain. *J. Mol. Biol.* **425**, 3073–3090.
- 28 Ferguson, K. M., Lemmon, M. A., Schlessinger, J. and Sigler, P. B. (1995) Structure of the high affinity complex of inositol trisphosphate with a phospholipase C pleckstrin homology domain. *Cell* **83**, 1037–1046.
- 29 Anand, K., Maeda, K. and Gavin, A.-C. (2012) Structural Analyses of the Slm1-PH Domain Demonstrate Ligand Binding in the Non-Canonical Site. *PLoS ONE* **7**, e36526.
- 30 Ceccarelli, D. F. J., Blasutig, I. M., Goudreault, M., Li, Z., Ruston, J., Pawson, T. and Sicheri, F. (2007) Non-canonical interaction of phosphoinositides with pleckstrin homology domains of Tiam1 and ArhGAP9. *J. of Biol. Chem.*, **282**, 13864–13874.
- 31 Kalli, A. C. and Sansom, M. S. P. (2014) Interactions of peripheral proteins with model membranes as viewed by molecular dynamics simulations. *Biochem. Soc. Trans.* **42**, 1418–1424.
- 32 Yamamoto, E., Kalli, A. C., Akimoto, T., Yasuoka, K. and Sansom, M. S. P. (2015) Anomalous Dynamics of a Lipid Recognition Protein on a Membrane Surface. *Sci. Rep.*, **5**, 18245.
- 33 Yamamoto, E., Kalli, A. C., Yasuoka, K. and Sansom, M. (2016) Interactions of Pleckstrin Homology Domains with Membranes: Adding Back the Bilayer via High-Throughput Molecular Dynamics. *Structure* **24**, 1421-1431.
- 34 Naughton, F. B., Kalli, A. C. and Sansom, M. S. P. (2016) Association of Peripheral Membrane Proteins with Membranes: Free Energy of Binding of GRP1 PH Domain with Phosphatidylinositol Phosphate-Containing Model Bilayers. *J. Phys. Chem. Lett.* **7**, 1219–1224.
- 35 Lumb, C. N., He, J., Xue, Y., Stansfeld, P. J., Stahelin, R. V., Kutateladze, T. G. and Sansom, M. S. P. (2011) Biophysical and Computational Studies of Membrane Penetration by the GRP1 Pleckstrin Homology Domain. *Structure* **19**, 1338–1346.

- 36 Orłowski, A., Kukkurainen, S., Pöyry, A., Rissanen, S., Vattulainen, I., Hytönen, V. P. and Róg, T. (2015) PIP2 and Talin Join Forces to Activate Integrin. *J. Phys. Chem. B* **119**, 12381–12389.
- 37 Schmidt, T., Suk, J.-E., Ye, F., Situ, A. J., Mazumder, P., Ginsberg, M. H. and Ulmer, T. S. (2015) Annular anionic lipids stabilize the integrin  $\alpha$ IIb $\beta$ 3 transmembrane complex. *J. Biol. Chem.* **290**, 8283–8293.
- 38 Borch, J. and Hamann, T. (2009) The nanodisc: a novel tool for membrane protein studies. *Biol. Chem.* **390**, 805–814.
- 39 Wang, X., Mu, Z., Li, Y., Bi, Y. and Wang, Y. (2015) Smaller Nanodiscs are Suitable for Studying Protein Lipid Interactions by Solution NMR. *Protein J.* **34**, 205–211.
- 40 Ellenbroek, W. G., Wang, Y.-H., Christian, D. A., Discher, D. E., Janmey, P. A. and Liu, A. J. (2011) Divalent Cation-Dependent Formation of Electrostatic PIP2 Clusters in Lipid Monolayers. *Biophys. J.* **101**, 2178–2184.
- 41 van den Bogaart, G., Meyenberg, K., Risselada, H. J., Amin, H., Willig, K. I., Hubrich, B. E., Dier, M., Hell, S. W., Grubmüller, H., Diederichsen, U., et al. (2011) Membrane protein sequestering by ionic protein-lipid interactions. *Nature* **479**, 552–555.
- 42 Vonkova, I., Saliba, A.-E., Deghou, S., Anand, K., Ceschia, S., Doerks, T., Galih, A., Kugler, K. G., Maeda, K., Rybin, V., et al. (2015) Lipid Cooperativity as a General Membrane-Recruitment Principle for PH Domains. *Cell Rep* **12**, 1519–1530.
- 43 Ferguson, K. M., Kavran, J. M., Sankaran, V. G., Fournier, E., Isakoff, S. J., Skolnik, E. Y. and Lemmon, M. A. (2000) Structural Basis for Discrimination of 3-Phosphoinositides by Pleckstrin Homology Domains. *Molecular Cell* **6**, 373–384.
- 44 Pilling, C., Landgraf, K. E. and Falke, J. J. (2011) The GRP1 PH domain, like the AKT1 PH domain, possesses a sentry glutamate residue essential for specific targeting to plasma membrane PI(3,4,5)P(3). *Biochemistry* **50**, 9845–9856.
- 45 Moravcevic, K., Oxley, C. L. and Lemmon, M. A. (2012) Conditional Peripheral Membrane Proteins: Facing up to Limited Specificity. *Structure* **20**, 15–27.
- 46 van Meer, G., Voelker, D. R. and Feigenson, G. W. (2008) Membrane lipids: where they are and how they behave. *Nat. Rev. Mol. Cell Biol.* **9**, 112–124.
- 47 Bayburt, T. H. and Sligar, S. G. (2010) Membrane protein assembly into Nanodiscs. *FEBS Letters* **584**, 1721–1727.
- 48 Borhani, D. W., Rogers, D. P. and Engler, J. A. (1997) Crystal structure of truncated human apolipoprotein AI suggests a lipid-bound conformation, *Proc. Natl. Acad. Sci.* **94**, 12291-12296
- 49 Lemmon, M. A. (2008) Membrane recognition by phospholipid-binding domains. *Nat. Rev. Mol. Cell Biol.* **9**, 99–111.

- 50 Brown, D. A. (2015) PIP2Clustering: From model membranes to cells. *Chemistry and Physics of Lipids* **192**, 33–40.
- 51 Jian, X., Tang, W.-K., Zhai, P., Roy, N. S., Luo, R., Gruschus, J. M., Yohe, M. E., Chen, P.-W., Li, Y., Byrd, R. A., et al. (2015) Molecular Basis for Cooperative Binding of Anionic Phospholipids to the PH Domain of the Arf GAP ASAP1. *Structure* **23**, 1977–1988.
- 52 Goñi, G. M., Epifano, C., Boskovic, J., Camacho-Artacho, M., Zhou, J., Bronowska, A., Martín, M. T., Eck, M. J., Kremer, L., Gräter, F., et al. (2014) Phosphatidylinositol 4,5-bisphosphate triggers activation of focal adhesion kinase by inducing clustering and conformational changes. *Proc. Natl. Acad. Sci.* **111**, E3177–86.
- 53 Calderwood, D. A., Zent, R., Grant, R., Rees, D. J., Hynes, R. O. and Ginsberg, M. H. (1999) The Talin head domain binds to integrin beta subunit cytoplasmic tails and regulates integrin activation. *J. Biol. Chem.* **274**, 28071–28074.
- 54 Goult, B. T., Bouaouina, M., Harburger, D. S., Bate, N., Patel, B., Anthis, N. J., Campbell, I. D., Calderwood, D. A., Barsukov, I. L., Roberts, G. C., et al. (2009) The Structure of the N-Terminus of Kindlin-1: A Domain Important for  $\alpha$ IIb $\beta$ 3 Integrin Activation. *J. Mol. Biol.* **394**, 944–956.
- 55 Heckman, K. L. and Pease, L. R. (2007) Gene splicing and mutagenesis by PCR-driven overlap extension. *Nat. Protoc.* **2**, 924–932.
- 56 Fairhead, M. and Howarth, M. (2014) Site-Specific Biotinylation of Purified Proteins Using BirA. *Methods Mol. Biol.* **1266**, 171–184.
- 57 Winter, G. (2009) xia2: an expert system for macromolecular crystallography data reduction. *J. Appl Crystallogr.* **43**, 186–190.
- 58 Adams, P. D., Afonine, P. V., Bunkóczi, G., Chen, V. B., Davis, I. W., Echols, N., Headd, J. J., Hung, L. W., Kapral, G. J., Grosse-Kunstleve, R. W., et al. (2010) PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213–221.
- 59 Cowtan, K.IUCr. (2006) The Buccaneer software for automated model building. 1. Tracing protein chains. *Acta Crystallogr. D Biol. Crystallogr.* **62**, 1002–1011.
- 60 Emsley, P., Cowtan, K.IUCr. (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr. D Biol. Crystallogr.* **60**, 2126–2132.
- 61 Ludtke, S. J., Baldwin, P. R. and Chiu, W. (1999) EMAN: Semiautomated Software for High-Resolution Single-Particle Reconstructions. *J. Struct. Biol.* **128**, 82–97.
- 62 van Heel, M., Harauz, G., Orlova, E. V., Schmidt, R. and Schatz, M. (1996) A New Generation of the IMAGIC Image Processing System. *J. Struct. Biol.* **116**, 17–24.
- 63 Frank, J., Radermacher, M., Penczek, P., Zhu, J., Li, Y., Ladjadj, M. and Leith, A.



- (1996) SPIDER and WEB: Processing and Visualization of Images in 3D Electron Microscopy and Related Fields. *J. Struct. Biol.* **116**, 190–199.
- 64 Hess, B., Kutzner, C., van der Spoel, D. and Lindahl, E. (2008) GROMACS 4: Algorithms for Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *J. Chem. Theory Comput.* **4**, 435–447.
- 65 Siewert J Marrink, H Jelger Risselada, Serge Yefimov, D Peter Tieleman, A. and de Vries, A. H. (2007) The MARTINI Force Field: Coarse Grained Model for Biomolecular Simulations. *J. Phys. Chem. B.* **111**, 7812–7824.
- 66 Monticelli, L., Kandasamy, S. K., Periole, X., Larson, R. G., Tieleman, D. P. and Marrink, S.-J. (2008) The MARTINI Coarse-Grained Force Field: Extension to Proteins. *J. Chem. Theory Comput.* **4**, 819–834.
- 67 Atilgan, A. R., Durell, S. R., Jernigan, R. L., Demirel, M. C., Keskin, O. and Bahar, I. (2001) Anisotropy of Fluctuation Dynamics of Proteins with an Elastic Network Model. *Biophys. J.* **80**, 505–515.
- 68 Hess, B., Bekker, H. and Berendsen, H. (1997) LINCS: a linear constraint solver for molecular simulations. *J. Comput. Chem.* **18**, 1463–1472.
- 69 Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A. and Haak, J. R. (1984) Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* **81**, 3684–3690.
- 70 Stansfeld, P. J. and Sansom, M. S. P. (2011) From Coarse Grained to Atomistic: A Serial Multiscale Approach to Membrane Protein Simulations. *J. Chem. Theory Comput.* **7**, 1157–1166.
- 71 Walter R P Scott, Philippe H Hünenberger, Ilario G Tironi, Alan E Mark, Salomon R Billeter, Jens Fennen, Andrew E Torda, Thomas Huber, Peter Krüger, A. and van Gunsteren, W. F. (1999) The GROMOS Biomolecular Simulation Program Package. *J. Phys. Chem. A* **103**, 3596–3607.
- 72 Bussi, G., Donadio, D. and Parrinello, M. (2007) Canonical sampling through velocity rescaling. *J. Chem. Phys* **126**, 014101.
- 73 Parrinello, M. and Rahman, A. (1981) Polymorphic transitions in single crystals: A new molecular dynamics method. *J. Applied Phys.* **52**, 7182–7190.
- 74 Hub, J.S., de Groot, B.L., and van der Spoel, D. (2010). g\_wham—A Free Weighted Histogram Analysis Implementation Including Robust Error and Autocorrelation Estimates. *J. Chem. Theory Comput.* **6**, 3713–3720.



**Table 1:** Data collection and structure refinement

Data collection	
Space group	<i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2
Cell dimensions	
<i>a</i> , <i>b</i> , <i>c</i> (Å)	132.04, 36.19, 52.69
$\alpha$ , $\beta$ , $\gamma$ (°)	90, 90, 90
Resolution (Å)	31.73-2.23 (2.31-2.23)
R <sub>merge</sub>	0.049 (0.432)
<i>I</i> / $\sigma$ <i>I</i>	23.7 (2.3)
Completeness (%)	96 (93)
Redundancy	3.1 (2.6)
CC half	0.998 (0.771)
Refinement	
Resolution (Å)	31.73-2.23 (2.31-2.23)
No. reflections	38032 (3014)
No. unique reflections	12441 (1155)
R <sub>work</sub> /R <sub>free</sub>	21.94 (27.07)
No. atoms	
Protein	2123
Ligand/ion	1
Water	13
B-factors (Å <sup>2</sup> )	
Protein	84.92
Ligand/ion	13.06
Water	61.66
R.m.s. deviations	
Bond lengths (Å)	0.007
Bond angles (°)	1.11
Ramachandran	
Favoured (outliers)	98% (0%)

**Table 2:** Summary of simulations

Simulation	Bilayer composition	Duration
<i>Coarse-grained</i>		
<i>Kin3PH-1</i>	POPC:POPS:PIP <sub>2</sub> (~73:20:7)	20 x 1.5 $\mu$ s
<i>Kin3PH-2</i>	POPC:POPS:PIP <sub>3</sub> (~73:20:7)	20 x 1.5 $\mu$ s
<i>Kin3PH-3</i>	POPC:POPS (~78:20) + 4 PIP <sub>3</sub> molecule per leaflet	20 x 1.5 $\mu$ s
<i>Kin3PH-4</i>	POPC:POPS (~79:20) + 2 PIP <sub>3</sub> molecule per leaflet	20 x 1.5 $\mu$ s
<i>Kin3PH-5</i>	POPC:POPS (~79:20) + 1 PIP <sub>3</sub> molecule per leaflet	20 x 1.5 $\mu$ s
<i>Kin3PH-6</i>	POPC:POPS (~73:20) + 5% PIP <sub>2</sub> and 4 PIP <sub>3</sub> molecule per leaflet	20 x 1.5 $\mu$ s
<i>Kit3mutPH-1</i>	POPC:POPS (~78:25) + 4 PIP <sub>3</sub> molecule per leaflet - K363A, K367A mutation	20 x 1.5 $\mu$ s
<i>Atomistic</i>		
<i>Kin3PH_AT-1</i>	No bilayer – GROMOS force-field	3 x 100 ns
<i>Kin3PH_AT-1</i>	No bilayer – OPLS force-field	3 x 100 ns
<i>Kin3PH_AT-3</i>	POPC:POPS:PIP <sub>2</sub> (~73:20:7)	6 x 100 ns
<i>Kin3PH_AT-4</i>	POPC:POPS (78:20) + 4 PIP <sub>3</sub> molecule per leaflet	4 x 100 ns

### **Figure 1: Overall structure of kindlin-3 PH domain**

(A) Crystal structure of kindlin-3 PH domain. The protein is coloured rainbow-wise from its amino (blue) to carboxy (red) terminus and the canonical inositol phosphate binding site is highlighted with an arrow (left). The hydrogen bonds within the open end of the PH domain are shown (right). (B) Surface electrostatic charge of kindlin-3 PH domain. (C) Average Root Mean Square Fluctuation (RMSF) of the kindlin-3 PH domain during atomistic MD simulations in water (i.e. in the absence of a bilayer) for two different forcefields (i.e. GROMOS and OPLS). The magnitude of the average RMSF of each residue was mapped on the kindlin-3 PH domain crystal structure.

### **Figure 2: Coarse-grained MD simulations of Kindlin-3 PH domain interacting with a PIP-containing lipid bilayer**

(A) Snapshots at the beginning and at the end of one of the simulations with 4 PIP<sub>3</sub> molecules in each leaflet (Kin3PH-3 in Table 2). PIP<sub>3</sub> molecules are shown in green, POPC and POPS phosphate atoms are shown in grey and the kindlin-3 PH domain is shown in ice blue.

(B) Rotation of the kindlin-3 PH domain (shown as the  $R_{zz}$  component of its rotational matrix) relative to the distance between the centre of mass of the protein and the centre of mass of the bilayer. (C) Final snapshot from one of the atomistic simulations that started using snapshots from the simulations shown in (A). (D) Number of PIP molecules interacting with the kindlin-3 PH domain. The number of PIP lipids is shown for Kin3PH-1, Kin3PH-2 and Kin3PH-3 simulations (Table 2) in orange, green and light green, respectively. For this calculation a PIP molecule is considered to be in contact with the protein if its phosphate is within 1 nm of the protein. The number of PIP molecules are shown only for the frames in which the protein is bound to the bilayer (i.e. the z component of the distance between the protein centre of mass and the bilayer centre of mass is smaller than 4.75 nm).

**Figure 3: Potential of mean force (PMF) calculations of Kindlin-3 PH domain with PIPs.**

(A) PMF calculations for the binding of the wild type and the mutated form of kindlin-3 PH domain to PIP<sub>2</sub> (black) and PIP<sub>3</sub> (cyan and red) molecules. (B) Kindlin-3 PH/bilayer complexes corresponding to wells 1 and 2 in (A). The PIP<sub>3</sub> molecules are shown in green and the PH domain in ice blue. For this calculation the initial structures of wild type or mutant kindlin-3 PH domains bound to PIP<sub>3</sub> or PIP<sub>2</sub> in a 80:20 POPC:POPS bilayer were generated by alignment with an existing Grp1-PH/PIP<sub>3</sub> complex. The PH domain was pulled from the bound PIP lipid and bilayer along the membrane normal (z-axis). See Methods for more details.

**Figure 4: SPR study of kindlin-3 PH domain with immobilised PIP2 and PIP3 headgroups.**

(A) Schematic representation of SPR experiments with immobilised lipid head group as ligand and PH domain as analyte. (B) SPR measurements of kindlin-3 PH domain WT and mutants with immobilised PtdIns(4,5)P<sub>2</sub> and PtdIns(3,4,5)P<sub>3</sub>, respectively. On the left in each case we show the background-subtracted raw sensogram data for the wild type protein, while on the right we show binding saturation curves for wild type, K363A K367A and IPRR insertion mutants. The highest concentration used is given on each set of sensogram curves; subsequent binding profiles derive from a series of twofold dilutions.

**Figure 5: SPR study of kindlin-3 PH domain with PtdIns(3,4,5)P<sub>3</sub> incorporated nanodiscs.**

(A) Schematic representation of SPR experiments with immobilised PH domain as ligand and lipid nanodiscs as analyte in solution. (B) SPR measurement between WT kindlin-3 PH domain and nanodiscs with 5% PtdIns(3,4,5)P<sub>3</sub>. The background-subtracted raw sensogram data for the wild type protein is shown on the left and the resulting binding saturation curve on the right. The highest concentration used is given on the sensogram graph; subsequent binding profiles derive from a series of twofold dilutions. See Materials and Methods for experimental details.

**Figure 6: SPR study of interaction between Grp1PH domain and PtdIns(3,4,5)P<sub>3</sub> in two experimental setups.**

(A) Immobilised PtdIns(3,4,5)P<sub>3</sub> headgroup as ligand and PH domain as analyte. (B) immobilised Grp1PH domain as ligand and lipid nanodiscs as analyte. The upper panel in each case gives the background-subtracted raw sensogram data and the lower panel the resulting binding saturation curve. The highest concentration used is given on the sensogram graphs (83µM Grp1 PH domain in A; 3.4µM nanodiscs in B); subsequent binding profiles derive from a series of twofold dilutions. See Materials and Methods for experimental details.

**Figure 7: A proposed model for integrin activation by kindlin-3 with recruitment by clustered inositol phosphates to a membrane.**

The activating conversion of integrins to an upright conformation is promoted by the binding of talin to the membrane-proximal NPxY and of kindlin to the membrane-distal NPxY on an integrin β-subunit cytoplasmic tail. Clustered PtdIns(3,4,5)P<sub>3</sub> interacts and recruits kindlin-3 from cytosol to membrane, thus facilitating the formation of integrin-talin-kindlin tertiary complex.



Figure 1






Figure 2











Figure 3

**A**



**B**







Figure 4

**A**



Analytes: K3PH

Ligand: PtdInsPs headgroups



**B**










Figure 5

**A**



Analytes: PtdInsPs nanodiscs

Ligand: K3PH



Nanodiscs



PtdInsPs

**B**

K3PH binding to **PtdIns(3,4,5)P<sub>3</sub>** nanodiscs








Figure 6

**A**

Grp1PH binding to **PtdIns(3,4,5)P<sub>3</sub>**



**B**

Grp1PH binding to **PtdIns(3,4,5)P<sub>3</sub>** nanodiscs








Figure 7




## Supporting Information



SI Figure S1 Sequence alignment of the PH domains of murine kindlins -1, -2 and -3.

Numbering is by reference to the complete protein sequence, as is the residue numbering in the main text.






SI Figure S2 Phylogenetic tree of representative PH domains from crystallographic studies.


Some PH domains are highlighted: orange, kindlins; blue, high-affinity specific PtdInsP binders; green, low affinity or promiscuous PtdInsP binders. The following proteins are used to generate the phylogenetic tree: Btk (1B55); PI3Kinase enhancer (2RLO);  $\alpha_1$  syntrophin (2ADZ); P63Rho GEF (2RGN); Dbs (1KZ7); intersectin (1KI1); PLC $\beta$ 2 (2FJU); PLC $\delta$  (1MAI); GPCR kinase 2 (1BAK); Unc89 (1FHO); PDK1 (1W1H); SOS1 (1AWE); pleckstrin (2I5F); PKB/Akt (1UNQ); Grb10 (3HK0); ARNO (1U27); Grp1 (1FGY); dynamin (3ZVR); SLM1 (3NSU); PEPP1 (1UPQ); kindlin-1(4BBK); kindlin-2 (4F7H); kindlin-3 (5L81 (this work)); OBPRP1 (2D9X); FAPP1 (3RCP); PLCG2 (2W2W); ArhGAP9 (2P0D); SKAP-Hom (1U5E); Tiam-2 (3A8P); RAL GEF (2DTC); B-spectrin (1BTN).






*SI Figure S3 Comparison of the canonical and non-canonical inositol phosphate binding sites of PH domains.*

(A) Surface electrostatic charge distribution of kindlin-2 and kindlin-3 PH domains. Kindlin-2 and -3 are placed in the same orientation with the open barrel at the top. (B) Surface electrostatic charge distribution of PH domains with canonical inositol phosphate binding pocket (PLC $\delta$ 1-PH and Grp1-PH) and non-canonical binding site (ArhGAP9 and Slm1). These molecules are placed in the same orientation as in (A). PH domains with a canonical inositol phosphate binding site typically exhibit a positively-charged barrel opening.




SI Figure S4: Hydrogen bond between Arg<sup>360</sup> and Gln<sup>396</sup> in simulation.

(A,B) Minimum distance between residues Arg<sup>360</sup> and Gln<sup>396</sup> during the simulations with the kindlin-3 PH domain in solution. The distance is shown separately for the three repeat simulations with the GROMOS (A) and the OPLS (B) force-fields.




SI Figure S5: Contacts between Kindlin-3 PH domain and PIPs

(A,B,C) Normalised average number of contacts of the interactions of the kindln-3 PH domain with PIP<sub>2</sub> or PIP<sub>3</sub> molecules. The contacts were calculated using a cutoff distance of 0.5 nm. The interactions with PIP<sub>2</sub> molecules (A) were calculated from the Kin3PH-1 simulation. The interactions of kindlin-3 with the PIP<sub>3</sub> molecules are shown as the average over all the simulations with PIP<sub>3</sub> in the bilayer (B; simulations Kin3PH-2 to 5) and the interactions with PIP<sub>2</sub> and PIP<sub>3</sub> molecules in (C) were calculated from Kin3PH-6 system. (D,E) Final snapshots from the atomistic simulations with the PIP<sub>2</sub>-containing bilayer showing the two binding modes of kindlin-3 PH domain to the bilayer.



*SI Figure S6: Orientation of kindlin-3 PH domain on the membrane bilayer.*


Orientation of the kindlin-3 PH domain relative to the bilayer shown as the  $R_{zz}$  component of the rotational matrix to align the kindlin-3 PH domain to a reference structure versus the distance between the centres of mass of the protein and the bilayer. The reference structure for each system was in the same orientation as in Figure 1C. In the figure we show the normalized density to aid comparison between the different systems.



SI Figure S7: Contact between the Kindlin-3 PH domain and PIP molecules in the free energy calculations:


(A,B) Normalised average number of contacts of the interactions of the wild type kindlin-3 PH domain with PIP<sub>3</sub> (A) or PIP<sub>2</sub> (B) molecules in the free energy calculations. The contacts were calculated using the first four windows of each simulation system (that correspond to the first well in Fig. 4A).







SI Figure S8: *Kin3mutPH-1* dissociates from membrane during MD simulations.

(A) Distance between the centre of mass of the protein and the centre of mass of the bilayer for one of the simulations with the mutated form of the protein (*Kin3mutPH-1*). (B) Snapshots from the same system shown at 0, 0.26, 0.65 and 1.5 μs. PIP<sub>3</sub> molecules are shown in green, POPE and POPC phosphate atoms are shown in grey and the *kin3mutPH-1* is shown in ice blue.



SI Figure S9 Preparation and characterisation of nanodiscs.

(A) Size-exclusion chromatography profile of nanodisc purification. The regions shadowed in light pink and green represent the peak fractions used for this study. (B) Analytical ultracentrifugation study of nanodiscs showing their homogeneous distribution in solution. (C) Dynamic light scattering measurement of two representative nanodisc species. (D) 3D cryo-EM reconstruction of POPC/POPS nanodiscs. (E) Size distribution of nanodiscs with various lipid compositions measured by dynamic light scattering.



SI Figure S10 Grp1 PH domain binding to PtdIns(3,5)P<sub>2</sub>

SPR study with immobilised PtdIns(3,5)P<sub>2</sub> on sensor chip and Grp1PH domain as analyte. The experimental setup was the same with that in Figure 6A.