



**UNIVERSITY OF LEEDS**

This is a repository copy of *p-mechanics as a physical theory: an introduction*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/111528/>

Version: Accepted Version

---

**Article:**

Kisil, VV [orcid.org/0000-0002-6593-6147](http://orcid.org/0000-0002-6593-6147) (2004) *p-mechanics as a physical theory: an introduction*. *Journal of Physics A: Mathematical and General*, 37 (1). pp. 183-204. ISSN 0305-4470

<https://doi.org/10.1088/0305-4470/37/1/013>

---

© 2004 IOP Publishing Ltd. This is an author produced version of an article published in *Journal of Physics A: Mathematical and General*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# $p$ -MECHANICS AS A PHYSICAL THEORY. AN INTRODUCTION

VLADIMIR V. KISIL

ABSTRACT. The paper provides an introduction into  $p$ -mechanics, which is a consistent physical theory suitable for a simultaneous description of classical and quantum mechanics.  $p$ -Mechanics naturally provides a common ground for several different approaches to quantisation (geometric, Weyl, coherent states, Berezin, deformation, Moyal, etc.) and has a potential for expansions into field and string theories. The backbone of  $p$ -mechanics is solely the representation theory of the Heisenberg group.

## CONTENTS

List of Figures	1
1. Introduction	2
2. The Heisenberg Group and Its Representations	2
2.1. Representations $\mathbb{H}^n$ and Method of Orbit	2
2.2. Structure and Topology of the Unitary Dual of $\mathbb{H}^n$	5
2.3. Fock Spaces $F_2(\mathcal{O}_h)$ and Coherent States	6
3. $p$ -Mechanics: Statics	8
3.1. Observables in $p$ -Mechanics, Convolutions and Commutators	8
3.2. Berezin Quantisation and Wavelet Transform	10
3.3. From Classical and Quantum Observables to $p$ -Mechanics	12
4. $p$ -Mechanics: Dynamics	14
4.1. $p$ -Mechanical Brackets on $\mathbb{H}^n$	15
4.2. $p$ -Mechanical Dynamic Equation	16
4.3. Symplectic Invariance from Automorphisms of $\mathbb{H}^n$	18
5. Conclusions	19
5.1. Discussion	19
5.2. Further Developments	20
Acknowledgements	21
References	21

## LIST OF FIGURES

1 The unitary dual object to the Heisenberg group	5
2 Quantisation and wavelet transforms	13
3 Dynamics of the harmonic oscillator	17

---

2000 *Mathematics Subject Classification*. Primary 81R05; Secondary 81R15, 22E27, 22E70, 43A65.

*Key words and phrases*. Classical mechanics, quantum mechanics, Moyal brackets, Poisson brackets, commutator, Heisenberg group, orbit method, deformation quantisation, symplectic group, representation theory, metaplectic representation, Berezin quantisation, Weyl quantisation, Segal–Bargmann–Fock space, coherent states, wavelet transform, contextual interpretation, string theory, field theory.

On leave from the Odessa University.

## 1. INTRODUCTION

This paper describes how classical and quantum mechanics are naturally united within a construction based on the Heisenberg group  $\mathbb{H}^n$  and the complete set of its unitary representations. There is a dynamic equation (4.9) on  $\mathbb{H}^n$  which generates both Heisenberg (4.10) and Hamilton (4.11) equations and corresponding classical and quantum dynamics. The standard assumption that observables constitute an algebra, which is discussed in [24, 26] and elsewhere, is not necessary for setting up a valid quantisation scheme.

*The paper outline* is as follows. In the next Section we recall the representation theory of the Heisenberg group based on the *orbit method* of Kirillov [21] and utilising Fock–Segal–Bargmann spaces [11, 15]. We emphasise the existence *and* usability of the family of one-dimensional representations: they play for classical mechanics exactly the same rôle as infinite dimensional representations do for quantum. In Section 3 we introduce the concept of observable in  $p$ -mechanics and describe their relations with quantum and classical observables. These links are provided by the representations of the Heisenberg group and *wavelet* transforms. In Section 4 we study  $p$ -mechanical brackets and the associated dynamic equation together with its classical and quantum representations. In conclusion we derive the symplectic invariance of dynamics from automorphisms of  $\mathbb{H}^n$ .

The notion of physical states in  $p$ -mechanics is introduced in subsequent publications [6, 7];  $p$ -mechanical approach to quantised fields is sketched in [28] with some further papers to follow.

## 2. THE HEISENBERG GROUP AND ITS REPRESENTATIONS

We start from the representation theory of the Heisenberg group  $\mathbb{H}^n$  based on the *orbit method* of Kirillov. Analysis of the unitary dual of  $\mathbb{H}^n$  in Subsection 2.2 suggests that the family of one-dimensional representations of  $\mathbb{H}^n$  forms the phase space of a classical system. Infinite dimensional representations in the Fock type space are described in Subsection 2.3.

**2.1. Representations  $\mathbb{H}^n$  and Method of Orbit.** Let  $(s, x, y)$ , where  $x, y \in \mathbb{R}^n$  and  $s \in \mathbb{R}$ , be an element of the Heisenberg group  $\mathbb{H}^n$  [11, 15]. We assign physical units to coordinates on  $\mathbb{H}^n$ . Let  $M$  be a unit of mass,  $L$ —of length,  $T$ —of time then we adopt the following

- Convention 2.1.** (1) Only physical quantities of the *same dimension* can be added or subtracted.
- (2) Therefore mathematical functions, e.g.  $\exp(u) = 1 + u + u^2/2! + \dots$  or  $\sin(u)$ , can be naturally constructed out of a dimensionless number  $u$  only. Thus Fourier dual variables, say  $x$  and  $q$ , should possess reciprocal dimensions because they have to form the expression  $e^{ixq}$ .
- (3) We assign to  $x$  and  $y$  components of  $(s, x, y)$  physical units  $1/L$  and  $T/(LM)$  respectively.

The Convention 2.1.3 is the only *a priori* assumption which we made about physical dimensions and it will be justified *a posteriori* as follows. From 2.1.2 we need dimensionless products  $qx$  and  $py$  in order to get the exponent in (2.15), where  $q$  and  $p$  represent the classical coordinates and momenta (in accordance to the main observation of  $p$ -mechanics). All other dimensions will be assigned strictly in agreement with the Convention 2.1.1 and 2.1.2.

The group law on  $\mathbb{H}^n$  is given as follows:

$$(s, x, y) * (s', x', y') = (s + s' + \frac{1}{2}\omega(x, y; x', y'), x + x', y + y'), \quad (2.1)$$

where the non-commutativity is solely due to  $\omega$ —the *symplectic form* [2, § 37] on the Euclidean space  $\mathbb{R}^{2n}$ :

$$\omega(x, y; x', y') = xy' - x'y. \quad (2.2)$$

Consequently the parameter  $s$  should be measured in  $T/(L^2M)$ —the product of units of  $x$  and  $y$ . The Lie algebra  $\mathfrak{h}^n$  of  $\mathbb{H}^n$  is spanned by the basis  $S, X_j, Y_j$ ,  $j = 1, \dots, n$ , which may be represented by either left- or right-invariant vector fields on  $\mathbb{H}^n$ :

$$S^{l(r)} = \pm \frac{\partial}{\partial s}, \quad X_j^{l(r)} = \pm \frac{\partial}{\partial x_j} - \frac{y_j}{2} \frac{\partial}{\partial s}, \quad Y_j^{l(r)} = \pm \frac{\partial}{\partial y_j} + \frac{x_j}{2} \frac{\partial}{\partial s}. \quad (2.3)$$

These fields satisfy the Heisenberg *commutator relations* expressed through the Kronecker delta  $\delta_{i,j}$  as follows:

$$[X_i^{l(r)}, Y_j^{l(r)}] = \delta_{i,j} S^{l(r)} \quad (2.4)$$

and all other commutators (including any between a left and a right fields) vanishing. Units to measure  $S^{l(r)}$ ,  $X_j^{l(r)}$ , and  $Y_j^{l(r)}$  are inverse to  $s, x, y$ —i.e.  $L^2M/T, L$ , and  $LM/T$  respectively—which are obviously compatible with (2.4).

The exponential map  $\exp : \mathfrak{h}^n \rightarrow \mathbb{H}^n$  respecting the multiplication (2.1) and Heisenberg commutators (2.4) is provided by the formula:

$$\exp : sS + \sum_{j=1}^n (x_j X_j + y_j Y_j) \mapsto (s, x_1, \dots, x_n, y_1, \dots, y_n).$$

The composition of the exponential map with representations (2.3) of  $\mathfrak{h}^n$  by the left(right)-invariant vector fields produces the right (left) regular representation  $\lambda_{r(l)}$  of  $\mathbb{H}^n$  by right (left) shifts. Linearised [19, § 7.1] to  $L_2(\mathbb{H}^n)$  they are:

$$\lambda_r(g) : f(h) \mapsto f(hg), \quad \lambda_l(g) : f(h) \mapsto f(g^{-1}h), \quad \text{where } f(h) \in L_2(\mathbb{H}^n). \quad (2.5)$$

As any group  $\mathbb{H}^n$  acts on itself by the conjugation automorphisms  $A(g)h = g^{-1}hg$ , which fix the unit  $e \in \mathbb{H}^n$ . The differential  $\text{Ad} : \mathfrak{h}^n \rightarrow \mathfrak{h}^n$  of  $A$  at  $e$  is a linear map which can be differentiated again to the representation  $\text{ad}$  of the Lie algebra  $\mathfrak{h}^n$  by the commutator:  $\text{ad}(A) : B \mapsto [B, A]$ . The adjoint space  $\mathfrak{h}_n^*$  of the Lie algebra  $\mathfrak{h}^n$  can be realised by the left invariant first order differential forms on  $\mathbb{H}^n$ . By the duality between  $\mathfrak{h}^n$  and  $\mathfrak{h}_n^*$  the map  $\text{Ad}$  generates the *co-adjoint representation* [19, § 15.1]  $\text{Ad}^* : \mathfrak{h}_n^* \rightarrow \mathfrak{h}_n^*$ :

$$\text{Ad}^*(s, x, y) : (h, q, p) \mapsto (h, q + hy, p - hx), \quad \text{where } (s, x, y) \in \mathbb{H}^n \quad (2.6)$$

and  $(h, q, p) \in \mathfrak{h}_n^*$  in bi-orthonormal coordinates to the exponential ones on  $\mathfrak{h}^n$ . These coordinates  $h, q, p$  should have units of an *action*  $ML^2/T$ , *coordinates*  $L$ , and *momenta*  $LM/T$  correspondingly. Again nothing in (2.6) violates the Convention 2.1.

There are two types of orbits for  $\text{Ad}^*$  (2.6): isomorphic to Euclidean spaces  $\mathbb{R}^{2n}$  and single points:

$$\mathcal{O}_h = \{(h, q, p) : \text{for a fixed } h \neq 0 \text{ and all } (q, p) \in \mathbb{R}^{2n}\}, \quad (2.7)$$

$$\mathcal{O}_{(q,p)} = \{(0, q, p) : \text{for a fixed } (q, p) \in \mathbb{R}^{2n}\}. \quad (2.8)$$

The *orbit method* of Kirillov [19, § 15], [21] starts from the observation that the above orbits parametrise all irreducible unitary representations of  $\mathbb{H}^n$ . All representations are *induced* [19, § 13] by the character  $\chi_h(s, 0, 0) = e^{2\pi ihs}$  of the centre of  $\mathbb{H}^n$  generated by  $(h, 0, 0) \in \mathfrak{h}_n^*$  and shifts (2.6) from the “left hand side” (i.e. by

$g^{-1}$ ) on orbits. Using [19, § 13.2, Prob. 5] we get a neat formula, which (unlike some other in literature, e.g. [33, Chap. 1, (2.23)]) respects the Convention 2.1 for all physical units:

$$\rho_h(s, x, y) : f_h(q, p) \mapsto e^{-2\pi i(hs + qx + py)} f_h\left(q - \frac{\hbar}{2}y, p + \frac{\hbar}{2}x\right). \quad (2.9)$$

Exactly the same formula is obtained if we apply the Fourier transform  $\hat{\cdot} : L_2(\mathbb{H}^n) \rightarrow L_2(\mathfrak{h}_n^*)$  given by:

$$\hat{\phi}(F) = \int_{\mathfrak{h}^n} \phi(\exp X) e^{-2\pi i\langle X, F \rangle} dX \quad \text{where } X \in \mathfrak{h}^n, F \in \mathfrak{h}_n^* \quad (2.10)$$

to the left regular action (2.5), see [21, § 2.3] for relations of the Fourier transform (2.10) and the orbit method.

The derived representation  $d\rho_h$  of the Lie algebra  $\mathfrak{h}^n$  defined on the vector fields (2.3) is:

$$d\rho_h(S) = -2\pi i\hbar I, \quad d\rho_h(X_j) = \frac{\hbar}{2}\partial_{p_j} - 2\pi i q_j I, \quad d\rho_h(Y_j) = -\frac{\hbar}{2}\partial_{q_j} - 2\pi i p_j I, \quad (2.11)$$

which clearly represents the commutation rules (2.4). The representation  $\rho_h$  (2.9) is reducible on whole  $L_2(\mathcal{O}_h)$  as can be seen from the existence of the set of ‘‘right-invariant’’, i.e. commuting with (2.11), differential operators:

$$d\rho_h^r(S) = 2\pi i\hbar I, \quad d\rho_h^r(X_j) = -\frac{\hbar}{2}\partial_{p_j} - 2\pi i q_j I, \quad d\rho_h^r(Y_j) = \frac{\hbar}{2}\partial_{q_j} - 2\pi i p_j I, \quad (2.12)$$

which also represent the commutation rules (2.4).

To obtain an irreducible representation defined by (2.9) we need to restrict it to a subspace of  $L_2(\mathcal{O}_h)$  where operators (2.12) acts as scalars, e.g. use a *polarisation* from the *geometric quantisation* [35]. For  $\hbar > 0$  consider the vector field  $-X_j + ic_i Y_j$  from the complexification of  $\mathfrak{h}^n$ , where the constant  $c_i$  has the dimension  $T/M$  in order to satisfy the Convention 2.1, the numerical value of  $c_i$  in given units can be assumed 1. We introduce operators  $D_h^j$ ,  $1 \leq j \leq n$  representing vectors  $-X_j + ic_i Y_j$ :

$$D_h^j = d\rho_h^r(-X_j + ic_i Y_j) = \frac{\hbar}{2}(\partial_{p_j} + c_i i \partial_{q_j}) + 2\pi(c_i p_j + i q_j)I = \hbar \partial_{\bar{z}_j} + 2\pi z_j I \quad (2.13)$$

where  $z_j = c_i p_j + i q_j$ . For  $\hbar < 0$  we define  $D_h^j = d\rho_h^r(-c_i Y_j + i X_j)$ . Operators (2.13) are used to give the following classical result in terms of orbits:

**Theorem 2.2** (Stone–von Neumann, cf. [11, Chap. 1, § 5], [19, § 18.4]). *All unitary irreducible representations of  $\mathbb{H}^n$  are parametrised up to equivalence by two classes of orbits (2.7) and (2.8) of co-adjoint representation (2.6) in  $\mathfrak{h}_n^*$ :*

- (1) *The infinite dimensional representations by transformation  $\rho_h$  (2.9) for  $\hbar \neq 0$  in Fock [11, 15] space  $F_2(\mathcal{O}_h) \subset L_2(\mathcal{O}_h)$  of null solutions to the operators  $D_h^j$  (2.13):*

$$F_2(\mathcal{O}_h) = \{f_h(q, p) \in L_2(\mathcal{O}_h) \mid D_h^j f_h = 0, 1 \leq j \leq n\}. \quad (2.14)$$

- (2) *The one-dimensional representations as multiplication by a constant on  $\mathbb{C} = L_2(\mathcal{O}_{(q,p)})$  which drop out from (2.9) for  $\hbar = 0$ :*

$$\rho_{(q,p)}(s, x, y) : c \mapsto e^{-2\pi i(qx + py)} c, \quad (2.15)$$

*with the corresponding derived representation*

$$d\rho_{(q,p)}(S) = 0, \quad d\rho_{(q,p)}(X_j) = -2\pi i q_j, \quad d\rho_{(q,p)}(Y_j) = -2\pi i p_j. \quad (2.16)$$




FIGURE 1. The structure of unitary dual object to  $\mathbb{H}^n$  appearing from the method of orbits. The space  $\mathfrak{h}_n^*$  is sliced into “horizontal” hyperplanes. Planes with  $h \neq 0$  form single orbits and correspond to different classes of unitary irreducible representation. The plane  $h = 0$  is a family of one-point orbits  $(0, q, p)$ , which produce one-dimensional representations. The topology on the dual object is the factor topology inherited from the  $\mathfrak{h}_n^*$  [21, § 2.2].

**2.2. Structure and Topology of the Unitary Dual of  $\mathbb{H}^n$ .** The structure of the unitary dual object to  $\mathbb{H}^n$ —the collection of all different classes of unitary irreducible representations—as it appears from the method of orbits is illustrated by the Figure 1, cf. [20, Chap. 7, Fig. 6 and 7]. The adjoint space  $\mathfrak{h}_n^*$  is sliced into “horizontal” hyperplanes. A plane with a parameter  $h \neq 0$  forms a single orbit (2.7) and corresponds to a particular class of unitary irreducible representation (2.9). The plane with parameter  $h = 0$  is a family of one-point orbits  $(0, q, p)$  (2.8), which produce one-dimensional representations (2.15). The topology on the dual object is the factor topology inherited from the adjoint space  $\mathfrak{h}_n^*$  under the above identification, see [21, § 2.2].

**Example 2.3.** A set of representations  $\rho_h$  (2.9) with  $h \rightarrow 0$  is dense in the whole family of one-dimensional representations (2.15), as can be seen either from the Figure 1 or the analytic expressions (2.9) and (2.15) for those representations.

Non-commutative representations  $\rho_h$ ,  $h \neq 0$  (2.9) are known to be connected with quantum mechanics [11] from its origin. This explains, for example, the name of the Heisenberg group. In the contrast commutative representations (2.15) are mostly neglected and only mentioned for sake of completeness in some mathematical formulations of the Stone–von Neumann theorem. The development of  $p$ -mechanics started [23] from the observation that the union of all representations  $\rho_{(q,p)}$ ,  $(q, p) \in \mathbb{R}^{2n}$  naturally acts as the classical *phase space*. The sensibleness of the single union

$$\mathcal{O}_0 = \bigcup_{(q,p) \in \mathbb{R}^{2n}} \mathcal{O}_{(q,p)} \tag{2.17}$$

rather than unrelated set of disconnected orbits manifests itself in several ways:

- (1) The topological position of  $\mathcal{O}_0$  as the limiting case (cf. Example 2.3) of quantum mechanics for  $h \rightarrow 0$  realises the *correspondence principle* between quantum and classical mechanics.
- (2) Symplectic automorphisms of the Heisenberg group (see Subsection 4.3) produce the metaplectic representation in quantum mechanics and *transitively* act by linear symplectomorphisms on the whole set  $\mathcal{O}_0 \setminus \{0\}$ .

- (3) We got the Poisson brackets (4.7) on  $\mathcal{O}_0$  from the same source (4.2) that leads to the correct Heisenberg equation in quantum mechanics.

The identification of  $\mathcal{O}_0$  with the classical phase space justifies that  $q$  and  $p$  are measured by the units of length and momentum respectively, which supports our choice of units for  $x$  and  $y$  in Convention 2.1.3.

*Remark 2.4.* Since unitary representations are classified up to a unitary equivalence one may think that their explicit realisations in particular Hilbert spaces are “the same”. However a suitable form of a representation can give many technical advantages. The classical illustration is the paper [15], where the comparison of the (unitary equivalent!) Schrödinger and Fock representations of  $\mathbb{H}^n$  is the principal tool of investigation.

Our form (2.9) of representations of  $\mathbb{H}^n$  given in Theorem 2.2 has at least two following advantages, which are rarely combined together:

- (1) There is the explicit physical meaning of all entries in (2.9) as will be seen below. In the contrast the formula (2.23) in [33, Chap. 1] contains terms  $\sqrt{\hbar}$  (in our notations), which could be hardly justified from a physical point of view.
- (2) The one-dimension representations (2.15) explicitly correspond to the case  $\hbar = 0$  in (2.9). The Schrödinger representation (the most used in quantum mechanics!) is handicapped in this sense: a transition  $\hbar \rightarrow 0$  from  $\rho_\hbar$  in the Schrödinger form to  $\rho_{(q,p)}$  requires a long discussion [20, Ex. 7.11].

We finish the discussion of the unitary dual of  $\mathbb{H}^n$  by a remark about negative values of  $\hbar$ . Since its position in the Heisenberg equation (4.10) a negative value of  $\hbar$  revert the flow of time. Thus representations  $\rho_\hbar$  with  $\hbar < 0$  seems to be suitable for a description of anti-particles. There is the explicit (cf. Figure 1) mirror symmetry between matter and anti-matter through classical mechanics. In this paper however we will consider only the case of  $\hbar > 0$ .

**2.3. Fock Spaces  $F_2(\mathcal{O}_\hbar)$  and Coherent States.** Our Fock type spaces (2.14) are not very different [25, Ex. 4.3] from the standard Segal–Bargmann spaces.

**Definition 2.5.** [11, 15] The Segal–Bargmann space (with a parameter  $\hbar > 0$ ) consists of functions on  $\mathbb{C}^n$  which are holomorphic in  $z$ , i.e.  $\partial_{\bar{z}_j} f(z) = 0$ , and square integrable with respect to the measure  $e^{-2|z|^2/\hbar} dz$  on  $\mathbb{C}^n$ :

$$\int_{\mathbb{C}^n} |f(z)|^2 e^{-2|z|^2/\hbar} dz < \infty.$$

Noticing the  $\partial_{\bar{z}_j}$  component in the operator  $D_\hbar^j$  (2.13) we obviously obtain

**Proposition 2.6.** *A function  $f_\hbar(q, p)$  is in  $F_2(\mathcal{O}_\hbar)$  (2.14) for  $\hbar > 0$  if and only if the function  $f_\hbar(z)e^{|z|^2/\hbar}$ ,  $z = p + iq$  is in the classical Segal–Bargmann space.*

The space  $F_2(\mathcal{O}_\hbar)$  can be also described in the language of *coherent states*, (also known as *wavelets*, *matrix elements* of representation, *Berezin transform*, etc., see [1, 25]). Since the representation  $\rho_\hbar$  is irreducible any vector  $v_0$  in  $F_2(\mathcal{O}_\hbar)$  is *cyclic*, i.e. vectors  $\rho_\hbar(g)v_0$  for all  $g \in G$  span the whole space  $F_2(\mathcal{O}_\hbar)$ . Even all vectors are equally good in principle, some of them are more equal for particular purposes (cf. Remark 2.4). For the harmonic oscillator the preferred vector is the dimensionless *vacuum state*:

$$v_0(q, p) = \exp\left(-\frac{2\pi}{\hbar} (c_1^{-1}q^2 + c_1p^2)\right), \quad (2.18)$$

which corresponds to the minimal level of energy. Here  $c_1$  as was defined before (2.13) has the dimensionality  $T/M$ . One can check directly the validity of

the both equation (2.14) and Convention 2.1 for (2.18), particularly the exponent is taken from a dimensionless pure number. Note also that  $v_0(q, p)$  is destroyed by the *annihilation operators* (sf. (2.11) and (2.13)):

$$A_h^j = d\rho_h(X_j + ic_1 Y_j) = \frac{\hbar}{2}(\partial_{p_j} - ic_1 \partial_{q_j}) + 2\pi(c_1 p_j - iq_j)I. \quad (2.19)$$

We introduce a dimensionless inner product on  $F(\mathcal{O}_h)$  by the formula:

$$\langle f_1, f_2 \rangle = \left(\frac{4}{\hbar}\right)^n \int_{\mathbb{R}^{2n}} f_1(q, p) \bar{f}_2(q, p) dq dp \quad (2.20)$$

With respect to this product the vacuum vector (2.18) is normalised:  $\|v_0\| = 1$ . For any observable  $A$  the formula

$$\langle Av_0, v_0 \rangle = \left(\frac{4}{\hbar}\right)^n \int_{\mathbb{R}^{2n}} Av_0(q, p) \bar{v}_0(q, p) dq dp$$

gives an expectation in the units of  $A$  since both the vacuum vector  $v_0(q, p)$  and the inner product (2.20) are dimensionless. The term  $\hbar^{-n}$  in (2.20) does not only normalise the vacuum and fix the dimensionality of the inner product; it is also related to the *Plancherel measure* [11, (1.61)], [33, Chap. 1, Th. 2.6] on the unitary dual of  $\mathbb{H}^n$ .

Elements  $(s, 0, 0)$  of the centre of  $\mathbb{H}^n$  trivially act in the representation  $\rho_h$  (2.9) as multiplication by scalars, e.g. any function is a common eigenvector of all operators  $\rho_h(s, 0, 0)$ . Thus the essential part [25, Defn. 2.5] of the operator  $\rho_h(s, x, y)$  is determined solely by  $(x, y) \in \mathbb{R}^{2n}$ . The *coherent states*  $v_{(x,y)}(q, p)$  are “left shifts” of the vacuum vector  $v_0(q, p)$  by operators (2.9):

$$\begin{aligned} v_{(x,y)}(q, p) &= \rho_h(0, x, y)v_0(q, p) \\ &= \exp\left(-2\pi i(qx + py) - \frac{2\pi}{\hbar}\left(c_1^{-1}\left(q - \frac{\hbar}{2}y\right)^2 + c_1\left(p + \frac{\hbar}{2}x\right)^2\right)\right). \end{aligned} \quad (2.21)$$

Now any function from the space  $F_2(\mathcal{O}_h)$  can be represented [25, Ex. 4.3] as a linear superposition of coherent states:

$$\begin{aligned} f(q, p) = [\mathcal{M}_h \check{f}](q, p) &= \hbar^n \int_{\mathbb{R}^{2n}} \check{f}(x, y)v_{(x,y)}(q, p) dx dy \\ &= \hbar^n \int_{\mathbb{R}^{2n}} \check{f}(x, y)\rho_h(x, y) dx dy v_{(0,0)}(q, p) \end{aligned} \quad (2.22)$$

where  $\check{f}(x, y)$  is the *wavelet* (or *coherent states*) *transform* [1, 25] of  $f(q, p)$ :

$$\begin{aligned} \check{f}(x, y) = [\mathcal{W}_h f](x, y) &= \langle f, v_{(x,y)} \rangle_{F_2(\mathcal{O}_h)} \\ &= \left(\frac{4}{\hbar}\right)^n \int_{\mathbb{R}^{2n}} f(q, p)\bar{v}_{(x,y)}(q, p) dq dp. \end{aligned} \quad (2.23)$$

The formula (2.22) can be regarded [25] as the *inverse wavelet transform*  $\mathcal{M}$  of  $\check{f}(x, y)$ . Note that all above integrals are dimensionless, thus both the wavelet transform and its inverse are measured in the same units.

The straightforward use of the basic formula:

$$\int_{-\infty}^{\infty} \exp(-ax^2 + bx + c) dx = \sqrt{\frac{\pi}{a}} \exp\left(\frac{b^2}{4a} + c\right), \quad \text{where } a > 0. \quad (2.24)$$

for the wavelet transform (2.22) leads to:

$$\check{v}_0(s, x, y) = \exp 2\pi \left( ihs - \frac{\hbar}{4} (c_1 x^2 + c_1^{-1} y^2) \right). \quad (2.25)$$

Since [25, Prop. 2.6] the wavelet transform  $\mathcal{W}_h$  (2.22) intertwines  $\rho_h$  (2.9) with the left regular representation  $\lambda_l$  (2.5):

$$\mathcal{W}_h \circ \rho_h(g) = \lambda_l(g) \circ \mathcal{W}_h, \quad \text{for all } g \in \mathbb{H}^n$$

the image of an arbitrary coherent state is:

$$\begin{aligned} \check{v}_{(s',x',y')}(s,x,y) &= \\ \exp 2\pi \left( ih \left( s - s' - \frac{1}{2}(x'y - xy') \right) - \frac{h}{4} (c_i(x-x')^2 + c_i^{-1}(y-y')^2) \right). \end{aligned} \quad (2.26)$$

Needless to say that these functions are obeying Convention 2.1.

We should mention however a problem related to coherent states (2.21): all their “classical limits” for  $h \rightarrow 0$  are functions with supports in neighbourhoods of  $(0,0)$ . In the contrast we may wish them be supported around different classical states  $(q,p)$ . This difficulty can be resolved through a replacement of the group action of  $\mathbb{H}^n$  in (2.21) by the “shifts” (4.8) generated by the  $p$ -mechanical brackets (4.3).

### 3. $p$ -MECHANICS: STATICS

We define  $p$ -mechanical observables to be convolutions on the Heisenberg group. The next Subsection describes their multiplication and commutator as well as quantum and classical representations. The Berezin quantisation in form of wavelet transform is considered in Subsection 3.2. This is developed in Subsection 3.3 into a construction of  $p$ -observables out of either quantum or classical ones.

**3.1. Observables in  $p$ -Mechanics, Convolutions and Commutators.** In line with the standard quantum theory we give the following definition:

**Definition 3.1.** *Observables* in  $p$ -mechanics ( $p$ -observables) are presented by operators on  $L_2(\mathbb{H}^n)$ .

Actually we will need here<sup>1</sup> only operators generated by convolutions on  $L_2(\mathbb{H}^n)$ . Let  $dg$  be a left invariant measure [19, § 7.1] on  $\mathbb{H}^n$ , which coincides with the standard Lebesgue measure on  $\mathbb{R}^{2n+1}$  in the exponential coordinates  $(s,x,y)$ . Then a function  $k_1$  from the linear space  $L_1(\mathbb{H}^n, dg)$  acts on  $k_2 \in L_2(\mathbb{H}^n, dg)$  by the convolution as follows:

$$\begin{aligned} (k_1 * k_2)(g) &= c_h^{n+1} \int_{\mathbb{H}^n} k_1(g_1) k_2(g_1^{-1}g) dg_1 \\ &= c_h^{n+1} \int_{\mathbb{H}^n} k_1(gg_1^{-1}) k_2(g_1) dg_1, \end{aligned} \quad (3.1)$$

where the constant  $c_h$  is measured in units of the action and can be assumed equal to 1. Then  $c_h^{n+1}$  has units inverse to  $dg$ . Thus the convolution  $k_1 * k_2$  is measured in units those are product of units for  $k_1$  and  $k_2$ . The composition of two convolution operators  $K_1$  and  $K_2$  with kernels  $k_1$  and  $k_2$  has the kernel defined by the same formula (3.1). Clearly two products  $K_1K_2$  and  $K_2K_1$  could have a different value due to non-commutativity of  $\mathbb{H}^n$  but always are measured in the same units. Thus we can find out how distinct they are from the difference  $K_1K_2 - K_2K_1$ , which does not violate the Convention 2.1. This also produces the *inner derivations*  $D_k$  of  $L_1(\mathbb{H}^n)$  by the *commutator*:

$$\begin{aligned} D_k : f \mapsto [k, f] &= k * f - f * k \\ &= c_h^{n+1} \int_{\mathbb{H}^n} k(g_1) (f(g_1^{-1}g) - f(gg_1^{-1})) dg_1. \end{aligned} \quad (3.2)$$

---

<sup>1</sup>More general operators are in use for a string-like version of  $p$ -mechanics, see Subsection 5.2.3.

Because we only consider observables those are convolutions on  $\mathbb{H}^n$  we can extend a unitary representation  $\rho_h$  of  $\mathbb{H}^n$  to a  $*$ -representation  $L_1(\mathbb{H}^n, dg)$  by the formula:

$$\begin{aligned} [\rho_h(k)f](q, p) &= c_h^{n+1} \int_{\mathbb{H}^n} k(g) \rho_h(g) f(q, p) dg \\ &= c_h^n \int_{\mathbb{R}^{2n}} \left( c_h \int_{\mathbb{R}} k(s, x, y) e^{-2\pi i h s} ds \right) \\ &\quad \times e^{-2\pi i (qx + py)} f \left( q - \frac{h}{2} y, p + \frac{h}{2} x \right) dx dy. \end{aligned} \quad (3.3)$$

The last formula in the Schrödinger representation defines for  $h \neq 0$  a *pseudodifferential operator* [11, 15, 32] on  $L_2(\mathbb{R}^n)$  (2.14), which are known to be *quantum observables* in the Weyl quantisation. For representations  $\rho_{(q,p)}$  (2.15) an expression analogous to (3.3) defines an operator of multiplication on  $\mathcal{O}_0$  (2.17) by the Fourier transform of  $k(s, x, y)$ :

$$\rho_{(q,p)}(k) = \hat{k}(0, q, p) = c_h^{n+1} \int_{\mathbb{H}^n} k(s, x, y) e^{-2\pi i (qx + py)} ds dx dy, \quad (3.4)$$

where the direct  $\hat{\cdot}$  and inverse  $\check{\cdot}$  Fourier transforms are defined by the formulae:

$$\hat{f}(v) = \int_{\mathbb{R}^m} f(u) e^{-2\pi i uv} du \quad \text{and} \quad f(u) = (\hat{f})\check{(u)} = \int_{\mathbb{R}^m} \hat{f}(v) e^{2\pi i vu} dv.$$

For reasons discussed in subsections 2.2 and 4.1 we regard the functions (3.4) on  $\mathcal{O}_0$  as *classical observables*. Again the both representations  $\rho_h(k)$  and  $\rho_{(q,p)}k$  are measured in the same units as the function  $k$  does.

From (3.3) follows that  $\rho_h(k)$  for a fixed  $h \neq 0$  depends only from  $\hat{k}_s(h, x, y)$ , which is the partial Fourier transform  $s \rightarrow h$  of  $k(s, x, y)$ . Then the representation of the composition of two convolutions depends only from

$$\begin{aligned} (k' * k)_s^\wedge &= c_h \int_{\mathbb{R}} e^{-2\pi i h s} c_h^{n+1} \int_{\mathbb{H}^n} k'(s', x', y') \\ &\quad \times k(s - s' + \frac{1}{2}(xy' - yx'), x - x', y - y') ds' dx' dy' ds \\ &= c_h^n \int_{\mathbb{R}^{2n}} e^{\pi i h (xy' - yx')} \cdot c_h \int_{\mathbb{R}} e^{-2\pi i h s'} k'(s', x', y') ds' \\ &\quad \times c_h \int_{\mathbb{R}} e^{-2\pi i h (s - s' + \frac{1}{2}(xy' - yx'))} \\ &\quad \times k(s - s' + \frac{1}{2}(xy' - yx'), x - x', y - y') ds dx' dy' \\ &= c_h^n \int_{\mathbb{R}^{2n}} e^{\pi i h (xy' - yx')} \hat{k}'_s(h, x', y') \hat{k}_s(h, x - x', y - y') dx' dy'. \end{aligned} \quad (3.5)$$

Note that if we apply the Fourier transform  $(x, y) \rightarrow (q, p)$  to the last expression (3.5) then we get the *star product* of  $\hat{k}'$  and  $\hat{k}$  known in *deformation* quantisation, cf. [36, (9)–(13)]. Consequently the representation  $\rho_h([k', k])$  of the commutator (3.2) depends only from:

$$\begin{aligned} [k', k]_s^\wedge &= c_h^n \int_{\mathbb{R}^{2n}} \left( e^{i\pi h (xy' - yx')} - e^{-i\pi h (xy' - yx')} \right) \\ &\quad \times \hat{k}'_s(-h, x', y') \hat{k}_s(-h, x - x', y - y') dx' dy' \\ &= 2ic_h^n \int_{\mathbb{R}^{2n}} \sin(\pi h (xy' - yx')) \hat{k}'_s(h, x', y') \hat{k}_s(h, x - x', y - y') dx' dy'. \end{aligned} \quad (3.6)$$

The integral (3.6) turns to be equivalent to the *Moyal brackets* [36] for the (full) Fourier transforms of  $k'$  and  $k$ . It is commonly accepted that the method of orbit is a mathematical side of the *geometric* quantisation [35]. Our derivation of the Moyal brackets in terms of orbits shows that deformation and geometric quantisations are closely connected and both are not very far from the original quantisation of Heisenberg and Schrödinger. Yet one more their close relative can be identified as the Berezin quantisation [4], see the next Subsection.

*Remark 3.2.* The expression (3.6) vanishes for  $h = 0$  as can be expected from the commutativity of representations (2.15). Thus it does not produce anything interesting on  $\mathcal{O}_0$ , that supports the common negligence to this set.

Summing up,  $p$ -mechanical observables, i.e. convolutions on  $L_2(\mathbb{H}^n)$ , are transformed

- (1) by representations  $\rho_h$  (2.9) into quantum observables (3.3) with the Moyal bracket (3.6) between them;
- (2) by representations  $\rho_{(q,p)}$  (2.15) into classical observables (3.4).

We did not get a meaningful brackets on classical observables yet, this will be done in Section 4.1.

**3.2. Berezin Quantisation and Wavelet Transform.** There is the following construction, known as the *Berezin quantisation* [3, 4], allowing us to assign a function to an operator (observable) and an operator to a function. The scheme is based on the construction of the *coherent states* and can be derived from different sources [29, 30]. We prefer the group-theoretic origin of Perelomov coherent states [30], which is realised in (2.21). Following [3] we introduce the *covariant* symbol  $a(g)$  of an operator  $A$  on  $F_2(\mathcal{O}_h)$  by the simple expression:

$$a(g) = \langle Av_g, v_g \rangle, \quad (3.7)$$

i.e. we get a map from the linear space of operators on  $F_2(\mathcal{O}_h)$  to a linear space of function on  $\mathbb{H}^n$ . A map in the opposite direction assigns to a function  $\check{a}(g)$  on  $\mathbb{H}^n$  the linear operator  $A$  on  $F_2(\mathcal{O}_h)$  by the formula

$$A = c_h^{n+1} \int_{\mathbb{H}^n} \mathring{a}(g) P_g dg, \quad \text{where } P_g \text{ is the projection } P_g v = \langle v, v_g \rangle v_g. \quad (3.8)$$

The function  $\mathring{a}(g)$  is called the *contravariant* symbol of the operator  $A$  (3.8).

The co- and contravariant symbols of operators are defined through the coherent states, in fact both types of symbols are realisations [25, § 3.1] of the direct (2.23) and inverse (2.22) wavelet transforms. Let us define a representation  $\rho_{bh}$  of the group  $\mathbb{H}^n \times \mathbb{H}^n$  in the space  $\mathcal{B}(F_2(\mathcal{O}_h))$  of operators on  $F_2(\mathcal{O}_h)$  by the formula:

$$\rho_{bh}(g_1, g_2) : A \mapsto \rho_h(g_1^{-1}) A \rho_h(g_2), \quad \text{where } g_1, g_2 \in \mathbb{H}^n. \quad (3.9)$$

According to the scheme from [25] for any state  $f_0$  on  $\mathcal{B}(F_2(\mathcal{O}_h))$  we get the wavelet transform  $\mathcal{W}_{f_0} : \mathcal{B}(F_2(\mathcal{O}_h)) \rightarrow C(\mathbb{H}^n \times \mathbb{H}^n)$ :

$$\mathcal{W}_{f_0} : A \mapsto \check{a}(g_1, g_2) = \langle \rho_{bh}(g_1, g_2) A, f_0 \rangle. \quad (3.10)$$

The important particular case is given by  $f_0$  defined through the vacuum vector  $v_0$  (2.18) by the formula  $\langle A, f_0 \rangle_{\mathcal{B}(F_2(\mathcal{O}_h))} = \langle Av_0, v_0 \rangle_{F_2(\mathcal{O}_h)}$ . Then the wavelet transform (3.10) produces the *covariant presymbol*  $\check{a}(g_1, g_2)$  of operator  $A$ . Its restriction  $a(g) = \check{a}(g, g)$  to the diagonal  $D$  of  $\mathbb{H}^n \times \mathbb{H}^n$  is exactly [25] the Berezin covariant symbol (3.7) of  $A$ . Such a restriction to the diagonal is done without a lost of information due to holomorphic properties of  $\check{a}(g_1, g_2)$  [3].

Another important example of the state  $f_0$  is given by the trace:

$$\langle A, f_0 \rangle = \text{tr } A = h^n \int_{\mathbb{R}^{2n}} \langle Av_{(x,y)}, v_{(x,y)} \rangle_{F_2(\mathcal{O}_h)} dx dy, \quad (3.11)$$

where coherent states  $v_{(x,y)}$  are again defined in (2.21). Operators  $\rho_{bh}(g, g)$  from the diagonal  $D$  of  $\mathbb{H}^n \times \mathbb{H}^n$  trivially act on the wavelet transform (3.10) generated by the trace (3.11) since the trace is invariant under  $\rho_{bh}(g, g)$ . According to the general scheme [25] we can consider *reduced wavelet transform* to the homogeneous space  $\mathbb{H}^n \times \mathbb{H}^n / D$  instead of the entire group  $\mathbb{H}^n \times \mathbb{H}^n$ . The space  $\mathbb{H}^n \times \mathbb{H}^n / D$  is isomorphic to  $\mathbb{H}^n$  with the embedding  $\mathbb{H}^n \rightarrow \mathbb{H}^n \times \mathbb{H}^n$  given by  $g \mapsto (g; 0)$ . Furthermore the centre  $Z$  of  $\mathbb{H}^n$  acts trivially in the representation  $\rho_{bh}$  as usual. Thus the only essential part of  $\mathbb{H}^n \times \mathbb{H}^n / D$  in the wavelet transform is the homogeneous space  $\Omega = \mathbb{H}^n / Z$ . A Borel section  $\mathbf{s} : \Omega \rightarrow \mathbb{H}^n \times \mathbb{H}^n$  in the principal bundle  $G \rightarrow \Omega$  can be defined as  $\mathbf{s} : (x, y) \mapsto ((0, x, y); (0, 0, 0))$ . We got the reduced realisation  $\mathcal{W}_r$  of the wavelet transform (3.10) in the form:

$$\begin{aligned} \mathcal{W}_r : A \mapsto \check{a}_r(x, y) &= \langle \rho_{bh}(\mathbf{s}(x, y))A, f_0 \rangle \\ &= \text{tr}(\rho_h((0, x, y)^{-1})A) \end{aligned} \quad (3.12)$$

$$\begin{aligned} &= h^n \int_{\mathbb{R}^{2n}} \langle \rho_h((0, x, y)^{-1})Av_{(x',y')}, v_{(x',y')} \rangle_{F_2(\mathcal{O}_h)} dx' dy' \\ &= h^n \int_{\mathbb{R}^{2n}} \langle Av_{(x',y')}, v_{(x,y) \cdot (x',y')} \rangle_{F_2(\mathcal{O}_h)} dx' dy'. \end{aligned} \quad (3.13)$$

The formula (3.12) is the principal ingredient of the *inversion formula* for the Heisenberg group [11, Chap. 1, (1.60)], [33, Chap. 1, Th. 2.7], which reconstructs kernels of convolutions  $k(g)$  out of operators  $\rho_h(k)$ . Therefore if we define a mother wavelet to be the identity operator  $I$  the inverse wavelet transform (cf. (2.22)) will be

$$\begin{aligned} \mathcal{M}_r a &= h^n \int_{\mathbb{R}^{2n}} a(x, y) \rho_{bh}(\mathbf{s}((0, x, y)^{-1})) I dx dy \\ &= h^n \int_{\mathbb{R}^{2n}} a(x, y) \rho_h(0, x, y) dx dy. \end{aligned} \quad (3.14)$$

The inversion formula for  $\mathbb{H}^n$  insures that

**Proposition 3.3.** *The composition  $\mathcal{M}_r \circ \mathcal{W}_r$  is the identity map on the representations  $\rho_h(k)$  of convolution operators on  $\mathcal{O}_h$ .*

**Example 3.4.** The wavelet transform  $\mathcal{W}_r$  (3.13) applied to the quantum coordinate  $Q = d\rho_h(X)$ , momentum  $P = d\rho_h(Y)$  (see (2.11)), and the energy function of the harmonic oscillator  $(c_1 Q^2 + c_2 P^2)/2$  produces the distributions on  $\mathbb{R}^{2n}$ :

$$\begin{aligned} Q &\mapsto \frac{1}{2\pi i} \delta^{(1)}(x) \delta(y), \\ P &\mapsto \frac{1}{2\pi i} \delta(x) \delta^{(1)}(y), \\ \frac{1}{2} (c_1 Q^2 + c_2 P^2) &\mapsto -\frac{1}{8\pi^2} \left( c_1 \delta^{(2)}(x) \delta(y) + c_2 \delta(x) \delta^{(2)}(y) \right), \end{aligned}$$

where  $\delta^{(1)}$  and  $\delta^{(2)}$  are the first and second derivatives of the Dirac *delta function*  $\delta$  respectively. The constants  $c_1$  and  $c_2$  have units  $M/T^2$  and  $1/M$  correspondingly. We will use these distributions later in Example 3.7.

**3.3. From Classical and Quantum Observables to  $p$ -Mechanics.** It is commonly accepted that we cannot deal with quantum mechanics directly and thus classical dynamics serves as an unavoidable intermediate step. A passage from classical observables to quantum ones—known as a *quantisation*—is the huge field with many concurring approaches (geometric, deformation, Weyl, Berezin, etc. quantisations) each having its own merits and demerits. Similarly one has to construct  $p$ -mechanical observables starting from classical or quantum ones by some procedure (should it be named “ $p$ -mechanisation”?), which we about to describe now.

The transition from a  $p$ -mechanical observable to the classical one is given by the formula (3.4), which in a turn is a realisation of the inverse wavelet transform (2.22):

$$\rho_{(q,p)}k = \hat{k}(0, q, p) = c_h^{n+1} \int_{\mathbb{H}^n} k(s, x, y) e^{-2\pi i(qx+py)} ds dx dy. \quad (3.15)$$

Similarly to a case of quantisation the classical image  $\rho_{(q,p)}k$  (3.15) contains only a partial information about  $p$ -observable  $k$  unless we make some additional assumptions. Let us start from a classical observable  $c(q, p)$  and construct the corresponding  $p$ -observable. From the general consideration (see [25] and Section 2.3) we can partially invert the formula (3.15) by the wavelet transform (2.23):

$$\check{c}(x, y) = [\mathcal{W}_0 c](x, y) = \langle cv_{(0,0)}, v_{(x,y)} \rangle = c_h^n \int_{\mathbb{R}^{2n}} c(q, p) e^{2\pi i(qx+py)} dq dp, \quad (3.16)$$

where  $v_{(x,y)} = \rho_{(q,p)}v_{(0,0)} = e^{-2\pi i(qx+py)}$ .

However the function  $\check{c}(x, y)$  (3.16) is not defined on the entire  $\mathbb{H}^n$ . The natural domain of  $\check{c}(x, y)$  according to the construction of the reduced wavelet transform [25] is the homogeneous space  $\Omega = G/Z$ , where  $G = \mathbb{H}^n$  and  $Z$  is its normal subgroup of central elements  $(s, 0, 0)$ . Let  $\mathbf{s} : \Omega \rightarrow G$  be a Borel section in the principal bundle  $G \rightarrow \Omega$ , which is used in the construction of induced representation, see [19, § 13.1]. For the Heisenberg group [25, Ex. 4.3] it can be simply defined as  $\mathbf{s} : (x, y) \in \Omega \mapsto (0, x, y) \in \mathbb{H}^n$ . One can naturally transfer functions from  $\Omega$  to the image  $\mathbf{s}(\Omega)$  of the map  $\mathbf{s}$  in  $G$ . However the range  $\mathbf{s}(\Omega)$  of  $\mathbf{s}$  has oftenly (particularly for  $\mathbb{H}^n$ ) a zero Haar measure in  $G$ . Probably two simplest possible ways out are:

- (1) To increase the “weight” of function  $\check{c}(s, x, y)$  vanishing outside of the range  $\mathbf{s}(\Omega)$  of  $\mathbf{s}$  by a suitable Dirac delta function on the subgroup  $Z$ . For the Heisenberg group this can be done, for example, by the map:

$$\mathcal{E} : \check{c}(x, y) \mapsto \tilde{c}(s, x, y) = \delta(s)\check{c}(x, y), \quad (3.17)$$

where  $\check{c}(x, y)$  is given by the inverse wavelet (Fourier) transform (3.16). As we will see in Proposition 3.6 this is related to the Weyl quantisation and the Moyal brackets.

- (2) To extend the function  $\check{c}(x, y)$  to the entire group  $G$  by a tensor product with a suitable function on  $Z$ , for example  $e^{-s^2}$ :

$$\check{c}(x, y) \mapsto \tilde{c}(s, x, y) = e^{-s^2}\check{c}(x, y).$$

In order to get the *correspondence principle* between classical and quantum mechanics (cf. Example 2.3) the function on  $Z$  has to satisfy some additional requirements. For  $\mathbb{H}^n$  it should vanish for  $s \rightarrow \pm\infty$ , which fulfils for both  $e^{-s^2}$  and  $\delta(s)$  from the previous item. In this way we get infinitely many essentially different quantisations with non-equivalent *deformed* Moyal brackets between observables.

There are other more complicated possibilities not mentioned here, which can be of some use if some additional information or assumptions are used to extend functions




FIGURE 2. The relations between:  
 $\mathcal{Q}_h$ —the Weyl quantisation from classical mechanics to quantum;  
 $\mathcal{C}_{h \rightarrow 0}$ —the classical limit  $h \rightarrow 0$  of quantum mechanics;  
 $\rho_h$  and  $\rho_{(q,p)}$ —unitary representations of Heisenberg group  $\mathbb{H}^n$ ;  
 $\mathcal{W}_r$  and  $\mathcal{W}_0$ —wavelet transforms defined in (3.12) and (3.16);  
 $\mathcal{E}$ —extension of functions from  $\Omega = \mathbb{H}^n/Z$  to the whole group  $\mathbb{H}^n$ .  
 Note the relations  $\mathcal{Q}_h = \rho_h \circ \mathcal{E} \circ \mathcal{W}_0$  and  $\mathcal{C}_{h \rightarrow 0} = \rho_{(q,p)} \circ \mathcal{E} \circ \mathcal{W}_r$ .

from  $\Omega$  to  $G$ . We will focus here only on the first “minimalistic” approach from the two listed above.

**Example 3.5.** The composition of the wavelet transform  $\mathcal{W}_0$  (3.16) and the map  $\mathcal{E}$  (3.17) applied to the classical coordinate, momentum, and the energy function of a harmonic oscillator produces the distributions on  $\mathbb{H}^n$ :

$$q \mapsto \frac{1}{2\pi i} \delta(s) \delta^{(1)}(x) \delta(y), \tag{3.18}$$

$$p \mapsto \frac{1}{2\pi i} \delta(s) \delta(x) \delta^{(1)}(y), \tag{3.19}$$

$$\frac{1}{2} (c_1 q^2 + c_2 p^2) \mapsto -\frac{1}{8\pi^2} \left( c_1 \delta(s) \delta^{(2)}(x) \delta(y) + c_2 \delta(s) \delta(x) \delta^{(2)}(y) \right), \tag{3.20}$$

where  $\delta^{(1)}$ ,  $\delta^{(2)}$ ,  $c_1$ , and  $c_2$  are defined in Example 3.4. We will use these distributions later in the Example 4.3.

If we apply the representation  $\rho_h$  (3.3) to the function  $\check{c}(s, x, y)$  (3.17) we will get the operator on  $F_2(\mathcal{O}_h)$ :

$$\begin{aligned} \mathcal{Q}_h(c) &= c_h^{n+1} \int_{\mathbb{H}^n} \check{c}(s, x, y) \rho_h(s, x, y) ds dx dy \\ &= c_h^{n+1} \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}} \delta(s) \check{c}(x, y) e^{s \cdot d\rho_h(S) + x \cdot d\rho_h(X) + y \cdot d\rho_h(Y)} ds dx dy \\ &= c_h^{n+1} \int_{\mathbb{R}} \delta(s) e^{-2\pi i s h} ds \int_{\mathbb{R}^{2n}} \check{c}(x, y) e^{x \cdot d\rho_h(X) + y \cdot d\rho_h(Y)} dx dy \\ &= c_h^n \int_{\mathbb{R}^{2n}} \check{c}(x, y) e^{x \cdot d\rho_h(X) + y \cdot d\rho_h(Y)} dx dy, \end{aligned} \tag{3.21}$$

where the last expression is exactly the *Weyl quantisation* (the *Weyl correspondence* [11, § 2.1]) if the Schrödinger realisation with  $d\rho_h(X) = q$  and  $d\rho_h(Y) = ih\partial_q$  on  $L_2(\mathbb{R}^n)$  is chosen for  $\rho_h$ . Thus we demonstrate that

**Proposition 3.6.** *The Weyl quantisation  $\mathcal{Q}_h$  (3.21) is the composition of the wavelet transform (3.16), the extension  $\mathcal{E}$  (3.17), and the representation  $\rho_h$  (2.9):*

$$\mathcal{Q}_h = \rho_h \circ \mathcal{E} \circ \mathcal{W}_0. \quad (3.22)$$

The similar construction can be carried out if we have a quantum observable  $A$  and wish to recover a related  $p$ -mechanical object. The wavelet transform  $\mathcal{W}_r$  (3.12) maps  $A$  into the function  $a(x, y)$  defined on  $\Omega$  and we again face the problem of extension of  $a(x, y)$  to the entire group  $\mathbb{H}^n$ . If it will be once more solved as in the classical case by the tensor product with the delta function  $\delta(s)$  then we get the following formula:

$$A \mapsto a(s, x, y) = \mathcal{E} \circ \mathcal{W}_r(A) = h^n \delta(s) \int_{\mathbb{R}^{2n}} \langle Av_{(x', y')}, v_{(x, y) \cdot (x', y')} \rangle_{F_2(\mathcal{O}_h)} dx' dy'.$$

We can apply to this function  $a(s, x, y)$  the representation  $\rho_{(q, p)}$  and obtain a classical observables  $\rho_{(q, p)}(a)$ . For a reasonable quantum observable  $A$  its classical image  $\rho_{(q, p)} \circ \mathcal{E} \circ \mathcal{W}_r(A)$  will coincide with its classical limit  $C_{h \rightarrow 0} A$ :

$$C_{h \rightarrow 0} = \rho_{(q, p)} \circ \mathcal{E} \circ \mathcal{W}_r, \quad (3.23)$$

which is expressed here through integral transformations and does not explicitly use any limit transition for  $h \rightarrow 0$ . The Figure 2 illustrates various transformations between quantum, classical, and  $p$ -observables. Besides the mentioned decompositions (3.22) and (3.23) there are presentations of identity maps on classical and quantum spaces correspondingly:

$$\mathcal{I}_c = \rho_{(q, p)} \circ \mathcal{E} \circ \mathcal{W}_0, \quad \mathcal{I}_h = \rho_h \circ \mathcal{E} \circ \mathcal{W}_h.$$

**Example 3.7.** The wavelet transform  $\mathcal{W}_r$  applied to the quantum coordinate  $Q$ , momentum  $P$ , and the energy function of a harmonic oscillator  $(c_1 Q^2 + c_2 P^2)/2$  was calculated in Example 3.4. The composition with the above map  $\mathcal{E}$  yields the distributions:

$$\begin{aligned} Q &\mapsto \frac{1}{2\pi i} \delta(s) \delta^{(1)}(x) \delta(y), \\ P &\mapsto \frac{1}{2\pi i} \delta(s) \delta^{(1)}(x) \delta(y), \\ \frac{1}{2} (c_1 Q^2 + c_2 P^2) &\mapsto -\frac{1}{2\pi^2} \left( c_1 \delta(s) \delta^{(2)}(x) \delta(y) + c_2 \delta(s) \delta(x) \delta^{(2)}(y) \right), \end{aligned}$$

which are exactly the same as in the Example 3.5.

#### 4. $p$ -MECHANICS: DYNAMICS

We introduce the  $p$ -mechanical brackets, which suit to all essential physical requirements and have a non-trivial classical representation coinciding with the Poisson brackets. A consistent  $p$ -mechanical dynamic equations is given in Subsection 4.2 and is analysed for the harmonic oscillator. Symplectic automorphisms of the Heisenberg group produce symplectic symmetries of  $p$ -mechanical, quantum, and classical dynamics in Subsection 4.3.

4.1.  **$p$ -Mechanical Brackets on  $\mathbb{H}^n$ .** Having observables as convolutions on  $\mathbb{H}^n$  we need a dynamic equation for their time evolution. To this end we seek a time derivative generated by an observable associated with energy.

*Remark 4.1.* The first candidate is the derivation coming from commutator (3.2). However the straight commutator has at least two failures:

- (1) It cannot produce any dynamics on  $\mathcal{O}_0$  (2.17), see Remark 3.2.
- (2) It violates the Convention 2.1 as indicated bellow.

As well known the classical energy is measured in  $ML^2/T^2$  so does the  $p$ -mechanical energy  $E$ . Consequently the commutator  $[E, \cdot]$  (3.2) with the  $p$ -energy has units  $ML^2/T^2$  whereas the time derivative should be measured in  $1/T$ , i.e. the mismatch is in the units of action  $ML^2/T$ .

Fortunately, there is a possibility to fix the both above defects of the straight commutator at the same time. Let us define a multiple  $\mathcal{A}$  of a right inverse operator to the vector field  $S$  (2.3) on  $\mathbb{H}^n$  by its actions on exponents—characters of the centre  $Z \in \mathbb{H}^n$ :

$$S\mathcal{A} = 4\pi^2 I, \quad \text{where} \quad \mathcal{A}e^{2\pi ihs} = \begin{cases} \frac{2\pi}{i\hbar} e^{2\pi ihs}, & \text{if } h \neq 0, \\ 4\pi^2 s, & \text{if } h = 0. \end{cases} \quad (4.1)$$

An alternative definition of  $\mathcal{A}$  as the convolution with a distribution is given in [27].

We can extend  $\mathcal{A}$  by the linearity to the entire space  $L_1(\mathbb{H}^n)$ . As a multiplier of a right inverse to  $S$  the operator  $\mathcal{A}$  is measured in  $T/(ML^2)$ —exactly that we need to correct the second of above mentioned defects of the straight commutator. Thus we introduce [27] the modified convolution operation  $\star$  on  $L_1(\mathbb{H}^n)$ :

$$k' \star k = (k' * k)\mathcal{A} \quad (4.2)$$

and the associated modified commutator ( $p$ -mechanical brackets):

$$\{[k', k]\} = [k', k]\mathcal{A} = k' \star k - k \star k'. \quad (4.3)$$

Obviously (4.3) is a bilinear antisymmetric form on the convolution kernels. It was also demonstrated in [27] that  $p$ -mechanical brackets satisfy to the Leibniz and Jacoby identities. They are all important for a consistent dynamics [8] along with the dimensionality condition given in the beginning of this Subsection.

From (3.3) one gets  $\rho_h(\mathcal{A}k) = \frac{2\pi}{i\hbar}\rho_h(k)$  for  $h \neq 0$ . Consequently the modification of the commutator for  $h \neq 0$  is only slightly different from the original one:

$$\rho_h \{[k', k]\} = \frac{1}{i\hbar} [\rho_h(k'), \rho_h(k)], \quad \text{where } \hbar = \frac{h}{2\pi} \neq 0. \quad (4.4)$$

The integral representation of the modified commutator kernel become (cf. (3.6)):

$$\{[k', k]\}_s^\wedge = c_h^n \int_{\mathbb{R}^{2n}} \frac{4\pi}{h} \sin(\pi h(xy' - yx')) \hat{k}'_s(h, x', y') \hat{k}_s(h, x - x', y - y') dx' dy', \quad (4.5)$$

where we can understand the expression under the integral as

$$\frac{4\pi}{h} \sin(\pi h(xy' - yx')) = 4\pi^2 \sum_{k=1}^{\infty} (-1)^{k+1} (\pi h)^{2(k-1)} \frac{(xy' - yx')^{2k-1}}{(2k-1)!} \quad (4.6)$$

This makes the operation (4.5) for  $h = 0$  significantly distinct from the vanishing integral (3.6). Indeed it is natural to assign the value  $4\pi^2(xy' - yx')$  to (4.6) for  $h = 0$ . Then the integral in (4.5) becomes the Poisson brackets for the Fourier transforms of  $k'$  and  $k$  defined on  $\mathcal{O}_0$  (2.17):

$$\rho_{(q,p)} \{[k', k]\} = \frac{\partial \hat{k}'(0, q, p)}{\partial q} \frac{\partial \hat{k}(0, q, p)}{\partial p} - \frac{\partial \hat{k}'(0, q, p)}{\partial p} \frac{\partial \hat{k}(0, q, p)}{\partial q}. \quad (4.7)$$

The same formula is obtained [27, Prop. 3.5] if we directly calculate  $\rho_{(q,p)} \{[k', k]\}$  rather than resolve the indeterminacy for  $h = 0$  in (4.6). This means the continuity of our construction at  $h = 0$  and represents the *correspondence principle* between quantum and classical mechanics.

We saw that the remedy of the second failure of commutator in Remark 4.1 (which was our duty according to Convention 2.1) by the antiderivative (4.1) improves the first defect as well (which is a very pleasant and surprising bonus). There are probably much simpler ways to fix the dimensionality of commutator “by hands”. However not all of them obviously would produce the Poisson brackets on  $\mathcal{O}_0$  as the antiderivative (4.1).

We arrived to the following observation: *Poisson brackets and inverse of the Planck constant  $1/h$  have the same dimensionality because they are image of the same object (anti-derivative (4.1)) under different representations (2.9) and (2.15) of the Heisenberg group.*

Note that functions  $X = \delta(s)\delta^{(1)}(x)\delta(y)$  and  $Y = \delta(s)\delta(x)\delta^{(1)}(y)$  (see (3.18) and (3.19)) on  $\mathbb{H}^n$  are measured in units  $L$  and  $ML^2/T$  (inverse to  $x$  and  $y$ ) correspondingly because they are respective derivatives of the dimensionless function  $\delta(s)\delta(x)\delta(y)$ . Then the  $p$ -mechanical brackets  $\{[X, \cdot]\}$  and  $\{[Y, \cdot]\}$  with these functions have dimensionality of  $T/(ML^2)$  and  $1/L$  correspondingly. Their representation  $\rho_* \{[X, \cdot]\}$  and  $\rho_* \{[Y, \cdot]\}$  (for both type of representations  $\rho_h$  and  $\rho_{(q,p)}$ ) are measured by  $L$  and  $ML^2/T$  and are simple derivatives:

$$\rho_* \{[X, \cdot]\} = \frac{\partial}{\partial p}, \quad \rho_* \{[Y, \cdot]\} = \frac{\partial}{\partial q}. \quad (4.8)$$

Thus  $\rho_* \{[X, \cdot]\}$  and  $\rho_* \{[Y, \cdot]\}$  are generators of shifts on both types of orbits  $\mathcal{O}_h$  and  $\mathcal{O}_0$  independent from value of  $h$ .

**4.2.  $p$ -Mechanical Dynamic Equation.** Since the modified commutator (4.3) with a  $p$ -mechanical energy has the dimensionality  $1/T$ —the same as the time derivative—we introduce the dynamic equation for an observable  $f(s, x, y)$  on  $\mathbb{H}^n$  based on that modified commutator as follows

$$\frac{df}{dt} = \{[f, E]\}. \quad (4.9)$$

*Remark 4.2.* It is a general tendency to make a Poisson brackets or quantum commutator out of any two observables and say that they form a Lie algebra. However there is a physical meaning to do that if at least one of two observables is an energy, coordinate or momentum: in these cases the brackets produce the time derivative (4.9) or corresponding shift generators (4.8) [16] of the other observable.

A simple consequence of the previous consideration is that the  $p$ -dynamic equation (4.9) is reduced

- (1) by the representation  $\rho_h$ ,  $h \neq 0$  (2.9) on  $F_2(\mathcal{O}_h)$  (2.7) to Moyal’s form of Heisenberg equation [36, (8)] based on the formulae (4.4) and (4.5):

$$\frac{d\rho_h(f)}{dt} = \frac{1}{i\hbar}[\rho_h(f), H_h], \quad \text{where the operator } H_h = \rho_h(E); \quad (4.10)$$

- (2) by the representations  $\rho_{(q,p)}$  (2.15) on  $\mathcal{O}_0$  (2.17) to Poisson’s equation [2, § 39] based on the formula (4.7):

$$\frac{d\hat{f}}{dt} = \{\hat{f}, H\} \quad \text{where the function } H(q, p) = \rho_{(q,p)}E = \hat{E}(0, q, p). \quad (4.11)$$

The same connections are true for the solutions of the three equations (4.9)–(4.11).




FIGURE 3. Dynamics of the harmonic oscillator in the adjoint space  $\mathfrak{h}_n^*$  is given by the identical linear symplectomorphisms of all orbits  $\mathcal{O}_h$  and  $\mathcal{O}_0$ . The vertical dotted string is uniformly rotating in the “horisontal” plane around the  $h$ -axis without any dynamics along the “vertical” direction.

**Example 4.3** (harmonic oscillator, of course ;-). [27] Let the  $p$ -mechanical energy function of a harmonic oscillator be as obtained in Examples 3.5 and 3.7:

$$E(s, x, y) = -\frac{1}{8\pi^2} \left( c_1 \delta(s) \delta^{(2)}(x) \delta(y) + c_2 \delta(s) \delta(x) \delta^{(2)}(y) \right), \quad (4.12)$$

Then the  $p$ -dynamic equation (4.9) on  $\mathbb{H}^n$  obeying the Convention 2.1 is

$$\frac{d}{dt} f(t; s, x, y) = \sum_{j=1}^n \left( c_2 x_j \frac{\partial}{\partial y_j} - c_1 y_j \frac{\partial}{\partial x_j} \right) f(t; s, x, y). \quad (4.13)$$

Solutions to the above equation is well known to be rotations in each of  $(x_j, y_j)$  planes given by:

$$f(t; s, x, y) = f_0 \left( s, x \cos(\sqrt{c_1 c_2} t) - \sqrt{\frac{c_1}{c_2}} y \sin(\sqrt{c_1 c_2} t), \right. \\ \left. \sqrt{\frac{c_2}{c_1}} x \sin(\sqrt{c_1 c_2} t) + y \cos(\sqrt{c_1 c_2} t) \right). \quad (4.14)$$

This expression respects the Convention 2.1. Since the dynamics on  $L_2(\mathbb{H}^n)$  is given by a symplectic linear transformation of  $\mathbb{H}^n$  its Fourier transform (2.10) to  $L_2(\mathfrak{h}_n^*)$  is the adjoint symplectic linear transformation of orbits  $\mathcal{O}_h$  and  $\mathcal{O}_0$  in  $\mathfrak{h}_n^*$ , see Figure 3.

The representation  $\rho_h$  transforms the energy function  $E$  (4.12) into the operator

$$H_h = -\frac{1}{8\pi^2} (c_1 Q^2 + c_2 P^2), \quad (4.15)$$

where  $Q = d\rho_h(X)$  and  $P = d\rho_h(Y)$  are defined in (2.11). The representation  $\rho_{(q,p)}$  transforms  $E$  into the classical Hamiltonian

$$H(q, p) = \frac{c_1}{2} q^2 + \frac{c_2}{2} p^2. \quad (4.16)$$

The  $p$ -dynamic equation (4.9) in form (4.13) is transformed by the representations  $\rho_h$  into the Heisenberg equation

$$\frac{d}{dt} f(t; Q, P) = \frac{1}{i\hbar} [f, H_h], \quad \text{where} \quad \frac{1}{i\hbar} [f, H_h] = c_1 p \frac{\partial f}{\partial q} - c_2 q \frac{\partial f}{\partial p}, \quad (4.17)$$




FIGURE 4. Automorphisms of  $\mathbb{H}^n$  generated by the symplectic group  $Sp(n)$  do not mix representations  $\rho_h$  with different Planck constants  $h$  and act by the metaplectic representation inside each of them. In the contrast those automorphisms of  $\mathbb{H}^n$  act transitively on the set of one-dimensional representations  $\rho_{(q,p)}$  joining them into the tangent space of the classical phase space  $\mathbb{R}^{2n}$ .

defined by the operator  $H_h$  (4.15). The representation  $\rho_{(q,p)}$  produces the Hamilton equation

$$\frac{d}{dt}f(t; q, p) = c_1 p \frac{\partial f}{\partial q} - c_2 q \frac{\partial f}{\partial p} \quad (4.18)$$

defined by the Hamiltonian  $H(q, p)$  (4.16). Finally, to get the solution for equations (4.17) and (4.18) it is enough to apply representations  $\rho_h$  and  $\rho_{(q,p)}$  to the solution (4.14) of  $p$ -dynamic equation (4.13).

Summing up we can rephrase the title of [36]: *quantum and classical mechanics live and work **together** on the Heisenberg group and are separated only in irreducible representations of  $\mathbb{H}^n$ .*

**4.3. Symplectic Invariance from Automorphisms of  $\mathbb{H}^n$ .** Let  $A : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$  be a linear *symplectomorphism* [2, § 41], [11, § 4.1], i.e. a map defined by  $2n \times 2n$  matrix:

$$A : \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix}$$

preserving the symplectic form (2.2):

$$\omega(A(x, y); A(x', y')) = \omega(x, y; x', y'). \quad (4.19)$$

All such transformations form the symplectic group  $Sp(n)$ . The Convention 2.1 implies that sub-blocks  $a$  and  $d$  of  $A$  have to be dimensionless while  $b$  and  $c$  have to be of reciprocal dimensions  $M/T$  and  $T/M$  respectively.

It follows from the identities (4.19) and (2.1) that the linear transformation  $\alpha : \mathbb{H}^n \rightarrow \mathbb{H}^n$  such that  $\alpha(s, x, y) = (s, A(x, y))$  is an automorphism of  $\mathbb{H}^n$ . Let us also denote by  $\tilde{\alpha} = \tilde{\alpha}_A$  a unitary transformation of  $L_2(\mathbb{H}^n)$  in the form

$$\tilde{\alpha}(f)(s, x, y) = \sqrt{\det a} f(s, A(x, y)),$$

which is well defined [11, § 4.2] on the double cover  $\tilde{Sp}(n)$  of the group  $Sp(n)$ . The correspondence  $A \mapsto \tilde{\alpha}_A$  is a linear unitary representation of the symplectic group

in  $L_2(\mathbb{H}^n)$ . One can also check the intertwining property

$$\lambda_{l(r)}(g) \circ \tilde{\alpha} = \tilde{\alpha} \circ \lambda_{l(r)}(\alpha(g)) \quad (4.20)$$

for the left (right) regular representations (2.5) of  $\mathbb{H}^n$ .

Because  $\alpha$  is an automorphism of  $\mathbb{H}^n$  the map  $\alpha^* : k(g) \mapsto k(\alpha(g))$  is an automorphism of the convolution algebra  $L_1(\mathbb{H}^n)$  with the multiplication  $*$  (3.1), i.e.  $\alpha^*(k_1) * \alpha^*(k_2) = \alpha^*(k_1 * k_2)$ . Moreover  $\alpha^*$  commutes with the antiderivative  $\mathcal{A}$  (4.1), thus  $\alpha^*$  is an automorphism of  $L_1(\mathbb{H}^n)$  with the modified multiplication  $\star$  (4.2) as well, i.e.  $\alpha^*(k_1) \star \alpha^*(k_2) = \alpha^*(k_1 \star k_2)$ . By the linearity we can extend the intertwining property (4.20) to the convolution operator  $K$  as follows:

$$\alpha^* K \circ \tilde{\alpha} = \tilde{\alpha} \circ K. \quad (4.21)$$

Since  $\alpha$  is an automorphism of  $\mathbb{H}^n$  it fixes the unit  $e$  of  $\mathbb{H}^n$  and its differential  $d\alpha : \mathfrak{h}^n \rightarrow \mathfrak{h}^n$  at  $e$  is given by the same matrix as  $\alpha$  in the exponential coordinates. Obviously  $d\alpha$  is an automorphism of the Lie algebra  $\mathfrak{h}^n$ . By the duality between  $\mathfrak{h}^n$  and  $\mathfrak{h}_n^*$  we obtain the adjoint map  $d\alpha^* : \mathfrak{h}_n^* \rightarrow \mathfrak{h}_n^*$  defined by the expression

$$d\alpha^* : (h, q, p) \mapsto (h, A^t(q, p)), \quad (4.22)$$

where  $A^t$  is the transpose of  $A$ . Obviously  $d\alpha^*$  preserves any orbit  $\mathcal{O}_h$  (2.7) and maps the orbit  $\mathcal{O}_{(q,p)}$  (2.8) to  $\mathcal{O}_{A^t(q,p)}$ .

Identity (4.22) indicates that both representations  $\rho_h$  and  $(\rho_h \circ \alpha)(s, x, y) = \rho_h(s, A(x, y))$  for  $h \neq 0$  correspond to the same orbit  $\mathcal{O}_h$ . Thus they should be equivalent, i.e. there is an intertwining operator  $U_A : F_2(\mathcal{O}_h) \rightarrow F_2(\mathcal{O}_h)$  such that  $U_A^{-1} \rho_h U_A = \rho_h \circ \alpha$ . Then the correspondence  $\sigma : A \mapsto U_A$  is a linear unitary representation of the double cover  $\widetilde{Sp}(n)$  of the symplectic group called the *metaplectic representation* [11, § 4.2], [13].

Thus we have

**Proposition 4.4.** *The  $p$ -mechanical brackets are invariant under the symplectic automorphisms of  $\mathbb{H}^n$ :  $\{[\tilde{\alpha}k_1, \tilde{\alpha}k_2]\} = \tilde{\alpha} \{[k_1, k_2]\}$ . Consequently the dynamic equation (4.9) has symplectic symmetries which are reduced*

- (1) by  $\rho_h$ ,  $h \neq 0$  on  $\mathcal{O}_h$  (2.7) to the metaplectic representation in quantum mechanics;
- (2) by  $\rho_{(q,p)}$  on  $\mathcal{O}_0$  (2.17) to the symplectic symmetries of classical mechanics [2, § 38].

Combining intertwining properties of all three components (3.22) in the Weyl quantisation we get

**Corollary 4.5.** *The Weyl quantisation  $\mathcal{Q}_h$  (3.21) is the intertwining operator between classical and metaplectic representations.*

## 5. CONCLUSIONS

**5.1. Discussion.** Our intention is to demonstrate that the complete representation theory of the Heisenberg group  $\mathbb{H}^n$ , which includes one-dimensional commutative representations, is a sufficient language for *both* classical and quantum theory.

It is natural to describe the complete set of unitary irreducible representations by the orbit method of Kirillov. The analysis carried out in Subsection 2.2 and illustrated in Figure 1 shows that the position of one-dimensional representations  $\rho_{(q,p)}$  within the unitary dual of  $\mathbb{H}^n$  relates them to classical mechanics. Various connections of infinite dimensional representations  $\rho_h$  of  $\mathbb{H}^n$  to quantum mechanics has been known for a long time.

Convolutions operators on  $\mathbb{H}^n$  is a natural class to be associated with physical observables. They are reduced by infinite dimensional representations  $\rho_h$  to the

pseudodifferential operators, which are observables in the Weyl quantisation. The one-dimensional representations  $\rho_{(q,p)}$  map convolutions into classical observables—functions on the phase space. The wavelets technique allows us to transform these three types of observables into each other, which is illustrated by Figure 2.

A nontrivial dynamics in the phase space—the space of one-dimensional representations of  $\mathbb{H}^n$ —could be obtained from the commutator on  $\mathbb{H}^n$  with a help of the anti-derivative operator  $\mathcal{A}$  (4.1). The  $p$ -mechanical dynamic equation (4.9) based on the operator  $\mathcal{A}$  possesses all desirable properties for description of a physical time evolution and its solution gives both classic and quantum dynamics. See Figure 3 for the familiar dynamics of the harmonic oscillator.

Finally, the symplectic automorphisms of the Heisenberg group preserve the dynamic equation (4.9) and all its solutions. In representations of the Heisenberg group this reduces to the symplectic invariance of classical mechanics and the metaplectic invariance of the quantum description. Moreover the symplectic transformations act transitively on the set  $\mathcal{O}_0$  (2.17) of one-dimensional representations supporting its  $p$ -mechanical interpretation as the classical phase space, see Figure 4.

**5.2. Further Developments.** The present paper deals only with elementary aspects of  $p$ -mechanics. Notion of physical states in  $p$ -mechanics is considered in [6, 7], where its usefulness for a forced oscillator is demonstrated. Paper [7] discusses also connection of  $p$ -mechanics and *contextual interpretation* [18]. Our study is a part of the Erlangen-type approach [24, 26] in non-commutative geometry. It could be extended in several directions:

**5.2.1. Quantum-Classical Interaction.** The long standing discussion [8, 31] about quantum-classical interaction can be treated as follows. Let  $\mathbb{B}$  be a nilpotent step two Heisenberg-like group of elements  $(s_1, s_2; x_1, y_1; x_2, y_2)$  with the only non-trivial commutators in the Lie algebra (cf. (2.4)) as follows:

$$[X_i, Y_j] = \delta_{ij} S_i.$$

Thus  $\mathbb{B}$  has the two dimensional centre  $(s_1, s_2, 0, 0, 0, 0)$  and the adjoint space of characters of  $\mathbb{B}$  is also two dimensional. We can regard it as being spanned by two different Planck constants  $h_1$  and  $h_2$ . There is a possibility to study the case  $h_1 \neq 0$  and  $h_2 = 0$ , which correspond to a quantum behaviour of coordinates  $(x_1, y_1)$  and a classical dynamics in  $(x_2, y_2)$ . This study was initiated in [31] but oversaw some homological aspects of the construction and is not satisfactory completed yet.

**5.2.2. Quantum Field Theory.** Mathematical formalism of quantum mechanics uses complex numbers in order to provide unitary infinite dimensional representations of the Heisenberg group  $\mathbb{H}^n$ . In a similar way the De Donder–Weyl formalism for classical field theories [17] requires *Clifford numbers* [14] for their quantisation. It was recently realised [9] that the appearance of Clifford algebras is induced by the Galilean group—a nilpotent step two Lie group with multidimensional centre. In the one-dimensional case an element of the Galilean group is  $(s_1, \dots, s_n, x, y_1, \dots, y_n)$  with corresponding the Lie algebra described by the non-vanishing commutators:

$$[X, Y_j] = S_j, \quad j = 1, 2, \dots, n.$$

This corresponds to several momenta  $y_1, y_2, \dots, y_n$  adjoint to a single field coordinate  $x$  [17]. For field theories it is worth [28] to consider Clifford valued representations induced by Clifford valued “characters”  $e^{2\pi i(e_1 h_1 s_1 + \dots + e_n h_n s_n)}$  of the centre, where  $e_1, \dots, e_n$  are imaginary units spanning the Clifford algebra. The associated Fock spaces were described in [9]. In [28] we quantise the De Donder–Weyl field equations (similarly to our consideration in Subsection 4.1) with the help of

composed antiderivative operator  $\mathcal{A} = \sum_1^n e_i \mathcal{A}_i$ , where  $S_i \mathcal{A}_i = 4\pi^2 I$ . There are important mathematical and physical questions related to the construction, notably the rôle of the Dirac operator [12], which deserve further careful considerations.

5.2.3. *String Theory.* There is a possibility to use  $p$ -mechanical picture for a string-like theory. Indeed the  $p$ -dynamics of a harmonic oscillator as presented in the Example 4.3 and Figure 3 consists of the uniform rotation of lines around the  $h$ -axis—one can say *strings*—with the same  $(q, p)$  coordinates but different values of the Planck constant  $h$ .

In case of a more general energy, which is still however given by a *convolution* on  $\mathbb{H}^n$ , the dynamics can be more complicated. For example, it may not correspond to a point transformation of the adjoint space  $\mathfrak{h}_n^*$ . Alternatively a generic point transformation may transform a straight line  $(h, q_0, p_0)$  with fixed  $(q_0, p_0) \in \mathbb{R}^{2n}$  and variable  $h$  into a generic curve transversal to all  $(q, p)$ -planes. However all spaces  $F_2(\mathcal{O}_h)$  are invariant under any  $p$ -dynamics generated by a convolution on  $\mathbb{H}^n$ .

However if an energy is given by an arbitrary operator on  $L_2(\mathbb{H}^n)$  [10, 22] then spaces  $F_2(\mathcal{O}_h)$  for different  $h$  are no longer invariant during the evolution and could be mixed together. This opens a possibility of longitudinal dynamics of strings along the  $h$ -axis as well. It may seem strange to have a dynamics along  $h$  which is a *constant*, not a variable. However there is a duality [34] between the “Planck constant”  $h$  and the “tension of string”  $\alpha'$ . Dualities and symmetries between  $h$  and  $\alpha'$  can be reflected in a dynamics which mixes spaces  $F_2(\mathcal{O}_h)$  with different  $h$ .

#### ACKNOWLEDGEMENTS

I am grateful to A. Brodlie for many useful discussions and comments. Anonymous referees made numerous critical remarks and suggestions, which resulted into the paper’s improvements.

#### REFERENCES

1. Syed Twareque Ali, Jean-Pierre Antoine, and Jean-Pierre Gazeau, *Coherent states, wavelets and their generalizations*, Graduate Texts in Contemporary Physics, Springer-Verlag, New York, 2000. MR **2002m**:81092
2. V. I. Arnol’d, *Mathematical methods of classical mechanics*, Graduate Texts in Mathematics, vol. 60, Springer-Verlag, New York, 1991, Translated from the 1974 Russian original by K. Vogtmann and A. Weinstein, Corrected reprint of the second (1989) edition. MR **96c**:70001
3. F. A. Berezin, *Covariant and contravariant symbols of operators*, Izv. Akad. Nauk SSSR Ser. Mat. **36** (1972), 1134–1167, Reprinted in [5, pp. 228–261]. MR 50 #2996
4. ———, *General concept of quantization*, Comm. Math. Phys. **40** (1975), 153–174. MR 53 #15186
5. ———, *Metod vtorichnogo kvantovaniya*, second ed., “Nauka”, Moscow, 1986, Edited and with a preface by M. K. Polivanov. MR **89c**:81001
6. Alastair Brodlie, *Classical and quantum coherent states*, Internat. J. Theoret. Phys. (2003), 20 p., E-print: [arXiv:quant-ph/0303142](https://arxiv.org/abs/quant-ph/0303142) (To appear).
7. Alastair Brodlie and Vladimir V. Kisil, *Observables and sates in p-mechanics*, to appear, Advances in Mathematics Research, Nova Sci., 2003, E-print: [arXiv:quant-ph/0304023](https://arxiv.org/abs/quant-ph/0304023).
8. J. Caro and L. L. Salcedo, *Impediments to mixing classical and quantum dynamics*, Phys. Rev. **A60** (1999), 842–852, E-print: [arXiv:quant-ph/9812046](https://arxiv.org/abs/quant-ph/9812046).
9. Jan Cnops and Vladimir V. Kisil, *Monogenic functions and representations of nilpotent Lie groups in quantum mechanics*, Mathematical Methods in the Applied Sciences **22** (1998), no. 4, 353–373, E-print: [arXiv:math/9806150](https://arxiv.org/abs/math/9806150). MR **2000b**:81044. **Zbl** # 1005.22003.
10. A. S. Dynin, *Pseudodifferential operators on the Heisenberg group*, Dokl. Akad. Nauk SSSR **225** (1975), no. 6, 1245–1248. MR 54 #11410
11. Gerald B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies, vol. 122, Princeton University Press, Princeton, NJ, 1989. MR **92k**:22017
12. Ezra Getzler, *A short proof of the local Atiyah-Singer index theorem*, Topology **25** (1986), no. 1, 111–117. MR **87h**:58207

13. K. C. Hannabuss, *Characters and contact transformations*, Math. Proc. Cambridge Philos. Soc. **90** (1981), no. 3, 465–476. MR **83e**:81035
14. David Hestenes, *New foundations for classical mechanics*, second ed., Kluwer Academic Publishers Group, Dordrecht, 1999. MR **2001h**:70004
15. Roger Howe, *Quantum mechanics and partial differential equations*, J. Funct. Anal. **38** (1980), no. 2, 188–254. MR **83b**:35166
16. R. L. Hudson and S. N. Peck, *Canonical Fourier transforms*, J. Math. Phys. **20** (1979), no. 1, 114–119. MR **81j**:46107
17. Igor V. Kanatchikov, *Precanonical quantum gravity: quantization without the space-time decomposition*, Internat. J. Theoret. Phys. **40** (2001), no. 6, 1121–1149, E-print: [arXiv:gr-qc/0012074](https://arxiv.org/abs/gr-qc/0012074). MR **2002m**:83038
18. Andrei Khrennikov, *Växjö interpretation of quantum mechanics*, (2002), E-print: [arXiv:quant-ph/0202107](https://arxiv.org/abs/quant-ph/0202107).
19. A. A. Kirillov, *Elements of the theory of representations*, Springer-Verlag, Berlin, 1976, Translated from the Russian by Edwin Hewitt, Grundlehren der Mathematischen Wissenschaften, Band 220. MR 54 #447
20. ———, *Introduction to the theory of representations and noncommutative harmonic analysis* [ MR **90a**:22005], Representation Theory and Noncommutative Harmonic Analysis, I, Springer, Berlin, 1994, MR 1311488., pp. 1–156, 227–234. MR 1 311 488
21. ———, *Merits and demerits of the orbit method*, Bull. Amer. Math. Soc. (N.S.) **36** (1999), no. 4, 433–488. MR **2000h**:22001
22. Vladimir V. Kisil, *Local algebras of two-sided convolutions on the Heisenberg group*, Mat. Zametki **59** (1996), no. 3, 370–381, 479, MR **97h**:22006.
23. ———, *Plain mechanics: Classical and quantum*, J. Natur. Geom. **9** (1996), no. 1, 1–14, MR **96m**:81112. E-print: [arXiv:funct-an/9405002](https://arxiv.org/abs/funct-an/9405002).
24. ———, *Two approaches to non-commutative geometry*, Complex Methods for Partial Differential Equations (H. Begehr, O. Celebi, and W. Tutschke, eds.), Kluwer Academic Publishers, Netherlands, 1999, E-print: [arXiv:funct-an/9703001](https://arxiv.org/abs/funct-an/9703001), MR **2001a**:01002, pp. 219–248.
25. ———, *Wavelets in Banach spaces*, Acta Appl. Math. **59** (1999), no. 1, 79–109, E-print: [arXiv:math/9807141](https://arxiv.org/abs/math/9807141). MR **2001c**:43013.
26. ———, *Meeting Descartes and Klein somewhere in a noncommutative space*, Highlights of Mathematical Physics (A. Fokas, J. Halliwell, T. Kibble, and B. Zegarlinski, eds.), AMS, 2002, E-print: [arXiv:math-ph/0112059](https://arxiv.org/abs/math-ph/0112059), pp. 165–189.
27. ———, *Quantum and classical brackets*, Internat. J. Theoret. Phys. **41** (2002), no. 1, 63–77, E-print: [arXiv:math-ph/0007030](https://arxiv.org/abs/math-ph/0007030). MR **2003b**:81105
28. ———, *p-Mechanics and De Donder–Weyl theory*, The Fifth International Conference “Symmetry in Nonlinear Mathematical Physics”, Inst. of Math., NAS of Ukraine, 2003, E-print: [arXiv:quant-ph/0306101](https://arxiv.org/abs/quant-ph/0306101) (To appear).
29. John R. Klauder, *Coherent states and coordinate-free quantization*, Internat. J. Theoret. Phys. **33** (1994), no. 3, 509–522. MR **95a**:81105
30. A. Perelomov, *Generalized coherent states and their applications*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1986. MR **87m**:22035
31. Oleg V. Prezhdo and Vladimir V. Kisil, *Mixing quantum and classical mechanics*, Phys. Rev. A (3) **56** (1997), no. 1, 162–175, MR **99j**:81010. E-print: [arXiv:quant-ph/9610016](https://arxiv.org/abs/quant-ph/9610016).
32. M. A. Shubin, *Pseudodifferential operators and spectral theory*, second ed., Springer-Verlag, Berlin, 2001, Translated from the 1978 Russian original by Stig I. Andersson. MR **2002d**:47073
33. Michael E. Taylor, *Noncommutative harmonic analysis*, Mathematical Surveys and Monographs, vol. 22, American Mathematical Society, Providence, RI, 1986. MR **88a**:22021
34. Edward Witten, *Reflections on the fate of spacetime*, Phys. Today **49** (1996), no. 4, 24–30. MR **97i**:81003
35. N. M. J. Woodhouse, *Geometric quantization*, second ed., Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1992, Oxford Science Publications. MR **94a**:58082
36. Cosmas Zachos, *Deformation quantization: quantum mechanics lives and works in phase-space*, Internat. J. Modern Phys. A **17** (2002), no. 3, 297–316, E-print: [arXiv:hep-th/0110114](https://arxiv.org/abs/hep-th/0110114). MR 1 888 937

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UK

E-mail address: [kisilv@maths.leeds.ac.uk](mailto:kisilv@maths.leeds.ac.uk)

URL: <http://maths.leeds.ac.uk/~kisilv/>