UNIVERSITY of York

This is a repository copy of *Enhanced ozone loss by active inorganic bromine chemistry in the tropical troposphere*.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/112138/</u>

Version: Accepted Version

Article:

Breton, Michael Le, Bannan, Thomas J., Shallcross, Dudley E. et al. (14 more authors) (2017) Enhanced ozone loss by active inorganic bromine chemistry in the tropical troposphere. Atmospheric Environment. pp. 1-20. ISSN 1352-2310

https://doi.org/10.1016/j.atmosenv.2017.02.003

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Accepted Manuscript

Enhanced ozone loss by active inorganic bromine chemistry in the tropical troposphere

Michael Le Breton, Thomas J. Bannan, Dudley E. Shallcross, M. Anwar Khan, Mathew J. Evans, James Lee, Richard Lidster, Stephen Andrews, Lucy J. Carpenter, Johan Schmidt, Daniel Jacob, Neil R.P. Harris, Stephane Bauguitte, Martin Gallagher, Asan Bacak, Kimberley E. Leather, Carl J. Percival



DOI: 10.1016/j.atmosenv.2017.02.003

Reference: AEA 15176

To appear in: Atmospheric Environment

Received Date: 25 November 2016

Revised Date: 26 January 2017

Accepted Date: 2 February 2017

Please cite this article as: Le Breton, M., Bannan, T.J., Shallcross, D.E., Khan, M.A., Evans, M.J., Lee, J., Lidster, R., Andrews, S., Carpenter, L.J., Schmidt, J., Jacob, D., Harris, N.R.P., Bauguitte, S., Gallagher, M., Bacak, A., Leather, K.E., Percival, C.J., Enhanced ozone loss by active inorganic bromine chemistry in the tropical troposphere, *Atmospheric Environment* (2017), doi: 10.1016/j.atmosenv.2017.02.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



- 1 Enhanced ozone loss by active inorganic bromine chemistry in the tropical troposphere
- 2 Michael Le Breton^{1a}, Thomas J. Bannan¹, Dudley E. Shallcross², M. Anwar Khan², Mathew
- 3 J. Evans³, James Lee³, Richard Lidster³, Stephen Andrews³, Lucy J. Carpenter³, Johan
- 4 Schmidt⁴, Daniel Jacob⁴, Neil R. P. Harris⁵, Stephane Bauguitte⁶, Martin Gallagher¹, Asan
- 5 Bacak¹, Kimberley E. Leather¹, Carl J. Percival^{1b}
- 6 ¹The Centre for Atmospheric Science, School of Earth, Atmospheric and Environmental Sciences, University of
- 7 Manchester, Simon Building, Brunswick Street, Manchester, M13 9PL, UK
- 8 ²School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, UK
- 9 ³National Centre for Atmospheric Science (NCAS), Department of Chemistry, University of York, York, YO10 5DD, UK
- ⁴Harvard University, School of Engineering and Applied Sciences, 29 Oxford St, Cambridge, MA02138, USA
- 11 And Department of Chemistry, University of Copenhagen, 2100 Copenhagen, Denmark
- 12 ⁵Department of Chemistry, University of Cambridge, Cambridge, UK
- ⁵*Facility for Airborne Atmospheric Measurements (FAAM), Building 125, Cranfield University, Cranfield, Bedford,*
- 14 MK43 OAL, UK
- ^a now at Department of Chemistry and Molecular Biology, Medicinaregatan 9 C, 40530, Gothenburg, Sweden
- ^b now at Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, CA 91109.
- 18 *Authors to whom correspondence should be addressed. Email carl.j.percival@jpl.nasa.gov
- 19 d.e.shallcross@bristol.ac.uk
- 20

17

21 Abstract: Bromine chemistry, particularly in the tropics, has been suggested to play an important 22 role in tropospheric ozone loss (Theys et al., 2011)) although a lack of measurements of active bromine species impedes a quantitative understanding of its impacts. Recent modelling and 23 measurements of bromine monoxide (BrO) by Wang et al. (2015) have shown current models 24 under predict BrO concentrations over the Pacific Ocean and allude to a missing source of BrO. 25 26 Here, we present the first simultaneous aircraft measurements of atmospheric bromine monoxide, 27 BrO (a radical that along with atomic Br catalytically destroys ozone) and the inorganic Br 28 precursor compounds HOBr, BrCl and Br_2 over the Western Pacific Ocean from 0.5 to 7 km. The presence of 0.17-1.64 pptv BrO and 3.6-8 pptv total inorganic Br from these four species 29 30 throughout the troposphere causes 10-20% of total ozone loss, and confirms the importance of 31 bromine chemistry in the tropical troposphere; contributing to a 6 ppb decrease in ozone levels due to halogen chemistry. Observations are compared with a global chemical transport model and find 32

that the observed high levels of BrO, BrCl and HOBr can be reconciled by active multiphase
oxidation of halide (Br⁻ and Cl⁻) by HOBr and ozone in cloud droplets and aerosols. Measurements
indicate that 99% of the instantaneous free Br in the troposphere up to 8 km originates from
inorganic halogen photolysis rather than from photolysis of organobromine species.

37 Keywords

38 BrO, CIMS, ozone, troposphere

39 **1. Introduction**

Ozone (O₃) is one of the most important short-lived gases contributing to greenhouse radiative 40 forcing (RF) and plays a critical role in atmospheric chemistry and air quality (Crutzen et al., 41 1988). Therefore, it is vital to determine the mechanisms that control its regional and global levels. 42 In the troposphere O₃ has two sources: in situ photochemical production from anthropogenic 43 44 precursors such as volatile organic compounds (VOCs) and nitrogen oxides, and transport from the stratosphere (Liu et al., 2002). Subsequently, the temporal and spatial resolution of O₃ is rather 45 complex and our current understanding cannot account for the observed O₃ variation in the tropical 46 47 troposphere (Prather et al., 2001).

48

The primary loss route of tropospheric ozone is photolysis in the presence of water vapour. 49 However, recent studies have indicated that reactive halogens in the troposphere could make an 50 51 important contribution to photochemical O_3 depletion. Especially in tropical regions where higher 52 concentrations of halogenated species (assuming a biological source) are found due to warmer 53 waters, higher biological activity, and fast convective transport into the free troposphere (FT) (Liang et al., 2010, Holmes et al., 2010, Parella et al., 2012, Wang et al., 2015). Inorganic bromine 54 radicals (BrO and Br) are known to be efficient catalysts for O₃ destruction. 55 Sources of tropospheric bromine are poorly constrained (Parella et al., 2012) and include photochemical 56 57 breakdown of halocarbons (e.g. CHBr₃ and CH₂Br₂), debromination of sea salt aerosol, and input of inorganic bromine (Bry) from the stratosphere (Read et al., 2008). Model calculations have 58 59 indicated that bromine chemistry may be responsible for a reduction in the zonal mean O_3 mixing ratio by up to 18% and locally even up to 40% (Holmes et al., 2006) and account for the majority 60 of global oxidation of elemental mercury (Read et al., 2008). 61

62

63 Only a few pptv (parts per trillion by volume) of BrO are required to have a significant impact on 64 tropospheric chemistry (Holmes *et al.*, 2006). There is recent observational evidence for the

ubiquitous presence of BrO at these levels in the global troposphere, although reported BrO 65 background levels vary widely (Volkammer et al., 2015, Yang et al., 2005, Wang et al., 2015, 66 Mahajan et al., 2010) whilst global models under predict these levels of BrO. Wang et al. (2015) 67 have recently measured and modelled BrO over the tropical Eastern Pacific Ocean and shown 68 69 concentrations are up to 2-4 times greater than that predicted in the tropical free troposphere. Their modelling results also show a bias at the lower tropical transition layer, indicating possible missing 70 71 sources supplying activated inorganic bromine, which current models are unable to account for. 72 Inorganic precursors to BrO such as BrCl, Br₂ and HOBr have previously been measured in the 73 Arctic at significant concentrations to account for high BrO concentrations (Liao et al., 2011). To date, there have been no previous simultaneous measurements of these inorganic halogens in the 74 75 tropics. Model simulations of their concentrations and importance to ozone loss are currently 76 constrained by experimental observations of BrO concentrations.

77

78 2. Flight Campaign

The NERC CAST (Co-ordinated Airborne Studies in the Tropics) campaign took place in the West 79 80 Pacific in January/February 2014. The field campaign had two components: (i) the NERC FAAM BAe-146 research aircraft based in Guam (13.5°N, 144.8°E); and (ii) ground-based and ozone 81 sonde measurements in Manus, Papua New Guinea (2.1°S, 147.4°E). CAST was part of an 82 83 international collaboration involving the NASA ATTREX (Airborne Tropical Tropopause Experiment) project based around the Global Hawk, the NCAR-led CONTRAST (Convective 84 85 Transport of Active Species in the Tropics) campaign based around the Gulfstream V (HIAPER) aircraft, and the Japanese SOWER project (Soundings of Ozone and Water in the Equatorial 86 87 Region). Together, detailed measurements of atmospheric structure and composition were made from the ocean surface to 20 km. An overview of this campaign can is given by Harris et al. (2016), 88 89 describing the experimental setup, flying conditions and general meteorological and core chemistry 90 results.

91

In this coordinated campaign, the FAAM BAe-146 made measurements in the lower tropical troposphere below ~8 km altitude with an emphasis on the boundary layer. The flights were made to the south of Guam and occasionally penetrated the southern hemisphere (See figure S1 in supplementary). In total, 25 flights were made between January 25th and 18th 2014 with 90 hours of measurements. Other composition measurements included H₂O, CO, CO₂, CH₄, N₂O, VSLS (very Short Lived Substances) including many bromocarbons, NO, IO (although never above the limit of

detection), black carbon and aerosol. NOx concentrations averaged at 6 ppt, indicating no influenceof pollution outflow.

100

101 **3. Materials and Methods**

A chemical ionisation mass spectrometer (CIMS) was used for real-time detection of Bro, Br₂, 102 103 HOBr and BrCl. The CIMS instrument employed here was built by the Georgia Institute of Technology and was set up as in previous studies (Le Breton et al., 2013, 2014). The inlet consisted 104 of 9.5 mm outer diameter PFA tubing of length 580 mm, which was heated to 40°C to reduce 105 surface losses. The fast inlet pump (Picolino VTE-3, Gardner Denver Thomas) delivered ambient 106 107 air to the IMR (Ion Molecule reaction) chamber at a flow rate of 5.8 SLM which was subsampled using an orifice of diameter 0.2 mm positioned at the front of the inlet to restrict the flow into the 108 IMR to 0.8 SLM. The pressure in the ionisation region was maintained at 19 Torr (133.3 Pa) 109 110 throughout the flight by controlling the flow of nitrogen into the ionisation region using a mass flow 111 meter.

112

113 The I ionization scheme was used for the detection of BrO, Br₂, BrCl and HOBr as previously utilised by Liao et al., (2011, 2012). To generate I a flow of 20 SCCM flow of the ionization gas 114 mix (15 Torr CH₃I, 20 Torr H₂O and 47.3 PSI N₂) combined with a 2 SLM flow of N₂ and passed 115 through a ²¹⁰Po Nuclecel ionizer (NRD inc.). The reagent ion was then carried into the 116 ion-molecule region where Br₂ and BrO were detected as an I⁻ adduct producing I.BrO⁻ and I.Br₂⁻. 117 BrCl was detected as the ionized ion BrCl due to ion transfer. HOBr was detected as an adduct of 118 119 both I and H₂O as I.HOBr.H₂O. Contrary to Liao et al., (2012) under our conditions it is not observed at mass 223 and 225 (I.HOBr) even with up to 12 ppb of HOBr calibrant and therefore 120 cannot interfere with the BrO signal. A variation of voltage tuning through the collisional 121 122 dissociation chamber (CDC) and quadrupole can drastically change the clustering abilities of the 123 produced ions and possibly explain this difference. An example in the literature of such variation 124 can be viewed for N_2O_5 , where it has been reported as the adduct $I.N_2O_5$ (Kecher *et al.*, 2009) and NO₃⁻ declustered by the CDC (Le Breton *et al.*, 2014). 125

126

127 Calibration and gas preparation

The CIMS was calibrated to attain a sensitivity for BrO, HOBr, Br_2 , BrCl and formic acid in the laboratory before and after the campaign. Here, we describe the BrO setup, whereas the calibrations for other compounds can be found in the supplementary material. BrO was produced in a flow tube *via* a moveable injector and produced from the reaction

132 $Br_2 + O \rightarrow BrO + Br$ (R1)

Oxygen atoms were generated using a Beenakker microwave discharge cavity operating at 60W. A 134 1 SLM flow of He (99.999 %) purified using a molecular sieve trap cooled to 78 K was combined 135 with a 1 SCCM flow of 0.1 % O_2 (>99.9995 %) and passed through quartz tubing within the 136 microwave discharge cavity. A 1-10 SCCM flow of a 0.5 % gas mixture of bromine was added into 137 the side arm of the sliding injector, combined with a 3 SLM flow of He, downstream of the 138 microwave discharge cavity reacting with O atoms to produce BrO preceding entrance to the 139 CIMS, which was identified at mass m224; I.BrO⁻.

140 The BrO signal calibration was achieved by adding NO to the flow tube via the moveable injector 141 at a constant contact time of 20 ms and by monitoring the resultant NO₂ formed by reaction with 142 BrO, using a Thermo Fisher, model 42i NO-NO₂-NO_x Analyser.

143 BrO + NO \rightarrow NO₂ + Br (R2)

NO (Technical grade 98.5%, Sigma-Aldrich) was purified by freeze-pump-thaw cycles, and selective freezing of NO₂ impurities. NO₂ (Technical grade, Air Products 99.5+%, Sigma-Aldrich) was purified by freeze-pump-thaw cycles. He (CP Grade, BOC) was passed through N₂ (l) containing a molecular sieve (5Å 4-8 mesh bead, Sigma-Aldrich) before entering the Beenakker cell. All gas mixtures entering the microwave cavities were made up in electronic grade He (99.9999 %, BOC).

150 Complete removal of BrO was ensured by adding sufficient NO, confirmed by a constant NO₂ 151 signal with further increases in (NO). The sensitivity was estimated for BrO to be 50 ion counts ppt⁻¹ 152 1 s⁻¹. Validation of the BrO production was confirmed using a flow tube chemical ionization mass 153 spectrometer implementing the SF₆⁻ ionization scheme, as reported previously by Shallcross *et al.* 154 (2012).

155

A time series and mass scan of the BrO signal can be seen in figure 1 where the raw 1 Hz data and
30 second averaged data are displayed to show how the limit of detection (0.1 ppt) was achieved.
The increased standard deviation during the calibration is as a result of the increased noise from use

of the microwave discharge in the production of BrO in the flow tube and not an instrumental 159 factor. This would increase a limit of detection (LOD) to 0.36 ppt if these data were used for the 160 calculation, although this would not change any of the analysis as this value is below the 5th 161 percentile. The uncertainty of the BrO data is therefore +15%. The low LOD can be attributed to 162 the tuning of the instrument which has a signal of 0 Hz during background periods. The instrument 163 was tuned to maximise sensitivity and minimize the LOD by optimisation of the most well 164 characterised species the CIMS detects, formic acid. Simultaneous tuning of the formic acid peak 165 166 area and minimisation of non-formic acid peaks during a formic acid calibration ensured an increase in counts was not due to increased MCP voltage noise or instrumental interference. This 167 168 ensures minimal signal at the BrO mass is observed during background periods and attributes. To validate this limit of detection for field measurements, dry nitrogen was introduced into the inlet 169 before and after a flight to simulate zero ppt BrO ambient air. 170



171

173 Fig. 1a. BrO time series and mas scan during a calibration and background period. The standard deviations are

174 reported for both 1 Hz and 30 second averaged data.



175



178

All calibrations were performed relative to in-flight formic acid calibrations and pre-flight ground calibrations, previously described by Le Breton *et al.* (2012). Calibrations of BrO, Br_2 , HOBr and BrCl were performed alongside formic acid calibrations in the laboratory, under varying water vapour conditions, to assess relative changes in the instrument's sensitivity. The sensitivity of the instrument were found to have no dependence on relative humidity. This is attributed to a tuning of excess reagent ions and the $I.(H_2O)^-$ cluster which ensures IMR water availability and sensitivity is not dependent on atmospheric relative humidity.

186 Inlet losses

The loss of HOBr onto instrument and inlet walls was quantified using a perfluoroalkoxy (PFA) flow tube inlet system which varied in length from 0.2 m to 2 m. Up to 15 % of HOBr was seen to convert to Br_2 on the inlet walls, although this value did not vary with inlet length. Therefore, it was assumed that 15 % of the Br_2 signal resulted from HOBr. This was factored into every data point collected during the campaign.

192

193 4. Measurement Results

Here, we report the first simultaneous airborne real-time observations of BrO, Br₂, BrCl and HOBr,
in tropical marine air with minimal influence from anthropogenic activity. The 1 Hz inorganic
halogen data were averaged to 30 seconds in order to facilitate comparison with other in-flight

197 measurements and model outputs. A time series of these data can be seen in figure 2, displaying the concentration of BrO, HOBr, BrCl and Br₂ during flight B830 with the time displayed as local time, 198 199 together with O₃ and altitude. In figure 2 Br₂, BrCl and HOBr are all above the limit of detection threshold throughout the flight. O_3 and altitude measurements are also displayed in the figure 2. 200 The measurement uncertainties are 10%, 10%, 10% and 15% respectively. BrO concentrations 201 throughout the campaign ranged from below the limit of detection (0.1 ppt for 30 second averaged 202 203 data) up to 1.71 ppt, with a mean of 0.69 ppt. These concentrations are within the range of that 204 reported at other equatorial mid-ocean sites such as Cape Verde (Mahajan et al., (2010) and the 205 Canary Islands (Leser et al., (2003). Our results further confirm that BrO is present in the tropical 206 troposphere at globally significant levels.



207

Fig. 2. A time series during flight B841 (14/02/14) of BrO (blue), Br₂ (green), HOBr (red) and BrCl (black) 30
second averaged concentrations from the CIMS and O₃ (light blue). The aircraft altitude (black) is also
displayed.

211

The altitude profile of BrO and O₃ can be seen in figure 3A for 4 flights during the campaign. A 212 broad anti-correlation was observed between BrO and O₃ during these flights, even though 213 214 individual profiles varied from flight to flight. Tracer-tracer analysis of all the flight data over 4 215 altitude bins (figure 3B) is consistent with surface deposition being the dominant route for O_3 loss 216 although at higher altitude bins, a larger anti correlation is observed. Further analysis of 2 altitude bins (0-2 km and 2-8 km), where the BrO is split into two samples; above and below the 50th 217 percentile, shows a near identical level of O₃ below and above the 50th percentile in the 0-2 km bin. 218 The 2-8 km bin shows an 8 ppb reduction in O_3 compared with the low BrO sample (the green bar). 219

- 220 This is consistent with the model results, i.e. Br driven ozone loss is small in the MBL (most likely
- due to the already very short life time of O_3 at the surface), but is significant above the MBL in the Free Troposphere. It must be noted that this correlation was not observed during every profile. The
- possibility that observed features were of stratospheric origin was dismissed after analysis of N_2O
- measured by a Quantum Cascade Laser absorption spectrometer (QCL) (Pitt *et al.*, 2015) exhibited
- 225 no concentrations characteristic of that from a stratospheric source.



Fig.3 Figure A shows altitude profile box plots of BrO and O_3 for 4 flights. The median, mean, 25th, 75th, 5th and 95th quartile ranges are indicated. Graph B shows a correlation plot of ozone and BrO split into 4 altitude bins (red > 6 km, green 4 -6 km, blue 2 - 4 km, black 0 - 2 km) and also a bar chart of binned BrO data by altitude split into BrO concentrations above the 50th percentile (red) and below the 50th percentile (green).

- 235
- 236 5. Global model simulations

237 A GEOS-Chem (www.geos-chem.org) global chemical transport model is utilised to interpret observed inorganic bromine mixing ratios. The model is driven by GEOS-5 assimilated 238 239 meteorological data from the NASA Global Modeling and Assimilation Office with 1/2°x2/3° horizontal resolution and 47 vertical layers extending from the surface up to about 80 km. The 240 horizontal resolution is degraded to 4°x5°. The model includes a detailed ozone-NOx-VOC-241 aerosol-Br-Cl tropospheric chemistry mechanism as described in great detail by Schmidt et al., 242 243 (2016) building on the previous bromine mechanism of Parrella et al. (2012) by including chlorine 244 chemistry and more extensive halogen multiphase chemistry. Model sources of inorganic bromine include sea salt debromination, oxidation of bromocarbon (CHBr₃, CH₂Br₂, CH₃Br) and input from 245 246 the stratosphere. In simulating the source of bromine from sea salt aerosol (SSA) we assume that 247 50% of bromide in newly emitted SSA is activated and released as Br₂. Model CHBr₃ and CH₂Br₂ emissions are based on emission fields of Hossaini et al. (2013) and modelled bromocarbon mixing 248 249 ratios are consistent with aircraft observations from an ensemble of field campaigns (Parella et al., 250 2012). Stratospheric concentrations (monthly and diurnally varying) of Bry (Br, BrO, HOBr, HBr, 251 BrNO₃ and BrCl) are taken from a Goddard Earth Observing System Chemistry-Climate Model 252 (GEOSCCM) simulation that has been demonstrated to yield stratospheric BrO profiles consistent with balloon observations (Holmes et al., 2006). The halogen and heterogeneous chemistry 253 254 mechanism yields BrO column densities and mixing ratios broadly consistent with ground based and recent aircraft observations (Mahajan et al., 2010 Volkamer et al., 2015, Wang et al., 2015). 255

256

Simulated BrO and HOBr (figure 4) show fair agreement (over 55% and 75% of the data points 257 258 agreeing within error respectively) with observations in the MBL and FT. Agreement is only 259 achieved when heterogeneous chemistry is included, as shown in figure 4. Bromine driven O_3 loss 260 takes place by cycling between Br, BrO and HOBr. Bromine driven oxidation of NO_x (a key ozone precursor) also reduces the level of ozone in the troposphere. The BrO measurements in the Pacific 261 262 by Wang et al., (2015) suggested that their consistent high BrO concentrations were due to a missing inorganic source. Here, the measured inorganic halogen concentrations, similar to that by 263 264 Wang et al., (2015) further support their hypothesis that inorganic halogen concentrations may often be several ppt in the Pacific. The increase in BrO from heterogeneous chemistry results in a 265 266 further 6 ppb O₃ loss in the model.

The model predicts higher levels of BrCl than Br_2 in direct contrast with observations. The inconsistent pattern in simulated levels of BrCl and Br_2 may be due to multiphase halide exchange reactions (BrCl + Br⁻ \rightarrow Br₂ + Cl⁻) that are not accounted for in the simulations, which has

270 previously been reported to initiate O_3 depletion reactions (Wachsmuth *et al.*, 2002). The model 271 results are unable to account for the observed high levels of Br_2 throughout the campaign, seen in 272 figure 4 where the measured sum of BrCl and Br_2 is on average 0.5 ppt higher than the model. It 273 must be stated that this value represents absolute reported concentrations and does not account for 274 experimental uncertainty. Extensive laboratory studies have assessed a possible inlet loss of HOBr to form Br₂ (Liao et al. 2012 and Neuman et al., 2010). Our results indicate that a maximum of 275 15% of HOBr is lost on the walls, which could not account for the discrepancy between model and 276 277 measurement. O_3 may play in halogen activation on the inlet, although no sign of this interference was found; i.e. no correlation between observed Br2 and O3 is observed (see figure S4 in the 278 supporting material). Furthermore, there remains the possibility that our inlet walls could become 279 280 coated in sea salt and thus increase the Br₂ production along the inlet which would be difficult to 281 take into account. However, Huey and co-workers (see e.g. Neuman et al., 2010) experimentally 282 determined the efficiency of inlet conversion of HOBr on the inlet on various PFA in comparison 283 with a pure salt substrate and showed that there is a factor of 5 increase in Br_2 production. For the 284 data presented in the manuscript there would have to be a factor of 20 increase in wall loss 285 efficiency to explain our observed Br₂, thus we are confident that wall loss can not explain the observed Br_2 . Liao et al., (2012) found that NaBr deliquescence in the presence of OH and the 286 photolysis of O_3 could produce significant levels of molecular bromine. This source of bromine is 287 288 not represented in the model and could therefore account for its underestimation compared to the 289 measurements.



Fig. 4. Altitude profile box plot of the measured (CIMS) and modelled (GEOS-Chem) concentration of BrO (upper left), HOBr (upper right), BrCl (lower left, dashed), Br₂ (lower left, full), and ozone (lower right)

throughout the campaign for 16 flights. Black lines represent the measurements. The red line is output from our GEOS-Chem simulation. The green curve is model output from a simulation without multiphase halogen chemistry (no oxidation of Br⁻ and Cl⁻ and no halogen release from SSA). The cyan curve shows simulated ozone when halogen chemistry is not included. The horizontal bars on the observed data lines correspond to the maximum and minimal values in the data set

298 299

300 6. Contribution of inorganic bromine to ozone loss

301 CAST aircraft measurements of organobromine compounds taken by Whole Air Samples (WAS) (Lidster et al., 2014) and measured by GC-MS (Hopkins et al., 2011) coupled with modelled 302 303 photooxidation rates show that their contribution to the Br loading is minimal. Our results indicate that at least 99% of Br radicals originates from inorganic halogen species, in contrast to other 304 model studies (Liu et al., 2002). Non-halogen related O₃ loss, from photolysis and reactions with 305 OH and HO₂ radicals (Yang et al., 2005) was estimated using a steady state approximation 306 implementing all other inorganic halogen measurements as described in the SOM. The average 307 308 HO_2 concentration calculated in the steady state approximation in the upper free troposphere was 6 309 ppt, consistent with Jagle et al., (1997) and Tin et al, (2001). The results from this analysis imply that Br from inorganic halogen photolysis can account for up to 20% of the O₃ loss in the FT and 310 MBL as shown in figure 5. The figure also suggests that HO₂ can account for up to 80% loss of O₃ 311 312 in the upper troposphere.



Fig. 5. The graph represents the contribution to O_3 loss in the FT and MBL from reaction with HO₂ (yellow), O¹D (blue), Br from inorganic halogens measured by the CIMS (purple) and OH (red). These values have been

- 317 calculated from the steady state approximation explained in the SOM.
- 318

There remains the possibility that we have overestimated the impact of Br_2 photolysis as a result of inlet conversion of HOBr. However, even if we assume that all the Br_2 that we observe is resultant from HOBr, that would not alter the conclusions of this work. Inorganic Bromine would still be the major source (99%) of Br atoms in the MBL, the source being either from Br_2 or HOBr.

323

324 7. Conclusions

325

326 The recent work by Wang et al., (2015) showing the capacity for inorganic halogen chemistry to 327 destroy ozone in the Pacific region and suggestioning that such high concentrations of BrO may 328 result from high concentrations of inorganic halogens in the region have been further supported by 329 the work presented here. The first simultaneous airborne measurements of BrO, Br₂, HOBr and 330 BrCl suggest inorganic halogen photochemistry is the overwhelmingly dominant source of Br in 331 the FT and MBL in comparison with that of organobromine compounds. Analysis of data from this 332 work reveals that Br radicals generated from inorganic halogen photolysis can account for up to 333 20% of O_3 in the FT and to a lesser extent in the MBL as shown in figure 5. Previous model studies suggested that there could be up to 20% less ozone in the tropics as a result of bromine 334 335 chemistry (von Glasow et al., 2004, Long et al., 2014) and this study has provided the direct evidence of the inorganic sources of bromine. This study also builds on the advancements in this 336 field by Wang et al. (2015) and further supports the hypothesis that a missing source of BrO 337 338 production may propagate through the debromination of sea salt.

339

340

341 Acknowledgments

342 The experiments were conceived by NH, LJC, MJE, JDL, CJP and DES; designed by CJP, DES,

MLB and AB, and carried out by CJP, DES, MLB, AB, KEL and TB and funded by NERC. Model

344 studies were carried out by JAS, MAHK, DES, ME and DJJ. SJA, RTL and LJC contributed

345 organohalogen measurements. All authors participated in the data analysis and interpretation and

346 contributed to the manuscript. JAS acknowledges support from The Danish Council for

347 Independent Research | Natural Sciences and the NASA Atmospheric Composition Modelling and

348 Analysis Program. CAST data is available via NERC BADC

349 (http://catalogue.ceda.ac.uk/uuid/565b6bb5a0535b438ad2fae4c852e1b3) and the use of data

produced in the context of the CAST project is governed by the protocols outlined at

351 https://badc.nerc.ac.uk/conditions/cast.html.

352

354 **References**

- 355 Crutzen, P. (1988), Tropospheric ozone: an overview. In: Isaksen, I.S.A. (Ed.), Tropospheric
- 356 Ozone: Regional and Global Scale Interactions, Reidel Publishing, Boston, MA, 3 11.

Harris, N., Carpenter, L. J., Lee, J. D., Vaughan, G., Filus, M.T., Jones, R. L., Ouyang, B., Pyle, J.

- A., Robinson, A. D., Andrews, S., Lewis, A. C., Minaeian, J., Vaughan, A., Dorsey, J., Gallagher,
- 359 M. W., Le Breton, M., Newton, R., Percival, C. J., Ricketts, H., Bauguitte, S. J-B., Nott, G.,
- Wellpott, A., Ashfold, M., Flemming, J., Butler, R., Palmer, P., Stopford, C., Chemel, C., Kaye, P.,
- Boesch, H., Humpage, N., Vick, A., Mackenzie, A. R., Hyde, R., Angelov, P., Manning, A. and
- 362 Meneguz E., 2016. Co-ordinated Airborne Studies in the Tropics (CAST), Submitted to Bulletin
- 363 Amer. Met. Soc., doi: 10.1175/BAMS-D-14-00290.1

364

- Holmes, C., Jacob, D. and Yang, X., 2006. Global lifetime of elemental mercury against oxidation
- by atomic bromine in the free troposphere, Geophys. Res. Lett., 33, L20808,
- doi:10.1029/2006GL027176.

368

- Holmes, C., Jacob, D. J., Corbitt, E. S., Mao, J., Yang, X., Talbot, R. and Slemr, F., 2010. Global
- atmospheric model for mercury including oxidation by bromine atoms, Atmos. Chem. Phys., 10,
- 371 12037–12057, 2010, www.atmos-chem-phys.net/10/12037/2010/doi:10.5194/acp-10-12037-2010.

372

- Hopkins, J. R., Jones, C. E., and Lewis, A. C., 2011. A dual channel gas chromatograph for
- atmospheric analysis of volatile organic compounds including oxygenated and monoterpene
 compounds, J. Environ. Monit., doi:10.1039/c1em10050e.
- 376
- Hossaini, R., Mantle, H., Chipperfield, M. P., Montzka, S. A., Hamer, P., Ziska, F., Quack, B.,
- 378 Krüger, K., Tegtmeier, S., Atlas, E., Sala, S., Engel, A., Bönisch, H., Keber, T., Oram, D., Mills,
- 379 G., Ordóñez, C., Saiz-Lopez, A., Warwick, N., Liang, Q., Feng, W., Moore, F., Miller, B. R.,
- 380 Marécal, V., Richards, N. A. D., Dorf, M., and Pfeilsticker, K., 2013. Evaluating global emission
- inventories of biogenic bromocarbons, Atmos. Chem. Phys., 13, 11819-11838, doi:10.5194/acp-13-
- **382** 11819-2013.

383

- Jaegle, L., Jacob, D. J., Wennberg, P. O., Spivakowsky, C. M., Hanisco, T. F., Lanzendorf, E. J.,
- Hinsta, E. J., Fahey, D. W., Keim, E. R., Proffitt, M. H., Atlas, E. L., Flocke, F., Schauffler, S.,
- McElroy, C. T., Midwinter, C., Pfister, L. and Wilson, J. C., 1997. Geophys. Res. Lett, 24, 24, 3181
- **387** − **3**184.

- 389 Kercher, J. P., Riedel, T. P. and Thornton, J. A., 2009. Chlorine activation by N₂O₅: simultaneous,
- in situ detection of ClNO₂ and N_2O_5 by chemical ionization mass spectrometry, Atmos. Meas.
- 391 Tech., 2, 193–204.

- Le Breton, M., McGillen, M. R., Muller, J. B. A., Shallcross, D. E., Bacak, A., Xiao, P., Huey, L.
 G., Tanner, D., Coe, H. and Percival, C. J., 2012. Airborne observations of formic acid using a
- chemical ionisation mass spectrometer, Atmos. Meas. Tech., 5, 3029–3039.
- 396
- Le Breton, M., Bacak, A., Muller, J.B. A., Bannan, T. J., Kennedy, O., Ouyang, B., Xiao, P.,
- Bauguitte, S. J-B., Jones, R. L., Daniels, M. J. S., Ball, S. M. and Percival, C. J., 2014. The first
- airborne comparison of N_2O_5 measurements over the UK using a CIMS and BBCEAS during the RONOCO campaign, Anal. Methods, 6, 9731-9743
- 401
- Le Breton, M., McGillen, M. R., Muller, J. B. A., Shallcross, D. E., Batt, R., Cooke, M. C.,
- Bauguitte, S. J-B. and Percival, C. J., 2014b. Simultaneous airborne nitric acid and formic acid
- 404 measurements using a chemical ionization mass spectrometer around the UK: Analysis of primary
- and secondary production pathways, *Atmos. Environ.*, 83, 166–175,
- 406 doi:10.1016/j.atmosenv.2013.10.008.
- 407
- Leser, H., Honninger, G., Platt, U. (2003). MAX-DOAS measurements of BrO and NO₂ in the
 marine boundary layer. *Geophys. Res. Lett.*, **30**, Art No. 1537; DOI: 10.1029/2002GL015811.
- Liang, Q., Stolarski, R. S., Kawa, S. R., Nielsen, J. E., Douglass, A. R., Rodriguez, J. M., Blake, D.
- 412 R., Atlas, E. L. and Ott, L. E., 2010. Finding the missing stratospheric Bry: a global modeling study
- 413 of CHBr₃ and CH₂Br₂, Atmos. Chem. Phys., 10, 2269-2286.
- 414
- Lidster, R. T., Hamilton, J. F., Lee, J. D., Lewis, A. C., Hopkins, J. R., Punjabi, S., Rickard, A.
- 416 R., and Young, J. C., 2014. The impact of monoaromatic hydrocarbons on OH reactivity in the
- 417 coastal UK boundary layer and free troposphere, Atmos. Chem. Phys., 14, 6677-6693,
- 418 doi:10.5194/acp-14-6677-2014.
- 419
- 420 Liao, J., H. Sihler, L. G. Huey, J. A. Neuman, D. J. Tanner, U. Friess, U. Platt, F. M. Flocke, J. J.
- Orlando, P. B. Shepson, H. J. Beine, A. J. Weinheimer, S. J. Sjostedt, J. B. Nowak, D. J. Knapp, R.
 M. Staebler, W. Zheng, R. Sander, S. R. Hall, and K. Ullman (2011), A comparison of Arctic BrO
- 423 measurements by chemical ionization mass spectrometry and long path-differential optical
- 424 absorption spectroscopy, J. Geophys. Res, 116, D14, 10.1029/2010JD014788.
- 425
- 426 Liao, J., Huey, L. G., Scheuer, E., Dibbi, J. E., Stickel, R. R., Tanner, D. J., Neuman, J. A., Nowak,
- 427 J. B., Choi, S., Wang, Y., Salawitch, R. J., Canty, T., Chance, K., Jurosu, T., Suleiman, R.,
- 428 Weinheimer, A. J., Shetter, R. R, Fired, A., Brune, W., Anderson, B., Zhang, X., Chen, G.,

429 Crawford, J., Hecobia, A. and Ingall, E. D. 2012. Characterization of soluble bromide

430 measurements and a case study of BrO observations during ARCTAS, Atmos. Chem. Phys., 12,

- 431 1327–1338.
- 432

Liu, H., Jacob, D. J., Chan, L. Y., Oltmans, S. J., Bey, I., Yantosca, R. M., Harris, J. M., Duncan,

- B. N. and Martin, R. V., 2002. Sources of tropospheric ozone along the Asian Pacific Rim: An
- analysis of ozonesonde observations, J. Geophys. Res., 107, doi:10.1029/2001JD002005.
- 436

437 Long, M. S., Keene, W. C., Easter, R. C., Sander, R., Liu, X., Kerkweg, A. and Erickson, D., 2014.

438 Sensitivity of tropospheric chemical composition to halogen-radical chemistry using a fully

439 coupled size-resolved multiphase chemistry–global climate system: halogen distributions, aerosol

440 composition, and sensitivity of climate-relevant gases, Atmos. Chem. Phys., 14, 3397–3425,

441 doi:10.5194/acp-14-3397-2014.

442

443 Mahajan, A. S., Plane, J. M. C., Oetjen, H., Mendes, L., Saunders, R. W., Saiz-Lopez, A., Jones, C.

E., Carpenter, L. J. and McFiggans, G. B., 2010. Measurement and modelling of tropospheric

reactive halogen species over the tropical pacific Atlantic Ocean, Atmos. Chem. Phys., 10, 46114624, doi:10.5194/acp-10-4611-2010.

447

448 Neuman, J. A., Nowak, J. B., Huey, L. G., Burkholder, J. B., Dibb, J. D., Holloway, J. S., Liao, J.,

Peischl, J., Robert, J. M., Ryerson, T. B., Scheur, E., Stark, H., Stickel, R. E., Tanner, D. J. and

450 Weinheimer, A (2010), Bromine measurements in ozone depleted air over the Arctic Ocean,

451 Atmos. Chem. Phys., 10, 6503-6514.

452

453 Parella, J. P, Jacob, D. J., Liang, Q., Zhang, Y., Mickeley, L. J., Miller, B., Evans, M. J., Yang, X.,

454 Pyle, J. A., Theys, N. and Van Roozendael, M., 2012. Tropospheric bromine chemistry:

455 implications for present and pre-industrial ozone and mercury, Atmos. Chem. Phys., 12, 6723–

456 6740, doi:10.5194/acp-12-6723-2012.

457

458 Prather, M., Ehhalt, D., Dentener, F., Derwent, R., Dlugokencky, E., Holland, E., Isaksen, I.,

459 Katima, J., Kirchhoff, V., Matson, P., Midgley, P., and Wang, M., 2001. Atmospheric Chemistry

460 and Greenhouse Gases. In: Climate Change 2001: The Scientific Basis. Contribution of Working

461 Group I to the Third Assessment Report of the Intergovermental Panel on Climate Change.

462 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

463

464 Pitt, J. R., Le Breton, M., Allen, G., Percival, C. J., Gallagher, M. W., Bauguitte, S. J-B., O'Shea,

465 S. J., Muller, J. B., Zahniser, M. S., Pyle, J. and Palmer, P. I., 2015. The development and

evaluation of airborne in situ N₂O and CH₄ sampling using a Quantum Cascade Laser Absorption
Spectrometer (QCLAS), Atmos. Meas. Tech. Discuss., 8, 8859-8902, doi:10.5194/amtd-8-8859-

468 2015.

469

- 470 Read, K. A., Mahajan, A. S., Carpenter, L. J., Evans, M. J., Faria, B. V. E., Heard, D. E., Jopkins, J.
- 471 R., Lee, J. D., Moller, S. J., Lewis, A. C., Mendes, L., McQuaid, J. B., Oetjen, H., Saiz-Lopez, A.,
- Pilling, M. J. and Plane J. M. C., 2008. Extensive halogen-mediated ozone destruction over the
- tropical Atlantic Ocean, *Nature* 453, 1232-1235 (26 June 2008) doi:10.1038/nature07035.
- 474
- 475 Saiz-Lopez, A., Lamarque, J-F., Kinnison, D. E., Tilmes, S., Ordóñez, C., Orlando, J.
- 476 J., Conley, A. J., Plane, J. M. C., Mahajan, A. S., Sousa Santos, G., Atlas, E. L., Blake, D.
- 477 R., Sander, S. P., Schauffler, S., Thompson, A. M. and Brasseur, G., 2012. Estimating the climate
- significance of halogen-driven ozone loss in the tropical marine troposphere, Atmos. Chem. Phys.,
- 479 12, 3939-3949, doi:10.5194/acp-12-3939-2012.
- 480
- 481 Schmidt, J. A., Jacob, D. J., Horowitz, H. M., Hu, L., Sherwen, T., Evans, M. J., Lang, Q.,
- 482 Suleiman, R. M., Oram, D. E., Le Breton, M., Percival, C. J., Wang, S., Dix, B. and Volkamer, R.,
- 483 2016. Modeling the observed tropospheric BrO background: Importance of multiphase chemistry
- and implications for ozone, OH, and mercury, accepted by J. Geophs. Res.
- 485
- Shallcross, D. E., Leather, K. E., Bacak, A., Xiao, P., Lee, E. P., Ng, M., Mok, D. K. W., Dyke, J.
 M., Hossaini, R., Chipperfield, M. P., Khan, M. A. H. and Percival, C. J., 2015. Reaction between
- CH_3O_2 and BrO Radicals: A New Source of Upper Troposphere Lower Stratosphere Hydroxyl
- 489 Radicals, J. Phys. Chem. A, 2015, 119 (19), pp 4618–4632, DOI: 10.1021/jp5108203.

490

- 491 Theys, N., Van Roozendael, M., Hendrick, F., Yang, X., De Smedt, I., Richter, A., Begoin, M.,
- 492 Errera, Q., Johnston, P. V., Kreher, K. and De Mazière, M., 2011. Global observations of
- tropospheric BrO columns using GOME-2 satellite data, Atmos. Chem. Phys., 11, 1791-1811,
- doi:10.5194/acp-11-1791-2011.

495

- Tin, D, I. Faloona, J.B. Sampas, W. Brune, J. Olson, J. Crawford, M. Avery, G. Sachse, S. vay, S.
- 497 Sandholm, H-W Guan, T. Vaughn, J. Mastromarino, B., Heikes, J. Snow, J. Podolski and H. Singh
- 498 (2001), J. Geophys. Res, 106, D23, 32,667-32,681.

- 500 Volkamer, R., Baidar, S., Campos, T. L., Coburn, S., DiGangi, J. P., Dix, B., Eloranta, E. W.,
- 501 Koenig, T. K., Morley, B., Ortega, I., Pierce, B. R., Reeves, M., Sinreich, R., Wang, S., Zondlo, M.
- A., and Romashkin, P. A., 2015. Aircraft measurements of BrO, IO, glyoxal, NO₂, H₂O, O₂–O₂ and
- aerosol extinction profiles in the tropics: comparison with aircraft-/ship-based in situ and lidar
- 504 measurements, Atmos. Meas. Tech., 8, 2121-2148, doi:10.5194/amt-8-2121-2015.

505

- Von Glasow, R., von Kuhlman, R., Lawrence, M. G., Platt, U. and Crutzen, P. J., 2004. Impact of
 reactive bromine chemistry in the troposphere, Atmos. Chem. Phys., 4, 2481–2497, 16807324/acp/2004-4-2481.
- 509
- 510 Wachsmith, M., Gaggler, H. W., von Glasow, R. and Ammann, M., 2002. Accommodation
- 511 coefficient of HOBr on deliquescent sodium bromide aerosol particles, Atmos. Chem. Phys., 2,
- 512 121–131.

513

- 514 Wang, S.Y., Schmidt, J.A., Baidar, S., Coburn, S., Dix, B., Koenig, T.K. Apel, E., Bowdalo, D.,
- 515 Campos, T.L. Eloranta, E., Evans, M.J., DiGangi, J.P., Zondlo, M.A., Gao, R.S., Haggerty, J.A.,
- Hall, S.R., Hornbrook, R.S., Jacob, D., Morley, B., Pierce, B., Reeves, M., Romashkin, P., ter
- 517 Schure, A., Volkamer, R. (2015). Active and widespread halogen chemistry in the tropical and
- subtropical free troposphere. *Proc. Natl, Acad. Sci.*, **112**, 9281-9286.
- 519
- 520 Yang, X., Cox, R. A., Warwick, N. J., Pyle, J. A., Carver, G. D., O'Connor, F. M. and Savage, N.
- 521 H., 2005. Tropospheric bromine chemistry and its impacts on ozone: A model study, J. Geophys.
- 522 Res., 110, D23,311, doi:10.1029/2005JD006244.

Enhanced ozone loss by active inorganic bromine chemistry in the tropical troposphere

Highlights

Ozone (O_3) is one of the pivotal species in the lower atmosphere, initiating production of key radicals OH, NO₃ and Criegee Intermediates, as well as being a powerful oxidizing agent in its own right. Hence, O_3 plays a vital role in controlling atmospheric chemistry and air quality and in addition is a key greenhouse gas. The role of O_3 in the tropical troposphere is none more important where its photochemistry drives high OH production in the lower marine boundary layer through the availability of high solar fluxes and water vapour. This region is where significant removal of pollutants occurs and is, dynamically, the gateway to the stratosphere. Therefore, the O_3 budget in the region is critical and yet there are significant gaps in our knowledge. For example, it is currently assumed that low concentrations of O_3 observed in the tropical marine boundary are due to higher photolysis rates in the presence of high water vapour concentrations. However, recent measurements of extremely low O_3 concentrations (<10 ppb) indicate that another unexplained mechanism is responsible for O_3 loss in the tropics, which models cannot explain.

In this work we present results from simultaneous measurements of a number of gaseous inorganic bromine species in the tropics onboard the FAAM 146 Research Aircraft. The results show that BrO, HOBr, BrCl and Br₂ represent 99% of the free Br, contrary to current theory in which organobromine species dominate the Br production pathway. This suite of inorganic bromine species can rapidly recycle Br during sunlit hours and provide an efficient catalytic cycle that removes significant levels of ozone. Good agreement with GEOS Chem, a global 3-D chemical transport model which implemented a new multi phase chemistry scheme, reveals the underestimation of halogen chemistry within standard models and accounts for their under prediction of O_3 loss in the tropical troposphere. Model results show that the concentrations of inorganic bromine species presented here can contribute up to 6 ppb O_3 loss in the tropical troposphere, representing 20% of the total O_3 destruction which in this region is a significant additional loss. Implementation of this inorganic halogen chemistry into future models is imperative to accurately quantify O_3 variability in the tropics and therefore the oxidizing power of the atmosphere.