



UNIVERSITY OF LEEDS

This is a repository copy of *A stochastic partially reversible investment problem on a finite time-horizon: free-boundary analysis.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/112931/>

Version: Accepted Version

Article:

De Angelis, T and Ferrari, G (2014) A stochastic partially reversible investment problem on a finite time-horizon: free-boundary analysis. *Stochastic Processes and their Applications*, 124. pp. 4080-4119. ISSN 0304-4149

<https://doi.org/10.1016/j.spa.2014.07.008>

© 2014 Elsevier B.V. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

THE DIVIDEND PROBLEM WITH A FINITE HORIZON

TIZIANO DE ANGELIS AND ERIK EKSTRÖM¹

ABSTRACT. We characterise the value function of the optimal dividend problem with a finite time horizon as the unique classical solution of a suitable Hamilton-Jacobi-Bellman equation. The optimal dividend strategy is realised by a Skorokhod reflection of the fund's value at a time-dependent optimal boundary. Our results are obtained by establishing for the first time a new connection between singular control problems with an absorbing boundary and optimal stopping problems on a diffusion reflected at 0 and created at a rate proportional to its local time.

1. INTRODUCTION

The dividend problem is a foundational problem in actuarial mathematics whose formulation dates back to De Finetti's work [11]. The model addresses the question of how a fund or an insurance company should distribute dividends to its beneficiaries prior to the time of ruin. After De Finetti's seminal work, the dividend problem has attracted the interest of many mathematicians and economists who produced a substantial body of literature on the subject. An extensive review of existing models and related mathematical results was published by Avanzi [1] in 2009, and the list of papers relative to the topic has continued to increase since.

Here we consider a canonical formulation of the problem in a simple diffusive setting that was proposed by Radner and Shepp [28] and later considered also by [20] among many others. The value of a fund after dividends have been paid out evolves according to

$$X_t^D = x + \mu t + \sigma B_t - D_t, \quad t \geq 0,$$

where μ and $\sigma > 0$ are constants, B is a Brownian motion and D_t is the cumulative amount of dividends paid out up to time t . The objective of the fund manager is to maximise the expected present value of future dividends up to the fund's default time $\gamma^D := \inf\{t \geq 0 : X_t^D \leq 0\}$. In addition, we also assume that the manager has a finite time horizon T for the investment plan. The assumption of a finite horizon is the main difference between our model and the vast majority of the existing literature (including [28] and [20]). From the financial point of view, this restriction on the set of

Date: January 14, 2017.

Key words and phrases. The dividend problem; singular control; optimal stopping.

¹ Support from the Swedish Research Council (VR) is gratefully acknowledged.

admissible dividend strategies is very natural and it simply means that an investment fund is liquidated at a pre-specified future date.

If the fund's value at time $t \in [0, T]$ is $x > 0$, the optimisation problem that the fund manager is faced with may be stated as follows:

$$(1) \quad \text{Find } D^* \text{ that maximises } \mathcal{J}(t, x; D) := \mathbb{E} \left[\int_{0-}^{\gamma^D \wedge (T-t)} e^{-rs} dD_s \right]$$

where we integrate from $0-$ to account for the possibility of a jump of D at time zero. From the mathematical point of view this is a problem of singular stochastic control (SSC) on a finite time horizon in which the underlying process is absorbed at zero. It is important to notice that zero is a regular point for the uncontrolled process $R_t = x + \mu t + \sigma B_t$ so that default may occur prior to T with positive probability even if no dividends are distributed.

In this work we solve (1) by constructing an optimal dividend strategy and by proving that the corresponding value function $V(t, x) := \mathcal{J}(t, x; D^*)$ is a classical solution of the Hamilton-Jacobi-Bellman equation, i.e. in particular $V \in \mathcal{C}^{1,2}([0, T] \times (0, \infty))$. The optimal dividend strategy is shown to be the solution of a Skorokhod reflection problem at an appropriate time-dependent optimal boundary.

To accomplish our task we develop a self-contained, fully probabilistic proof that hinges on a new type of connection between SSC problems and optimal stopping. Indeed, we show that $V_x = U$, where U is the value function of an optimal stopping problem whose underlying process is a Brownian motion with drift μ and variance σ^2 , which gets *reflected at zero and created at a rate proportional to its local time*. Although links between optimal stopping and singular control have been known for many years (see for example [2–10, 15, 19, 21–23, 32] among others) our result makes a fundamental forward leap in this field. For the first time we establish that an absorbing boundary in SSC translates to a reflecting boundary with creation in optimal stopping. This new characterisation proves to be a powerful tool to tackle problem (1) in an effective way.

We remark that, despite the vast existing literature on SSC, the study of problems that combine absorbing boundary behaviour with a finite time horizon is still a major theoretical challenge. For example, we observe that in the literature on optimal dividend problems an analytical characterisation of the optimal strategy and of the value function can only be found in models with infinite time horizon (for a theoretical study of problems of this kind one may refer to [31]). These models are substantially easier to deal with compared to (1) because they give rise to variational problems in the form of ordinary differential equations whereas our problem is associated to a parabolic one.

To the best of our knowledge, an analytical study of the problem in (1) has only been addressed very recently by Grandits in a series of two papers, [16] and [17], followed by a third one [18] containing an extension of the

canonical model. In these papers, Grandits uses methods from PDE and free-boundary analysis that rely on several transformations of the variational problem associated to (1). In [16], the author obtains ϵ -optimal boundaries, whereas in [17] he manages to pass to the limit as $\epsilon \rightarrow 0$ and shows that the optimal strategy is of barrier type. Moreover, Grandits proves that the optimal boundary is continuous and that the value function V is continuous with locally bounded weak derivatives V_t and V_{xx} . Under strong assumptions on the regularity of the boundary he also derives asymptotic estimates for $t \rightarrow T$.

Our new connection between the SSC problem (1) and optimal stopping enables us to use powerful methods from optimal stopping theory in the study of the dividend problem. This leads to a self-contained probabilistic analysis that complements and improves results in [16] and [17]. We obtain spatial concavity and $\mathcal{C}^{1,2}$ -regularity of the value function, monotonicity of the boundary and, without further assumptions, the boundary's asymptotic behaviour at T along with its characterisation as the unique continuous solution of an integral equation (both these properties are actually consequences of results relative to the Russian option, see the last remark in Section 3 and results in Section 8).

The paper is organised as follows. In Section 2 we introduce the dividend problem with a finite time horizon in some further detail, and we provide a verification theorem. In Section 3 we introduce a related optimal stopping problem with a peculiar boundary condition at 0, and we state our main result, Theorem 3.1, which shows the connection between these two problems. To prove Theorem 3.1, we begin our study of the optimal stopping problem in Section 4 by proving continuity of the value function as well as existence and continuity of the optimal stopping boundary. To apply a verification result, however, additional regularity of the optimal stopping problem is needed, which is the main contribution of Sections 5 (spatial regularity) and 6 (regularity in time). The proof of Theorem 3.1 is instead contained in Section 7. Finally, Section 8 gives a couple of concluding remarks concerning additional properties of the optimal boundary.

2. THE OPTIMAL DIVIDEND PROBLEM

Denote by $X^D = (X_s^D)_{s \in [0, \infty)}$ the value of a fund after dividends have been paid out according to a strategy D . We assume that

$$(2) \quad X_s^D = x + \mu s + \sigma B_s - D_s,$$

where $x \geq 0$, μ and $\sigma > 0$ are constants, B is a standard Brownian motion and D is a non-negative, non-decreasing and right-continuous process (adapted to the filtration generated by B) with the interpretation that D_s represents the accumulated dividends paid out until time s . In particular, if $D_0 > 0$, then a lump sum D_0 is paid out at time 0. We only consider dividend strategies that satisfy $D_s - D_{s-} \leq X_{s-}^D$ at all times $s \in [0, \infty)$

(with a convention that $D_{0-} = 0$), and we denote the set of such dividend strategies \mathcal{A} .

For a given dividend strategy $D \in \mathcal{A}$, denote by

$$\gamma^D := \inf\{s \geq 0 : X_s^D \leq 0\}$$

the (possibly infinite) default time of the firm, and consider the stochastic control problem

$$(3) \quad V(t, x) = \sup_{D \in \mathcal{A}} \mathbb{E}_x \left[\int_{0-}^{\gamma^D \wedge (T-t)} e^{-rs} dD_s \right].$$

Here $\mathbb{E}_x[\cdot] = \mathbb{E}[\cdot | X_{0-}^D = x]$, $T > 0$ is a given time horizon, and we refer to problem (3) as the **dividend problem with finite horizon**.

Remark The integral in (3) is interpreted in the Riemann-Stieltjes sense. In particular, the lower limit $0-$ of integration accounts for the contribution from an initial dividend payment $D_0 > 0$. We also point out that choosing a strategy with $D_0 = x$ in (3) yields the trivial inequality $V \geq x$.

Remark Notice that in (3) we consider the optimisation problem as if it were started at time $0-$ (i.e. before dividends are paid) but with a time horizon equal to $T - t$. This is justified because X^D is time-homogeneous and, if at time t the fund's value before dividends are paid is $x > 0$, then the residual time of the optimisation is $T - t$. Moreover we also point out that in the rest of the paper we use the time-space process $(t + s, X_s^D)_{s \in [0, T-t]}$ under the measure \mathbb{P}_x . The latter is equivalent to the process $(s, X_s^D)_{s \in [t, T]}$ under the measure $\mathbb{P}_{t,x}(\cdot) = \mathbb{P}(\cdot | X_{t-}^D = x)$.

Denote by \mathcal{L} the differential operator

$$(4) \quad \mathcal{L} = \partial_t + \frac{\sigma^2}{2} \partial_x^2 + \mu \partial_x - r.$$

We have the following verification theorem.

Theorem 2.1. (Verification). *Let a function $v \in \mathcal{C}([0, T] \times [0, \infty)) \cap \mathcal{C}^{1,2}([0, T] \times (0, \infty))$ be given. Assume that*

- (i) $\max\{\mathcal{L}v, 1 - v_x\} = 0$ on $[0, T] \times (0, \infty)$;
- (ii) $v(t, 0) = 0$ for all $t \in [0, T]$;
- (iii) $v(T, x) = x$ for $x \in [0, \infty)$.

Further assume that there exists a continuous function $a : [0, T] \rightarrow [0, \infty)$ with $a(T) = 0$ such that

- (iv) $\mathcal{L}v = 0$ for $(t, x) \in [0, T] \times (0, \infty)$ with $0 < x < a(t)$;
- (v) $v_x = 1$ for $(t, x) \in [0, T] \times (0, \infty)$ with $x \geq a(t)$.

Then $V = v$. Moreover,

$$(5) \quad D_s^a := \sup_{0 \leq u \leq s} (x + \mu u + \sigma B_u - a(t + u))^+$$

is an optimal strategy in the sense that

$$V(t, x) = \mathbb{E}_x \left[\int_{0-}^{\gamma^{D^a} \wedge (T-t)} e^{-rs} dD_s^a \right].$$

Proof. Fix $(t, x) \in [0, T) \times (0, \infty)$. For a given dividend strategy $D \in \mathcal{A}$, denote by D^c the continuous part of D , and let, for $\epsilon > 0$,

$$\gamma^\epsilon := \inf\{s \geq 0 : X_s^D \leq \epsilon\} \wedge (T - t - \epsilon).$$

Then

$$\begin{aligned} e^{-r\gamma^\epsilon} v(t + \gamma^\epsilon, X_{\gamma^\epsilon}^D) &= v(t, x) + \int_0^{\gamma^\epsilon} e^{-rs} \mathcal{L}v(t + s, X_{s-}^D) ds \\ &\quad - \int_0^{\gamma^\epsilon} e^{-rs} v_x(t + s, X_{s-}^D) dD_s^c \\ &\quad + \sum_{0 \leq s \leq \gamma^\epsilon} e^{-rs} (v(t + s, X_s^D) - v(t + s, X_{s-}^D)) \\ &\quad + \int_0^{\gamma^\epsilon} e^{-rs} \sigma v_x(t + s, X_{s-}^D) dB_s. \end{aligned}$$

Note that v_x is bounded on the set $[0, T - \epsilon] \times [\epsilon, \infty)$ (recall (v)), so the last integral is a (stopped) martingale. Taking expected values of both sides gives

$$\begin{aligned} \mathbb{E}_x [e^{-r\gamma^\epsilon} v(t + \gamma^\epsilon, X_{\gamma^\epsilon}^D)] &= v(t, x) + \mathbb{E}_x \left[\int_0^{\gamma^\epsilon} e^{-rs} \mathcal{L}v(t + s, X_{s-}^D) ds \right] \\ (6) \quad &\quad - \mathbb{E}_x \left[\int_0^{\gamma^\epsilon} e^{-rs} v_x(t + s, X_{s-}^D) dD_s^c \right] \\ &\quad + \mathbb{E}_x \left[\sum_{0 \leq s \leq \gamma^\epsilon} e^{-rs} (v(t + s, X_s^D) - v(t + s, X_{s-}^D)) \right]. \end{aligned}$$

We notice that (i) implies $v_x \geq 1$ and therefore $v \geq 0$ follows from (ii). Using also $\mathcal{L}v \leq 0$ and rewriting

$$v(t + s, X_s^D) - v(t + s, X_{s-}^D) = - \int_0^{\Delta D_s} v_x(t + s, X_{s-}^D - y) dy$$

yields

$$v(t, x) \geq \mathbb{E}_x \left[\int_{0-}^{\gamma^\epsilon} e^{-rs} dD_s \right].$$

Letting $\epsilon \rightarrow 0$ gives

$$v(t, x) \geq \mathbb{E}_x \left[\int_{0-}^{\gamma^D \wedge (T-t)} e^{-rs} dD_s \right],$$

and since $D \in \mathcal{A}$ is arbitrary, $v \geq V$.

To prove the opposite inequality, let D^a be the strategy given by

$$D_s^a = \sup_{0 \leq u \leq s} (R_u - a(t+u))^+,$$

where $R_u = x + \mu u + \sigma B_u$. Then $X_s^{D^a} = R_s - D_s^a$, and we notice that (X^{D^a}, D^a) is the solution of the Skorokhod reflection problem at the boundary $a(\cdot)$. Therefore (iv), (v) and (6) give

$$v(t, x) = \mathbb{E}_x [e^{-r\gamma^\epsilon} v(t + \gamma^\epsilon, X_{\gamma^\epsilon}^{D^a})] + \mathbb{E}_x \left[\int_{0-}^{\gamma^\epsilon} e^{-rs} dD_s^a \right].$$

Recall that a is continuous with $a(T) = 0$. By continuity of v , and since $x \mapsto v(t, x)$ is increasing due to (i), we have by Dini's theorem that $0 \leq v(t + \gamma^\epsilon, X_{\gamma^\epsilon}^{D^a}) \leq h(\epsilon)$ for some function h with $h(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$. Thus

$$v(t, x) \leq h(\epsilon) + \mathbb{E}_x \left[\int_{0-}^{\gamma^\epsilon} e^{-rs} dD_s^a \right] \rightarrow \mathbb{E}_x \left[\int_{0-}^{\gamma^{D^a} \wedge (T-t)} e^{-rs} dD_s^a \right]$$

as $\epsilon \rightarrow 0$. Consequently, we also have the inequality $v \leq V$, and the strategy D^a is optimal. \square

Remark As noted above, the inequality $V \geq x$ always holds. Moreover, if $\mu \leq 0$, then actually $V = x$, and the optimal strategy is to immediately distribute the whole capital as dividends. To see this, notice that with $v(t, x) = x$ and $a(t) = 0$ we have $\mathcal{L}v = \mu - rx \leq 0$, so that (i)-(v) of Theorem 2.1 are satisfied. From now on, we only consider the case $\mu > 0$.

3. AN OPTIMAL STOPPING PROBLEM WITH LOCAL TIME

Our approach to solving the optimisation problem given by (3) is to connect it to a suitable problem of optimal stopping. In order to find the correct candidate for the latter we begin by making some useful heuristic observations.

If V satisfies the variational problem cast in Theorem 2.1, and if in addition V_t is continuous everywhere, then $V_t(t, 0) = 0$ for all $t \in [0, T)$ due to (ii). It follows that (iv) gives the boundary condition

$$(7) \quad \frac{\sigma^2}{2} V_{xx}(t, 0+) = -\mu V_x(t, 0+), \quad \text{for } t \in [0, T).$$

Setting $u := V_x$ we now notice that u should solve, at least formally,

$$(8) \quad \begin{cases} \mathcal{L}u = 0 & \text{for } (t, x) \in [0, T) \times (0, \infty) \text{ with } 0 < x < a(t), \\ u \geq 1 & \text{for } (t, x) \in [0, T) \times (0, \infty), \\ u = 1 & \text{for } (t, x) \in [0, T) \times (0, \infty) \text{ with } x \geq a(t), \\ u(T, x) = 1 & \text{for all } x \in [0, \infty), \end{cases}$$

with the additional boundary condition

$$(9) \quad u_x(t, 0+) = -\frac{2\mu}{\sigma^2} u(t, 0+) \quad \text{for } t \in [0, T).$$

The boundary value problem (8) is reminiscent of the one associated to an optimal stopping problem with payoff of immediate stopping equal to 1. Moreover, (9) is similar to the so-called *Feller's elastic boundary condition* at zero for a diffusion that lives on \mathbb{R}_+ (except for the minus sign on the right-hand side of (9)). This observation is the key to finding the right connection to optimal stopping.

Recall that (see for example [29, Ch. X, Exercise 1.15]), given a real-valued Markov process $(X_t)_{t \geq 0}$ and an additive functional A , one can construct the killed process

$$\tilde{X}_t = \begin{cases} X_t & \text{on } \{A_t < e\} \\ \Delta & \text{on } \{A_t \geq e\}. \end{cases}$$

Here e is an exponentially distributed random variable with parameter 1 which is independent of X , Δ is a cemetery state, and this process has the associated semigroup $(P_t)_{t \geq 0}$ given by

$$P_t f(x) = \mathbb{E}_x [e^{-A_t} f(X_t)].$$

Moreover, if $A_t = \lambda L_t^0$, where $(L_t^0)_{t \geq 0}$ is the local time of X at 0, then the process \tilde{X} is a diffusion with elastic behaviour at zero, the associated semigroup is given by

$$(10) \quad P_t f(x) = \mathbb{E}_x [e^{-\lambda L_t^0} f(X_t)],$$

and the infinitesimal generator coincides with the generator of X on the space of functions $u(x)$ with $u'(0+) - u'(0-) = 2\lambda u(0)$.

It is then clear that our problem boils down to finding the appropriate process X . Since (9) expresses a one-sided condition, and recalling the expression for \mathcal{L} in (4), we expect that in our case the process X should be a *reflected Brownian motion with drift μ and variance σ^2* , killed at a rate r . Hence, letting X be a reflected Brownian motion with drift μ and variance σ^2 , and L^0 its local time at zero, we are naturally led by (8) and (10) to consider the optimal stopping problem

$$(11) \quad U(t, x) = \sup_{0 \leq \tau \leq T-t} \mathbb{E}_x [e^{\lambda L_\tau^0 - r\tau}]$$

where $\lambda = 2\mu/\sigma^2$.

Remark Note that in problem (11) the presence of the local time corresponds to a *creation* of the process (rather than killing) at 0.

We notice that the reflected Brownian motion with drift is traditionally defined in terms of analytical properties of its infinitesimal generator. However, it is known that a useful equivalence in law holds between $(X, L^0(X))$ and a more explicit process. In fact, given a standard Brownian motion B , setting $Y_t = -\mu t + \sigma B_t$ and

$$S_t = \sup_{0 \leq s \leq t} Y_s,$$

it is shown in [27] that

$$(12) \quad (X_t^x, L^0(X^x)) \stackrel{\text{law}}{=} (x \vee S_t - Y_t, x \vee S_t - x).$$

From now on we identify

$$(13) \quad X_t^x := x \vee S_t - Y_t \quad \text{and} \quad L_t^0(X^x) = x \vee S_t - x.$$

Using (12)-(13) we can rewrite (11) in a more tractable form as

$$(14) \quad U(t, x) = \sup_{0 \leq \tau \leq T-t} \mathbb{E}_x \left[e^{\lambda L_\tau^0(X) - r\tau} \right] = \sup_{0 \leq \tau \leq T-t} \mathbb{E} \left[e^{\lambda(x \vee S_\tau - x) - r\tau} \right],$$

where the supremum is taken over all stopping times of B .

The following theorem describes the main findings of this article.

Theorem 3.1. (Connection) *Let U be the value function of the optimal stopping problem in (14), and denote by $b : [0, T] \rightarrow \mathbb{R}$ the corresponding optimal stopping boundary from Proposition 4.2 below. Then the value V of the dividend problem satisfies*

$$V(t, x) = \int_0^x U(t, y) dy,$$

and the dividend strategy $D_s^b = \sup_{0 \leq u \leq s} (x + \mu u + \sigma B_u - b(t+u))^+$ is optimal.

The proof of Theorem 3.1 is given in Section 7 below and it builds on the results of Sections 4-6 concerning U and b . Additional properties of V and b may also be deduced from the study in Sections 4-6 and we summarise some of them as follows.

Theorem 3.2. (Properties of the value function and the optimal boundary)

- (i) *The value function V belongs to $\mathcal{C}^{0,1}([0, T] \times [0, \infty)) \cap \mathcal{C}^{1,2}([0, T) \times (0, \infty))$.*
- (ii) *V satisfies $\max\{\mathcal{L}V, 1 - V_x\} = 0$ on $[0, T) \times (0, \infty)$.*
- (iii) *$x \rightarrow V(t, x)$ is concave, and $t \mapsto V(t, x)$ is non-increasing.*
- (iv) *The boundary $b : [0, T) \rightarrow (0, \infty)$ describing the optimal dividend strategy is non-increasing, continuous and satisfies $b(T-) = 0$.*

Remark It is worth observing that we may consider a more general controlled dynamic for the value of the fund, say

$$(15) \quad dX_t^D = \mu(X_t^D)dt + \sigma(X_t^D)dB_t - dD_t$$

for some functions μ and σ . Then the verification result Theorem 2.1 may be stated in a similar form, but with \mathcal{L} being the second order operator associated to $\mu(x)$ and $\sigma(x)$, and with the process D^a realising the Skorokhod downwards reflection of X^D at the boundary $a(\cdot)$. In this setting, the formal derivation of (8) requires to replace \mathcal{L} by $\tilde{\mathcal{L}} := \partial_t + \frac{\sigma^2}{2}\partial_{xx} + (\mu + \sigma\sigma')\partial_x - (r - \mu')$ and (9) by $u_x(t, 0+) = -\frac{2\mu(0)}{\sigma^2(0)}u(t, 0+)$.

It follows that (11) is again the natural candidate optimal stopping problem to be linked to (3), but now L^0 is the local time at zero of a diffusion

X associated to $\tilde{\mathcal{L}}$ and reflected at zero, $\lambda = 2\mu(0)/\sigma^2(0)$ and the constant killing rate r should be replaced by a level-dependent rate $r - \mu'$. Unfortunately, however, the general version of (11) cannot be reduced to a simpler form (as in (14)) because there seems to be no analogue of (12). In Sections 5 and 6 below we exploit the explicitness of (12) to derive sufficient regularity of U so that the verification result Theorem 2.1 can be applied to $v(t, x) = \int_0^x U(t, y) dy$. While we conjecture that $U = V_x$ holds also for more general absorbed diffusion processes, a full treatment of the general case is technically more demanding and is therefore left for future studies.

Remark One may add a discounted running cost $e^{-rt}f(t, X_t^D)$ in the formulation for (3). This would simply give rise to an additional running cost in (11) of the form $e^{\lambda L_t - rt}f_x(t, X_t)$.

Remark The optimal stopping problem (14) is closely connected to the Russian option with a finite horizon, see [12], [13], or [26]. In fact,

$$U(t, x) = e^{-\lambda x} \tilde{U}(t, x),$$

where

$$\tilde{U}(t, x) := \sup_{0 \leq \tau \leq T-t} \mathbb{E} \left[e^{\lambda(x \vee S_\tau) - r\tau} \right]$$

is the value of a Russian option written on a stock with current price 1, a historic maximal price $e^{\lambda x}$, volatility $\lambda\sigma$ and drift 0. While some parts of our analysis in Section 4 below can be deduced from studies of the Russian option, we choose, for the convenience of the reader, to include a detailed study. It should be noticed, however, that we go substantially beyond the regularity results contained in [12], [13], or [26] by proving that indeed U is \mathcal{C}^1 on $[0, T) \times (0, \infty)$.

4. ANALYSIS OF THE OPTIMAL STOPPING PROBLEM

Choosing $\tau = 0$ in (14) gives $U(t, x) \geq 1$. Denote by

$$\mathcal{C} := \{(t, x) \in [0, T] \times [0, \infty) : U(t, x) > 1\}$$

and

$$\mathcal{D} := \{(t, x) \in [0, T] \times [0, \infty) : U(t, x) = 1\}$$

the continuation region and the stopping region, respectively. We notice that assumptions in [25, Appendix D, Theorem D.12] are satisfied in our setting. Hence, from standard optimal stopping theory the stopping time

$$(16) \quad \begin{aligned} \tau^* &:= \tau_{t,x}^* &:= \inf\{s \geq 0 : U(t+s, X_s^x) = 1\} \\ & &= \inf\{s \geq 0 : (t+s, X_s^x) \in \mathcal{D}\} \end{aligned}$$

is optimal for the problem (14) in the sense that

$$U(t, x) = E \left[e^{\lambda(x \vee S_{\tau^*} - x) - r\tau^*} \right].$$

Moreover, for any $x \in \mathbb{R}_+$, the process

$$(17) \quad e^{-rs + \lambda L_s^0(X)} U(t+s, X_s), \quad s \geq 0,$$

is a \mathbb{P}_x -supermartingale, and

$$(18) \quad e^{-rs \wedge \tau^* + \lambda L_t^0(X)} U(t + s \wedge \tau^*, X_{s \wedge \tau^*}), \quad s \geq 0,$$

is a \mathbb{P}_x -martingale. In Proposition 4.2 below, we provide some qualitative properties of the shape of \mathcal{C} . First, however, we list a few properties of the value function U .

Proposition 4.1. (Properties of U) *The function $U : [0, T] \times [0, \infty) \rightarrow [0, \infty)$ is*

- (i) *equal to one at all points (T, x) , $x \in [0, \infty)$;*
- (ii) *non-increasing in t ;*
- (iii) *non-increasing and convex in x ;*
- (iv) *continuous on $[0, T] \times [0, \infty)$.*

Proof. The first property follows directly from the definition of U , and (ii) is obvious since the set of stopping times is decreasing in t .

For (iii), note that the function $e^{\lambda(x \vee S_\tau - x) - r\tau}$ is a.s. non-increasing and convex in x for any fixed τ . It follows that $\mathbb{E}[e^{\lambda(x \vee S_\tau - x) - r\tau}]$ is non-increasing and convex, and hence also U .

Finally, for (iv) we let $x_2 > x_1 \geq 0$ and $t \in [0, T]$. Then, by (iii),

$$\begin{aligned} 0 &\leq U(t, x_1) - U(t, x_2) \\ &\leq \sup_{0 \leq \tau \leq T-t} \mathbb{E} \left[e^{-r\tau} \left(e^{\lambda(x_1 \vee S_\tau - x_1)} - e^{\lambda(x_2 \vee S_\tau - x_2)} \right) \right] \\ &\leq \lambda(x_2 - x_1) \sup_{0 \leq \tau \leq T-t} \mathbb{E} \left[e^{\lambda S_\tau} \right] \\ &\leq \lambda(x_2 - x_1) \mathbb{E} \left[e^{\lambda S_T} \right], \end{aligned}$$

where the second last inequality follows from the fact that $e^{\lambda(x \vee s - x)}$ is Lipschitz continuous in x with constant $\lambda e^{\lambda s}$. This proves that U is Lipschitz continuous in x , uniformly in $t \in [0, T]$. Thus, to prove continuity of U it suffices to check that $t \mapsto U(t, x)$ is continuous for any fixed $x \in [0, \infty)$. To do that, fix x and let $t_1 \leq t_2$. Let $\tau^* = \tau_{t_1, x}^*$ be optimal for (t_1, x) , and define $\hat{\tau} := \tau^* \wedge (T - t_2)$. Then

$$\begin{aligned} 0 &\leq U(t_1, x) - U(t_2, x) \\ &\leq \mathbb{E} \left[e^{\lambda(x \vee S_{\tau^*} - x) - r\tau^*} - e^{\lambda(x \vee S_{\hat{\tau}} - x) - r\hat{\tau}} \right] \\ &= \mathbb{E} \left[\left(e^{\lambda(x \vee S_{\tau^*} - x) - r\tau^*} - e^{\lambda(x \vee S_{T-t_2} - x) - r(T-t_2)} \right) \mathbf{1}_{\{\tau^* \in (T-t_2, T-t_1)\}} \right] \\ &\leq e^{-r(T-t_2)} \mathbb{E} \left[e^{\lambda(x \vee S_{T-t_1} - x)} - e^{\lambda(x \vee S_{T-t_2} - x)} \right], \end{aligned}$$

which tends to 0 as $t_2 - t_1 \rightarrow 0$ by dominated convergence. This completes the proof of (iv). \square

Proposition 4.2. *There exists a boundary function $b : [0, T] \rightarrow (0, \infty)$ such that*

- (i) $\mathcal{C} = \{(t, x) \in [0, T] \times [0, \infty) : 0 \leq x < b(t)\}$;
- (ii) b is non-increasing;
- (iii) b is continuous on $[0, T]$ if one sets $b(T) := 0$.

Remark It follows that the optimal stopping time from (16) satisfies

$$\tau_{t,x}^* = \inf\{s \geq 0 : X_s^x \geq b(t+s)\}.$$

The function b is therefore referred to as **the optimal stopping boundary**.

Proof. The existence of a function $b : [0, T] \rightarrow [0, \infty]$ satisfying (i) is obvious from the fact that $x \mapsto U(t, x)$ is non-increasing. To prove that $b > 0$ on $[0, T]$, it suffices to check that $U(t, 0) > 1$ for $t \in [0, T]$. Choosing $\tau = \epsilon \leq T - t$ yields

$$U(t, 0) \geq \mathbb{E} \left[e^{\lambda S_\epsilon - r\epsilon} \right] \geq e^{-(r + \frac{2\mu^2}{\sigma^2})\epsilon} \mathbb{E} \left[e^{\sup_{0 \leq s \leq \epsilon} \frac{2\mu}{\sigma} B_s} \right].$$

By the reflection principle,

$$\mathbb{P} \left(\sup_{0 \leq s \leq \epsilon} B_s \in dz \right) = 2\mathbb{P}(B_\epsilon \in dz) = \sqrt{\frac{2}{\pi\epsilon}} e^{-\frac{z^2}{2\epsilon}} dz$$

for $z \geq 0$, so

$$\begin{aligned} \mathbb{E} \left[e^{\sup_{0 \leq s \leq \epsilon} \frac{2\mu}{\sigma} B_s} \right] &= \sqrt{\frac{2}{\pi\epsilon}} \int_0^\infty e^{\frac{2\mu z}{\sigma}} e^{-\frac{z^2}{2\epsilon}} dz = \sqrt{\frac{2}{\pi}} e^{\frac{2\mu^2\epsilon}{\sigma^2}} \int_{-\frac{2\mu\sqrt{\epsilon}}{\sigma}}^\infty e^{-z^2/2} dz \\ &\geq e^{\frac{2\mu^2\epsilon}{\sigma^2}} + \frac{2\mu}{\sigma} \sqrt{\frac{2\epsilon}{\pi}}. \end{aligned}$$

Consequently,

$$U(t, 0) \geq 1 + \frac{2\mu}{\sigma} \sqrt{\frac{2\epsilon}{\pi}} + \mathcal{O}(\epsilon)$$

as $\epsilon \downarrow 0$. This proves that $U(t, 0) > 1$, so $b(t) > 0$ for $t \in [0, T]$.

Next, note that (ii) is immediate from (ii) of Proposition 4.1. To prove that $b(t) < \infty$ we assume, to reach a contradiction, that $b(t) = \infty$ for some $t \in (0, T)$ (by time-homogeneity of the model, the assumption that $t > 0$ is without loss of generality). Then, since b is non-increasing, $b(s) = \infty$ for all $s \in [0, t]$, so given a starting point $(0, x)$, the optimal stopping time $\tau^* = \tau_{0,x}^*$ in (16) satisfies $\tau^* \geq t$. However,

$$\mathbb{E} \left[e^{\lambda(x \vee S_{\tau^*} - x) - r\tau^*} \right] \leq e^{-rt} \mathbb{E} \left[e^{\lambda(x \vee S_T - x)} \right] \rightarrow e^{-rt} < 1$$

as $x \rightarrow \infty$ by dominated convergence, since $e^{\lambda(x \vee S_T - x)} \leq e^{\lambda S_T} \in L^1(\mathbb{P})$. The latter inequality contradicts the optimality of τ^* . Thus $b(t) < \infty$ for all times $t \in [0, T]$.

From the continuity of U , the function b is lower semi-continuous, and thus (ii) implies that it is right-continuous. To finish the proof of (iii) it thus suffices to prove left-continuity on $(0, T]$. For this, assume (to reach a contradiction) that $b(t-) > b(t)$ for some $t \in (0, T]$. For $\epsilon \in (0, t)$, consider

the starting point (t_ϵ, x) , where $t_\epsilon = t - \epsilon$ and $x = (b(t-) + b(t))/2$, and define

$$\gamma_\epsilon := \inf\{s \geq 0 : X_s^x \notin (b(t), b(t-))\} \wedge \epsilon.$$

Notice that $\mathbb{P}_{t_\epsilon, x}(L_{\gamma_\epsilon}^0(X) = 0) = 1$, so that $x \vee S_{\gamma_\epsilon} - x = 0$ almost surely as well. Moreover $\mathbb{P}_{t_\epsilon, x}(\gamma_\epsilon \leq \tau^*) = 1$, and therefore the martingale property (18) yields

$$\begin{aligned} U(t_\epsilon, x) &= \mathbb{E} \left[e^{\lambda(x \vee S_{\gamma_\epsilon} - x) - r\gamma_\epsilon} U(t_\epsilon + \gamma_\epsilon, X_{\gamma_\epsilon}^x) \right] \\ &\leq \mathbb{E} \left[e^{-r\epsilon} 1_{\{\gamma_\epsilon = \epsilon\}} \right] + \mathbb{E} \left[U(t_\epsilon + \gamma_\epsilon, X_{\gamma_\epsilon}^x) 1_{\{\gamma_\epsilon < \epsilon\}} \right] \\ &\leq e^{-r\epsilon} + U(0, 0) \mathbb{P}(\gamma_\epsilon < \epsilon) \\ &= 1 - r\epsilon + o(\epsilon) \end{aligned}$$

as $\epsilon \rightarrow 0$, which contradicts $U \geq 1$. Thus $b(t-) = b(t)$, which completes the proof of (iii). \square

We end this section by stating that the value function U is a classical solution to a parabolic equation below the boundary. The proof of this fact is standard (e.g., see [25, Theorem 2.7.7]) and is therefore left out.

Proposition 4.3. *The value function U belongs to $\mathcal{C}^{1,2}$ separately in the interior of the continuation set \mathcal{C} and in the interior of the stopping set \mathcal{D} . Moreover it satisfies*

$$(19) \quad \mathcal{L}U(t, x) = 0 \quad \text{for } 0 < x < b(t) \text{ and } t \in [0, T),$$

$$(20) \quad \mathcal{L}U(t, x) = -r \quad \text{for } x > b(t) \text{ and } t \in [0, T).$$

5. FURTHER REGULARITY OF U : THE SPATIAL DERIVATIVE

In this section we prove that the value function U is continuously differentiable in the spatial variable, see Theorem 5.3.

Lemma 5.1. *Let*

$$\tau'_{t,x} := \inf\{s \geq 0 : X_s^x > b(t+s)\} \wedge (T-t).$$

Then $\tau'_{t,x} = \tau_{t,x}^$ a.s.*

Remark The proof of this follows the proof of [14, Lemma 6.2].

Proof. The claim is trivial for (t, x) such that $x > b(t)$ so we fix $(t, x) \in [0, T) \times (0, \infty)$ with $x \leq b(t)$. Since

$$\tau^* := \tau_{t,x}^* = \inf\{s \geq 0 : X_s^x \geq b(t+s)\} \wedge (T-t),$$

we have $\tau' := \tau'_{t,x} \geq \tau^*$. Moreover we notice that

$$(21) \quad \tau'_{t,b(t)} = 0 \quad \mathbb{P}\text{-a.s.}$$

due to the monotonicity of b and well-known properties of Brownian motion.

To prove also that $\tau'_{t,x} \leq \tau^*$ we introduce $Z_{s,s'} = \sup_{s \leq u \leq s'} (X_u^x - b(t+u))$ for any $0 < s < s' \leq T-t$. We claim that for arbitrary but fixed $0 < s_1 < s_2 \leq T-t$ one has

$$(22) \quad \mathbb{P}(Z_{s_1,s_2} = 0) = 0.$$

Let $\tau_1 := \inf\{u \geq s_1 : X_u^x = b(t+u)\} \wedge (T-t)$. Then

$$(23) \quad \mathbb{P}(Z_{s_1,s_2} = 0) = \mathbb{P}(Z_{s_1,s_2} = 0, \tau_1 \in [s_1, s_2]) + \mathbb{P}(Z_{s_1,s_2} = 0, \tau_1 = s_2)$$

because on the event $\{Z_{s_1,s_2} = 0\}$ it must be $\tau_1 \in [s_1, s_2]$.

For the first term of the expression on the right-hand side of (23) we have by continuity of X and b that

$$\begin{aligned} & \mathbb{P}(Z_{s_1,s_2} = 0, \tau_1 \in [s_1, s_2]) \\ &= \mathbb{P}(Z_{s_1,s_2} = 0, \tau_1 \in [s_1, s_2], X_{\tau_1}^x = b(t+\tau_1)) \\ &= \mathbb{E} \left[\mathbf{1}_{\{\tau_1 \in [s_1, s_2], X_{\tau_1}^x = b(t+\tau_1)\}} \mathbb{P}(Z_{s_1,s_2} = 0 | \mathcal{F}_{\tau_1}) \right] \\ &= \mathbb{E} \left[\mathbf{1}_{\{\tau_1 \in [s_1, s_2], X_{\tau_1}^x = b(t+\tau_1)\}} \mathbb{P}(Z_{s_1,\tau_1} \vee Z_{\tau_1,s_2} = 0 | \mathcal{F}_{\tau_1}) \right] \\ &= \mathbb{E} \left[\mathbf{1}_{\{\tau_1 \in [s_1, s_2], X_{\tau_1}^x = b(t+\tau_1)\}} \mathbb{P}(Z_{\tau_1,s_2} = 0 | X_{\tau_1}^x = b(t+\tau_1)) \right] = 0 \end{aligned}$$

where we have used the strong Markov property, the fact that $Z_{s_1,\tau_1} \mathbf{1}_{\{\tau_1 < s_2\}} = 0$ and that $\mathbb{P}(Z_{\tau_1,s_2} = 0 | X_{\tau_1}^x = b(t+\tau_1)) = 0$ due to (21).

For the second term on the right-hand side of (23) we simply have

$$\mathbb{P}(Z_{s_1,s_2} = 0, \tau_1 = s_2) \leq \mathbb{P}(X_{s_2}^x = b(t+s_2)) = 0$$

since $X_{s_2}^x$ has a continuous distribution, which finishes the proof of (22).

Now, if $\tau^* \in [s_1, s_2]$, then $Z_{s_1,s_2} \geq X_{\tau^*}^x - b(t+\tau^*) \geq 0$, so

$$\begin{aligned} \mathbb{P}(\tau^* \in [s_1, s_2]) &= \mathbb{P}(\tau^* \in [s_1, s_2], Z_{s_1,s_2} \geq 0) \\ &= \mathbb{P}(\tau^* \in [s_1, s_2], Z_{s_1,s_2} > 0) \\ &= \mathbb{P}(\tau^* \in [s_1, s_2], \tau' \in [s_1, s_2]). \end{aligned}$$

Consequently,

$$\mathbb{P}_{t,x}(s_1 \leq \tau^* \leq s_2, \tau' > s_2) = 0,$$

and since this holds for any rational s_1 and s_2 , we have $\tau^* = \tau'$ a.s. \square

Proposition 5.2. *The optimal stopping time $\tau_{t,x}^*$ is continuous in $(t, x) \in [0, T] \times [0, \infty)$.*

Proof. First notice that $x \mapsto X_t^x$ is a.s. Lipschitz continuous in x from (13), uniformly in $t \in [0, T]$. Fix $(t, x) \in [0, T] \times [0, \infty)$ and ω , and take a sequence $[0, T] \times [0, \infty) \ni (t_n, x_n) \rightarrow (t, x)$ as $n \rightarrow \infty$.

If $X_s^x > b(t+s)$ for some $s \in [0, T-t]$, i.e. $\tau'_{t,x} \leq s$, then continuity implies

$$X_s^{x_n} > b(t_n + s)$$

for n large, so $\tau'_{t_n, x_n} \leq s$ for any such n . Therefore $\limsup_n \tau'_{t_n, x_n} \leq \tau'_{t, x}$, since s was arbitrary and

$$(24) \quad \limsup_{n \rightarrow \infty} \tau_{t_n, x_n}^* = \limsup_{n \rightarrow \infty} \tau'_{t_n, x_n} \leq \tau'_{t, x} = \tau_{t, x}^*.$$

Next, if $s \in [0, T - t)$ is such that $X_u^x < b(t + u)$ for all $u \in [0, s]$, then $\inf_{u \in [0, s]} b(t + u) - X_u^x =: \delta > 0$ by continuity of b and X^x in time. By Lipschitz continuity of $x \mapsto X_u^x$, $\inf_{u \in [0, s]} b(t + u) - X_u^{x_n} \geq \delta/2$ for n large enough. By continuity (in time), this implies that $\inf_{u \in [0, s]} b(t_n + u) - X_u^{x_n} > 0$ for large n . Consequently, $\tau_{t_n, x_n}^* \geq s$, so

$$\liminf_{n \rightarrow \infty} \tau_{t_n, x_n}^* \geq \tau_{t, x}^*$$

since s was arbitrary. Together with (24), this yields $\lim_{n \rightarrow \infty} \tau_{t_n, x_n}^* = \tau_{t, x}^*$. \square

Theorem 5.3. *The spatial derivative $U_x(t, x)$ exists at all points $(t, x) \in [0, T] \times [0, \infty)$ and is continuous on $[0, T] \times [0, \infty)$. Moreover, it satisfies*

$$(25) \quad U_x(t, x) = -\lambda \mathbb{E} \left[\mathbf{1}_{\{S_{\tau^*} > x\}} e^{\lambda(S_{\tau^*} - x) - r\tau^*} \right].$$

Proof. We first show that the function

$$g(t, x) := -\lambda \mathbb{E} \left[\mathbf{1}_{\{S_{\tau^*} > x\}} e^{\lambda(S_{\tau^*} - x) - r\tau^*} \right]$$

is continuous on $[0, T] \times [0, \infty)$. To do that, assume that $[0, T] \times [0, \infty) \ni (t_n, x_n) \rightarrow (t, x) \in [0, T] \times [0, \infty)$ as $n \rightarrow \infty$. By Proposition 5.2, $\tau_{t_n, x_n}^* \rightarrow \tau_{t, x}^*$ a.s. Moreover, notice that

$$(26) \quad \mathbb{P}(S_{\tau_{t, x}^*} = x) = 0.$$

Indeed, define $\hat{\tau}_{t, x} := \inf\{s \geq 0 : S_s = x\} \wedge (T - t)$, so that $\hat{\tau}_{t, x}$ is also the first time that X^x equals zero. Since $s \mapsto S_s$ is increasing and $\mathbb{P}(S_{\hat{\tau}_{t, x} + u} > x) = 1$ for all $u > 0$, we have that $\mathbb{P}(S_{\tau_{t, x}^*} = x) = \mathbb{P}(\tau_{t, x}^* = \hat{\tau}_{t, x})$. However, since $b(t + s) > 0$ for $s \in [0, T - t)$ and $X_{\hat{\tau}_{t, x}}^x = 0$,

$$\mathbb{P}(\tau_{t, x}^* = \hat{\tau}_{t, x}) = \mathbb{P}(\tau_{t, x}^* = T - t) \leq \mathbb{P}(X_{T-t}^x = 0) = 0,$$

and hence (26) holds. By (26),

$$\mathbf{1}_{\{S_{\tau_{t_n, x_n}^*} > x_n\}} \rightarrow \mathbf{1}_{\{S_{\tau_{t, x}^*} > x\}}$$

a.s. as $n \rightarrow \infty$, and consequently,

$$\begin{aligned} g(t_n, x_n) &= -\lambda \mathbb{E} \left[\mathbf{1}_{\{S_{\tau_{t_n, x_n}^*} > x_n\}} e^{\lambda(S_{\tau_{t_n, x_n}^*} - x_n) - r\tau_{t_n, x_n}^*} \right] \\ &\rightarrow -\lambda \mathbb{E} \left[\mathbf{1}_{\{S_{\tau_{t, x}^*} > x\}} e^{\lambda(S_{\tau_{t, x}^*} - x) - r\tau_{t, x}^*} \right] = g(t, x) \end{aligned}$$

by dominated convergence. This shows that g is continuous on $[0, T] \times [0, \infty)$.

Now note that (25) holds for $t = T$ since $U(x, T) = 1$ and τ^* in that case equals 0.

Next, since $x \mapsto U(t, x)$ is convex for $t \in [0, T)$, its right derivative exists everywhere on $[0, \infty)$ and its left derivative exists everywhere on $(0, \infty)$, and the set of points where the right and the left derivative differ (for a fixed $t \in [0, T)$) is at most countable. Fix $(t, x) \in [0, T) \times (0, \infty)$ such that the right and the left (spatial) derivatives agree at (t, x) , and let $\epsilon > 0$. Denote $\tau^* = \tau_{t,x}^*$. Then

$$\begin{aligned} U(t, x) - U(t, x + \epsilon) &\leq \mathbb{E} \left[e^{-r\tau^*} \left(e^{\lambda(x \vee S_{\tau^*} - x)} - e^{\lambda((x+\epsilon) \vee S_{\tau^*} - (x+\epsilon))} \right) \right] \\ &= \mathbb{E} \left[e^{-r\tau^*} \left(e^{\lambda(S_{\tau^*} - x)} - 1 \right) \mathbf{1}_{\{x < S_{\tau^*} \leq x+\epsilon\}} \right] \\ &\quad + \mathbb{E} \left[e^{-r\tau^*} \left(e^{\lambda(S_{\tau^*} - x)} - e^{\lambda(S_{\tau^*} - (x+\epsilon))} \right) \mathbf{1}_{\{S_{\tau^*} > x+\epsilon\}} \right] \\ &\leq (e^{\lambda\epsilon} - 1) \mathbb{P}(x < S_{\tau^*} \leq x + \epsilon) \\ &\quad + (1 - e^{-\lambda\epsilon}) \mathbb{E} \left[e^{-r\tau^*} e^{\lambda(S_{\tau^*} - x)} \mathbf{1}_{\{S_{\tau^*} > x+\epsilon\}} \right]. \end{aligned}$$

Dividing by ϵ and using that

$$\lim_{\epsilon \downarrow 0} \mathbb{P}(x < S_{\tau^*} \leq x + \epsilon) = 0,$$

we obtain that the right (spatial) derivative at (t, x) satisfies

$$(27) \quad \lim_{\epsilon \downarrow 0} \frac{U(t, x + \epsilon) - U(t, x)}{\epsilon} \geq -\lambda \mathbb{E} \left[\mathbf{1}_{\{S_{\tau^*} > x\}} e^{\lambda(S_{\tau^*} - x) - r\tau^*} \right].$$

Similarly, for $\epsilon \in (0, x)$,

$$\begin{aligned} U(t, x) - U(t, x - \epsilon) &\leq \mathbb{E} \left[e^{-r\tau^*} \left(e^{\lambda(x \vee S_{\tau^*} - x)} - e^{\lambda((x-\epsilon) \vee S_{\tau^*} - (x-\epsilon))} \right) \right] \\ &= \mathbb{E} \left[e^{-r\tau^*} \left(1 - e^{\lambda(S_{\tau^*} - (x-\epsilon))} \right) \mathbf{1}_{\{x-\epsilon < S_{\tau^*} \leq x\}} \right] \\ &\quad + \mathbb{E} \left[e^{-r\tau^*} \left(e^{\lambda(S_{\tau^*} - x)} - e^{\lambda(S_{\tau^*} - (x-\epsilon))} \right) \mathbf{1}_{\{S_{\tau^*} > x\}} \right] \\ &\leq (1 - e^{\lambda\epsilon}) \mathbb{E} \left[e^{\lambda(S_{\tau^*} - x) - r\tau^*} \mathbf{1}_{\{S_{\tau^*} > x\}} \right], \end{aligned}$$

so the left (spatial) derivative satisfies

$$(28) \quad \lim_{\epsilon \downarrow 0} \frac{U(t, x) - U(t, x - \epsilon)}{\epsilon} \leq -\lambda \mathbb{E} \left[\mathbf{1}_{\{S_{\tau^*} > x\}} e^{\lambda(S_{\tau^*} - x) - r\tau^*} \right].$$

Since the derivative exists at (t, x) , it follows from (27) and (28) that U_x satisfies (25) at (t, x) . Moreover, since $U_x(t, \cdot)$ coincides with the continuous function $g(t, \cdot)$ outside a countable set, and since the right (left) derivative of a convex function is right (left) continuous, it follows that $U_x = g$ on $[0, T) \times [0, \infty)$, which completes the proof. \square

Corollary 5.4. (Creation condition) *The value function U satisfies the boundary condition $U_x(t, 0) + \lambda U(t, 0) = 0$ for $t < T$.*

Proof. For $t \in [0, T)$ we have $\tau_{0,t}^* > 0$ a.s., so $S_{\tau_{0,t}^*} > 0$ a.s. Thus

$$U_x(t, 0) = -\lambda \mathbb{E} \left[e^{\lambda S_{\tau_{0,t}^*} - r\tau_{0,t}^*} \right] = -\lambda U(t, 0).$$

□

Corollary 5.5. *The smooth fit condition holds, i.e. $U_x(t, b(t)) = 0$ for $t \in [0, T)$.*

Proof. This follows since $U(t, x) = 1$ for $x \geq b(t)$ and U_x is continuous. □

6. FURTHER REGULARITY OF U : THE TIME DERIVATIVE

In this section we show that the time derivative of U is continuous, see Theorem 6.2.

Lemma 6.1. *The function U is Lipschitz continuous in t on $[0, T_1] \times [0, \infty)$, uniformly with respect to x , for any $T_1 \in (0, T)$.*

Proof. Let $t_1, t_2 \in [0, T_1]$ with $t_1 < t_2$, let $x \in [0, \infty)$, and denote $\tau := \tau_{t_1, x}^*$. Then, recalling that $U(\cdot, x)$ is decreasing, we get

$$\begin{aligned} 0 &\leq U(t_1, x) - U(t_2, x) \\ &\leq \mathbb{E} \left[e^{\lambda(x \vee S_\tau - x) - r\tau} - e^{\lambda(x \vee S_{\tau \wedge (T-t_2)} - x) - r(\tau \wedge (T-t_2))} \right] \\ &= \mathbb{E} \left[\left(e^{\lambda(x \vee S_\tau - x) - r\tau} - e^{\lambda(x \vee S_{T-t_2} - x) - r(T-t_2)} \right) \mathbf{1}_{\{T-t_2 < \tau \leq T-t_1\}} \right] \\ &\leq e^{-\lambda x} e^{-r(T-t_2)} \mathbb{E} \left[e^{\lambda(x \vee S_{T-t_1})} - e^{\lambda(x \vee S_{T-t_2})} \right] \\ &\leq \mathbb{E} \left[e^{\lambda S_{T-t_1}} - e^{\lambda S_{T-t_2}} \right], \end{aligned}$$

where we for the last inequality used

$$e^{\lambda(x \vee S_{T-t_1})} - e^{\lambda(x \vee S_{T-t_2})} = \mathbf{1}_{\{x < S_{T-t_1}\}} \left(e^{\lambda S_{T-t_1}} - e^{\lambda(x \vee S_{T-t_2})} \right).$$

By explicit formulas, see e.g. [24, Section 3.5.C],

$$\mathbb{P}(S_t \geq z) = \int_0^t \frac{z}{\sigma \sqrt{2\pi s^3}} e^{-\frac{(z+\mu)^2}{2\sigma^2 s}} ds,$$

so

$$\mathbb{P}(S_t \in dz) = \int_0^t \frac{1}{\sigma \sqrt{2\pi s^3}} \left(\frac{z+\mu}{\sigma^2 s} z - 1 \right) e^{-\frac{(z+\mu)^2}{2\sigma^2 s}} ds dz.$$

Thus

$$\begin{aligned} \mathbb{E} \left[e^{\lambda S_t} \right] &= \int_0^\infty e^{\lambda z} \int_0^t \frac{1}{\sigma \sqrt{2\pi s^3}} \left(\frac{z+\mu}{\sigma^2 s} z - 1 \right) e^{-\frac{(z+\mu)^2}{2\sigma^2 s}} ds dz \\ &=: f(t) \end{aligned}$$

and

$$f'(t) = \int_0^\infty e^{\lambda z} \frac{1}{\sigma \sqrt{2\pi t^3}} \left(\frac{z+\mu}{\sigma^2 t} z - 1 \right) e^{-\frac{(z+\mu)^2}{2\sigma^2 t}} dz.$$

Since $f'(t)$ is bounded for $t \in [T - T_1, T]$, the function U is Lipschitz on $[0, T_1] \times [0, \infty)$ \square

The following theorem is the main result of the current section.

Theorem 6.2. *The time derivative U_t is continuous on $[0, T) \times (0, \infty)$.*

Proof. Fix $T_0 \in (0, T)$, let $t_1, t_2 \in [0, T_0]$ with $t_1 < t_2$ and $T_0 + t_2 - t_1 < T$, and let $x \in [0, \infty)$. Define $\tau := \tau_{t_1, x}^* \wedge (T_0 - t_2)$, and note that $U(t_1 + \tau, X_\tau^x) = U(t_2 + \tau, X_\tau^x) = 1$ on the set where $\tau = \tau_{t_1, x}^*$ thanks to (ii) in Proposition 4.1. Consequently, using (17) and (18), we have

$$\begin{aligned} 0 &\leq U(t_1, x) - U(t_2, x) \\ &\leq \mathbb{E} \left[e^{\lambda(x \vee S_\tau - x) - r\tau} (U(t_1 + \tau, X_\tau^x) - U(t_2 + \tau, X_\tau^x)) \right] \\ &= \mathbb{E} \left[e^{\lambda(x \vee S_\tau - x) - r\tau} (U(t_1 + \tau, X_\tau^x) - U(t_2 + \tau, X_\tau^x)) \mathbf{1}_{\{\tau = T_0 - t_2\}} \right] \\ &\leq \mathbb{E} \left[e^{\lambda S_{T_0 - t_2} - r(T_0 - t_2)} \left(U(T_0 + t_1 - t_2, X_{T_0 - t_2}^x) \right. \right. \\ &\quad \left. \left. - U(T_0, X_{T_0 - t_2}^x) \right) \mathbf{1}_{\{\tau = T_0 - t_2\}} \right] \\ &\leq C_0(t_2 - t_1) \mathbb{E} \left[e^{\lambda S_T} \mathbf{1}_{\{\tau = T_0 - t_2\}} \right], \end{aligned}$$

where C_0 is a Lipschitz constant of $t \mapsto U(t, x)$, see Lemma 6.1, which depends on T_0 . In particular, for $(t, x) \in \mathcal{C}$ with $t \leq T_0$, we have

$$\begin{aligned} 0 &\leq U_t(t, x) \leq C_0 \mathbb{E} \left[e^{\lambda S_T} \mathbf{1}_{\{\tau_{t, x}^* \geq T_0 - t\}} \right] \\ &\leq C_0 \sqrt{\mathbb{E} [e^{2\lambda S_T}]} \sqrt{\mathbb{P}(\tau_{t, x}^* \geq T_0 - t)} \end{aligned}$$

by the Cauchy-Schwarz inequality. By Proposition 5.2, $\tau_{t, x}^* \rightarrow \tau_{t_0, b(t_0)}^* = 0$ a.s. as $(t, x) \rightarrow (t_0, b(t_0))$ for $t_0 < T_0$. Consequently, $\mathbb{P}(\tau_{t, x}^* \geq T_0 - t) \rightarrow 0$, and hence $U_t(t, x) \rightarrow 0$ as $(t, x) \rightarrow (t_0, b(t_0))$. Since U_t is continuous in the interior of \mathcal{D} and in \mathcal{C} , this shows that U_t is continuous on $[0, T_0) \times (0, \infty)$. Consequently, since T_0 is arbitrary, this shows that U_t is continuous on $[0, T) \times (0, \infty)$. \square

Remark The C^1 -differentiability of the value function is typically referred to as the ‘smooth fit’ condition in optimal stopping theory. This is a well-known condition that can be utilized in perpetual problems to produce a candidate solution, which then can be verified to equal the value function. While the smooth fit condition is generally believed to hold also for time-dependent problems (with sufficiently smooth underlying data), a formal verification of this fact is often lacking. In fact, in most studies of time-dependent optimal stopping problems it is only shown that smooth fit holds in the *spatial* variable for each fixed time, see e.g. [12], [13] and [25, Lemma 2.7.8].

In that respect, Theorems 5.3 and 6.2 go beyond established theory for optimal stopping problems.

7. PROOF OF THEOREM 3.1

Define a new function $v : [0, T] \times [0, \infty) \rightarrow [0, \infty)$ by

$$v(t, x) = \int_0^x U(t, y) dy.$$

From Theorem 5.3 and Theorem 6.2 it is immediate to see that $v \in \mathcal{C}^{1,2}([0, T] \times (0, \infty))$ and it is continuous everywhere. Moreover $t \mapsto v(t, x)$ is decreasing and $x \mapsto v(t, x)$ is concave due to Proposition 4.1.

Now we want to show that (v, b) solves (i) to (v) of Theorem 2.1 so that $v = V$ and D^b as in (5) is optimal.

Since $U \geq 1$ and $U(T, x) = 1$ we have $v_x \geq 1$ and $v(T, x) = x$. For $t \in [0, T)$ and $0 < x < b(t)$ and recalling (19) we obtain

$$\begin{aligned} & (v_t + \mu v_x + \frac{\sigma^2}{2} v_{xx} - rv)(t, x) \\ &= \int_0^x (U_t - rU)(t, y) dy + \mu U(t, x) + \frac{\sigma^2}{2} U_x(t, x) \\ &= - \int_0^x (\frac{\sigma^2}{2} U_{xx} + \mu U_x)(t, y) dy + \mu U(t, x) + \frac{\sigma^2}{2} U_x(t, x) \\ &= \frac{\sigma^2}{2} U_x(t, 0) + \mu U(t, 0) = 0 \end{aligned}$$

where the last equality follows from the creation condition at zero of Corollary 5.4. Repeating the same calculation for $t \in [0, T)$ and $x \geq b(t)$ and using (20) we then find $v_t + \mu v_x + \frac{\sigma^2}{2} v_{xx} - rv \leq 0$. Therefore we have verified (i) to (v) of Theorem 2.1, since (ii) is obviously true.

8. CONCLUDING REMARKS

The connection established in Theorem 3.1 enables the use of techniques from optimal stopping theory in the study of the dividend problem. For example, the precise asymptotic behaviour of the boundary can be derived, and the boundary can be characterised in terms of an integral equation. Both these results have been derived in the context of Russian options (recall the remark at the end of Section 3) and we only provide their statements.

8.1. Asymptotic behaviour of the boundary ([13]). The optimal stopping boundary satisfies

$$\lim_{t \uparrow T} \frac{b(t)}{\sqrt{(T-t) \ln \frac{1}{T-t}}} = \sigma,$$

where we notice that our $b(t)$ is equal to $\lambda^{-1} \ln a_t$ with a_t as in [13]. This asymptotic behaviour is the first term in an expansion that was found in [17]. It is important to remark that in [17] the author needs an a priori assumption regarding additional regularity of b . Here, on the other hand, relying on the asymptotic formula from [13], together with the established

connection between the Russian option and the dividend problem, we do not require any additional assumptions.

8.2. An integral equation for the boundary ([26]). The boundary b solves the integral equation

$$(29) \quad 1 = \mathbb{E} \left[e^{\lambda(b(t) \vee S_{T-t} - b(t)) - r(T-t)} \right] + r \int_0^{T-t} e^{-rs} \mathbb{P} \left(X_s^{b(t)} \geq b(t+s) \right) ds.$$

The above formula may be deduced from [26] via algebraic transformations. Alternatively, it can be directly verified by applying Dynkin's formula to $e^{\lambda L_t^b(X) - rt} U(t, X_t)$, since $U \in \mathcal{C}^1$ with U_{xx} bounded, and using (19), (20) and $U(t, b(t)) = 1$. Moreover, b is the unique solution of (29) in the class of continuous and positive functions.

8.3. The infinite horizon case. Finally, we remark that the connection suggested by Theorem 3.1 between the infinite horizon dividend problem (see [28]) and the perpetual optimal stopping problem

$$U(x) = \sup_{\tau \geq 0} \mathbb{E} \left[e^{\lambda(x \vee S_\tau - x) - r\tau} \right]$$

(which is closely related to the value function of a Russian option, see [30]) also holds. Indeed, one way to see this is to do the obvious changes in the scheme of the current paper. Alternatively, since these problems can be solved explicitly, it is straightforward to check that $V = U'$ by explicit calculations. Notably, however, this connection seems unnoticed even in the perpetual setting.

REFERENCES

- [1] Avanzi, B. Strategies for dividend distribution: a review. *N. Am. Actuar. J.* 13, no. 2 (2009) 217-251.
- [2] Bather, J.A. and Chernoff, H. Sequential decisions in the control of a spaceship. *Proc. Fifth Berkeley Symposium on Mathematical Statistics and Probability* 3 (1966) 181-207.
- [3] Beneš, V.E., Shepp, L.A. and Witsenhausen, H.S. Some Solvable Stochastic Control Problems. *Stochastics* 4 (1980) 39-83.
- [4] Benth, F.E. and Reikvam, K. A connection between singular stochastic control and optimal stopping. *Appl. Math. Optim.* 49 (2004) 27-41.
- [5] Boetius, F. and Kohlmann, M. Connections between optimal stopping and singular stochastic control. *Stochastic Process. Appl.* 77 (1998) 253-281.
- [6] Budhiraja, A. and Ross, K. Optimal stopping and free boundary characterizations for some Brownian control problems. *Ann. Appl. Probab.* 18 (6) (2008) 2367-2391.
- [7] De Angelis, T. and Ferrari, G. A stochastic partially reversible investment problem on a finite time-horizon: free-boundary analysis. *Stochastic Process. Appl.* 124 (2014), 4080-4119.

- [8] De Angelis, T. and Ferrari, G. Stochastic nonzero-sum games: a new connection between singular control and optimal stopping. arXiv: (2016).
- [9] De Angelis, T., Ferrari, G. and Moriarty, J. A nonconvex singular stochastic control problem and its related optimal stopping boundaries. *SIAM J. Control Optim.* 53 (3) (2015) 1199-1223.
- [10] De Angelis, T., Ferrari, G. and Moriarty, J. A solvable two-dimensional degenerate singular stochastic control problem with non-convex costs. arXiv:1411.2428 (2015).
- [11] De Finetti, B. Su un'impostazione alternativa della teoria collettiva del rischio. *Transactions of the XVth international congress of Actuaries.* Vol. 2. No. 1 (1957), 433-443.
- [12] Duistermaat, J., Kyprianou, A., and van Schaik, K. Finite expiry Russian options. *Stochastic Process. Appl.* 115 (2005), 609-638.
- [13] Ekström, E. Russian options with a finite time horizon. *J. Appl. Probab.* 41 (2004), 313-326.
- [14] Ekström, E. and Janson, S. The inverse first-passage problem and optimal stopping. *Ann. Appl. Probab.* 26 (2016), 3154-3177.
- [15] El Karoui, N. and Karatzas, I. Probabilistic aspects of finite-fuel, reflected follower problems. *Acta Applicandae Math.* 11 (1988) 223-258.
- [16] Grandits, P. Optimal consumption in a Brownian model with absorption and finite time horizon. *Appl. Math. Optim.* 67 (2) (2013), 197-241.
- [17] Grandits, P. Existence and asymptotic behavior of an optimal barrier for an optimal consumption problem in a Brownian model with absorption and finite time horizon. *Appl. Math. Optim.* 69 (2) (2014), 233-271.
- [18] Grandits, P. An optimal consumption problem in finite time with a constraint on the ruin probability. *Finance Stoch.* 19 (4) (2015), 791-847.
- [19] Guo, X. and Tomecek, P. Connections between singular control and optimal switching. *SIAM J. Control Optim.* 47 (1) (2008) 421-443.
- [20] Jeanblanc-Piqué, M. and Shiryaev, A. Optimization of the flow of dividends. *Russian Math. Surveys* 50 (1995), no. 2, 257-277.
- [21] Karatzas, I. Probabilistic aspects of finite-fuel stochastic control. *Proc. Natl. Acad. Sci. USA* 82 (1985) 5579-5581.
- [22] Karatzas, I. and Shreve, S. Connections between optimal stopping and singular stochastic control I. Monotone follower problems, *SIAM J. Control Optim.* 22 (1984) 856-877.
- [23] Karatzas, I. and Shreve, S. Equivalent models for finite-fuel stochastic control. *Stochastics* 18 (1986) 245-276.
- [24] Karatzas, I. and Shreve, S. *Brownian motion and stochastic calculus.* Second edition. Graduate Texts in Mathematics, 113. Springer-Verlag, New York, 1991.
- [25] Karatzas, I. and Shreve, S. *Methods of mathematical finance. Applications of Mathematics (New York),* 39. Springer-Verlag, New York,

- 1998.
- [26] Peskir, G. The Russian option: finite horizon. *Finance Stoch.* 9 (2005), 251-267.
 - [27] Peskir, G. On reflecting Brownian motion with drift. *Proc. 37th ISCIE Int. Symp. Stoch. Syst. Theory Applic., Inst. Systems Control Inform. Engrs., Kyoto*, (2006), 1-5.
 - [28] Radner, R. and Shepp, L. Risk vs. profit potential: A model for corporate strategy. *J. Econom. Dynam. Control* 20, no. 8 (1996), 1373-1393.
 - [29] Revuz, D. and Yor, M. Continuous martingales and Brownian motion, Third edition. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 293. Springer-Verlag, Berlin, 1999.
 - [30] Shepp, L. and Shiryaev, A. The Russian option: reduced regret. *Ann. Appl. Probab.* 3 (1993), no. 3, 631-640.
 - [31] Shreve, S.E., Lehoczky, J.P. and Gaver, D.P. Optimal consumption for general diffusions with absorbing and reflecting barriers. *SIAM J. Control Optim.* 22 (1) (1984), 55-75.
 - [32] Taksar, M.I. Average optimal singular control and a related stopping problem. *Math. Oper. Res.* 10 (1) (1985) 63-81.

T. DE ANGELIS: SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LS2 9JT LEEDS, UK.

E. EKSTRÖM: DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY, BOX 480, 75106 UPPSALA, SWEDEN.

E-mail address: t.deangelis@leeds.ac.uk, ekstrom@math.uu.se