

This is a repository copy of *Sums of reciprocals of fractional parts and multiplicative Diophantine approximation*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/114575/>

Version: Accepted Version

Article:

Beresnevich, Victor orcid.org/0000-0002-1811-9697, Haynes, Alan
orcid.org/0000-0001-6077-8162 and Velani, Sanju orcid.org/0000-0002-4442-6316
(Accepted: 2017) Sums of reciprocals of fractional parts and multiplicative Diophantine approximation. *Memoirs of the American Mathematical Society*. ISSN 0065-9266 (In Press)

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Sums of reciprocals of fractional parts and multiplicative Diophantine approximation

V. Beresnevich* A. Haynes† S. Velani‡

Dedicated to Wolfgang M. Schmidt on N°80

Abstract

There are two main interrelated goals of this paper. Firstly we investigate the sums

$$S_N(\alpha, \gamma) := \sum_{n=1}^N \frac{1}{n \|n\alpha - \gamma\|}$$

and

$$R_N(\alpha, \gamma) := \sum_{n=1}^N \frac{1}{\|n\alpha - \gamma\|},$$

where α and γ are real parameters and $\|\cdot\|$ is the distance to the nearest integer. Our theorems improve upon previous results of W.M. Schmidt and others, and are (up to constants) best possible. Related to the above sums, we also obtain upper and lower bounds for the cardinality of

$$\{1 \leq n \leq N : \|n\alpha - \gamma\| < \varepsilon\},$$

valid for all sufficiently large N and all sufficiently small ε . This first strand of the work is motivated by applications to multiplicative Diophantine approximation, which are also considered. In particular, we obtain complete Khintchine type results for multiplicative simultaneous Diophantine approximation on fibers in \mathbb{R}^2 . The divergence result is the first of its kind and represents an attempt of developing the concept of ubiquity to the multiplicative setting.

Mathematics subject classification: 11K60, 11J71, 11A55, 11J83, 11J20, 11J70, 11K38, 11J54, 11K06, 11K50

Keywords: Multiplicative Diophantine approximation, Ostrowski expansion, uniform distribution

*VB: Research supported by EPSRC grant EP/J018260/1.

†AH: Research supported by EPSRC grants EP/F027028/1, EP/J00149X/1 and EP/M023540/1.

‡SV: Research supported by EPSRC grants EP/E061613/1, EP/F027028/1 and EP/J018260/1.

Contents

I	Problems and main results	5
1	Sums of reciprocals	5
1.1	Background	5
1.2	Homogeneous results and corollaries	10
1.2.1	Corollaries to Theorem 1.1 regarding $S_N(\alpha, 0)$	11
1.2.2	Corollaries to Theorem 1.2 regarding $R_N(\alpha, 0)$	12
1.2.3	Homogeneous lower bounds via Minkowski's Theorem	14
1.2.4	Almost sure behaviour of $R_N(\alpha, 0)$	17
1.2.5	The homogeneous estimates for algebraic numbers	20
1.3	Inhomogeneous results and corollaries	21
1.3.1	Upper bounds	21
1.3.2	Lower bounds	22
1.4	Linear forms revisited	24
2	Multiplicative Diophantine approximation	25
2.1	Background	25
2.2	Problems and main results	27
II	Developing techniques and establishing the main results	32
3	Ostrowski expansion	32
3.1	Continued fractions: the essentials	32

3.2	The Ostrowski expansion	33
4	Estimates for $\ n\alpha - \gamma\$	35
4.1	The homogeneous case	35
4.2	The inhomogeneous case	36
5	A gaps lemma and its consequences	41
6	Counting solutions to $\ n\alpha - \gamma\ < \varepsilon$	46
6.1	The homogeneous case	46
6.2	The inhomogeneous case	50
7	Establishing the homogeneous results on sums of reciprocals	51
7.1	Proof of Theorem 1.2	52
7.1.1	The upper bound in (1.20)	52
7.1.2	The lower bound in (1.20)	54
7.1.3	Proof of (1.21) and (1.22)	55
7.2	Proof of Theorem 1.1	56
8	Establishing the inhomogeneous results on sums of reciprocals	58
8.1	Proof of Theorem 1.6	59
8.1.1	The case $m > K$	59
8.1.2	The case $ \Sigma \leq 1/2, m \leq K$	60
8.1.3	The case $ \Sigma > 1/2, m \leq K$	63
8.2	Proof of Theorem 1.4	65
8.3	Proof of Theorem 1.5	66

8.4	Proof of Theorem 1.7	67
8.5	Proof of Theorem 1.8	69
9	Proofs of Theorems 2.2, 2.3 and 2.4	74
9.1	Preliminaries	74
9.2	Proof of Theorem 2.2	76
9.3	Proof of Theorem 2.3	76
9.4	Proof of Theorem 2.4	79
10	Proof of Theorem 2.1	80
10.1	A zero-one law and quasi-independence on average	80
10.2	Setting up a limsup set	81
10.3	Choosing the indexing set \mathcal{T}^*	84
10.4	Overlaps estimates for $E_{t,k}$	87
10.4.1	Preliminary analysis	87
10.4.2	Further analysis	89
10.4.3	The remaining case	91
10.5	The finale	95

Part I

Problems and main results

Notation

To simplify notation the Vinogradov symbols \ll and \gg will be used to indicate an inequality with an unspecified positive multiplicative constant. If $a \ll b$ and $a \gg b$ we write $a \asymp b$, and say that the quantities a and b are *comparable*. For a real number x , the quantity $\{x\}$ will denote the fractional part of x . Also, $[x] := x - \{x\}$ denotes the largest integer not greater than x and $\lceil x \rceil := -\lfloor -x \rfloor := \min\{m \in \mathbb{Z} : m \geq x\}$ denotes the smallest integer not less than x . Given a real number x , $\text{sgn}(x)$ will stand for 1 if $x > 0$, -1 if $x < 0$ and 0 if $x = 0$. Finally, $\|x\| = \min\{|x - m| : m \in \mathbb{Z}\}$ will denote the distance from $x \in \mathbb{R}$ to the nearest integer, and $\mathbb{R}^+ = (0, +\infty)$.

1 Sums of reciprocals

1.1 Background

Let $\alpha, \gamma \in \mathbb{R}$ and suppose that

$$\|n\alpha - \gamma\| > 0 \quad \forall n \in \mathbb{N}. \quad (1.1)$$

For $N \in \mathbb{N}$ consider the sums

$$S_N(\alpha, \gamma) := \sum_{n=1}^N \frac{1}{n\|n\alpha - \gamma\|}$$

and

$$R_N(\alpha, \gamma) := \sum_{n=1}^N \frac{1}{\|n\alpha - \gamma\|}.$$

Note that for any real number x we have that $\|x\|$ is the minimum of the fractional parts $\{x\}$ and $\{-x\}$. Hence, it is legitimate that the sums $S_N(\alpha, \gamma)$ and $R_N(\alpha, \gamma)$ are referred to as the ‘sums of reciprocals of fractional parts’.

The above sums can be related by the following well known partial summation formula: given sequences (a_n) and (b_n) with $n \in \mathbb{N}$

$$\sum_{n=1}^N a_n b_n = \sum_{n=1}^N (a_n - a_{n+1})(b_1 + \cdots + b_n) + a_{N+1}(b_1 + \cdots + b_N). \quad (1.2)$$

In particular, it is readily seen that

$$S_N(\alpha, \gamma) = \sum_{n=1}^N \frac{R_n(\alpha, \gamma)}{n(n+1)} + \frac{R_N(\alpha, \gamma)}{N+1}. \quad (1.3)$$

Motivated by a wide range of applications, bounds for the above sums have been extensively studied over a long period of time, in particular, in connection with counting lattice points in polygons, problems in the theory of uniform distribution, problems in the metric theory of Diophantine approximation, problems in dynamical systems and problems in electronics engineering (see, for example, [2], [3], [4], [21], [28], [29], [30], [38, pp.108-110], [41], [40], [48], [49], [50], [55],[56]). Schmidt has shown in [49] that for any $\gamma \in \mathbb{R}$ and for any $\varepsilon > 0$

$$(\log N)^2 \ll S_N(\alpha, \gamma) \ll (\log N)^{2+\varepsilon}, \quad (1.4)$$

for almost all $\alpha \in \mathbb{R}$. In other words, the set of α for which (1.4) fails is a set of Lebesgue measure zero. It should be emphasised that (1.4) is actually a simple consequence of a much more general result, namely Theorem 2 in [49], established in higher dimensions for sums involving linear forms of integral polynomials with real coefficients. To a large extent Schmidt's interest in understanding the behavior of sums such as $S_N(\alpha, \gamma)$ lies in applications to metric Diophantine approximation, more specifically to obtaining Khintchine type theorems. Our motivation is somewhat similar.

In the homogeneous case, that is to say when $\gamma = 0$, the inequalities (1.4) are known to be true with $\varepsilon = 0$ for any badly approximable α . This result was originally proved by Hardy and Littlewood [29, Lemma 3]. Today this classical statement can be found as a set exercise in the monograph [41, Exercise 3.12] of Kuipers and Niederreiter. However, there is a downside in that the set of badly approximable numbers is of Lebesgue measure zero and therefore we are only guaranteed comparability in (1.4) for α in a thin set.

Walfisz [55, (48_{II}) on p. 571] proved that for any $\varepsilon > 0$ we have that $S_N(\alpha, 0) \ll (\log N)^{3+\varepsilon}$ for almost every $\alpha \in \mathbb{R}$. In another paper [56, p. 787], he showed that for any $\varepsilon > 0$ we have that $R_N(\alpha, 0) \ll N(\log N)^{1+\varepsilon}$ for almost every $\alpha \in \mathbb{R}$. This

together with (1.3) implies the right hand side of (1.4) was essentially known to Walfisz. Indeed, such upper bounds can be deduced from the even earlier work of Behnke [4] and metric properties of continued fractions appearing in [37].

More generally (see [41, Exercise 3.12]) if $\alpha \in \mathbb{R}$ and $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a nondecreasing function such that

$$\inf_{n \in \mathbb{N}} nf(n) \|n\alpha\| > 0, \quad (1.5)$$

then

$$S_N(\alpha, 0) \ll (\log N)^2 + f(N) + \sum_{1 \leq n \leq N} \frac{f(n)}{n}. \quad (1.6)$$

A simple consequence of the Borel-Cantelli Lemma in probability theory (or equivalently the trivial convergence part of Khintchine's Theorem [32, Theorem 2.2] – see also §2), implies that if the sum

$$\sum_{n=1}^{\infty} \frac{1}{nf(n)} \quad (1.7)$$

converges then (1.5), and so (1.6), holds for almost all α . However, when (1.7) converges the last term of (1.6) grows at least as fast as $(\log N)^2 \log \log N$. This means that even when $\gamma = 0$, for almost all α the lower bound of $(\log N)^2$ in (1.4) does not coincide with the upper bound given by (1.6).

The true magnitude of $S_N(\alpha, 0)$ for almost every number was eventually discovered by Kruse [40, Theorem 6(b)]. He proved that for almost every $\alpha \in \mathbb{R}$

$$S_N(\alpha, 0) \asymp (\log N)^2. \quad (1.8)$$

Beyond this ‘almost sure’ result, Kruse [40, Theorem 1(g)] showed that for any irrational α

$$S_N(\alpha, 0) \gg \log q_K \cdot \log(N/q_K) + \sum_{n=1}^{K-1} \log q_n \cdot \log a_{n+1} + \sum_{n=1}^{K+1} a_n, \quad (1.9)$$

$$S_N(\alpha, 0) \ll \log q_K \cdot \log(N/q_K) + \sum_{n=1}^{K-1} \log q_n \cdot (1 + \log a_{n+1}) + \sum_{n=1}^{K+1} a_n \quad (1.10)$$

where $[a_0; a_1, a_2, \dots]$ is the continued fraction expansion of α and K is the largest integer such that the denominator q_K of $[a_0; a_1, a_2, \dots, a_K]$ satisfies $q_K \leq N$ (see

below for further details of continued fractions). Kruse also provides the following simplification of (1.10)

$$S_N(\alpha, 0) \ll (\log N)(\log q_K) + \sum_{n=1}^{K+1} a_n. \quad (1.11)$$

Observe that the estimate (1.9) is not always sharp. For example, if $\alpha = [1; 1, 1, \dots]$, the golden ratio, and $N = q_K$, the lower bound (1.9) becomes $S_N(\alpha, 0) \gg \log N$ and this is significantly smaller than the truth – see Theorem 1.1 below. In fact, based on this example it is simple to construct many other real numbers for which (1.9) is not optimal. In short, they correspond to real numbers that contain sufficiently large blocks of 1’s in their continued fraction expansion. In this paper we will rectify this issue. Moreover, we shall prove that the two sides of (1.11) are comparable and provide explicit constants.

Turning our attention to $R_N(\alpha, 0)$, we have already mentioned Walfisz’s result that $R_N(\alpha, 0) \ll N(\log N)^{1+\varepsilon}$ for almost every $\alpha \in \mathbb{R}$. Beyond this ‘almost sure’ result, Behnke [4, pp. 289-290] showed that for any irrational α

$$R_N(\alpha, 0) \ll N \log N + \sum_{\substack{n=1 \\ n \equiv 0 \pmod{q_k}}^N \frac{1}{\|n\alpha\|}, \quad (1.12)$$

where K is the same as in (1.9). The sum appearing on the right hand side of (1.12) may result in substantial spikes and needs to be analysed separately. Lang [43, Theorem 2, p. 37] has shown that if, for some increasing function $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, the inequality $q_{K+1} \ll q_K g(q_K)$ holds for all sufficiently large $K \in \mathbb{N}$, where q_K are the denominators of the principal convergents of the continued fraction expansion of α , then

$$R_N(\alpha, 0) \ll N \log N + Ng(N). \quad (1.13)$$

As is mentioned in [43, Remark 1, p. 40] both terms on the right of (1.13) are necessary. This should be interpreted in the following sense: there are functions g , irrationals α and arbitrarily large N such that the inequality in (1.13) can be reversed.

Although (1.13) is not optimal for all choices of α and N , for badly approximable α the result is precise. Indeed, when α is badly approximable, (1.13) becomes

$$N \log N \ll R_N(\alpha, 0) \ll N \log N. \quad (1.14)$$

for all $N > 1$. The latter was originally obtained by Hardy and Littlewood [28, 29] in connection with counting integer points in certain polygons in \mathbb{R}^2 . The upper bound of (1.14) also appears in the work [21, p.546] of Chowla. Recently, Lê and Vaaler [44] have investigated a much more general problem that involves the sums $R_N(\alpha, 0)$ and their higher dimensional generalisations. In particular, they prove in [44, Theorem 1.1] that

$$R_N(\alpha, 0) \gg N \log N \quad (1.15)$$

for all N irrespective of the properties of the irrational number α . Indeed, the implicit constant within \gg can be taken to be 1 provided N is sufficiently large. In the same paper, Lê and Vaaler show that inequality (1.15) is best possible for a large class of α . More precisely, they prove in [44, Theorem 1.3] that for every sufficiently large N and every $0 < \varepsilon < 1$ there exists a subset $X_{\varepsilon;N}$ of $\alpha \in [0, 1]$ of Lebesgue measure $\geq 1 - \varepsilon$ such that

$$R_N(\alpha, 0) \ll_{\varepsilon} N \log N \quad \forall \alpha \in X_{\varepsilon;N}.$$

The above results for $R_N(\alpha, 0)$ can in fact be derived from the work of Kruse [40]. More precisely, inequalities (75) and (76) in [40] state that for any irrational α and N sufficiently large:

$$\sum_{\substack{n=1 \\ n \not\equiv 0 \pmod{q_K}}^N \frac{1}{\|n\alpha\|} \asymp N \log q_K + q_K a_{K+1} \log \frac{a_{K+1}}{\max\{1, a_{K+1} - N/q_K\}}, \quad (1.16)$$

$$\sum_{\substack{n=1 \\ n \equiv 0 \pmod{q_K}}^N \frac{1}{\|n\alpha\|} \asymp q_K a_{K+1} (1 + \log(N/q_K)), \quad (1.17)$$

where K is the same as in (1.9). These can then be combined to give (see formula (77) in [40]) the estimate

$$R_N(\alpha, 0) \asymp N \log N + q_K a_{K+1} (1 + \log(N/q_K)). \quad (1.18)$$

The techniques developed in this paper allow us to re-establish the above estimates of Kruse with fully explicit constants.

One of our principle goals is to undertake an in-depth investigation of the sums $S_N(\alpha, \gamma)$ and $R_N(\alpha, \gamma)$, especially in the currently fragmented inhomogeneous case ($\gamma \neq 0$). The intention is to establish (up to constants) best possible upper and lower bounds for the sums in question. We begin with the homogeneous case.

1.2 Homogeneous results and corollaries

In the homogeneous case ($\gamma = 0$) we are pretty much able to give exact bounds for

$$S_N(\alpha, 0) := \sum_{1 \leq n \leq N} \frac{1}{n \|n\alpha\|} \quad \text{and} \quad R_N(\alpha, 0) := \sum_{n=1}^N \frac{1}{\|n\alpha\|}$$

that are valid for all irrational α . In order to state our results we require notions from the theory of continued fractions, which will be recalled in §3. For now let $q_k = q_k(\alpha)$ denote the denominators of the principal convergents and let $a_k = a_k(\alpha)$ denote the partial quotients of (the continued fraction expansion of) α . Also, given $k \in \mathbb{N}$, let

$$A_k = A_k(\alpha) := \sum_{i=1}^k a_i$$

denote the sum of the first k partial quotients of α . Our first result concerns the sum $S_N(\alpha, 0)$.

Theorem 1.1. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $N \in \mathbb{N}$ and let $K = K(N, \alpha)$ denote the largest integer satisfying $q_K \leq N$. Then, for all sufficiently large N*

$$\max \left\{ \frac{1}{2}(\log N)^2, A_{K+1} \right\} \leq S_N(\alpha, 0) \leq 33(\log N)^2 + 10A_{K+1}. \quad (1.19)$$

Remark 1.1. The term A_{K+1} appearing in (1.19) is natural since

$$\frac{1}{2q_i \|q_i \alpha\|} - 1 < a_{i+1} < \frac{1}{q_i \|q_i \alpha\|}$$

(see (3.5) below) and thus we have that

$$A_{K+1} \asymp \sum_{i=1}^K \frac{1}{q_i \|q_i \alpha\|}.$$

Note that it is possible to quantify explicitly the meaning of ‘sufficiently large N ’ in Theorem 1.1. The upper bound within (1.19) is a consequence of the following statement for the ‘wilder’ behaving sum $R_N(\alpha, 0)$. Note that the sum $R_N(\alpha, 0)$ is split into two subsets which are treated separately.

Theorem 1.2. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $N \geq q_3$ and let $K = K(N, \alpha)$ be the largest integer satisfying $q_K \leq N$. Then*

$$\frac{1}{24}N \log q_K - \left(\frac{1}{3} \log q_2 + \frac{1}{2}\right)N \leq \sum_{\substack{1 \leq n \leq N \\ n \neq 0, q_{K-1} \pmod{q_K}}} \frac{1}{\|n\alpha\|} \leq 64N \log q_K + 2q_3N, \quad (1.20)$$

and

$$q_{K+1} \log(1 + N/q_K) \leq \sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \frac{1}{\|n\alpha\|} \leq 4q_{K+1}(1 + \log(1 + N/q_K)). \quad (1.21)$$

Furthermore, let $c > 0$. Then

$$\sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \min \left\{ cN, \frac{1}{\|n\alpha\|} \right\} \leq 12 N (ca_{K+1})^{\frac{1}{2}}. \quad (1.22)$$

Remark 1.2. Note that the upper and lower bounds in (1.20) and (1.21) are comparable and are thus (up to constants) best possible. The absolute constants appearing in the above results (and indeed elsewhere) can be improved. For the sake of clarity, during the course of proving our results, we do not attempt to obtain the sharpest possible constants, let alone asymptotic formulae. Instead, we aim to minimize technical details.

1.2.1 Corollaries to Theorem 1.1 regarding $S_N(\alpha, 0)$

First of all, we have the following straightforward consequence of Theorem 1.1.

Corollary 1.1. *Suppose $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ satisfies the condition*

$$A_{k+1} = o(k^2). \quad (1.23)$$

Then, for all sufficiently large N

$$\frac{1}{2} (\log N)^2 \leq S_N(\alpha, 0) \leq 34 (\log N)^2. \quad (1.24)$$

Proof. This result follows from (1.19), (1.23), the fact that $q_K \leq N$, and the estimate $K \ll \log q_K$ (see (3.10) below). \square

The set of $\alpha \in \mathbb{R}$ satisfying condition (1.23) is of full Lebesgue measure. In fact, for any $\varepsilon > 0$ the set of $\alpha \in \mathbb{R}$ such that $A_k \leq k^{1+\varepsilon}$ for all sufficiently large k is of full Lebesgue measure (see [22]). Thus we have the following result.

Corollary 1.2. *The upper bound in (1.24) holds for almost every real number α , and the lower bound holds for all α .*

1.2.2 Corollaries to Theorem 1.2 regarding $R_N(\alpha, 0)$

We now discuss various consequences of Theorem 1.2. By definition, $\log q_K \leq \log N$. Therefore, the sum in (1.20) is always $\ll N \log N$. In fact, we shall see that the sum is actually comparable to $N \log N$ unless α is a Liouville number. Throughout the paper \mathfrak{L} will denote the set of Liouville numbers; i.e. $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ such that

$$\liminf_{n \rightarrow \infty} n^w \|n\alpha\| = 0 \quad \forall w > 0.$$

It follows from the Jarník-Besicovitch Theorem [17, 35] (alternatively see [9]) that

$$\dim \mathfrak{L} = 0; \tag{1.25}$$

that is, \mathfrak{L} has zero Hausdorff dimension.

Corollary 1.3. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Then $\alpha \notin \mathfrak{L}$ if and only if for all sufficiently large N*

$$\sum_{\substack{1 \leq n \leq N \\ n \neq 0, q_{K-1} \pmod{q_K}}} \frac{1}{\|n\alpha\|} \asymp N \log N. \tag{1.26}$$

To establish Corollary 1.3, we will make use of well known facts about the growth of the denominators q_k associated with Liouville numbers. Recall that the exponent of approximation of $\alpha \in \mathbb{R}$ is defined as

$$w(\alpha) := \sup\{w > 0 : \|q\alpha\| < q^{-w} \text{ for i.m. } q \in \mathbb{N}\}. \tag{1.27}$$

Note that, for an irrational α , by definition, $w(\alpha) < \infty$ if and only if α is not Liouville. Also, by Dirichlet's Theorem, $w(\alpha) \geq 1$ for all α .

Lemma 1.1. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and let q_k denote the denominators of the principal convergents of α . Then*

$$w(\alpha) = \limsup_{k \rightarrow \infty} \frac{\log q_{k+1}}{\log q_k}. \tag{1.28}$$

In particular, $\alpha \notin \mathfrak{L}$ if and only if $\log q_{k+1} \ll \log q_k$ for all k .

Proof. This result is well known, however the proof is short and we give it for completeness. It easily follows from the definition of $w(\alpha)$ that

$$w(\alpha) = \limsup_{q \rightarrow \infty} \frac{\log \|q\alpha\|^{-1}}{\log q}.$$

Recall that the principal convergents of the continued fraction expansion of α are best approximations, that is $\|q_k\alpha\| \leq \|q\alpha\|$ whenever $q_k \leq q < q_{k+1}$. Hence

$$w(\alpha) = \limsup_{k \rightarrow \infty} \frac{\log \|q_k\alpha\|^{-1}}{\log q_k}.$$

Since $(2q_{k+1})^{-1} < \|q_k\alpha\| < q_{k+1}^{-1}$, this expression for $w(\alpha)$ immediately implies the required result. \square

Proof of Corollary 1.3. Assume that $\alpha \notin \mathfrak{L}$. Then, by Lemma 1.1, $\log q_K \asymp \log q_{K+1}$. Since, by definition, $q_K \leq N < q_{K+1}$, we also have that $\log N \asymp \log q_K$. By Theorem 1.2, the sum in (1.20) is comparable to $N \log q_K \asymp N \log N$. This proves one direction of Corollary 1.3. Now assume that $\alpha \in \mathfrak{L}$. Then, by Lemma 1.1, there is a sequence of K_i such that $\log q_{K_i+1} / \log q_{K_i} \rightarrow \infty$ as $i \rightarrow \infty$. Taking $N = N_i := q_{K_i+1} - 1$ implies that the sum in (1.20) is $\asymp N_i \log q_{K_i} = o(N_i \log N_i)$ as $i \rightarrow \infty$. This shows that (1.26) does not hold for Liouville numbers and thereby completes the proof. \square

Another consequence of (1.20) is that the sum in (1.26) can become $o(N \log N)$, along a subsequence of N , when α is a Liouville number. Nevertheless, on combining the estimates (1.20) and (1.21) we are able to recover the lower bound (1.15) obtained by L e and Vaaler in [44]. At this point it is worth formally restating (1.15) as a result in its own right.

Corollary 1.4. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Then, for all sufficiently large N*

$$R_N(\alpha, 0) \gg N \log N. \tag{1.29}$$

Proof. If $N \leq q_K^2$ then $\log q_K \geq \frac{1}{2} \log N$ and Corollary 1.4 is a consequence of (1.20) appearing within Theorem 1.2. Otherwise $N/q_K > N^{1/2}$, and it follows that $\log(1 + N/q_K) \geq \frac{1}{2} \log N$. In addition, by the definition of q_K , we have that $q_{K+1} > N$ and so in this case Corollary 1.4 is a consequence of (1.21) appearing within Theorem 1.2. \square

The proof of (1.29) given in [44] has its basis in harmonic analysis. However, we will demonstrate in the next section that such homogeneous lower bound estimates do not require either the full power of Theorem 1.2 or indeed the harmonic analysis tools utilized in [44].

1.2.3 Homogeneous lower bounds via Minkowski's Theorem

It turns out that appropriately exploiting Minkowski's Convex Body Theorem [20, p.71] from the geometry of numbers, leads to sharper lower bounds for $S_N(\alpha, 0)$ and $R_N(\alpha, 0)$ than those described above in Corollary 1.2 and Corollary 1.4. Furthermore, the approach provides lower bounds within the more general linear forms setting. To start with, we prove the following key statement, which only uses Minkowski's theorem for convex bodies and partial summation.

Theorem 1.3. *Let $A = (\alpha_1, \dots, \alpha_n)$ be any n -tuple of real numbers such that $1, \alpha_1, \dots, \alpha_n$ are linearly independent over \mathbb{Q} . Let T_1, \dots, T_n be any positive integers such that $T := T_1 \cdots T_n \geq 2$ and let $L \geq 2$ be a real number. Then*

$$\sum_{\substack{\mathbf{q} \in \mathbb{Z}^n \setminus \{\mathbf{0}\} \\ |q_j| \leq T_j \ (1 \leq j \leq n)}} \min \left\{ L, \|q_1 \alpha_1 + \cdots + q_n \alpha_n\|^{-1} \right\} \geq 2T \min\{\log L, \log T\} + (2^{n+1} - 2 - \log 4)T + 4. \quad (1.30)$$

Proof. Since both sides of (1.30) depend on L continuously, we can assume without loss of generality that $L > 2$. Also in the case $T = 2$ the left hand side is at least $2 \times \prod_{i=1}^n (2T_i) = 2^{n+2}$, while the right hand side is $4 \log 2 + (2^{n+1} - 2 - \log 4)2 + 4 = 2^{n+2}$. Thus, (1.30) holds for $T = 2$ and we can assume without loss of generality that

$$\min\{T, L\} > 2. \quad (1.31)$$

Fix any $b \in \mathbb{R}$ with $1 < b < \sqrt{2}$ such that for some $m \in \mathbb{Z}$,

$$b^m = \min\{T, L\}. \quad (1.32)$$

Let $m_0 \in \mathbb{N}$ satisfy

$$b^{m_0} \leq 2 < b^{m_0+1}. \quad (1.33)$$

Since $b < \sqrt{2}$, we have that $m_0 \geq 2$. By (1.31), $b^m > 2$ and so $2 \leq m_0 < m$. Next, given an integer $k \geq 1$, let

$$N_k := \left\{ \mathbf{q} \in \mathbb{Z}^n \setminus \{\mathbf{0}\} : |q_j| \leq T_j \ (1 \leq j \leq n), \ b^{-k-1} < \left\| \sum_{j=1}^n \alpha_j q_j \right\| \leq b^{-k} \right\}$$

and let $\Phi_k := \bigcup_{\ell=k}^{\infty} N_\ell$. Obviously the sets N_ℓ are disjoint and so $\#\Phi_k = \sum_{\ell=k}^{\infty} \#N_\ell$, where $\#X$ stands for the cardinality of X . Note that, by definition, Φ_k consists precisely of integer vectors $\mathbf{q} \in \mathbb{Z}^n \setminus \{\mathbf{0}\}$ such that $|q_j| \leq T_j$

($1 \leq j \leq n$) and $\|\sum_{j=1}^n \alpha_j q_j\| \leq b^{-k}$. Hence, by Minkowski's Convex Body Theorem we conclude that

$$\#\Phi_k \geq 2\lfloor b^{-k}T \rfloor \geq 2b^{-k}T - 2 \quad \text{when } b^k \leq T. \quad (1.34)$$

Furthermore, note that we trivially have that

$$\#\Phi_k = (2T_1 + 1) \cdots (2T_n + 1) - 1 \geq 2^n T \quad \text{when } b^k \leq 2; \quad (1.35)$$

that is, when $k \leq m_0$. Then, since $\#N_k = \#\Phi_k - \#\Phi_{k+1}$, it follows that

$$\begin{aligned} \sum_{\substack{\mathbf{q} \in \mathbb{Z}^n \setminus \{\mathbf{0}\} \\ |q_j| \leq T_j \ (1 \leq j \leq n)}} \min \left\{ L, \left\| \sum_{j=1}^n \alpha_j q_j \right\|^{-1} \right\} &\geq \sum_{k=1}^{\infty} \sum_{\mathbf{q} \in N_k} \min\{L, b^k\} \\ &\geq \sum_{k=1}^{\infty} \sum_{\mathbf{q} \in N_k} \min\{b^m, b^k\} \\ &= \sum_{k=1}^m \sum_{\mathbf{q} \in N_k} b^k + \sum_{k=m+1}^{\infty} \sum_{\mathbf{q} \in N_k} b^m \\ &= \sum_{k=1}^m \#N_k b^k + \#\Phi_{m+1} b^m \\ &= \sum_{k=1}^m (\#\Phi_k - \#\Phi_{k+1}) b^k + \#\Phi_{m+1} b^m \\ &= \sum_{k=1}^m \#\Phi_k b^k - \sum_{k=2}^{m+1} \#\Phi_k b^{k-1} + \#\Phi_{m+1} b^m \\ &= \#\Phi_1 b + \sum_{k=2}^m \#\Phi_k (b^k - b^{k-1}) \\ &= \#\Phi_1 b + (b-1) \sum_{k=2}^m b^{k-1} \#\Phi_k \\ &= \#\Phi_1 b + (b-1) \sum_{k=2}^{m_0} b^{k-1} \#\Phi_k + (b-1) \sum_{k=m_0+1}^m b^{k-1} \#\Phi_k \\ &\stackrel{(1.34) \ \& \ (1.35)}{\geq} 2^n T b + (b-1) 2^n T \sum_{k=2}^{m_0} b^{k-1} + (b-1) \sum_{k=m_0+1}^m b^{k-1} (2b^{-k}T - 2) \\ &= 2^n T b + 2^n T (b^{m_0} - b) + 2(b-1)b^{-1} \sum_{k=m_0+1}^m (T - b^k) \\ &\stackrel{(1.33)}{\geq} 2^{n+1} T b^{-1} + 2(b-1)b^{-1} \sum_{k=m_0+1}^m (T - b^k) \end{aligned}$$

$$\begin{aligned}
&= 2^{n+1}Tb^{-1} + 2(b-1)b^{-1}\left((m-m_0)T - \frac{b^{m+1} - b^{m_0+1}}{b-1}\right) \\
&= 2^{n+1}Tb^{-1} + 2(b-1)b^{-1}(m-m_0)T - 2b^m + 2b^{m_0} \\
&\stackrel{(1.32)}{=} 2^{n+1}Tb^{-1} + 2(b-1)b^{-1}(m-m_0)T - 2\min\{L, T\} + 2b^{m_0}.
\end{aligned}$$

Now since

$$m = \frac{\min\{\log L, \log T\}}{\log b}, \quad m_0 \leq \frac{\log 2}{\log b} \quad \text{and} \quad 2b^{-1} < b^{m_0},$$

we have that the above is

$$\geq 2^{n+1}Tb^{-1} + 2T\frac{b-1}{b\log b}\left(\min\{\log L, \log T\} - \log 2\right) - 2\min\{L, T\} + 4b^{-1}.$$

Since this estimate holds for b arbitrarily close to 1 and since $\lim_{b \rightarrow 1^+} \frac{b-1}{b\log b} = 1$, we obtain that the above sum is

$$\begin{aligned}
&\geq 2^{n+1}T + 2T\left(\min\{\log L, \log T\} - \log 2\right) - 2\min\{L, T\} + 4 \\
&= 2T\min\{\log L, \log T\} + (2^{n+1} - \log 4)T - 2\min\{L, T\} + 4,
\end{aligned}$$

whence the required estimate immediately follows. \square

Remark 1.3. It is worth mentioning that a similar argument can be given for the sums of products investigated in [44].

Theorem 1.3 together with formula (1.3), yields the following lower bound estimates for $R_N(\alpha, 0)$ and $S_N(\alpha, 0)$, that are sharper than those given by Corollary 1.2 and Corollary 1.4. We leave the details of the proof to the reader.

Corollary 1.5. *For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and any integer $N \geq 2$*

$$R_N(\alpha, 0) \geq N \log N + N \log(e/2) + 2$$

and

$$S_N(\alpha, 0) \geq \frac{1}{2}(\log N)^2.$$

Remark 1.4. Another consequence of Theorem 1.3 is that

$$\sum_{\substack{\mathbf{q} \in \mathbb{Z}^n \setminus \{\mathbf{0}\} \\ |q_j| \leq T_j \ (1 \leq j \leq n)}} \frac{1}{\|\Pi_+(\mathbf{q})\| \|q_1\alpha_1 + \cdots + q_n\alpha_n\|} \gg (\log T)(\log T_1) \cdots (\log T_n),$$

where $\alpha_1, \dots, \alpha_n$ and T_1, \dots, T_n are as above in Theorem 1.3 and

$$\Pi_+(\mathbf{q}) = \prod_{q_i \neq 0} |q_i|, \quad \text{where } \mathbf{q} = (q_1, \dots, q_n).$$

We omit the details of the proof since in §1.4 below we will prove a more general inhomogeneous statement.

By analogy with the one-dimensional case (see also [49, Theorem 2]), it makes sense to make the following upper bound conjecture.

Conjecture 1.1. *Let T_1, \dots, T_n be positive integer parameters and $T = T_1 \cdots T_n$. Then, as $T \rightarrow \infty$, for almost every $(\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$ we have that*

$$\sum_{\substack{\mathbf{q} \in \mathbb{Z}^n \setminus \{\mathbf{0}\} \\ |q_j| \leq T_j \ (1 \leq j \leq n)}} \frac{1}{\Pi_+(\mathbf{q}) \|q_1 \alpha_1 + \cdots + q_n \alpha_n\|} \ll (\log T)(\log T_1) \cdots (\log T_n). \quad (1.36)$$

Remark 1.5. Bounds such as (1.36) are instrumental in the theory of uniform distribution to study the discrepancy of Kronecker sequences $(n(\alpha_1, \dots, \alpha_n))_{n \in \mathbb{N}}$, see for example [2], [23, §1.4.2], [34] or [39, §6]. Note that Lemma 4.4 in [2] (see also [23, Lemma 1.95]) implies that the left hand side of (1.36) is $\ll (\log T)^{n+2}$. When $T_1 = \cdots = T_n$, this is $\log T$ times bigger than the conjectured bound of $(\log T)^{n+1}$. It may well be that the proof of Lemma 4.1 in [2] (see also [23, Lemma 1.93]) can be adapted to prove the conjecture in the symmetric case. It is worth highlighting the fact that compared to the sum in (1.36) the range of summation in [2, Lemma 4.1] is restricted by the addition condition that

$$\Pi_+(\mathbf{q}) \|q_1 \alpha_1 + \cdots + q_n \alpha_n\| \ll (\log T)^{20n}.$$

1.2.4 Almost sure behaviour of $R_N(\alpha, 0)$

Corollary 1.2 completely describes the almost sure behaviour of the sum $S_N(\alpha, 0)$; namely that for almost all $\alpha \in \mathbb{R}$ we have that

$$\frac{1}{2} (\log N)^2 \leq S_N(\alpha, 0) \leq 34 (\log N)^2,$$

for all sufficiently large $N \in \mathbb{N}$. In this section, we make use of Theorem 1.2 to investigate the almost sure behaviour of the sum $R_N(\alpha, 0)$.

In order to understand the almost sure behaviour of inequalities (1.21) and (1.22) associated with Theorem 1.2, we first recall from §1.1, that by the Borel-Cantelli Lemma, if $f : \mathbb{N} \rightarrow \mathbb{R}^+$ is any function such that the sum (1.7) converges, then for almost all $\alpha \in \mathbb{R}$ one has that $qf(q)\|q\alpha\| \geq 1$ for all sufficiently large q . In particular, taking $q = q_K$, where $q_K = q_K(\alpha)$ is as before the sequence of denominators of the principle convergents of α , we have that $\|q_K\alpha\| \geq (q_K f(q_K))^{-1}$ for sufficiently large K . In addition, we have the following well known inequalities from the theory of continued fractions [37]:

$$(2q_{K+1})^{-1} < \|q_K\alpha\| < q_{K+1}^{-1},$$

see also (3.5) below. Hence, whenever (1.7) converges, for almost every α we have that

$$q_{K+1} < q_K f(q_K)$$

for sufficiently large K .

On the other hand, the main substance of Khintchine's Theorem, the divergent part (namely, (2.2) below with $k = 1$), implies that if f is monotonic and the sum (1.7) diverges, then for almost all α we have that $qf(q)\|q\alpha\| < 1$ for infinitely many $q \in \mathbb{N}$. Since convergents are best approximations, for almost all α we have that $2q_K f(q_K)\|q_K\alpha\| < 1$ for infinitely many K . Thus, in the case f is monotonic and the sum (1.7) diverges, it follows that for almost all α we have that

$$q_{K+1} > q_K f(q_K)$$

for infinitely many K .

The above observations regarding the size of q_{K+1} together with (1.21) of Theorem 1.2 give rise to the following statement.

Corollary 1.6. *Let $f : \mathbb{N} \rightarrow (0, \infty)$ be any increasing function. If (1.7) converges, then for almost all $\alpha \in \mathbb{R}$ we have that*

$$\sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \frac{1}{\|n\alpha\|} \ll q_K f(q_K) \log(1 + N/q_K) \ll N f(N),$$

for all sufficiently large $N \in \mathbb{N}$. On the other hand, if (1.7) diverges, then for almost all α there are infinitely many $N \in \mathbb{N}$ such that

$$\sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \frac{1}{\|n\alpha\|} \gg N f(N).$$

Example 1.1. Let $f(q) = \log q \log \log q$. Then, Corollary 1.6 implies that for almost all $\alpha \in \mathbb{R}$

$$R_N(\alpha, 0) \gg N \log N \log \log N \quad \text{for infinitely many } N \in \mathbb{N}.$$

However, if $f(q) = \log q (\log \log q)^{1+\varepsilon}$, then Corollary 1.6 together with Corollary 1.3 (and the well known fact that the set of Liouville numbers are a set of measure zero), shows that for almost all $\alpha \in \mathbb{R}$

$$R_N(\alpha, 0) \ll N \log N (\log \log N)^{1+\varepsilon} \quad \text{for all sufficiently large } N \in \mathbb{N}.$$

Finally, we analyze the almost all behavior of the sum appearing in (1.22), in which the possible spikes of $\|n\alpha\|^{-1}$ are “trimmed” to be no more than cN . Once again, we appeal to the Borel-Cantelli Lemma, which implies that if (1.7) converges then, for almost all α , we have $q_{K+1} < q_K f(q_K)$ for sufficiently large K . Since $q_{K+1} = a_{K+1}q_K + q_{K-1} \geq a_{K+1}q_K$, we have that $a_{K+1} < f(q_K) \leq f(N)$ for large K . This observation with f replaced by $\frac{1}{144}f$, together with (1.22) implies the following statement.

Corollary 1.7. *Let $f : \mathbb{N} \rightarrow \mathbb{R}^+$ be an increasing function such that (1.7) converges. Then, for almost all $\alpha \in \mathbb{R}$ and for all sufficiently large N and $c > 0$*

$$\sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \min \left\{ cN, \frac{1}{\|n\alpha\|} \right\} \leq N (cf(N))^{1/2}. \quad (1.37)$$

Example 1.2. Taking $f(q) = \log q (\log \log q)^{1+\varepsilon}$ with $\varepsilon > 0$ in the above corollary, gives the upper bound estimate

$$\text{l.h.s. of (1.37)} \leq N (c \log N (\log \log N)^{1+\varepsilon})^{1/2}$$

for almost all $\alpha \in \mathbb{R}$, $c > 0$ and for all sufficiently large N .

Example 1.3. Take f as the previous example and $c = \frac{\log N}{(\log \log N)^{1+\varepsilon}}$ with $\varepsilon > 0$. Then Corollary 1.7 together with the upper bound appearing in (1.20) of Theorem 1.2 implies that

$$\sum_{1 \leq n \leq N} \min \left\{ \frac{N \log N}{(\log \log N)^{1+\varepsilon}}, \frac{1}{\|n\alpha\|} \right\} \ll N \log N \quad (1.38)$$

for almost all $\alpha \in \mathbb{R}$ and for all sufficiently large N .

1.2.5 The homogeneous estimates for algebraic numbers

In this section we investigate the implication of the estimates obtained above for $R_N(\alpha, 0)$ and $S_N(\alpha, 0)$ on specific classes of numbers. In particular, we consider the case that α is an algebraic irrational.

To begin with, observe that the task at hand is much simplified if we know the continued fraction expansion of the number α under consideration. For instance, it is well known that

$$e = [2; 1, 2, 1, 1, 4, 1, 1, 6, 1, 1, 8, 1, 1, 10, \dots].$$

Now, for any irrational number we have that

$$\begin{aligned} \text{r.h.s. of (1.21)} &\ll q_{K+1}(1 + \log(2N/q_K)) \leq 2a_{K+1}q_K(1 + \log(2N/q_K)) \\ &\ll Na_{K+1}q_K(1 + \log(2N/q_K))/N \ll Na_{K+1} \end{aligned}$$

and note that for e we also have that $a_k \ll k$. Hence, Theorem 1.2 implies that for all sufficiently large N

$$R_N(e, 0) := \sum_{1 \leq n \leq N} \frac{1}{\|ne\|} \asymp N \log N.$$

Similarly, it is readily seen from Corollary 1.1 that

$$S_N(e, 0) := \sum_{1 \leq n \leq N} \frac{1}{n\|ne\|} \asymp (\log N)^2.$$

The sums $R_N(\alpha, 0)$ and $S_N(\alpha, 0)$ are of course just as easily estimated when α is a quadratic irrational. One just exploits the fact that the continued fraction of α is periodic. The behaviour of the sums for quadratic irrationals is similar to that for $\alpha = e$.

The case of algebraic numbers α of degree ≥ 3 is problematic, although some bounds can be obtained using, for instance, Roth's Theorem. Indeed, Roth's Theorem implies that the denominators q_k of the convergents of an algebraic number α satisfy $q_{k+1} < q_k^{1+\varepsilon}$ for every $\varepsilon > 0$ and for all k sufficiently large. Hence, Theorem 1.2 implies that if α is an algebraic irrational, then

$$N \log N \ll R_N(\alpha, 0) \ll N^{1+\varepsilon}.$$

Lang’s Conjecture [42], dating back to 1965, implies that $q_{k+1} \ll q_k(\log q_k)^{1+\varepsilon}$ for any $\varepsilon > 0$. Together with Theorem 1.2, this would in turn imply the following strengthening of the previous statement: if α is an algebraic irrational, then

$$N \log N \ll R_N(\alpha, 0) \ll N(\log N)^{1+\varepsilon}.$$

Using (1.3), one can deduce similar inequalities in relation to $S_N(\alpha, 0)$ for algebraic irrational α . Furthermore, it is not unreasonable to believe in the truth of the following statement which, in view of Corollary 1.2, it implies that $S_N(\alpha, 0)$ for algebraic irrationals behaves in the same way as for almost all irrationals.

Conjecture 1.2. *For any algebraic $\alpha \in \mathbb{R} \setminus \mathbb{Q}$*

$$S_N(\alpha, 0) \asymp (\log N)^2.$$

In order to establish Conjecture 1.2, it is not necessary to know that all the partial quotients associated with an algebraic irrational grow relatively slowly. Instead, all that is required is that the growth “on average” is relatively slow. Indeed, Conjecture 1.2 is equivalent to the following statement.

Conjecture 1.3. *For any algebraic $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, there exists a constant $c_\alpha > 0$ such that for all $k \in \mathbb{N}$*

$$A_k(\alpha) \leq c_\alpha k^2.$$

1.3 Inhomogeneous results and corollaries

1.3.1 Upper bounds

We concentrate our attention on the sums $S_N(\alpha, \gamma)$. As we have seen from the homogeneous case, these sums behave more predictably than the sums $R_N(\alpha, \gamma)$.

Our first inhomogeneous result removes the ‘epsilon’ term in the upper bound appearing in (1.4) obtained by Schmidt.

Theorem 1.4. *For each $\gamma \in \mathbb{R}$ there exists a set $\mathcal{A}_\gamma \subset \mathbb{R}$ of full Lebesgue measure such that for all $\alpha \in \mathcal{A}_\gamma$ and all sufficiently large N*

$$S_N(\alpha, \gamma) := \sum_{1 \leq n \leq N} \frac{1}{n \|n\alpha - \gamma\|} \ll (\log N)^2. \quad (1.39)$$

In view of the lower bound given by (1.4), this theorem is best possible up to the implied constant. Furthermore, the set \mathcal{A}_γ cannot be made independent of γ , as demonstrated by the following result.

Theorem 1.5. *For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and any increasing function $f : \mathbb{N} \rightarrow \mathbb{R}^+$ satisfying $f(N) = o(N)$, there exist continuum many $\gamma \in \mathbb{R}$, satisfying (1.1), and such that*

$$\limsup_{N \rightarrow \infty} f(N)^{-1} \left(S_N(\alpha, \gamma) - \max_{1 \leq n \leq N} \frac{1}{n \|n\alpha - \gamma\|} \right) = \infty.$$

Despite Theorem 1.5, the main ‘bulk’ of the sum in (1.39) can be estimated on a set \mathcal{A} of α ’s independent of γ . The following can be viewed as the inhomogeneous analogue of the upper bound appearing in Corollary 1.1.

Theorem 1.6. *There exists a set $\mathcal{A} \subset \mathbb{R}$ of full Lebesgue measure such that for any $\alpha \in \mathcal{A}$, any $\gamma \in \mathbb{R}$ and any $c > 0$, we have that for all sufficiently large N ,*

$$\sum_{\substack{1 \leq n \leq N \\ n \|n\alpha - \gamma\| \geq c}} \frac{1}{n \|n\alpha - \gamma\|} \ll_{\alpha, \gamma, c} (\log N)^2.$$

We now move on to describing lower bound estimates.

1.3.2 Lower bounds

All our lower bound results will apply to any irrational number α which is not Liouville, that is $\alpha \notin \mathfrak{L}$. Recall, that \mathfrak{L} is a small set in the sense that it has Hausdorff dimension zero (see (1.25)).

Theorem 1.7. *Let $\alpha \in \mathbb{R} \setminus (\mathfrak{L} \cup \mathbb{Q})$ and let $0 < v < 1$. There exist constants r, C_1 , and N_0 , depending on α and v only, with $0 < r, C_1 < 1$ and $N_0 > 1$, and such that, for any $\gamma \in \mathbb{R}$ and all $N > N_0$,*

$$\sum_{\substack{rN < n \leq N \\ \|n\alpha - \gamma\| \geq N^{-v}}} \frac{1}{\|n\alpha - \gamma\|} \geq C_1 N \log N.$$

Using (1.3) we readily obtain the following result.

Corollary 1.8. *Let $\alpha \in \mathbb{R} \setminus (\mathfrak{L} \cup \mathbb{Q})$ and let $0 < u, v < 1$. There exist constants r, C_2 , and N_1 , depending on α, u and v only, with $0 < r, C_2 < 1 < N_1$, and such that, for any $\gamma \in \mathbb{R}$ and all $N > N_1$,*

$$\sum_{\substack{N^u \leq n \leq N \\ n^v \|n\alpha - \gamma\| \geq r^v}} \frac{1}{n \|n\alpha - \gamma\|} \geq C_2 (\log N)^2.$$

Proof. Let r be as in Theorem 1.7 and set

$$\ell_0 := \left\lfloor \frac{u \log N}{\log r} - 1 \right\rfloor.$$

Then, it follows that

$$\begin{aligned} \sum_{\substack{N^u \leq n \leq N \\ n^v \|n\alpha - \gamma\| \geq r^v}} \frac{1}{n \|n\alpha - \gamma\|} &\geq \sum_{\ell=0}^{\ell_0} \sum_{\substack{r^{\ell+1}N < n \leq r^\ell N \\ \|n\alpha - \gamma\| \geq (r^\ell N)^{-v}}} \frac{1}{n \|n\alpha - \gamma\|} \\ &\geq \sum_{\ell=0}^{\ell_0} \frac{1}{r^\ell N} \sum_{\substack{r^{\ell+1}N < n \leq r^\ell N \\ \|n\alpha - \gamma\| \geq (r^\ell N)^{-v}}} \frac{1}{\|n\alpha - \gamma\|} \\ &\geq \sum_{\ell=0}^{\ell_0} \frac{1}{r^\ell N} \cdot C_1 r^\ell N \log(r^\ell N) \\ &\geq C_1 \ell_0 \log(r^{\ell_0} N) \geq C_2 (\log N)^2, \end{aligned}$$

provided that $\ell_0 \geq 1$ and $rN^u > N_0$. These conditions determine the quantity N_1 appearing in the statement of the corollary. \square

Corollary 1.8 complements the upper bounds associated with Theorems 1.4 and 1.6. It also implies the following statement.

Corollary 1.9. *Let $\alpha \in \mathbb{R} \setminus (\mathfrak{L} \cup \mathbb{Q})$ and $c > 0$. Then, for all sufficiently large N and any $\gamma \in \mathbb{R}$, we have that*

$$S_N(\alpha, \gamma) \geq \sum_{\substack{1 \leq n \leq N \\ n \|n\alpha - \gamma\| \geq c}} \frac{1}{n \|n\alpha - \gamma\|} \gg (\log N)^2.$$

Remark 1.6. Recall, that Schmidt's inhomogeneous result (1.4) is true on a set of $\alpha \in \mathbb{R}$ of full measure, which may depend on γ . Corollary 1.9 shows that the lower bound holds for all $\alpha \in \mathbb{R}$ except possibly for Liouville numbers - an explicit set of Hausdorff dimension zero that is independent of γ .

Remark 1.7. We do not know if the condition that α is not a Liouville number is necessary for the conclusion of Corollary 1.9 to hold for all γ . However the next result, which can be viewed as the inhomogeneous analogue of Corollary 1.4, demonstrates the necessity of the condition that α is not a Liouville number in the statement of Theorem 1.7.

Theorem 1.8. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Then, $\alpha \notin \mathfrak{L}$ if and only if for any $\gamma \in \mathbb{R}$,*

$$R_N(\alpha, \gamma) := \sum_{1 \leq n \leq N} \frac{1}{\|n\alpha - \gamma\|} \gg N \log N \quad \text{for } N \geq 2.$$

1.4 Linear forms revisited

In this section we deduce a few further corollaries via the results of §1.3.2. These provide inhomogeneous generalisations of the results of §1.2.3 and also give an alternative proof of Theorem 1.3 and its corollaries, albeit with weaker constants. The constants are not explicitly given but can be computed from the proofs given below and in §1.3.2.

Theorem 1.9. *Let $A = (\alpha_1, \dots, \alpha_n)$ be any n -tuple of real numbers such that α_1 is irrational but not a Liouville number. Then, for any positive integers T_1, \dots, T_n such that T_1 is sufficiently large, and for any $\gamma \in \mathbb{R}$,*

$$\sum_{\substack{\mathbf{q} \in \mathbb{Z}^n \\ 1 \leq q_j \leq T_j \ (1 \leq j \leq n)}} \frac{1}{\|q_1 \alpha_1 + \dots + q_n \alpha_n - \gamma\|} \gg T_1 \cdots T_n \log T_1. \quad (1.40)$$

Proof. By Theorem 1.7, for any $T_1 > T_0$ and any $\gamma_1 \in \mathbb{R}$,

$$\sum_{1 \leq q_1 \leq T_1} \frac{1}{\|q_1 \alpha_1 - \gamma_1\|} \geq C_1 T_1 \log T_1,$$

where T_0 and C_1 do not depend on γ_1 . Applying this inequality with $\gamma_1 = \gamma - (q_2 \alpha_2 + \dots + q_n \alpha_n)$, and summing over $1 \leq q_i \leq T_i$ ($2 \leq i \leq n$), we obtain the desired statement. \square

Theorem 1.10. *Let $A = (\alpha_1, \dots, \alpha_n)$ be any n -tuple of real numbers such that α_1 is irrational but not a Liouville number. Then for any positive integers T_1, \dots, T_n such that T_1 is sufficiently large, and for any $\gamma \in \mathbb{R}$,*

$$\sum_{\substack{\mathbf{q} \in \mathbb{Z}^n \\ 1 \leq q_j \leq T_j \ (1 \leq j \leq n)}} \frac{1}{q_1 \cdots q_n \|q_1 \alpha_1 + \cdots + q_n \alpha_n - \gamma\|} \gg \log T_1 \prod_{i=1}^n \log T_i. \quad (1.41)$$

Proof. By Corollary 1.8, for any $T_1 > T_0$ and any $\gamma_1 \in \mathbb{R}$

$$\sum_{1 \leq q_1 \leq T_1} \frac{1}{q_1 \|q_1 \alpha_1 - \gamma_1\|} \geq C_2 (\log T_1)^2, \quad (1.42)$$

where T_0 and C_2 do not depend on γ_1 . Now consider (1.42) with $\gamma_1 = \gamma - (q_2 \alpha_2 + \cdots + q_n \alpha_n)$, where q_2, \dots, q_n are integers. Then, on multiplying the resulting inequality by $(q_2 \cdots q_n)^{-1}$ we obtain that

$$\sum_{1 \leq q_1 \leq T_1} \frac{1}{q_1 \cdots q_n \|q_1 \alpha_1 + \cdots + q_n \alpha_n - \gamma\|} \geq C_2 (\log T_1)^2 \frac{1}{q_2 \cdots q_n}.$$

Summing this over $1 \leq q_i \leq T_i$ ($2 \leq i \leq n$), we obtain the desired statement. \square

2 Multiplicative Diophantine approximation

2.1 Background

Many problems in multiplicative Diophantine approximation can be phrased in terms of the set

$$\mathcal{S}_k^\times(\psi) := \left\{ (x_1, \dots, x_k) \in \mathbb{R}^k : \prod_{i=1}^k \|n x_i\| < \psi(n) \text{ for i.m. } n \in \mathbb{N} \right\},$$

where $k \geq 1$ is an integer, ‘i.m.’ means ‘infinitely many,’ and $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ is a function which, for obvious reasons, is referred to as an approximating function. The famous conjecture of Littlewood from the nineteen thirties states that $\mathcal{S}_2^\times(n \mapsto \varepsilon n^{-1}) = \mathbb{R}^2$ for any $\varepsilon > 0$, or equivalently that

$$\liminf_{n \rightarrow \infty} n \|n\alpha\| \|n\beta\| = 0 \quad \text{for all } (\alpha, \beta) \in \mathbb{R}^2. \quad (2.1)$$

Littlewood's conjecture has attracted much attention— see [1, 25, 46, 54] and references within. Despite some recent remarkable progress, the Littlewood Conjecture remains very much open. For instance, we are unable to show that (2.1) is valid for the pair $(\sqrt{2}, \sqrt{3})$.

On the contrary, the measure theoretic description of $\mathcal{S}_2^\times(n \mapsto \varepsilon n^{-1})$, and indeed $\mathcal{S}_2^\times(\psi)$ is well understood. For $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ monotonic¹, a simple ‘volume’ argument together with the Borel-Cantelli Lemma from probability theory implies that

$$|\mathcal{S}_k^\times(\psi)| = 0 \quad \text{if} \quad \sum_{n=1}^{\infty} \psi(n) (\log n)^{k-1} < \infty.$$

Throughout, $|X|$ denotes the k -dimensional Lebesgue measure of the set $X \subset \mathbb{R}^k$.

For the case when the above sum diverges we have the following non-trivial result due to Gallagher [26].

Theorem (Gallagher). *Let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be monotonic. Then*

$$|\mathbb{R}^k \setminus \mathcal{S}_k^\times(\psi)| = 0 \quad \text{if} \quad \sum_{n=1}^{\infty} \psi(n) (\log n)^{k-1} = \infty. \quad (2.2)$$

Remark 2.1. In fact Gallagher [26] established the above statement for $k > 1$. The $k = 1$ case is the famous classical (divergent) result of Khintchine [36] dating back to 1924. The monotonicity condition in this classical statement cannot in general be relaxed, as was shown by Duffin and Schaeffer [24] in 1941. In short, they constructed a non-monotonic function ψ for which $\sum_{n=1}^{\infty} \psi(n)$ diverges but $|\mathcal{S}_1^\times(\psi)| = 0$. In the same paper they formulated the following alternative statement for arbitrary approximating functions.

The Duffin-Schaeffer Conjecture. *Let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$. Then for almost all $\alpha \in \mathbb{R}$ the inequality*

$$|n\alpha - r| < \psi(n)$$

holds for infinitely many coprime pairs $(n, r) \in \mathbb{N} \times \mathbb{Z}$ provided that

$$\sum_{n=1}^{\infty} \frac{\varphi(n)}{n} \psi(n) = \infty. \quad (2.3)$$

Here φ is the Euler phi function. Note for comparison that, for monotonic ψ , condition (2.3) is equivalent to $\sum_{n=1}^{\infty} \psi(n) = \infty$. Although various partial results

¹When ψ is not monotonic the convergence condition reads $\sum_{n=1}^{\infty} \psi(n) |\log \psi(n)|^{k-1} < \infty$; see [11] for more details.

have been obtained, this conjecture represents a key unsolved problem in number theory. For background and recent developments regarding this fundamental problem see [8, 32, 33]. However, since we shall make use of it later on, it is worth highlighting the following ‘partial’ result established in [24].

Theorem (Duffin & Schaeffer). *The above conjecture is true if, in addition to the divergent sum condition (2.3), we also have that*

$$\limsup_{N \rightarrow \infty} \left(\sum_{n=1}^N \frac{\varphi(n)}{n} \psi(n) \right) \left(\sum_{n=1}^N \psi(n) \right)^{-1} > 0 . \quad (2.4)$$

Note that condition (2.4) implies that the convergence/divergence behavior of the sum in (2.3) and the sum $\sum_{n=1}^{\infty} \psi(n)$ are equivalent.

Remark 2.2. In the case $k > 1$, we expect to be able to remove the condition that ψ is monotonic in the hypothesis of Gallagher’s Theorem (without imposing any other condition such as coprimality) if we replace the divergence condition appearing in (2.2) by $\sum_{n=1}^{\infty} \psi(n) |\log \psi(n)|^{k-1} = \infty$; see [11] for more details. For the multiplicative analogue of the Duffin-Schaeffer Theorem see [11, Theorem 2].

2.2 Problems and main results

With $k = 2$, observe that for almost all $(\alpha, \beta) \in \mathbb{R}^2$, Gallagher’s Theorem improves upon Littlewood’s Conjecture by a factor of $(\log n)^2$. More precisely, it implies that

$$\liminf_{n \rightarrow \infty} n(\log n)^2 \|n\alpha\| \|n\beta\| = 0 \quad \text{for almost all } (\alpha, \beta) \in \mathbb{R}^2. \quad (2.5)$$

Note that this is beyond the scope of what Khintchine’s Theorem (i.e., (2.2) with $k = 1$) can tell us; namely that

$$\liminf_{n \rightarrow \infty} n \log n \|n\alpha\| \|n\beta\| = 0 \quad \forall \alpha \in \mathbb{R} \quad \text{and} \quad \text{for almost all } \beta \in \mathbb{R}. \quad (2.6)$$

However, the extra log factor in (2.5) comes at a cost of having to sacrifice a set of measure zero on the α side. As a consequence, unlike with (2.6) which is valid for any α , we are unable to claim that the stronger ‘log squared’ statement (2.5) is true for example when $\alpha = \sqrt{2}$. Obviously, the role of α and β in (2.6) can be reversed. This raises the natural question of whether (2.5) holds for every α . If true, it would mean that for any α we still beat Littlewood’s Conjecture by ‘log squared’ for almost all β . In general, this line of thought leads to following problem.

Problem 2.1. *Let $k \geq 2$. Prove that for any real numbers $\alpha_1, \dots, \alpha_{k-1} \in \mathbb{R}$ one has that*

$$\liminf_{n \rightarrow \infty} n(\log n)^k \|n\alpha_1\| \cdots \|n\alpha_{k-1}\| \|n\alpha_k\| = 0 \quad \text{for almost all } \alpha_k \in \mathbb{R}.$$

Problems of this nature fall within the scope of the theory of multiplicative Diophantine approximation on manifolds [15, 16]. In short, the approximated points $(\alpha_1, \dots, \alpha_k) \in \mathbb{R}^k$ associated with Problem 2.1 are confined to lie on the manifold, or rather the line, given by $(\alpha_1, \dots, \alpha_{k-1}) \times \mathbb{R}$. More generally, one can pose a similar problem for arbitrary submanifolds \mathcal{M} of \mathbb{R}^k – see [15] for details in the case the manifold is a planar curve.

In this paper we resolve Problem 2.1 in the two-dimensional case by obtaining the following fiber version of Gallagher’s Theorem. Basically, given $\alpha \in \mathbb{R}$, the points $(\alpha_1, \alpha_2) \in \mathbb{R}^2$ of interest are forced to lie on the line given by $\{\alpha\} \times \mathbb{R}$.

Theorem 2.1. *Let $\alpha \in \mathbb{R}$ and let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be a monotonic function such that*

$$\sum_{n=1}^{\infty} \psi(n) \log n \tag{2.7}$$

diverges and such that for some $\varepsilon > 0$,

$$\liminf_{\ell \rightarrow \infty} q_\ell^{3-\varepsilon} \psi(q_\ell) \geq 1, \tag{2.8}$$

where $q_\ell = q_\ell(\alpha)$ denotes the denominators of the principal convergents of α . Then for almost all $\beta \in \mathbb{R}$, there exists infinitely many $n \in \mathbb{N}$ such that

$$\|n\alpha\| \|n\beta\| < \psi(n). \tag{2.9}$$

Remark 2.3. Condition (2.8) is not particularly restrictive. Irrespective of ψ , it holds for all α with exponent of approximation $w(\alpha) < 3$ and it follows from the Jarník-Besicovitch Theorem [17, 35] (alternatively see [9]) that the complement is of relatively small Hausdorff dimension; namely $\dim\{\alpha \in \mathbb{R} : w(\alpha) \geq 3\} = \frac{1}{2}$.

Remark 2.4. Theorem 2.1 is applicable with $\psi(n) = (n(\log n)^2 \log \log n)^{-1}$ and arbitrary irrational α and thus resolves Problem 2.1 for $k = 2$.

Remark 2.5. We have reason to believe that the approach taken in §10 to prove Theorem 2.1 could be the foundation for developing a general multiplicative framework of regular/ubiquitous systems. We hope to explore this in the near future. For details of the current systems and their role in establishing measure theoretic statements for a general class of lim sup sets associated with the simultaneous approximation see [7, 9].

Remark 2.6. For the sake of completeness, we mention that a strengthening of Khintchine's simultaneous theorem to fibers, akin to the above strengthening of Gallagher's multiplicative theorem, has recently been obtained. Recall that the basic object in the (homogeneous) theory of simultaneous Diophantine approximation is the set

$$\mathcal{W}_k(\psi) := \left\{ (x_1, \dots, x_k) \in \mathbb{R}^k : \max_{1 \leq i \leq k} \|nx_i\| < \psi(n) \text{ for i.m. } n \in \mathbb{N} \right\},$$

and under the assumption that $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ is monotonic, Khintchine's simultaneous theorem states that

$$|\mathbb{R}^k \setminus \mathcal{W}_k(\psi)| = 0 \quad \text{if} \quad \sum_{n=1}^{\infty} \psi(n)^k = \infty.$$

We refer the reader to [12, §4.5] and references within for further details.

We next consider the situation in which the sum (2.7) converges. In this case we are able to obtain the following inhomogeneous statement that naturally complements the above theorem. It can be viewed as the fiber analogue of the convergence results established in [13, 15] for (non-degenerate) planar curves.

Theorem 2.2. *Let $\gamma, \delta \in \mathbb{R}$ and $\alpha \in \mathbb{R}$ be any irrational real number and let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be such that the sum (2.7) converges. Furthermore, assume either of the following two conditions:*

(i) $n \mapsto n\psi(n)$ is decreasing and

$$S_N(\alpha; \gamma) \ll (\log N)^2 \quad \text{for all } N \geq 2; \quad (2.10)$$

(ii) $n \mapsto \psi(n)$ is decreasing and

$$R_N(\alpha; \gamma) \ll N \log N \quad \text{for all } N \geq 2. \quad (2.11)$$

Then for almost all $\beta \in \mathbb{R}$, there exist only finitely many $n \in \mathbb{N}$ such that

$$\|n\alpha - \gamma\| \|n\beta - \delta\| < \psi(n). \quad (2.12)$$

Remark 2.7. Recall that in view of Theorem 1.4, we know that condition (2.10) is satisfied for almost every real number. This is far from the truth regarding condition (2.11), as demonstrated by Example 1.1 in §1.2.4. Also recall that in the homogeneous case, we are able to give explicit examples of real numbers α satisfying (2.10) and indeed (2.11), such as quadratic irrationals (and more generally badly approximable numbers) and e (the base of natural logarithm).

Corollary 2.1. *Let α be a badly approximable number, $\gamma = 0$ and $\delta \in \mathbb{R}$. Let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be monotonic such that (2.7) converges. Then, for almost every $\beta \in \mathbb{R}$ inequality (2.12) holds for only finitely many $n \in \mathbb{N}$.*

Remark 2.8. Conjecture 1.2, if true, implies the validity of Corollary 2.1 for any irrational algebraic number α .

Returning to the case when the sum (2.7) diverges, we highlight the rather amazing fact that currently nothing is known concerning the natural inhomogeneous version of Gallagher's Theorem.

Conjecture 2.1 (Inhomogeneous Gallagher). *Let $\gamma, \delta \in \mathbb{R}$ and let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be monotonic such that the sum (2.7) is divergent. Then, for almost all $(\alpha, \beta) \in \mathbb{R}^2$ inequality (2.12) holds for infinitely many $n \in \mathbb{N}$.*

In this paper we establish the following partial result.

Theorem 2.3. *Let $\gamma \in \mathbb{R}$, $\delta = 0$ and $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ monotonic, and assume that the sum (2.7) is divergent. Then for almost all $(\alpha, \beta) \in \mathbb{R}^2$ inequality (2.12) holds for infinitely many $n \in \mathbb{N}$.*

It will be evident from the proof of Theorem 2.3 given in §9.3, that the same argument would establish Conjecture 2.1 in full if we had the inhomogeneous version of the Duffin-Schaeffer Theorem at hand. Unfortunately and somewhat surprisingly, such a statement does not seem to be in the existing literature and we have not been able to prove it.

Problem 2.2. *Let $\gamma \in \mathbb{R}$ and $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$. Prove, that for almost all $\alpha \in \mathbb{R}$ the inequality*

$$|n\alpha - r - \gamma| < \psi(n)$$

holds for infinitely many coprime pairs $(n, r) \in \mathbb{N} \times \mathbb{Z}$ provided that the divergent sum condition (2.3) and the limsup condition (2.4) hold.

Of course a version of Conjecture 2.1 in which the points (α, β) are restricted to a fiber in \mathbb{R}^2 or a (non-degenerate) curve is of interest but currently seems well out of reach. In particular, it is natural to consider the following two-dimensional inhomogeneous version of Problem 2.1.

Problem 2.3. *Let $\gamma, \delta \in \mathbb{R}$ and $\alpha \in \mathbb{R} \setminus (\mathfrak{L} \cup \mathbb{Q})$. Prove that*

$$\liminf_{n \rightarrow \infty} n (\log n)^2 \|n\alpha - \gamma\| \|n\beta - \delta\| = 0 \quad \text{for almost all } \beta \in \mathbb{R}. \quad (2.13)$$

Within this paper we obtain the following conditional partial result.

Theorem 2.4. *Let $\gamma \in \mathbb{R}, \delta = 0$ and $\alpha \in \mathbb{R} \setminus (\mathfrak{L} \cup \mathbb{Q})$. Then, under the assumption that the Duffin-Schaeffer Conjecture is true, (2.13) holds.*

Remark 2.9. It is worthwhile comparing Problem 2.3 with the inhomogeneous version of Littlewood's Conjecture. The latter states: *for any $\alpha, \beta \in \mathbb{R}$ such that $1, \alpha, \beta$ are linearly independent over \mathbb{Q}*

$$\liminf_{n \rightarrow \infty} n \|n\alpha - \gamma\| \|n\beta - \delta\| = 0 \quad \text{for any } \gamma, \delta \in \mathbb{R}^2. \quad (2.14)$$

The existence of a single pair (α, β) satisfying (2.14) was conjectured by Cassels in the 1950s and remained open until the recent work of Shapira [51]. He showed the validity of (2.14) for almost all pairs (α, β) and, in particular, for the basis $1, \alpha, \beta$ of any real cubic number field. A higher dimensional version of these findings have subsequently been established by Lindenstrauss and Shapira [45]. In short, Problem 2.3 implies a $(\log n)^2$ strengthening of the inhomogeneous Littlewood's Conjecture at a cost of 'losing' a null set of $\beta \in \mathbb{R}$, which almost certainly will depend on δ . Note that without the $(\log n)^2$ strengthening, Shapira's result shows that for almost every $\alpha \in \mathbb{R}$, the null set is independent of δ . Recently, Gorodnik and Vishe [27] have obtained a $(\log \log \log \log \log n)^\lambda$ strengthening of Shapira's result. Here λ is some positive constant.

Part II

Developing techniques and establishing the main results

3 Ostrowski expansion

The Ostrowski expansion of real numbers is very much at the heart of our approach towards obtaining good estimates for the size of $\|n\alpha - \gamma\|$. In short, the Ostrowski expansion naturally expresses real numbers in terms of the basic parameters associated with the theory of continued fraction.

3.1 Continued fractions: the essentials

To begin with we recall various standard facts from the theory of continued fractions, which can for instance be found in [37] or [47]. Let α be a real irrational number. Throughout, the expression

$$\alpha = [a_0; a_1, a_2, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}$$

will denote the *simple continued fraction expansion* of α . The integers a_k are called the *partial quotients* of α and for $k \geq 1$ satisfy $a_k \geq 1$. The reduced rationals

$$\frac{p_k}{q_k} = [a_0; a_1, \dots, a_k] \quad (k \geq 0)$$

obtained by truncating the infinite continued fraction expansion are called the *principal convergents* of α . We will often make use of the quantities

$$D_k = q_k \alpha - p_k \quad (k \geq 0). \quad (3.1)$$

The following well known relations can be found in [47]:

$$p_{k+1} = a_{k+1}p_k + p_{k-1}, \quad q_{k+1} = a_{k+1}q_k + q_{k-1} \quad (k \geq 1), \quad (3.2)$$

$$D_0 = \{\alpha\}, \quad D_k = (-1)^k \|q_k \alpha\| \quad (k \geq 1), \quad (3.3)$$

$$a_{k+1} D_k = D_{k+1} - D_{k-1} \quad (k \geq 1), \quad (3.4)$$

$$\frac{1}{2} \leq q_{k+1} |D_k| \leq 1 \quad (k \geq 0). \quad (3.5)$$

Using (3.3) and (3.4) it is easily verified that

$$a_{k+2} |D_{k+1}| + |D_{k+2}| = |D_k| \quad (k \geq 0), \quad (3.6)$$

$$\sum_{i=1}^{\infty} a_{k+2i} |D_{k+2i-1}| = |D_k| \quad (k \geq 0) \quad (3.7)$$

and therefore

$$\sum_{i=k+1}^{\infty} a_{i+1} |D_i| = |D_k| + |D_{k+1}| \quad (k \geq 0). \quad (3.8)$$

Recall that $a_0 = \lfloor \alpha \rfloor$, $p_1 = a_0 a_1 + 1$, $q_1 = a_1$, and therefore

$$D_1 = q_1 \alpha - p_1 = a_1 \{\alpha\} - 1.$$

In view of (3.3), $D_1 < 0$ and so it follows that $|D_1| = 1 - a_1 \{\alpha\}$. Now, since $|D_0| = \{\alpha\}$, we have that $a_1 |D_0| + |D_1| = 1$ and on using (3.8) with $k = 0$ we obtain that

$$\sum_{i=0}^{\infty} a_{i+1} |D_i| = |D_0| + 1. \quad (3.9)$$

We will also appeal to the following well known and useful consequence of (3.2), which can easily be verified by induction:

$$\max\{2^{\frac{m-1}{2}}, a_1 \cdots a_m\} \leq q_m \leq 2^{m-1} \cdot a_1 \cdots a_m \quad \forall m \geq 1. \quad (3.10)$$

Finally, it is easy to show that

$$|D_k| > |D_{k+1}| \quad (k \geq 0). \quad (3.11)$$

3.2 The Ostrowski expansion

The continued fractions framework offers a natural and convenient way of encoding real numbers (both integers and irrationals) via the continued fraction expansion of a given irrational number. The following lemma explicitly determines the expansion of a positive integer in terms of the denominators of the principal convergents of a given irrational number.

Lemma 3.1. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and let $a_k = a_k(\alpha)$ and $q_k = q_k(\alpha)$ be as above. Then, for every $n \in \mathbb{N}$ there is a unique integer $K \geq 0$ such that*

$$q_K \leq n < q_{K+1},$$

and a unique sequence $\{c_{k+1}\}_{k=0}^{\infty}$ of integers such that

$$n = \sum_{k=0}^{\infty} c_{k+1} q_k, \quad (3.12)$$

$$0 \leq c_1 < a_1 \quad \text{and} \quad 0 \leq c_{k+1} \leq a_{k+1} \quad \forall k \geq 1, \quad (3.13)$$

$$c_k = 0 \quad \text{whenever} \quad c_{k+1} = a_{k+1} \quad \text{with} \quad k \geq 1, \quad (3.14)$$

$$c_{k+1} = 0 \quad \forall k > K. \quad (3.15)$$

The representation given by Lemma 3.1 is called the *Ostrowski expansion* (or *Ostrowski numeration*) of the integers. In view of (3.15) the sum (3.12) is finite. The next lemma shows that if we work with the quantities D_k rather than simply the denominators q_k , then a similar representation is valid for irrational numbers.

Lemma 3.2. *Let $\alpha \in [0, 1) \setminus \mathbb{Q}$ and let $a_k = a_k(\alpha)$, $q_k = q_k(\alpha)$ and $D_k = D_k(\alpha)$ be as above, and suppose that $\gamma \in [-\alpha, 1 - \alpha)$. Then there is a unique sequence $\{b_{k+1}\}_{k=0}^{\infty}$ of integers such that*

$$\gamma = \sum_{k=0}^{\infty} b_{k+1} D_k, \quad (3.16)$$

$$0 \leq b_1 < a_1 \quad \text{and} \quad 0 \leq b_{k+1} \leq a_{k+1} \quad \forall k \geq 1, \quad (3.17)$$

$$b_k = 0 \quad \text{whenever} \quad b_{k+1} = a_{k+1} \quad \text{with} \quad k \geq 1. \quad (3.18)$$

Observe that in view of (3.9) and (3.17), the series in (3.16) is absolutely convergent.

Further details of the expansion of real numbers given by Lemmas 3.1 and 3.2 can be found in [47, pp. 24, 33]. As already mentioned, the Ostrowski expansion of real numbers is a key tool in our approach towards obtaining estimates for the size of $\|n\alpha - \gamma\|$, which will be the subject of the next section.

4 Estimates for $\|n\alpha - \gamma\|$

In this section we give estimates for the size of $\|n\alpha - \gamma\|$ using the Ostrowski expansions of n and γ . Before we proceed with the general case of this fairly elaborate task, we consider the much easier homogenous case; that is, the case when $\gamma = 0$. Note that if (1.1) is not satisfied, $\|n\alpha - \gamma\| = \|(n - l)\alpha\|$ for some fixed $l \in \mathbb{N}$ and so we are within the homogeneous setup.

4.1 The homogeneous case

We begin by stating a lemma which is a slightly simplified version of [47, Theorem II.4.1].

Lemma 4.1. *Let $\alpha \in [0, 1) \setminus \mathbb{Q}$, $n \in \mathbb{N}$, and, with reference to Lemma 3.1, let m be the smallest integer such that $c_{m+1} \neq 0$. If $m \geq 2$ then*

$$\|n\alpha\| = \left| \sum_{k=m}^{\infty} c_{k+1} D_k \right| = \operatorname{sgn}(D_m) \cdot \sum_{k=m}^{\infty} c_{k+1} D_k. \quad (4.1)$$

Also if $m = 1$ and $\{\alpha\} < 1/2$, then we have (4.1). In all other cases, $\|n\alpha\| \geq |D_2|$, which is a positive constant depending on α only.

Lemma 4.1 underpins the following two-sided estimate for $\|n\alpha\|$.

Lemma 4.2. *Let $\alpha \in [0, 1) \setminus \mathbb{Q}$ and $n \in \mathbb{N}$, and let m be as in Lemma 4.1. If $\|n\alpha\| < |D_2|$ then $m \geq 2$, and if $m \geq 2$ then*

$$(c_{m+1} - 1)|D_m| + (a_{m+2} - c_{m+2})|D_{m+1}| \leq \|n\alpha\| \leq (c_{m+1} + 1)|D_m|. \quad (4.2)$$

Proof. By Lemma 4.1, if $\|n\alpha\| < |D_2|$ or $m \geq 2$, we have that (4.1) holds and $m \geq 1$. Then using the fact that $\operatorname{sgn}(D_k) = (-1)^k$ we find that

$$\begin{aligned} \|n\alpha\| &= c_{m+1}|D_m| - c_{m+2}|D_{m+1}| + c_{m+3}|D_{m+2}| - c_{m+4}|D_{m+3}| + \cdots \quad (4.3) \\ &\geq c_{m+1}|D_m| + (a_{m+2} - c_{m+2})|D_{m+1}| \\ &\quad - a_{m+2}|D_{m+1}| - a_{m+4}|D_{m+3}| - \cdots \end{aligned}$$

$$\stackrel{(3.7)}{=} c_{m+1}|D_m| + (a_{m+2} - c_{m+2})|D_{m+1}| - |D_m| \quad (4.4)$$

where the above inequality is obtained by removing the non-negative terms $c_{m+3}|D_{m+2}|$, $c_{m+5}|D_{m+4}|$, \dots , from (4.3) and using (3.13). This establishes the l.h.s. of (4.2).

By (3.14) and the fact that $c_{m+1} \neq 0$, we have that $c_{m+2} < a_{m+2}$ and so if m was equal to 1, then the l.h.s. of (4.2) would imply the inequality $\|n\alpha\| \geq |D_2|$. Thus, $\|n\alpha\| < |D_2|$ implies that $m \geq 2$.

To obtain the r.h.s. of (4.2) we argue similarly to the proof of the lower bound, this time removing the negative terms appearing in (4.3). It follows that

$$\begin{aligned} \|n\alpha\| &\leq c_{m+1}|D_m| + c_{m+3}|D_{m+2}| + c_{m+5}|D_{m+4}| + \dots \\ &\stackrel{(3.7)\&(3.13)}{\leq} c_{m+1}|D_m| + |D_{m+1}| \stackrel{(3.11)}{\leq} (c_{m+1} + 1)|D_m|. \end{aligned}$$

The proof is now complete. \square

4.2 The inhomogeneous case

It is worth pointing out that the inhomogeneous estimates obtained in this section are valid with $\gamma = 0$. We begin by establishing an inhomogeneous analogue of Lemma 4.1.

Lemma 4.3. *Let $\alpha \in [0, 1) \setminus \mathbb{Q}$ and $\gamma \in [-\alpha, 1 - \alpha)$, and suppose that (1.1) holds. Further, let $n \in \mathbb{N}$ and, with reference to expansions (3.12) and (3.16), let*

$$\delta_{k+1} := c_{k+1} - b_{k+1} \quad (k \geq 0). \quad (4.5)$$

Then, there exists a smallest integer $m = m(n, \alpha, \gamma)$ such that $\delta_{m+1} \neq 0$ and

$$\Sigma = \Sigma(n, \alpha, \gamma) := \sum_{k=m}^{\infty} \delta_{k+1} D_k \quad (4.6)$$

satisfies the equations

$$\|n\alpha - \gamma\| = \|\Sigma\| = \min \left\{ |\Sigma|, 1 - |\Sigma| \right\} \quad (4.7)$$

and

$$|\Sigma| = \operatorname{sgn}(\delta_{m+1} D_m) \Sigma. \quad (4.8)$$

Proof. Since $\|\cdot\|$ is invariant under integer translation, the l.h.s. of (4.7) is a straightforward consequence of (3.1), (3.12) and (3.16), and the existence of m follows trivially from (1.1). Furthermore, by (3.13) and (3.17), we have that $|\delta_2| \leq a_2$ and $|\delta_1| \leq a_1 - 1$. Then,

$$|\Sigma| \leq \sum_{k=0}^{\infty} |\delta_{k+1} D_k| \leq \sum_{k=0}^{\infty} a_{k+1} |D_k| - |D_0| \stackrel{(3.9)}{=} 1. \quad (4.9)$$

This implies that $\|\Sigma\| = \min\{|\Sigma|, 1 - |\Sigma|\}$ and establishes the r.h.s. of (4.7).

In order to prove (4.8) first note that $|\delta_{k+1}| \leq a_{k+1}$ for all $k \geq 0$. Then, for any integer $l \geq 1$

$$\begin{aligned} \left| \sum_{k=l}^{\infty} \delta_{k+1} D_k \right| &\leq |\delta_{l+1}| \cdot |D_l| + \sum_{k=l+1}^{\infty} a_{k+1} |D_k| \\ &\stackrel{(3.8)}{=} |\delta_{l+1}| \cdot |D_l| + |D_l| + |D_{l+1}| \\ &\stackrel{(3.6)}{=} (|\delta_{l+1}| + 1 - a_{l+1}) |D_l| + |D_{l-1}|. \end{aligned} \quad (4.10)$$

First, consider the case when $|\delta_{m+2}| \leq a_{m+2} - 1$. Then, by (4.10) with $l = m + 1$, we have that $\left| \sum_{k=m+1}^{\infty} \delta_{k+1} D_k \right| \leq |D_m|$. On the other hand, since $\delta_{m+1} \neq 0$ and it is an integer, we have that $|\delta_{m+1} D_m| \geq |D_m|$. Therefore, the first term in the sum $\sum_{k=m}^{\infty} \delta_{k+1} D_k$ dominates and so (4.8) holds.

Now assume that $\delta_{m+2} = a_{m+2}$. Then, by (3.13), (3.17) and (4.5), we have that $c_{m+2} = a_{m+2}$ and $b_{m+2} = 0$. Consequently, by (3.14), we get $c_{m+1} = 0$, and so $\delta_{m+1} = c_{m+1} - b_{m+1} < 0$. Then $\text{sgn}(\delta_{m+1} D_m) = \text{sgn}(\delta_{m+2} D_{m+1})$, which implies that

$$|\delta_{m+1} D_m + \delta_{m+2} D_{m+1}| \geq |D_m| + |D_{m+1}|.$$

In turn, by (4.10), we have that

$$\left| \sum_{k=m+2}^{\infty} \delta_{k+1} D_k \right| \leq |D_{m+1}| + |D_{m+2}| < |D_m| + |D_{m+1}|.$$

Therefore the first two terms in the sum $\sum_{k=m}^{\infty} \delta_{k+1} D_k$ (which have the same sign) dominate and so (4.8) holds again.

The remaining case $\delta_{m+2} = -a_{m+2}$ is established in a similar manner to the case $\delta_{m+2} = a_{m+2}$. We leave the details to the reader. \square

Lemma 4.3 enables us to compute the value of $\|n\alpha - \gamma\|$ via $|\Sigma|$. The next two lemmas, akin to Lemma 4.2, provide accurate estimates for $|\Sigma|$ and $1 - |\Sigma|$.

Lemma 4.4. *Let $\alpha, \gamma, n, \delta_{k+1}, m$ and Σ be as in Lemma 4.3 and let K be as in Lemma 3.1. Then there exists some $\ell \in \mathbb{N}$, with $\ell \leq \max\{2, K - m + 1\}$, such that*

$$\delta_{m+2+i} = (-1)^i \operatorname{sgn}(\delta_{m+1}) a_{m+2+i} \quad (1 \leq i \leq \ell - 1), \quad (4.11)$$

$$\delta_{m+2+\ell} \neq (-1)^\ell \operatorname{sgn}(\delta_{m+1}) a_{m+2+\ell}, \quad (4.12)$$

and we have that

$$\begin{aligned} |\Sigma| &= (|\delta_{m+1}| - 1) |D_m| \\ &\quad + (a_{m+2} - 1 - \operatorname{sgn}(\delta_{m+1}) \delta_{m+2}) |D_{m+1}| \\ &\quad + (a_{m+2+\ell} - (-1)^\ell \operatorname{sgn}(\delta_{m+1}) \delta_{m+2+\ell}) |D_{m+1+\ell}| \\ &\quad + \Delta, \end{aligned} \quad (4.13)$$

where

$$0 \leq |\delta_1| - 1 \leq a_1 - 2 \quad (\text{if } m = 0), \quad (4.14)$$

$$0 \leq |\delta_{m+1}| - 1 \leq a_{m+1} - 1 \quad (\text{if } m \geq 1), \quad (4.15)$$

$$0 \leq a_{m+2} - 1 - \operatorname{sgn}(\delta_{m+1}) \delta_{m+2} \leq 2a_{m+2} - 1, \quad (4.16)$$

$$1 \leq a_{m+2+\ell} - (-1)^\ell \operatorname{sgn}(\delta_{m+1}) \delta_{m+2+\ell} \leq 2a_{m+2+\ell}, \quad \text{and} \quad (4.17)$$

$$0 \leq \Delta \leq 2|D_{m+1+\ell}| + 2|D_{m+2+\ell}| < 4|D_{m+1+\ell}|. \quad (4.18)$$

Remark 4.1. In the above statement ℓ can in principle be 1 in which case condition (4.11) is empty and thus automatically holds.

Proof. By (3.15) and (3.17), we have that $\delta_{m+2+\ell} = -b_{m+2+\ell} \leq 0$ for all $\ell \geq \ell_0 := \max\{1, K - m\}$. Obviously, the expression

$$(-1)^\ell \operatorname{sgn}(\delta_{m+1}) a_{m+2+\ell}$$

takes one positive and one negative value when $\ell \in \{\ell_0, \ell_0 + 1\}$. Therefore, (4.12) holds for some $\ell \leq \ell_0 + 1 = \max\{2, K - m + 1\}$. On letting ℓ to be the minimal positive integer satisfying (4.12) we also ensure the validity of (4.11).

Equations (4.14) and (4.15) follow immediately from (4.5) together with the assumption that $\delta_{m+1} \neq 0$, and properties (3.13) and (3.17) of the Ostrowski expansion.

Next, we consider (4.16). If $\delta_{m+1} > 0$, then $c_{m+1} = \delta_{m+1} + b_{m+1} > 0$ and, by (3.13), we have that $c_{m+2} \leq a_{m+2} - 1$. This means that $\text{sgn}(\delta_{m+1})\delta_{m+2} = \delta_{m+2} = c_{m+2} - b_{m+2} \leq a_{m+2} - 1$ and implies the l.h.s. of (4.16). Similarly, if $\delta_{m+1} < 0$, then $b_{m+1} = c_{m+1} - \delta_{m+1} > 0$ and, by (3.17), we have that $b_{m+2} \leq a_{m+2} - 1$. This means that $\text{sgn}(\delta_{m+1})\delta_{m+2} = -\delta_{m+2} = b_{m+2} - c_{m+2} \leq a_{m+2} - 1$ and again implies the l.h.s. of (4.16). In either case $|\text{sgn}(\delta_{m+1})\delta_{m+2}| \leq a_{m+2}$ and so the r.h.s. of (4.16) is straightforward.

Equation (4.17) is a consequence of (4.12) and the inequality $|\delta_{m+2+\ell}| \leq a_{m+2+\ell}$, which is valid for the same reason as the r.h.s. of (4.15).

What remains is to prove (4.18). By (4.6), (4.8) and the fact that $\text{sgn}(D_i) = (-1)^i$, we obtain that

$$\begin{aligned}
|\Sigma| &= \text{sgn}(\delta_{m+1}D_m) \cdot \sum_{k=m}^{\infty} \delta_{k+1}D_k \\
&= |\delta_{m+1}| |D_m| - \text{sgn}(\delta_{m+1})\delta_{m+2}|D_{m+1}| + \\
&\quad + \sum_{k=m+2}^{\infty} (-1)^{k-m} \text{sgn}(\delta_{m+1})\delta_{k+1}|D_k| \\
&\stackrel{(4.11)}{=} |\delta_{m+1}| |D_m| - \text{sgn}(\delta_{m+1})\delta_{m+2}|D_{m+1}| - \sum_{k=m+2}^{m+\ell} a_{k+1}|D_k| - \\
&\quad - (-1)^\ell \text{sgn}(\delta_{m+1})\delta_{m+\ell+2}|D_{m+\ell+1}| + \\
&\quad + \sum_{k=m+\ell+2}^{\infty} (-1)^{k-m} \text{sgn}(\delta_{m+1})\delta_{k+1}|D_k|.
\end{aligned}$$

Therefore, on combining the implicit expression for Δ from (4.13) with the above equation for $|\Sigma|$, we obtain that

$$\Delta = |D_m| + |D_{m+1}| - \sum_{k=m+1}^{m+\ell+1} a_{k+1}|D_k| + \sum_{k=m+\ell+2}^{\infty} (-1)^{k-m} \text{sgn}(\delta_{m+1})\delta_{k+1}|D_k|. \tag{4.19}$$

Therefore, since $|\delta_{k+1}| \leq a_{k+1}$, we get that

$$\Delta \geq |D_m| + |D_{m+1}| - \sum_{k=m+1}^{\infty} a_{k+1}|D_k| \stackrel{(3.8)}{=} 0$$

and

$$\begin{aligned}
\Delta &\leq |D_m| + |D_{m+1}| - \sum_{k=m+1}^{m+1+\ell} a_{k+1}|D_k| + \sum_{k=m+2+\ell}^{\infty} a_{k+1}|D_k| \\
&= \underbrace{|D_m| + |D_{m+1}| - \sum_{k=m+1}^{\infty} a_{k+1}|D_k|}_{\substack{\| (3.8) \\ 0}} + 2 \sum_{k=m+2+\ell}^{\infty} a_{k+1}|D_k| \\
&\stackrel{(3.8)}{=} 2|D_{m+1+\ell}| + 2|D_{m+2+\ell}| < 4|D_{m+1+\ell}|.
\end{aligned}$$

This completes the proof. \square

Corollary 4.1. *Under the conditions of Lemma 4.4, we have that*

$$\|n\alpha - \gamma\| < (|\delta_{m+1}| + 2)|D_m|.$$

Proof. By (4.13) and (4.18), it follows that

$$\begin{aligned}
\|n\alpha - \gamma\| &\leq (|\delta_{m+1}| - 1)|D_m| + (2a_{m+2} - 1)|D_{m+1}| + (2a_{m+3} + 4)|D_{m+2}| \\
&\stackrel{(3.6)}{=} (|\delta_{m+1}| + 1)|D_m| + |D_{m+1}| < (|\delta_{m+1}| + 2)|D_m|.
\end{aligned}$$

\square

Lemma 4.5. *Let $\alpha, \gamma, n, \delta_{k+1}, m$ and Σ be as in Lemma 4.3 and let K be as in Lemma 3.1. Then there exists a positive integer $L \leq K + 2$ such that*

$$\delta_{i+1} = (-1)^{i+m} \operatorname{sgn}(\delta_{m+1}) a_{i+1} \quad (1 \leq i \leq L-1), \quad (4.20)$$

$$\delta_{L+1} \neq (-1)^{L+m} \operatorname{sgn}(\delta_{m+1}) a_{L+1} \quad (4.21)$$

and

$$\begin{aligned}
1 - |\Sigma| &= (a_1 - 1 - (-1)^m \operatorname{sgn}(\delta_{m+1}) \delta_1) |D_0| \\
&\quad + (a_{L+1} - (-1)^{L+m} \operatorname{sgn}(\delta_{m+1}) \delta_{L+1}) |D_L| \\
&\quad + \tilde{\Delta},
\end{aligned} \quad (4.22)$$

where all the three terms in the r.h.s. of (4.22) are non-negative, and

$$0 \leq \tilde{\Delta} < 4|D_L|. \quad (4.23)$$

Proof. By (3.15) and (3.17), we have that $\delta_{i+1} \leq 0$ for all $i \geq K + 1$. This readily implies the existence of $L \in \mathbb{N}$ such that $L \leq K + 2$ and (4.20) and (4.21) hold – see the proof of Lemma 4.4 for a similar and more detailed argument. The non-negativity of the first two terms in the r.h.s. of (4.22) follows immediately from the facts that $|\delta_1| \leq a_1 - 1$ and $|\delta_{k+1}| \leq a_{k+1}$ for $k \geq 1$.

Now we proceed to the proof of (4.23). We have that

$$\begin{aligned}
1 - |\Sigma| &\stackrel{(4.6)\&(4.8)}{=} 1 - \operatorname{sgn}(\delta_{m+1}D_m) \cdot \sum_{k=m}^{\infty} \delta_{k+1}D_k \\
&\stackrel{(3.9)}{=} (a_1 - 1)|D_0| + \sum_{k=1}^{\infty} a_{k+1}|D_k| - \operatorname{sgn}(\delta_{m+1}D_m) \cdot \sum_{k=m}^{\infty} \delta_{k+1}D_k \\
&= (a_1 - 1 - (-1)^m \operatorname{sgn}(\delta_{m+1})\delta_1)|D_0| \\
&\quad + \sum_{k=1}^{\infty} (a_{k+1} - (-1)^{k+m} \operatorname{sgn}(\delta_{m+1})\delta_{k+1})|D_k| \\
&\stackrel{(4.20)}{=} (a_1 - 1 - \operatorname{sgn}(\delta_{m+1}D_m)\delta_1)|D_0| \\
&\quad + (a_{L+1} - (-1)^{L+m} \operatorname{sgn}(\delta_{m+1})\delta_{L+1})|D_L| \\
&\quad + \sum_{k=L+1}^{\infty} (a_{k+1} - (-1)^{k+m} \operatorname{sgn}(\delta_{m+1})\delta_{k+1})|D_k|.
\end{aligned}$$

The latter sum is by definition $\tilde{\Delta}$. Since $|\delta_{k+1}| \leq a_{k+1}$ for all k , we trivially have that

$$0 \leq \tilde{\Delta} \leq 2 \sum_{k=L+1}^{\infty} a_{k+1}|D_k| \stackrel{(3.8)}{=} 2(|D_L| + |D_{L+1}|) \leq 4|D_L|.$$

This establishes (4.23) and completes the proof. \square

5 A gaps lemma and its consequences

In the course of establishing Theorems 1.4 – 1.8 we will group together natural numbers n such that $\|n\alpha - \gamma\|$ is of comparable size. Akin to Lebesgue integration, this natural idea will prove to be extremely fruitful for our purposes. We shall see that the ‘grouping’ in question leads to investigating subsets

$A(d_1, \dots, d_{m+1})$ of natural numbers that have, up to certain point, pre-determined digits d_1, \dots, d_{m+1} in their Ostrowski expansions. The following lemma reveals the structure of such sets in terms of the size of the gap between consecutive integers in $A(d_1, \dots, d_{m+1})$.

Lemma 5.1 (Gaps Lemma). *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $m \geq 0$ and d_1, \dots, d_{m+1} be non-negative integers such that $d_1 < a_1$, $d_{k+1} \leq a_{k+1}$ for $k = 1, \dots, m$ and $d_k = 0$ whenever $d_{k+1} = a_{k+1}$ for $k \leq m$. Let $A = A(d_1, \dots, d_{m+1})$ denote the set of all $n \in \mathbb{N}$ with Ostrowski expansions of the form*

$$n = \sum_{k=0}^m d_{k+1}q_k + \sum_{k=m+1}^{\infty} c_{k+1}q_k. \quad (5.1)$$

Write $A = \{n_1 < n_2 < \dots\}$ in increasing order. Then, for every $i \geq 1$ we have that:

- (i) if $d_{m+1} > 0$ then $n_{i+1} - n_i \in \{q_{m+1}, q_{m+1} + q_m\}$, and
- (ii) if $d_{m+1} = 0$ then $n_{i+1} - n_i \in \{q_{m+1}, q_m\}$. Furthermore, if $n_{i+1} - n_i = q_m$ then $c_{m+2}(n_i) = a_{m+2}$ and the gap $n_{i+1} - n_i$ is preceded by a_{m+2} consecutive gaps of length q_{m+1} .

Proof. The Ostrowski expansion of a natural number n can be defined via the following well known “greedy algorithm”. First, choose the largest integer M satisfying $q_M \leq n$. This is possible since α is irrational and so the sequence q_k is unbounded. Then, select c_{M+1} to be the largest integer satisfying $c_{M+1}q_M \leq n$. Next, subtract $c_{M+1}q_M$ from n and then repeat this process on the remaining integer to determine c_M and so on. Formally, with $n_M := n$ we define

$$c_{k+1} := \left\lfloor \frac{n_k}{q_k} \right\rfloor \quad \text{and} \quad n_{k-1} := n_k - c_{k+1}q_k \quad \text{for } k = M, M-1, \dots, 0. \quad (5.2)$$

Since $q_0 = 1$, this process will terminate with the remainder equal to 0. Also we define $c_{k+1} = 0$ for $k > M$.

Using (3.2) and (5.2) one easily verifies properties (3.12)–(3.15); that is, the above algorithm produces the Ostrowski expansion of n . Consequently, the set $A(d_1, \dots, d_{m+1})$ can be put in increasing order by using the reverse lexicographic ordering of the Ostrowski coefficients. For the rest of the proof let

$$n' := \sum_{k=0}^m d_{k+1}q_k. \quad (5.3)$$

In view of the conditions imposed on d_1, \dots, d_{m+1} in the statement of the lemma, we have that $A(d_1, \dots, d_{m+1}) \neq \emptyset$. Furthermore, if $n' > 0$ then $n' \in A(d_1, \dots, d_{m+1})$ and in this case n' is the smallest element of $A(d_1, \dots, d_{m+1})$. Now we fix some $i \in \mathbb{N}$ and suppose that

$$n_{i+1} = n' + \sum_{k=m'}^{\infty} c_{k+1}q_k,$$

where $c_{k+1} = c_{k+1}(n_{i+1})$ are the Ostrowski coefficients of n_{i+1} and $m' \geq m + 1$ with $c_{m'+1} \neq 0$. Since $i + 1 > 1$, we have that $n_{i+1} > n_1 \geq n'$, and such an m' exists. We now consider three separate scenarios.

Case (1). Suppose that $m' = m + 1$. Then, by the reverse lexicographic ordering of $A(d_1, \dots, d_{m+1})$, we have that

$$n_i = n' + (c_{m+2} - 1)q_{m+1} + \sum_{k=m+2}^{\infty} c_{k+1}q_k.$$

In this case, we obviously have that $n_{i+1} - n_i = q_{m+1}$.

Case (2). Suppose that $m' - m$ is positive and even. Then again appealing to the reverse lexicographic ordering we find that

$$\begin{aligned} n_i = n' + a_{m+3}q_{m+2} + a_{m+5}q_{m+4} + \dots + a_{m'}q_{m'-1} \\ + (c_{m'+1} - 1)q_{m'} + \sum_{k=m'+1}^{\infty} c_{k+1}q_k. \end{aligned}$$

Using (3.2) one readily verifies that $n_{i+1} - n_i = q_{m+1}$ in this case.

Case (3). Suppose that $m' - m - 2$ is positive and odd. Then the form of n_i depends on whether or not $d_{m+1} > 0$. If $d_{m+1} > 0$, then

$$\begin{aligned} n_i = n' + (a_{m+2} - 1)q_{m+1} + a_{m+4}q_{m+3} + a_{m+6}q_{m+5} + \dots + a_{m'}q_{m'-1} \\ + (c_{m'+1} - 1)q_{m'} + \sum_{k=m'+1}^{\infty} c_{k+1}q_k. \end{aligned}$$

Again using (3.2) one verifies that $n_{i+1} - n_i = q_{m+1} + q_m$. Finally, if $d_{m+1} = 0$ then

$$\begin{aligned} n_i = n' + a_{m+2}q_{m+1} + a_{m+4}q_{m+3} + a_{m+6}q_{m+5} + \dots + a_{m'}q_{m'-1} \\ + (c_{m'+1} - 1)q_{m'} + \sum_{k=m'+1}^{\infty} c_{k+1}q_k, \end{aligned}$$

and using (3.2) gives $n_{i+1} - n_i = q_m$. The ‘furthermore’ part of (ii) easily follows from the above explicit form of n_i and the reverse lexicographic ordering of $A(d_1, \dots, d_{m+1})$. This completes the proof. \square

Using Lemma 5.1, we are able to prove various useful counting results.

Lemma 5.2. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $N \geq 3$ and $m \geq 0$. Furthermore, let d_1, \dots, d_{m+1} and $A(d_1, \dots, d_{m+1})$ be as in Lemma 5.1, let n' be given by (5.3) and let*

$$A_N(d_1, \dots, d_{m+1}) := A(d_1, \dots, d_{m+1}) \cap [1, N].$$

Then

$$\sum_{\substack{n \in A_N(d_1, \dots, d_{m+1}) \\ n \neq n'}} \frac{1}{n} \leq \frac{5 \log N}{q_{m+1}}.$$

Proof. We will assume that $A_N(d_1, \dots, d_{m+1})$ contains at least one element different from n' , as otherwise the sum under consideration is zero and there is nothing to prove. By Lemma 5.1, for any distinct $n_1, n_2 \in A_N(d_1, \dots, d_{m+1})$ with $c_{m+2}(n_i) < a_{m+2}$ ($i = 1, 2$) we have that $|n_1 - n_2| \geq q_{m+1}$. In this case we find that

$$\begin{aligned} \sum_{\substack{n \in A_N(d_1, \dots, d_{m+1}) \\ c_{m+2} < a_{m+2}, n \neq n'}} \frac{1}{n} &\leq \sum_{1 \leq \ell \leq N/q_{m+1}} \frac{1}{n' + \ell q_{m+1}} \\ &\leq \frac{1}{q_{m+1}} \sum_{1 \leq \ell \leq N/q_{m+1}} \frac{1}{\ell} \leq \frac{2 \log N}{q_{m+1}}. \end{aligned}$$

On the other hand, any distinct $n_1, n_2 \in A_N(d_1, \dots, d_{m+1})$ with $c_{m+2}(n_i) = a_{m+2}$ ($i = 1, 2$) lie in $A_N(d_1, \dots, d_{m+1}, a_{m+2})$. Then, by Lemma 5.1, it follows that $|n_1 - n_2| \geq q_{m+2}$ and so similarly

$$\begin{aligned} \sum_{\substack{n \in A_N(d_1, \dots, d_{m+1}) \\ c_{m+2} = a_{m+2}}} \frac{1}{n} &\leq \sum_{0 \leq \ell \leq N/q_{m+2}} \frac{1}{n' + a_{m+2} q_{m+1} + \ell q_{m+2}} \\ &\leq \frac{1}{a_{m+2} q_{m+1}} + \frac{2 \log N}{q_{m+2}} \leq \frac{3 \log N}{q_{m+1}}. \end{aligned}$$

This together with the previous displayed estimate implies the desired statement. \square

Lemma 5.3. *Under the conditions of Lemma 5.2, let us assume that*

$$\#A_N(d_1, \dots, d_{m+1}) \geq 1.$$

Then

$$\frac{N}{3q_{m+1}} \leq \#A_N(d_1, \dots, d_{m+1}) \leq \frac{3N}{q_{m+1}} + 1. \quad (5.4)$$

Proof. In order to establish the lower bound, first observe that since we are assuming that $\#A_N(d_1, \dots, d_{m+1}) \geq 1$, the lower bound inequality in (5.4) is trivial if $N \leq 3q_{m+1}$. Without loss of generality, assume that $N > 3q_{m+1}$. It is readily verified that the minimal element of $A(d_1, \dots, d_{m+1})$ is $\leq q_{m+1}$. Hence, by Lemma 5.1, any block of $2q_{m+1}$ consecutive positive integers contains at least one element of $A(d_1, \dots, d_{m+1})$. Therefore,

$$\#A_N(d_1, \dots, d_{m+1}) \geq 1 + \left\lfloor \frac{N - q_{m+1}}{2q_{m+1}} \right\rfloor \geq \frac{N - q_{m+1}}{2q_{m+1}} \stackrel{N > 3q_{m+1}}{\geq} \frac{N}{3q_{m+1}}.$$

This verifies the lower bound of (5.4).

For the upper bound write $A(d_1, \dots, d_{m+1}) = \{n_1 < n_2 < \dots\}$ in increasing order. Let $t = \#A_N(d_1, \dots, d_{m+1})$. Thus, $n_t \leq N < n_{t+1}$. Observe, that without loss of generality, we can assume that $t \geq 2$ as otherwise the upper bound in (5.4) is trivial. With this in mind, we consider separately the two cases appearing in the statement of Lemma 5.1.

- In case (i), we have that $n_{i+1} - n_i \geq q_{m+1}$ for all $i \geq 1$ and therefore $N \geq n_1 + (t - 1)q_{m+1}$. Hence

$$\frac{N}{\#A_N(d_1, \dots, d_{m+1})} = \frac{N}{t} \geq \frac{n_1 + (t - 1)q_{m+1}}{t} \stackrel{t \geq 2}{\geq} \frac{q_{m+1}}{2}, \quad (5.5)$$

and the upper bound in (5.4) readily follows.

- In case (ii), we have that any gap of length q_m among $\{n_1, n_2, \dots\}$ must be preceded by a_{m+2} consecutive gaps of length q_{m+1} . Therefore, if $t - 1 < a_{m+2} + 1$, then all the gaps $n_i - n_{i-1}$ with $i \leq t$ must be q_{m+1} and we have that $N \geq n_1 + (t - 1)q_{m+1}$. Then, by (5.5), we again conclude that the upper bound in (5.4) holds. It remains to consider the case $t - 1 \geq a_{m+2} + 1$. In this case, it is obvious that $t \geq 3$. Further, by the division algorithm, $t - 1 = Q(a_{m+2} + 1) + R$ with $0 \leq R < a_{m+2} + 1$. Then, there are at most Q gaps of length q_m , each preceded by a_{m+2} consecutive gaps of length q_{m+1} , plus R gaps of length q_{m+1} . Then, we have that

$$N \geq n_1 + \frac{t - 1 - R}{a_{m+2} + 1} (a_{m+2}q_{m+1} + q_m) + Rq_{m+1}$$

$$\begin{aligned}
&\geq (t-1-R)\frac{q_{m+1}}{2} + Rq_{m+1} \\
&\geq (t-1-R)\frac{q_{m+1}}{2} + R\frac{q_{m+1}}{2} = \frac{t-1}{2}q_{m+1} \\
&\stackrel{t \geq 3}{\geq} \frac{t}{3}q_{m+1} = \frac{q_{m+1} \cdot \#A_N(d_1, \dots, d_{m+1})}{3},
\end{aligned}$$

whence the upper bound in (5.4) follows. \square

6 Counting solutions to $\|n\alpha - \gamma\| < \varepsilon$

Given an $\alpha, \gamma \in \mathbb{R}$, $N \in \mathbb{N}$ and $\varepsilon > 0$, consider the set

$$N_\gamma(\alpha, \varepsilon) := \{n \in \mathbb{N} : \|n\alpha - \gamma\| < \varepsilon, n \leq N\}. \quad (6.1)$$

In the homogeneous case ($\gamma = 0$), we will simply write $N(\alpha, \varepsilon)$ for $N_0(\alpha, \varepsilon)$. The main goal of this section is to estimate the cardinality of $N_\gamma(\alpha, \varepsilon)$.

6.1 The homogeneous case

We begin by observing that when $\varepsilon N \geq 1$, Minkowski's Theorem for convex bodies, see [20, p.71], implies that

$$\#N(\alpha, \varepsilon) \geq \lfloor \varepsilon N \rfloor. \quad (6.2)$$

In fact, we have already used this estimate in the proof of Theorem 1.3. We shall investigate under which conditions this bound can be reversed with some positive multiplicative constant. The main result established in this section is the following statement.

Lemma 6.1. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $(q_\ell)_{\ell \geq 0}$ be the sequence of denominators of the principal convergents of α . Let $N \in \mathbb{N}$ and $\varepsilon > 0$ such that $0 < 2\varepsilon < \|q_2\alpha\|$. Suppose that*

$$\frac{1}{2\varepsilon} \leq q_\ell \leq N \quad (6.3)$$

for some integer ℓ . Then

$$\lfloor \varepsilon N \rfloor \leq \#N(\alpha, \varepsilon) \leq 32\varepsilon N. \quad (6.4)$$

Proof. Without loss of generality, we will assume that ℓ is the smallest integer satisfying (6.3). Since $2\varepsilon < \|q_2\alpha\|$, by (3.5) and (6.3), we have that $q_\ell > q_2$, so $\ell \geq 3$. By the minimality of ℓ ,

$$q_{\ell-1} < \frac{1}{2\varepsilon} \leq q_\ell. \quad (6.5)$$

Let $n \in \mathbb{N}$ satisfy the condition $\|n\alpha\| < \varepsilon$ and let m be the same as in Lemma 4.2. Then, by (3.5) and (4.2), we obtain that

$$\frac{c_{m+1} - 1}{2q_{m+1}} + \frac{a_{m+2} - c_{m+2}}{2q_{m+2}} \leq \|n\alpha\| \leq \frac{c_{m+1} + 1}{q_{m+1}}. \quad (6.6)$$

Since, by definition, $c_{m+1} \neq 0$, condition (3.14) implies that $a_{m+2} - c_{m+2} \geq 1$. Hence, by (6.6), we obtain that

$$\frac{1}{2q_{m+2}} \leq \|n\alpha\| < \varepsilon$$

and thus $q_{m+2} \geq 1/(2\varepsilon)$. Since ℓ is defined to be the smallest integer satisfying $q_\ell \geq 1/(2\varepsilon)$, we conclude that $m+2 \geq \ell$. Consider the following three cases.

Case (1): $m \geq \ell$. Then n lies in the set $A_N(d_1, \dots, d_\ell)$ with $d_1 = \dots = d_\ell = 0$. By Lemma 5.3,

$$\#A_N(\underbrace{0, \dots, 0}_\ell) \leq \frac{3N}{q_\ell} + 1 \stackrel{(6.5)}{\leq} 6\varepsilon N + 1.$$

By (6.3), we have that $2\varepsilon N \geq 1$. Therefore, the number of $n \in N(\alpha, \varepsilon)$ that correspond to this case is

$$\leq 8\varepsilon N.$$

Case (2): $m+1 = \ell$. Then $q_{m+1} = q_\ell \geq 1/(2\varepsilon)$ and consequently, by (6.6) and the assumption $\|n\alpha\| < \varepsilon$, we have that

$$c_{m+1} < 1 + 2\varepsilon q_{m+1} \leq 4\varepsilon q_{m+1}. \quad (6.7)$$

It follows that each $n \in N(\alpha, \varepsilon)$ with $m+1 = \ell$, must lie in $A_N(d_1, \dots, d_{m+1})$ with $d_1 = \dots = d_m = 0$ and $d_{m+1} = c_{m+1}$, for some c_{m+1} satisfying (6.7). By Lemma 5.3,

$$\#A_N(\underbrace{0, \dots, 0}_{\ell-1}, c_{m+1}) \leq \frac{3N}{q_{m+1}} + 1.$$

Since $q_{m+1} = q_\ell < N$, we have that

$$\#A_N(\underbrace{0, \dots, 0}_{\ell-1}, c_{m+1}) \leq \frac{4N}{q_{m+1}}.$$

Therefore, using (6.7), we conclude that the number of $n \in N(\alpha, \varepsilon)$ that correspond to this case is

$$\leq \frac{4N}{q_{m+1}} \cdot 4\varepsilon q_{m+1} = 16\varepsilon N.$$

Case (3): $m + 2 = \ell$. Then, by (6.5), we have that

$$q_{m+1} = q_{\ell-1} < 1/(2\varepsilon) \leq q_\ell = q_{m+2}.$$

Consequently, by (6.6) we obtain that $c_{m+1} = 1$ and

$$\frac{a_{m+2} - c_{m+2}}{2q_{m+2}} \leq \|n\alpha\| < \varepsilon.$$

Thus, we have that

$$q_{m+2} > \frac{1}{2\varepsilon} \quad \text{and} \quad a_{m+2} - c_{m+2} < 2\varepsilon q_{m+2}.$$

It follows that each $n \in N(\alpha, \varepsilon)$ with $m + 2 = \ell$ must lie in $A_N(d_1, \dots, d_{m+2})$ with $d_1 = \dots = d_m = 0$, $d_{m+1} = c_{m+1} = 1$ and $d_{m+2} = c_{m+2}$. By Lemma 5.3,

$$\#A_N(\underbrace{0, \dots, 0}_{\ell-2}, 1, c_{m+2}) \leq \frac{3N}{q_{m+2}} + 1 \leq \frac{4N}{q_{m+2}}.$$

Therefore, we conclude that the number of $n \in N(\alpha, \varepsilon)$ that correspond to this case is

$$\leq \frac{4N}{q_{m+2}} \times 2\varepsilon q_{m+2} = 8\varepsilon N.$$

On summing the estimates obtained in each of the above three cases yields the desired result. \square

The following corollary of Lemma 6.1 is phrased in terms of the exponent of approximation $w(\alpha)$ of $\alpha \in \mathbb{R}$, – see (1.27) for the definition. Recall that \mathcal{L} denotes the set of Liouville numbers; that is, the set of real numbers α such that $w(\alpha) = \infty$.

Corollary 6.1. *Let $\alpha \notin \mathcal{L} \cup \mathbb{Q}$ and let $\nu \in \mathbb{R}$ satisfy the inequalities*

$$0 < \nu < \frac{1}{w(\alpha)}. \quad (6.8)$$

Then, there exists a constant $\varepsilon_0 = \varepsilon_0(\alpha) > 0$ such that for any sufficiently large N and any ε with $N^{-\nu} < \varepsilon < \varepsilon_0$, estimate (6.4) is satisfied.

Proof. We will assume that $2\varepsilon_0 < \|q_2\alpha\|$. Let ℓ be the smallest integer such that $(2\varepsilon)^{-1} \leq q_\ell$. In particular, we have that $\ell \geq 3$ and (6.5) is satisfied. By Lemma 1.1 and the condition $w(\alpha) < 1/\nu$, we have that $q_{m+1} \leq q_m^{1/\nu}$ for all sufficiently large m . In particular, for sufficiently small ε_0 the parameter ℓ will be sufficiently large and hence we will have that $q_\ell \leq q_{\ell-1}^{1/\nu}$. Then, using (6.5), we readily verify that

$$q_\ell \leq q_{\ell-1}^{1/\nu} < 1/(2\varepsilon)^{1/\nu} < 1/\varepsilon^{1/\nu} < N.$$

Since, by the definition of ℓ , we also have that $(2\varepsilon)^{-1} \leq q_\ell$, the inequalities associated with (6.3) are satisfied. Therefore, Lemma 6.1 is applicable and the conclusion of the corollary follows. \square

Now we discuss the situation not covered by Lemma 6.1, namely when (6.3) is not satisfied. To begin with, we give another ‘trivial’ lower bound for $\#N(\alpha, \varepsilon)$. Suppose that K is the largest integer such that $q_K \leq N$. Then $q_{K+1} > N$ and, by (3.5), we have that

$$\|q_K\alpha\| \leq |D_K| < 1/q_{K+1}.$$

Observe that for any positive integer s we trivially have that $\|sq_K\alpha\| < s\|q_K\alpha\|$. Hence, as long as $sq_K \leq N$ and $s/q_{K+1} \leq \varepsilon$, we have that $sq_K \in N(\alpha, \varepsilon)$. Hence,

$$\#N(\alpha, \varepsilon) \geq \min \{ \lfloor \varepsilon q_{K+1} \rfloor, \lfloor N/q_K \rfloor \}. \quad (6.9)$$

The next result shows that, up to a constant multiple, this rather trivial lower bound combined with the other trivial lower bound given by (6.2) is best possible.

Lemma 6.2. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $(q_\ell)_{\ell \geq 0}$ be the sequence of denominators of the principal convergents of α , $N \in \mathbb{N}$ and $\varepsilon > 0$ such that $N^{-1} < 2\varepsilon < \|q_2\alpha\|$. Let K be the largest non-negative integer such that $q_K \leq N$, and let*

$$M := \max \left\{ \varepsilon N, \min \left\{ \varepsilon q_{K+1}, \frac{N}{2q_K} \right\} \right\} = \min \left\{ \varepsilon q_{K+1}, \max \left\{ \varepsilon N, \frac{N}{2q_K} \right\} \right\}. \quad (6.10)$$

Then

$$\lfloor M \rfloor \leq \#N(\alpha, \varepsilon) \leq 32M. \quad (6.11)$$

Proof. The equality in (6.10) between the min-max and the max-min expressions is a consequence of the fact that $\varepsilon N < \varepsilon q_{K+1}$, which follows from the definition of K . The lower bound in (6.11) is a consequence of (6.2) and (6.9). Thus, it remains to prove the upper bound. Suppose that $q_K < 1/(2\varepsilon)$. Then $\varepsilon < \varepsilon' := 1/(2q_K)$ and we have that $N(\alpha, \varepsilon) \subset N(\alpha, \varepsilon')$. Clearly, Lemma 6.1 is applicable to $N(\alpha, \varepsilon')$ and thus it follows that

$$\#N(\alpha, \varepsilon) \leq 32\varepsilon' N = 32N/2q_K.$$

Again, by Lemma 6.1, in the case $q_K \geq 1/(2\varepsilon)$ we have that $\#N(\alpha, \varepsilon) \leq 32\varepsilon N$. Hence

$$\#N(\alpha, \varepsilon) \leq 32 \max\{\varepsilon N, N/2q_K\}.$$

Next, let $N' = q_{K+1}$. Note that $N' > N$ and therefore $N(\alpha, \varepsilon) \subset N'(\alpha, \varepsilon)$. Also condition (6.3) is satisfied with $\ell = K + 1$. Hence, it follows via Lemma 6.1 that

$$\#N(\alpha, \varepsilon) \leq \#N'(\alpha, \varepsilon) \leq 32\varepsilon N' = 32q_{K+1}\varepsilon.$$

Thus

$$\#N(\alpha, \varepsilon) \leq 32 \min\{\varepsilon q_{K+1}, \max\{\varepsilon N, N/2q_K\}\}.$$

This is the upper bound in (6.11) and thereby completes the proof of the lemma. \square

6.2 The inhomogeneous case

We prove a statement that relates the cardinality of the inhomogeneous set $N_\gamma(\alpha, \varepsilon)$ to that of the homogeneous set $N(\alpha, \varepsilon)$. The upshot is that the estimates obtained in the previous section can be exploited to provide estimates for $\#N_\gamma(\alpha, \varepsilon)$.

Lemma 6.3. *For any $\varepsilon > 0$ and $N \in \mathbb{N}$, we have that*

$$\#N_\gamma(\alpha, \varepsilon) \leq \#N(\alpha, 2\varepsilon) + 1. \quad (6.12)$$

Furthermore, if $N'_\gamma(\alpha, \varepsilon') \neq \emptyset$, where $N' := \frac{1}{2}N$ and $\varepsilon' := \frac{1}{2}\varepsilon$, then

$$\#N_\gamma(\alpha, \varepsilon) \geq \#N'_\gamma(\alpha, \varepsilon') + 1. \quad (6.13)$$

Proof. First we prove the upper bound (6.12). Suppose that $\#N_\gamma(\alpha, \varepsilon) \geq 2$, as otherwise there is nothing to prove. Let n_0 be the smallest element of $N_\gamma(\alpha, \varepsilon)$. Then for any $n \in N_\gamma(\alpha, \varepsilon)$ such that $n > n_0$ we have that

$$\|(n - n_0)\alpha\| = \|(n\alpha - \gamma) - (n_0\alpha - \gamma)\| \leq \|n\alpha - \gamma\| + \|n_0\alpha - \gamma\| < 2\varepsilon.$$

Hence, $n - n_0 \in N(\alpha, 2\varepsilon)$ and (6.12) immediately follows.

Now we prove the lower bound (6.13). In view of the assumptions of the lemma, fix any $n_1 \in N'_\gamma(\alpha, \varepsilon')$. Then, for any $n \in N'(\alpha, \varepsilon')$ we obtain that

$$\|(n_1 + n)\alpha - \gamma\| \leq \|n_1\alpha - \gamma\| + \|n\alpha\| < \varepsilon' + \varepsilon' = \varepsilon.$$

Also, note that $1 \leq n_1 + n \leq N$. Therefore, for any $n \in N'(\alpha, \varepsilon')$ we have that $n_1 + n \in N_\gamma(\alpha, \varepsilon)$. The lower bound (6.13) now readily follows. \square

7 Establishing the homogeneous results on sums of reciprocals

In this section we prove Theorems 1.1 and 1.2. Before beginning the proofs, we establish various useful auxiliary inequalities. As before, $q_k = q_k(\alpha)$ denotes the denominators of the principal convergents of α , and $K = K(N, \alpha)$ denotes the largest non-negative integer such that $q_K \leq N$, so

$$q_K \leq N < q_{K+1}. \quad (7.1)$$

By (3.10) and (7.1), we also have that

$$K - 1 \leq 2 \log N / \log 2. \quad (7.2)$$

Furthermore, it is easily verified that

$$\max\{\log(\frac{a_1}{2} \cdots \frac{a_K}{2}), K\} \geq \frac{1}{3} \log q_K. \quad (7.3)$$

Indeed, by (3.10), we have that

$$q_K \leq 2^{K-1} \cdot a_1 \cdots a_K < 4^K \cdot \frac{a_1}{2} \cdots \frac{a_K}{2}.$$

On taking logarithms, we obtain that

$$\log q_K \leq K \log 4 + \log(\frac{a_1}{2} \cdots \frac{a_K}{2}) \leq 2 \log 4 \max\{K, \log(\frac{a_1}{2} \cdots \frac{a_K}{2})\}$$

whence (7.3) follows.

7.1 Proof of Theorem 1.2

7.1.1 The upper bound in (1.20)

For each pair of integers (u, a) satisfying $1 \leq u \leq K + 1$ and $1 \leq a \leq a_{u+1}$, let $B_N(u, a)$ be the set of integers n with $1 \leq n \leq N$ such that the Ostrowski coefficients $\{c_{k+1}\}_{k=0}^\infty$ of n satisfy

$$c_1 = \cdots = c_{u-1} = 0$$

and one of the following two conditions:

- (i) $c_u = 0$ and $c_{u+1} = a + 1$, or
- (ii) $c_u = 1$ and $c_{u+1} = a_{u+1} - a$.

In view of the uniqueness of Ostrowski expansion, the sets $B_N(u, a)$ are pairwise disjoint. By definition, we always have that $B_N(u, a) \subseteq \{1, \dots, N\}$. We claim that

$$\{n \in \mathbb{N} : n \leq N, \|n\alpha\| < |D_2|\} \subset \bigcup_{u=2}^{K+1} \bigcup_{a=1}^{a_{u+1}} B_N(u, a). \quad (7.4)$$

To see this, take any n from the l.h.s. of (7.4) and, with reference to the Ostrowski expansion of n , let m be the smallest integer such that $c_{m+1} \neq 0$. By (7.1) and Lemma 4.2, we have that $2 \leq m \leq K$. If $c_{m+1} > 1$ then n lies in the set $B_N(m, c_{m+1} - 1)$. If $c_{m+1} = 1$ then n lies in the set $B_N(m+1, a_{m+2} - c_{m+2})$. Note that, by (3.14) and (3.15), in this case $1 \leq a_{m+2} - c_{m+2} \leq a_{m+2}$.

By Lemma 4.2, if n lies in the l.h.s. of (7.4) then

$$\|n\alpha\| \geq a|D_u| \quad \text{whenever } n \in B_N(u, a) \quad (7.5)$$

for some u and a from the r.h.s. of (7.4). Clearly, the set $B_N(u, a)$ is the union of

$$A_N(\underbrace{0, \dots, 0}_{u \text{ times}}, a + 1) \quad \text{and} \quad A_N(\underbrace{0, \dots, 0}_{u-1 \text{ times}}, 1, a_{u+1} - a), \quad (7.6)$$

where the sets $A_N(\dots)$ are defined in §5. Then, by Lemma 5.3 and the inequalities given by (7.1), we find that

$$\#B_N(u, a) \leq \frac{8N}{q_{u+1}} \quad \text{if } u \leq K - 1. \quad (7.7)$$

Putting this together with (7.5) implies that

$$\begin{aligned}
\sum_{u=2}^{K-1} \sum_{a=1}^{a_{u+1}} \sum_{n \in B_N(u,a)} \frac{1}{\|n\alpha\|} &\leq \sum_{u=2}^{K-1} \sum_{a=1}^{a_{u+1}} \frac{8N}{aq_{u+1}|D_u|} \\
&\stackrel{(3.5)}{\leq} 16N \sum_{u=2}^{K-1} \sum_{a=1}^{a_{u+1}} \frac{1}{a} \\
&\leq 16N \sum_{u=2}^{K-1} (1 + \log a_{u+1}) \\
&\leq 16N(K - 2 + \log(a_1 \dots a_K)) \\
&\stackrel{(3.10)}{\leq} 64N \log q_K. \tag{7.8}
\end{aligned}$$

Considering the case $\|n\alpha\| \geq |D_2|$, we get that

$$\sum_{\substack{n \leq N \\ \|n\alpha\| \geq |D_2|}} \frac{1}{\|n\alpha\|} \leq N|D_2|^{-1} \stackrel{(3.5)}{\leq} 2q_3N. \tag{7.9}$$

It remains to note that if

$$n \in \bigcup_{u=K}^{K+1} \bigcup_{a=1}^{a_{u+1}} B_N(u, a)$$

then

$$n \equiv 0 \text{ or } q_{K-1} \pmod{q_K}. \tag{7.10}$$

Indeed, if $n \in B_N(K, a)$ then, since $q_{K+1} > N \geq n$, by definition, the Ostrowski expansion of n is either $(a+1)q_K$ or $q_{K-1} + (a_{K+1} - a)q_K$. In both cases (7.10) is satisfied. Similarly, if $n \in B_N(K+1, a)$ then, by definition, the Ostrowski expansion of n can only be $q_K + (a_{K+2} - a)q_{K+1}$ and since $n \leq N < q_{K+1}$, we have that $n = q_K$. Clearly, (7.10) is satisfied again. The upshot is that the n 's which fall into $B_N(K, a)$ or $B_N(K+1, a)$ are irrelevant in estimating (1.20) and therefore on combining (7.4), (7.8) and (7.9) implies the upper bound in (1.20).

7.1.2 The lower bound in (1.20)

For each $m, a \in \mathbb{N}$, define $B_N^*(m, a)$ to be the collection of positive integers $n \leq N$ with Ostrowski expansions satisfying $c_{k+1} = 0$ for all $k < m$ and $c_{m+1} = a$. If $m \leq K - 1$ and $1 \leq a \leq a_{m+1}$, the set $B_N^*(m, a)$ is clearly non-empty. Then, by Lemma 5.3, it follows that

$$\#B_N^*(m, a) \geq \frac{N}{3q_{m+1}}. \quad (7.11)$$

Assume that $n \in B_N^*(m, a)$ and that one of the following is satisfied

- $1 \leq a \leq a_{m+1}$ and $m \leq K - 2$, or
- $2 \leq a \leq a_{m+1}$ and $m = K - 1$.

Then it is easily seen that $n \not\equiv 0, q_{K-1} \pmod{q_K}$. Therefore, the sum in (1.20) is bounded below by

$$X := \sum_{m=2}^{K-2} \sum_{a=1}^{a_{m+1}} \sum_{n \in B_N^*(m, a)} \frac{1}{\|n\alpha\|} + \sum_{a=2}^{a_K} \sum_{n \in B_N^*(K-1, a)} \frac{1}{\|n\alpha\|}.$$

By Lemma 4.2, $\|n\alpha\| \leq (a+1)|D_m|$ for every $n \in B_N^*(m, a)$ with $m \geq 2$. Then this together with (7.11) implies that

$$\begin{aligned} X &\geq \sum_{m=2}^{K-2} \sum_{a=1}^{a_{m+1}} \frac{N}{3q_{m+1}(a+1)|D_m|} + \sum_{a=2}^{a_K} \frac{N}{3q_K(a+1)|D_{K-1}|} \\ &\stackrel{(3.5)}{\geq} \frac{1}{3}N \left(\sum_{m=2}^{K-2} \sum_{a=1}^{a_{m+1}} + \sum_{a=2}^{a_K} \right) \frac{1}{a+1} = \frac{1}{3}N \sum_{m=2}^{K-1} \sum_{a=1}^{a_{m+1}} \frac{1}{a+1} - \frac{1}{6}N \\ &\geq \frac{1}{3}N \sum_{m=2}^{K-1} \log \frac{a_{m+1}+2}{2} - \frac{1}{6}N \\ &\geq \frac{1}{3}N \sum_{m=0}^{K-1} \log \frac{a_{m+1}+2}{2} - \frac{1}{3}N (\log \frac{a_1+2}{2} + \log \frac{a_2+2}{2}) - \frac{1}{6}N. \end{aligned} \quad (7.12)$$

Since $3a_i \geq a_i + 2 > a_i$,

$$\log \frac{3a_{m+1}}{2} \geq \log \frac{a_{m+1}+2}{2} \geq \max\{\log(3/2), \log \frac{a_{m+1}}{2}\},$$

and we obtain from (7.12) that

$$X > \frac{1}{3}N \max\{K \log(3/2), \log(\frac{a_1}{2} \dots \frac{a_K}{2})\} - \frac{1}{3}N \log(9a_1a_2/4) - \frac{1}{6}N. \quad (7.13)$$

Trivially, $\frac{1}{3} \log(3/2) > \frac{1}{8}$. Then, using (3.10) and (7.3) we obtain from (7.13) that

$$\begin{aligned} X &\geq \frac{1}{24}N \log q_K - \frac{1}{3}N \log q_2 - \frac{1}{3}N \log(9/4) - \frac{1}{6}N \\ &\geq \frac{1}{24}N \log q_K - \left(\frac{1}{3} \log q_2 + \frac{1}{2}\right)N. \end{aligned}$$

This completes the proof of (1.20).

7.1.3 Proof of (1.21) and (1.22)

First observe that since $N \geq q_3$ we have that $K \geq 3$. Furthermore, $1 \leq n_1 \leq N$ and $n_1 \equiv 0 \pmod{q_K}$ if and only if $n_1 = aq_K$ for some a with $1 \leq a \leq N/q_K$. Since $N < q_{K+1} = a_{K+1}q_K + q_{K-1}$, we also have that $a \leq a_{K+1}$. Thus, the expression $n_1 = aq_K$ is the Ostrowski expansion of such an integer n_1 and then, by Lemma 4.1 (with $m = K \geq 3$), we have that

$$\|n_1\alpha\| = a|D_K|.$$

Similarly, $1 \leq n_2 \leq N$ and $n_2 \equiv q_{K-1} \pmod{q_K}$ if and only if $n_2 = q_{K-1} + (a_{K+1} - b)q_K$ for some b with $a_{K+1} - (N - q_{K-1})/q_K \leq b \leq a_{K+1}$. Again since $N < a_{K+1}q_K + q_{K-1}$, we have that $a_{K+1} - (N - q_{K-1})/q_K > 0$, which implies that $b \geq 1$. By Lemma 4.1 (with $m = K - 1 \geq 2$), we have that

$$\|n_2\alpha\| = |D_{K-1}| - (a_{K+1} - b)|D_K| \stackrel{(3.6)}{=} |D_{K+1}| + b|D_K|.$$

By (3.5),

$$\frac{q_{K+1}}{a} \leq \frac{1}{\|n_1\alpha\|} \leq \frac{2q_{K+1}}{a} \quad \text{and} \quad \frac{q_{K+1}}{b+1} \leq \frac{1}{\|n_2\alpha\|} \leq \frac{2q_{K+1}}{b}.$$

Consequently,

$$\begin{aligned} \sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \frac{1}{\|n\alpha\|} &\leq \sum_{1 \leq a \leq N/q_K} \frac{2q_{K+1}}{a} + \sum_{a_{K+1} - (N - q_{K-1})/q_K \leq b \leq a_{K+1}} \frac{2q_{K+1}}{b} \\ &\leq 4q_{K+1} \sum_{1 \leq a \leq 1 + N/q_K} \frac{1}{a} \leq 4q_{K+1}(1 + \log(1 + N/q_K)) \end{aligned}$$

and

$$\sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \frac{1}{\|n\alpha\|} \geq \sum_{1 \leq a \leq N/q_K} \frac{q_{K+1}}{a} \geq q_{K+1} \log(1 + N/q_K).$$

These upper and lower bound inequalities prove (1.21). Similarly,

$$\begin{aligned} \sum_{\substack{1 \leq n \leq N \\ n \equiv 0, q_{K-1} \pmod{q_K}}} \min \left\{ cN, \frac{1}{\|n\alpha\|} \right\} &\leq 2 \sum_{1 \leq a \leq 2N/q_K} \min \left\{ cN, \frac{2q_{K+1}}{a} \right\} \\ &\leq 2 \sum_{1 \leq a \leq 2N/q_K} \min \left\{ cN, \frac{4a_{K+1}q_K}{a} \right\}. \end{aligned} \quad (7.14)$$

Here we have used the fact that $q_{K+1} = a_{K+1}q_K + q_{K-1} \leq 2a_{K+1}q_K$. Now split the sum on the r.h.s. of (7.14) into two sub-sums: one with $a \leq 2(a_{K+1}/c)^{\frac{1}{2}}$ and the other with $a > 2(a_{K+1}/c)^{\frac{1}{2}}$. The first sub-sum is trivially bounded by

$$2 \cdot cN \cdot 2(a_{K+1}/c)^{\frac{1}{2}} = 4N(ca_{K+1})^{\frac{1}{2}}$$

while the second sub-sum is bounded by

$$2 \cdot \frac{2N}{q_K} \cdot \frac{4a_{K+1}q_K}{2(a_{K+1}/c)^{\frac{1}{2}}} = 8N(ca_{K+1})^{\frac{1}{2}}.$$

On combining the previous two estimates with (7.14) we obtain (1.22).

7.2 Proof of Theorem 1.1

Let N be a sufficiently large integer and $K = K(N, \alpha)$ be as in statement of Theorem 1.1. By the definition of K , for any integer $0 \leq i \leq K$ we have that $q_i \leq N$. By (3.5), we have that $q_{i+1}\|q_i\alpha\| < 1$ and since $q_{i+1} = a_{i+1}q_i + q_{i-1} > a_{i+1}q_i$ ($q_{-1} := -1$), it follows that $a_{i+1}q_i\|q_i\alpha\| < 1$. Hence

$$S_N(\alpha, 0) \geq \sum_{i=0}^K \frac{1}{q_i\|q_i\alpha\|} \geq \sum_{i=0}^K a_{i+1} = A_{K+1}.$$

Combining this with Corollary 1.5 establishes the lower bound in (1.19). It remains to prove the upper bound.

By the partial summation formula (1.2), for any function $g : \mathbb{N} \rightarrow \mathbb{R}$, we have that

$$\sum_{n=1}^N \frac{g(n)}{n} = \sum_{n=1}^N \frac{G(n)}{n(n+1)} + \frac{G(N)}{N+1} \quad \text{where} \quad G(n) := \sum_{k=1}^n g(k). \quad (7.15)$$

Now consider the specific function g defined as follows: given an integer $k \geq 0$, for $q_k \leq n < q_{k+1}$, let

$$g(n) := \begin{cases} \|n\alpha\|^{-1} & \text{if } n \not\equiv 0, q_{k-1} \pmod{q_k} \\ 0 & \text{otherwise,} \end{cases}$$

where $q_{-1} := 0$. Theorem 1.2 implies that

$$G(m) \leq 64m \log m + O(m), \quad (7.16)$$

where the implied constant depends only on α . Indeed, to see that this is so, let M be the largest integer such that $q_M \leq m$. Then for $n < q_M$, we have that $n \not\equiv 0 \pmod{q_M}$ while $n \equiv q_{M-1} \pmod{q_M}$ means that $n = q_{M-1}$ and so $g(n) = 0$. Hence, on making use of (1.20), we have that

$$\begin{aligned} G(m) &= \sum_{k=0}^{M-1} \sum_{\substack{q_k \leq n < q_{k+1} \\ n \not\equiv 0, q_{k-1} \pmod{q_k}}} \frac{1}{\|n\alpha\|} + \sum_{\substack{q_M \leq n \leq m \\ n \not\equiv 0, q_{M-1} \pmod{q_M}}} \frac{1}{\|n\alpha\|} \\ &\leq \sum_{\substack{1 \leq n \leq m \\ n \not\equiv 0, q_{M-1} \pmod{q_M}}} \frac{1}{\|n\alpha\|} \leq 64m \log q_M + O(m) \\ &\leq 64m \log m + O(m) \end{aligned}$$

which is precisely (7.16). On combining (7.15) and (7.16), it follows that

$$\begin{aligned} \sum_{\substack{1 \leq n \leq N \\ g(n) \neq 0}} \frac{1}{n \|n\alpha\|} &= \sum_{n=1}^N \frac{g(n)}{n} \leq 64 \sum_{n=1}^N \frac{\log n}{n} + O(\log N) \\ &\leq 32(\log N)^2 + O(\log N). \end{aligned} \quad (7.17)$$

It remains to consider the sum

$$\sum_{\substack{1 \leq n \leq N \\ g(n) = 0}} \frac{1}{n \|n\alpha\|}.$$

We will use similar arguments to those appearing in §7.1.3. Fix any integer $k \geq 3$. A positive integer $n_1 \in [q_k, q_{k+1})$ satisfies $n_1 \equiv 0 \pmod{q_k}$ if and only if $n_1 = aq_k$ for some a with $1 \leq a \leq (q_{k+1} - 1)/q_k$. Similarly, a positive integer $n_2 \in [q_k, q_{k+1})$ satisfies $n_2 \equiv q_{k-1} \pmod{q_k}$ if and only if $n_2 = q_{k-1} + (a_{k+1} - b)q_k$

for some b with $a_{k+1} - (q_{k+1} - q_{k-1} - 1)/q_k \leq b < a_{k+1}$. It is easily seen that $b \geq 1$. Then, by Lemma 4.1, we have that

$$\|n_1\alpha\| = a|D_k|$$

and

$$\|n_2\alpha\| = |D_{k-1}| - (a_{k+1} - b)|D_k| \stackrel{(3.6)}{=} |D_{k+1}| + b|D_k|.$$

By (3.5), we have that

$$\frac{1}{\|n_1\alpha\|} \leq \frac{2q_{k+1}}{a} \quad \text{and} \quad \frac{1}{\|n_2\alpha\|} \leq \frac{2q_{k+1}}{b}.$$

These together with the inequality $q_{k+1} < 2a_{k+1}q_k$, imply that

$$\frac{1}{\|n_1\alpha\|} \leq \frac{4a_{k+1}q_k}{a} \quad \text{and} \quad \frac{1}{\|n_2\alpha\|} \leq \frac{4a_{k+1}q_k}{b}.$$

Consequently, it follows that

$$\begin{aligned} \sum_{\substack{q_k \leq n < q_{k+1} \\ g(n)=0}} \frac{1}{n\|n\alpha\|} &\leq \sum_{a \geq 1} \frac{4a_{k+1}q_k}{q_k a^2} + \sum_{1 \leq b \leq a_{k+1}-1} \frac{4a_{k+1}q_k}{b(a_{k+1} - b)q_k} \\ &= 4a_{k+1} \sum_{a \geq 1} \frac{1}{a^2} + 4 \sum_{1 \leq b \leq a_{k+1}-1} \left(\frac{1}{b} + \frac{1}{a_{k+1} - b} \right) \\ &= 4a_{k+1} \sum_{a \geq 1} \frac{1}{a^2} + 8 \sum_{1 \leq b \leq a_{k+1}-1} \frac{1}{b} \\ &\leq \frac{2\pi^2}{3} a_{k+1} + 1 + \log a_{k+1} < 9a_{k+1}. \end{aligned}$$

Hence, for N (and therefore K) sufficiently large we have that

$$\sum_{\substack{1 \leq n \leq N \\ g(n)=0}} \frac{1}{n\|n\alpha\|} \leq \sum_{k=0}^K \sum_{\substack{q_k \leq n < q_{k+1} \\ g(n)=0}} \frac{1}{n\|n\alpha\|} \leq 9A_{K+1} + O(1) \leq 10A_{K+1}.$$

This together with (7.17) establishes the upper bound appearing in (1.19) and thus completes the proof of Theorem 1.1.

8 Establishing the inhomogeneous results on sums of reciprocals

In this section we prove Theorems 1.4, 1.5, 1.6 and 1.8. Since $\|x\|$ is invariant under integer translations of x , without loss of generality, we can assume that

$\alpha \in [0, 1) \setminus \mathbb{Q}$ and that $\gamma \in [-\alpha, 1 - \alpha)$. Throughout, we will use the notation and language introduced in §3 – §5. Also, we assume that (1.1) holds.

8.1 Proof of Theorem 1.6

Take $0 < \varepsilon < 1$ and let

$$\mathcal{A} := \left\{ \alpha \in [0, 1) \setminus \mathbb{Q} : \log(a_1 \cdots a_n) \ll n, \sum_{i=1}^n a_i \ll n^{1+\varepsilon} \text{ for all } n \in \mathbb{N} \right\},$$

where the implied constants may depend on α . It follows from the Khintchine-Levy Theorem [47, Chapter V], Khintchine’s Theorem [32, Theorem 2.2], and a theorem of Diamond and Vaaler [22] that the set \mathcal{A} has Lebesgue measure one. We shall show that Theorem 1.6 holds with this particular choice of \mathcal{A} .

In order to estimate the relevant sum we will make use of expression (4.7) for $\|n\alpha - \gamma\|$. Namely,

$$\|n\alpha - \gamma\| = \min \left\{ |\Sigma|, 1 - |\Sigma| \right\},$$

where $\Sigma = \Sigma(n, \alpha, \gamma)$ is given by (4.6) and $m = m(n, \alpha, \gamma)$ is defined as in Lemma 4.3. We will use the estimates for $|\Sigma|$ and $1 - |\Sigma|$ given by Lemma 4.4 and Lemma 4.5.

8.1.1 The case $m > K$

Unlike our treatment of the corresponding homogeneous case, where m was always bounded by K , in the inhomogeneous case the parameter m (which is determined by the b_{k+1} ’s as well as by the c_{k+1} ’s) can be arbitrarily large if the Ostrowski expansion of γ contains a large number of consecutive zeros. More precisely, if $b_{K+1} = \cdots = b_{K+T} = 0$, $b_{K+T+1} \neq 0$, then for $n = \sum_{k=0}^K b_{k+1} q_k$ we will have that $m = K + T$. However, as is clear from the definition of m and the coefficients δ_{k+1} given by (4.5), whenever $m > K$, we have that $c_{k+1} = b_{k+1}$ for all $k \leq K$ and the associated integer n is uniquely defined by γ . Hence

$$\sum_{\substack{n \leq N : m > K \\ c \leq n \|n\alpha - \gamma\|}} \frac{1}{n \|n\alpha - \gamma\|} \leq \frac{1}{c}. \quad (8.1)$$

8.1.2 The case $|\Sigma| \leq 1/2$, $m \leq K$

By Lemma 4.3, in the case $|\Sigma| \leq 1/2$, we have that $\|n\alpha - \gamma\| = |\Sigma|$ and therefore we can call upon Lemma 4.4 when required. With this in mind, for each pair of integers (u, a) satisfying

$$0 \leq u \leq K + 1 \quad \text{and} \quad 1 \leq a \leq 2a_{u+1} - 1, \quad (8.2)$$

let $C_N(u, a)$ be the set of all positive integers $n \leq N$ such that $|\Sigma| \leq 1/2$ and one of the following two conditions is satisfied:

- (i) $\delta_1 = \dots = \delta_u = 0$, $|\delta_{u+1}| - 1 = a$, or
- (ii) $u \geq 1$, $\delta_1 = \dots = \delta_{u-1} = 0$, $|\delta_u| = 1$, $a_{u+1} - 1 - \text{sgn}(\delta_u)\delta_{u+1} = a$.

In the first case we have that $u = m$ while in the second case $u = m + 1$, where $m = m(n, \alpha, \gamma)$ is defined as in Lemma 4.3. Since we are dealing with the case $m \leq K$, the condition $u \leq K + 1$ within (8.2) is natural and non-restrictive. Then, applying Lemmas 4.3 and 4.4, we obtain that

$$\|n\alpha - \gamma\| \geq a|D_u| \quad (8.3)$$

for any $n \in C_N(u, a)$. It is easily seen that $C_N(u, a)$ lies within the union of the following four sets:

$$A_N(b_1, \dots, b_u, b_{u+1} \pm (a + 1)) \quad (8.4)$$

and

$$A_N(b_1, \dots, b_{u-1}, b_u \pm 1, b_{u+1} \pm (a_{u+1} - a - 1)), \quad (8.5)$$

where b_1, b_2, \dots are the Ostrowski coefficients of γ (see Lemma 3.2) and are fixed.

Let $M_N(u, a)$ be the collection of the minimal elements of the non-empty sets given by (8.4) and (8.5). Trivially, we have that $\#M_N(u, a) \leq 4$. Further, using Lemma 5.2, we obtain

$$\sum_{n \in C_N(u, a) \setminus M_N(u, a)} \frac{1}{n} \ll \frac{\log N}{q_{u+1}}.$$

On combining this with (8.3), we obtain that

$$\sum_{u=0}^{K+1} \sum_{a=1}^{2a_{u+1}-1} \sum_{\substack{n \in C_N(u, a) \\ c \leq n \|n\alpha - \gamma\|}} \frac{1}{n \|n\alpha - \gamma\|}$$

$$\begin{aligned}
&\leq \sum_{u=0}^{K+1} \sum_{a=1}^{2a_{u+1}-1} \left(\sum_{n \in C_N(u,a) \setminus M_N(u,a)} \frac{1}{na|D_u|} + \sum_{\substack{n \in M_N(u,a) \\ c \leq n \|n\alpha - \gamma\|}} \frac{1}{n \|n\alpha - \gamma\|} \right) \\
&\ll \log N \cdot \sum_{u=0}^{K+1} \frac{1}{q_{u+1}|D_u|} \sum_{a=1}^{2a_{u+1}-1} \frac{1}{a} + \sum_{u=0}^{K+1} \sum_{a=1}^{2a_{u+1}-1} \frac{4}{c} \\
&\ll \log N \cdot \sum_{u=0}^{K+1} (1 + \log a_{u+1}) + \sum_{u=0}^{K+1} a_{u+1}. \tag{8.6}
\end{aligned}$$

The last inequality is a consequence of (3.5) and the trivial fact that

$$\sum_{a=1}^{2a_{u+1}-1} \frac{1}{a} \ll 1 + \log a_{u+1}.$$

Recall that, by (3.10), we have that $K \ll \log q_K \leq \log N$ and, by the definition of \mathcal{A} , we have that

$$\sum_{u=0}^{K+1} \log a_{u+1} = \log(a_1 \cdots a_{K+2}) \ll K + 2 \ll K \ll \log N,$$

and

$$\sum_{u=0}^{K+1} a_{u+1} \ll K^{1+\varepsilon} \ll (\log N)^{1+\varepsilon}.$$

Hence, the above two estimates together with (8.6) imply that

$$\sum_{u=0}^{K+1} \sum_{a=1}^{2a_{u+1}-1} \sum_{\substack{n \in C_N(u,a) \\ c \leq n \|n\alpha - \gamma\|}} \frac{1}{n \|n\alpha - \gamma\|} \ll (\log N)^2. \tag{8.7}$$

Observe that this upper bound estimate is no bigger than the upper bound appearing in Theorem 1.6. However, we are not yet done with the case under consideration, since there may be integers n with $1 \leq n \leq N$ and $|\Sigma| \leq 1/2$ which do not fall in any of the sets $C_N(u, a)$. As is clear from the definition of $C_N(u, a)$ and the range of (u, a) given by (8.2), the remaining numbers n correspond to the situation when the first two terms of (4.13) vanish. In order to take these numbers into account, for each triple (m, a, ℓ) such that

$$0 \leq m \leq K, \quad 1 \leq \ell \leq \max\{2, K - m + 1\}, \quad 1 \leq a \leq 2a_{m+2+\ell},$$

we define the set $C_N^+(m, a, \ell)$ to consist of integers n with $1 \leq n \leq N$ such that $|\Sigma| \leq 1/2$ and their Ostrowski expansions satisfy the following set of equalities:

$$\left\{ \begin{array}{l} \delta_1 = \cdots = \delta_m = 0 \\ |\delta_{m+1}| = 1, \\ \operatorname{sgn}(\delta_{m+1})\delta_{m+2} = a_{m+2} - 1, \\ \delta_{m+2+i} = (-1)^i \operatorname{sgn}(\delta_{m+1})a_{m+2+i} \quad (1 \leq i \leq \ell - 1) \\ a_{m+2+\ell} - (-1)^\ell \operatorname{sgn}(\delta_{m+1})\delta_{m+2+\ell} = a. \end{array} \right. \quad (8.8)$$

Let $t = m + \ell + 1$. Then we have that

$$2 \leq t \leq K + 3. \quad (8.9)$$

Since we are assuming that $|\Sigma| \leq 1/2$, by Lemmas 4.3 and 4.4, we have that

$$\|n\alpha - \gamma\| \geq a\|q_t\alpha\| \quad \text{for all } n \in C_N^+(m, a, \ell).$$

In view of properties (3.13), (3.14), (3.17) and (3.18), the equalities associated with (8.8) can only occur if the coefficients b_{m+1}, \dots, b_t of the Ostrowski expansion of γ have one of the following two patterns:

b_{m+1}	b_{m+2}	b_{m+3}	b_{m+4}	b_{m+5}	b_{m+6}	\dots	
$< a_{m+1}$	0	a_{m+3}	0	a_{m+5}	0	\dots	(8.10)
> 0	$a_{m+2} - 1$	0	a_{m+4}	0	a_{m+6}	\dots	

Therefore, for each t satisfying (8.9), there are at most two corresponding pairs (m, ℓ) such that (8.8) is satisfied for some $n \leq N$. In other words, for each such t there are at most two values of (m, ℓ) such that $C_N^+(m, a, \ell)$ is non-empty. Next, in view of (8.8), observe that $C_N^+(m, a, \ell)$ is the union of two sets $A_N(d_1, \dots, d_{t+1})$ with

$$\left\{ \begin{array}{l} d_k = b_k \quad (1 \leq k \leq m) \\ d_{m+1} = b_{m+1} \pm 1, \\ d_{m+2} = b_{m+2} \pm (a_{m+2} - 1), \\ d_{m+2+i} = b_{m+2+i} \pm (-1)^i a_{m+2+i} \quad (1 \leq i \leq \ell - 1) \\ d_{m+2+\ell} = b_{m+2+\ell} \pm (-1)^\ell (a_{m+2+\ell} - a). \end{array} \right.$$

Then, on exploiting the same arguments used to establish (8.6) and (8.7), we obtain that

$$\begin{aligned} & \sum_{t=2}^{K+3} \sum_{\substack{(m,\ell) \\ m+\ell+1=t}} \sum_{a=1}^{2a_{t+1}} \sum_{\substack{n \in C_N^+(m,a,\ell) \\ c \leq n \|n\alpha - \gamma\|}} \frac{1}{n \|n\alpha - \gamma\|} \\ & \ll (\log N) \sum_{t=2}^{K+3} (1 + \log a_t) + \sum_{t=1}^{K+2} a_t \ll (\log N)^2. \end{aligned} \quad (8.11)$$

Again, observe that this upper bound estimate is no bigger than the upper bound appearing in Theorem 1.6. On using Lemma 4.4, it can be readily verified that the sets $C_N(u, a)$ and $C_N^+(m, a, \ell)$ considered above account for all integers n with $1 \leq n \leq N$ such that $|\Sigma| \leq 1/2$ and $m = m(n, \alpha, \gamma) \leq K$.

8.1.3 The case $|\Sigma| > 1/2$, $m \leq K$

By Lemma 4.3, when $|\Sigma| > 1/2$ we have that $\|n\alpha - \gamma\| = 1 - |\Sigma|$ and therefore we can call upon Lemma 4.5 when required.

We consider two subcases. In subcase one, we assume that

$$\delta_1 \neq (-1)^m \operatorname{sgn}(\delta_{m+1})(a_1 - 1). \quad (8.12)$$

Then, by (4.22), $1 - |\Sigma| \geq |D_0| = \{\alpha\}$ and

$$\sum_{\substack{1 \leq n \leq N \\ |\Sigma| \geq 1/2, (8.12) \text{ holds}}} \frac{1}{n \|n\alpha - \gamma\|} \ll \sum_{n=1}^N \frac{1}{n} \ll \log N. \quad (8.13)$$

In subcase two, we assume that

$$\delta_1 = (-1)^m \operatorname{sgn}(\delta_{m+1})(a_1 - 1). \quad (8.14)$$

In view of properties (3.13), (3.14), (3.17) and (3.18), the equalities associated with (4.20) can only occur if the coefficients b_1, \dots, b_L of the Ostrowski expansion of γ have one of the following two patterns:

b_1	b_2	b_3	b_4	b_5	b_6	\dots
$a_1 - 1$	0	a_3	0	a_5	0	\dots
0	a_{m+2}	0	a_4	0	a_6	\dots

(8.15)

For positive integers $L \leq K + 2$ and for $1 \leq a \leq 2a_{L+1}$, let $F_N(L, a)$ be the set of positive integers $n \leq N$ such that $|\Sigma| > 1/2$, $m = m(n, \alpha, \gamma) \leq K$ and

$$\begin{cases} \delta_1 = (-1)^m \operatorname{sgn}(\delta_{m+1})(a_1 - 1), \\ \delta_{i+1} = (-1)^{i+m} \operatorname{sgn}(\delta_{m+1})a_{i+1} & (1 \leq i \leq L - 1), \\ a_{L+1} - (-1)^{L+m} \operatorname{sgn}(\delta_{m+1})\delta_{L+1} = a. \end{cases} \quad (8.16)$$

It is easily verified that the sets $F_N(L, a)$ account for the remaining positive integers $n \leq N$, not accounted for elsewhere in the proof of the theorem. By Lemmas 4.3 and 4.5, for $n \in F_N(L, a)$ we have that

$$\|n\alpha - \gamma\| \geq a|D_L|.$$

Next, on using (8.16), it is readily verified that $F_N(L, a)$ is the set $A_N(d_1, \dots, d_{L+1})$ defined within Lemma 5.2, with

$$\begin{cases} d_1 = b_1 + (-1)^m \operatorname{sgn}(\delta_{m+1})(a_1 - 1), \\ d_{i+1} = b_{i+1} + (-1)^{i+m} \operatorname{sgn}(\delta_{m+1})a_{i+1} & (1 \leq i \leq L - 1), \\ d_{L+1} = b_{L+1} - (-1)^{L+m} \operatorname{sgn}(\delta_{m+1})(a - a_{L+1}). \end{cases}$$

On exploiting the same arguments used to establish (8.6) and (8.7), we obtain that

$$\begin{aligned} & \sum_{L=1}^{K+2} \sum_{a=1}^{2a_{L+1}} \sum_{\substack{n \in F_N(L, a) \\ c \leq n \|n\alpha - \gamma\|}} \frac{1}{n \|n\alpha - \gamma\|} \\ & \ll (\log N) \sum_{L=1}^{K+2} (1 + \log a_L) + \sum_{L=1}^{K+2} a_L \ll (\log N)^2. \end{aligned} \quad (8.17)$$

Combining the lower bound estimates given by (8.1), (8.7), (8.11), (8.13) and (8.17) completes the proof of Theorem 1.6.

8.2 Proof of Theorem 1.4

Fix $\gamma \in \mathbb{R}$. Theorem 1.4 will follow from Theorem 1.6 if we can show that

$$\sum_{\substack{1 \leq n \leq N \\ n \|n\alpha - \gamma\| \leq 1}} \frac{1}{n \|n\alpha - \gamma\|} \ll (\log N)^2 \quad \text{for almost all } \alpha \in [0, 1). \quad (8.18)$$

Using the Borel-Cantelli Lemma in a standard way (e.g., see [9] or [52]) one can easily see that for almost all $\alpha \in \mathbb{R}$ the inequality

$$\|n\alpha - \gamma\| \leq \frac{1}{n(\log n)^2}$$

has only finitely many solutions $n \in \mathbb{N}$. Therefore, (8.18) will follow on showing that for some $\varepsilon \in (0, 1)$

$$\sum_{\substack{1 \leq n \leq N \\ 1/(\log n)^2 < n \|n\alpha - \gamma\| \leq 1}} \frac{1}{n \|n\alpha - \gamma\|} \ll (\log N)^{1+\varepsilon} \quad \text{for almost all } \alpha \in [0, 1).$$

Given an integer $n \geq 2$, define

$$\mathcal{A}_n := \left\{ \alpha \in [0, 1) : \frac{1}{n(\log n)^2} < \|n\alpha - \gamma\| \leq \frac{1}{n} \right\}$$

and

$$J(n) = \int_{\mathcal{A}_n} \frac{d\alpha}{\|n\alpha - \gamma\|}.$$

It is elementary to check that

$$\begin{aligned} J(n) &= \sum_{a=1}^n \left(\int_{(a+\gamma)/n-1/n^2}^{(a+\gamma)/n-1/(n \log n)^2} + \int_{(a+\gamma)/n+1/(n \log n)^2}^{(a+\gamma)/n+1/n^2} \right) \frac{d\alpha}{|n\alpha - a - \gamma|} \\ &\sim 4 \log \log n \quad \text{as } n \rightarrow \infty. \end{aligned}$$

This together with Fatou's lemma, implies that

$$\int_0^1 \left(\sum_{\substack{n=2 \\ (\log n)^{-2} < n \|n\alpha - \gamma\| \leq 1}}^{\infty} \frac{1}{n(\log n)^{1+\varepsilon} \|n\alpha - \gamma\|} \right) d\alpha \leq \sum_{n=2}^{\infty} \frac{J(n)}{n(\log n)^{1+\varepsilon}} < \infty.$$

The upshot is that the above integrand must be finite almost everywhere, therefore, for almost all α we have that

$$\sum_{\substack{1 \leq n \leq N \\ (\log n)^{-2} < n \|n\alpha - \gamma\| \leq 1}} \frac{1}{n \|n\alpha - \gamma\|} \leq (\log N)^{1+\varepsilon} \sum_{\substack{1 \leq n \leq N \\ (\log n)^{-2} < n \|n\alpha - \gamma\| \leq 1}} \frac{1}{n (\log n)^{1+\varepsilon} \|n\alpha - \gamma\|}$$

$$\ll (\log N)^{1+\varepsilon}.$$

This proves (8.18) and thereby completes the proof of Theorem 1.4.

8.3 Proof of Theorem 1.5

Let α and ψ be as in the statement of Theorem 1.5. Without loss of generality, we will assume that $\alpha \in (0, 1)$ and we let p_k/q_k denote the principal convergents of α . Let $(\varepsilon_i)_{i \in \mathbb{N}}$ be a sequence of elements from the set $\{0, 1\}$ and define, inductively a strictly increasing sequence of integers $(k_i)_{i \in \mathbb{N}}$ as follows. Let $k_1 \geq 1$ be the smallest integer satisfying

$$|D_{k_1-1}| + |D_{k_1}| < \min\{\alpha, 1 - \alpha\}. \quad (8.19)$$

For $i \geq 1$, assuming k_1, \dots, k_i are given, let k_{i+1} be taken large enough that

$$k_{i+1} \geq k_i + 5, \quad (8.20)$$

$$k_{i+1} - k_i \equiv \varepsilon_i \pmod{2}, \quad (8.21)$$

$$\frac{q_{k_{i+1}+1}}{\psi(q_{k_{i+1}+1})} \geq (i+1) \cdot q_{k_i+1} \quad (8.22)$$

and

$$\frac{q_{k_{i+1}}}{q_{k_{i-1}+1}} \leq \frac{q_{k_{i+1}+1}}{q_{k_i+1}}. \quad (8.23)$$

Note that it is possible to choose k_{i+1} in this manner because of the assumption that $f(N) = o(N)$ as $N \rightarrow \infty$. Now let

$$b_{k+1} := \begin{cases} 1 & \text{if } k = k_i \text{ for some } i \in \mathbb{N}, \\ 0 & \text{otherwise} \end{cases} \quad (8.24)$$

and

$$\gamma := \sum_{k=0}^{\infty} b_{k+1} D_k = \sum_{i=1}^{\infty} D_{k_i}.$$

It is readily seen that $\gamma \in (-\alpha, 1 - \alpha)$. Indeed, using (3.8) and (8.19) we have that

$$|\gamma| \leq \sum_{k=k_1}^{\infty} a_{k+1}|D_k| = |D_{k_1-1}| + |D_{k_1}| < \min\{\alpha, 1 - \alpha\}.$$

Thus, $(b_{k+1})_{k \geq 0}$ is the sequence of Ostrowski coefficients associated with the expansion of γ . In view of the uniqueness property of the Ostrowski expansion of a real number, it follows that two different sequences $(k_i)_{i \in \mathbb{N}}$ and $(k'_i)_{i \in \mathbb{N}}$ give rise to two different numbers γ and γ' . Note that, by (8.21), there are at least as many sequences $(k_i)_{i \in \mathbb{N}}$ as sequences $(\varepsilon_i)_{i \in \mathbb{N}}$. Hence, there is a set of γ as above of the cardinality of continuum satisfying (1.1). Now fix any one of them and for each $i \in \mathbb{N}$ let

$$n_i = \sum_{k=0}^{k_i} b_{k+1}q_k = \sum_{j=1}^i q_{k_j}.$$

Clearly, $n_i \leq q_{k_{i+1}}$. Using Corollary 4.1 (recall the results of §4.2 are valid even in the case $\gamma = 0$) and (3.5) we find that

$$\|n_i\alpha - \gamma\| \ll \frac{1}{q_{k_{i+1}+1}}.$$

Thus, it follows that

$$\begin{aligned} S_{n_i}(\alpha, \gamma) - \max_{1 \leq n \leq n_i} \frac{1}{n\|n\alpha - \gamma\|} &\geq \min \left\{ \frac{1}{n_{i-1}\|n_{i-1}\alpha - \gamma\|}, \frac{1}{n_i\|n_i\alpha - \gamma\|} \right\} \\ &\gg \min \left\{ \frac{q_{k_i+1}}{q_{k_{i-1}+1}}, \frac{q_{k_{i+1}+1}}{q_{k_i+1}} \right\} \stackrel{(8.23)}{=} \frac{q_{k_i+1}}{q_{k_{i-1}+1}} \\ &\stackrel{(8.22)}{\geq} i\psi(q_{k_{i+1}}), \end{aligned}$$

and thereby completes the proof of the theorem since i can be taken arbitrarily large.

8.4 Proof of Theorem 1.7

As before, without loss of generality we assume that $\gamma \in [-\alpha, 1 - \alpha)$. We can assume that (1.1) holds since, if this were not the case, there would exist $n_0 \in \mathbb{N}$ and $m_0 \in \mathbb{Z}$ such that $\gamma = n_0\alpha + m_0$. Then, on making the substitution $n' = n - n_0$ we would have that

$$\sum_{\substack{rN < n \leq N \\ \|n\alpha - \gamma\| \geq N^{-v}}} \frac{1}{\|n\alpha - \gamma\|} = \sum_{\substack{rN < n \leq N \\ \|(n - n_0)\alpha\| \geq N^{-v}}} \frac{1}{\|(n - n_0)\alpha\|}.$$

Since $n \in [rN, N]$ takes at least $\lfloor (1-r)N \rfloor - 1$ values the range of $|n'| = |n - n_0|$ will contain all positive integers up to $M := \lfloor \frac{1}{2} \lfloor (1-r)N \rfloor - 1 \rfloor$. Hence, assuming that N is large enough so that $N^{-v} \geq M^{-v'}$, where $v < v' := (v+1)/2 < 1$, we have that

$$\sum_{\substack{rN < n \leq N \\ \|n\alpha - \gamma\| \geq N^{-v}}} \frac{1}{\|n\alpha - \gamma\|} \geq \sum_{\substack{rM < n' \leq M \\ \|n'\alpha\| \geq M^{-v'}}} \frac{1}{\|n'\alpha\|},$$

and the desired result would follow from the homogeneous case.

Let $(b_{k+1})_{k \geq 0}$ be the Ostrowski coefficients of γ given within Lemma 3.2. Let K be the largest integer such that $q_K \leq N$, and let

$$n = \begin{cases} \sum_{k=0}^{K-1} b_{k+1} q_k & \text{if it is non-zero,} \\ q_K & \text{otherwise.} \end{cases}$$

It is readily seen that $1 \leq n \leq q_K \leq N$ and that the integer $m = m(n, \alpha, \gamma)$ defined in Lemma 4.3 satisfies $m \geq K$. Then, by Corollary 4.1

$$\|n\alpha - \gamma\| < (|\delta_{m+1}| + 2)|D_m| \leq \frac{4a_{m+1}}{q_{m+1}} < \frac{4a_{m+1}}{a_{m+1}q_m} = \frac{4}{q_m} \leq \frac{4}{q_K}.$$

Let $0 < \nu < \min\{v, w(\alpha)^{-1}\}$, where $w(\alpha)$ is the exponent of approximation of α – see (1.27). Note that $w(\alpha) < \infty$ for $\alpha \notin \mathcal{L}$. Then, by Lemma 1.1, $q_K^{-1} < \frac{1}{8}q_{K+1}^{-\nu} < \frac{1}{8}N^{-\nu}$ if N is bigger than some sufficiently large N_0 depending on α . Thus, for any $N > N_0$ and any γ satisfying (1.1), there exists a positive integer $n \leq N$ such that

$$\|n\alpha - \gamma\| < \frac{1}{2}N^{-\nu}.$$

Thus, for all $N > N_0$ and all γ satisfying (1.1), we have that

$$N_\gamma(\alpha, \frac{1}{2}N^{-\nu}) \neq \emptyset$$

were $N_\gamma(\alpha, \varepsilon)$ is given by (6.1). By Corollary 6.1

$$\lfloor N\varepsilon \rfloor \leq N(\alpha, \varepsilon) \leq 32N\varepsilon \quad \text{for } N^{-\nu} < \varepsilon < \varepsilon_0 \text{ and } N > N_0,$$

provided $N_0 = N_0(\alpha)$ is sufficiently large and $\varepsilon_0 = \varepsilon_0(\alpha)$ is sufficiently small. Then, by Lemma 6.3 it follows that

$$\frac{1}{4}N\varepsilon \leq \#N_\gamma(\alpha, \varepsilon) \leq 65N\varepsilon \tag{8.25}$$

provided that $4N^{-\nu} < \varepsilon < \frac{1}{2}\varepsilon_0$.

Now let $R > 1$ be an integer such that $R^\nu > 1040$. For $k \in \mathbb{N}$, consider the sets

$$A_k := \{n \in \mathbb{N} : NR^{-1} < n \leq N, R^{-(k+1)\nu} \leq \|n\alpha - \gamma\| < R^{-k\nu}\}.$$

Then,

$$\begin{aligned} \#A_k &\geq \#N_\gamma(\alpha, R^{-k\nu}) - \#(N/R)_\gamma(\alpha, R^{-k\nu}) - \#N_\gamma(\alpha, R^{-(k+1)\nu}) \\ &\geq \frac{1}{4}NR^{-k\nu} - 65(N/R)R^{-k\nu} - 65NR^{-(k+1)\nu} \\ &\geq \left(\frac{1}{4} - 130R^{-\nu}\right)NR^{-k\nu} \geq \frac{1}{8}NR^{-k\nu}, \end{aligned}$$

provided that $4N^{-\nu} < R^{-k\nu} < \frac{1}{2}\varepsilon_0$. The latter holds when $k_0 \leq k \leq \log N / \log R - 1$ for some fixed $k_0 \in \mathbb{N}$. Clearly, the sets A_k are disjoint and on taking $r = R^{-1}$ we obtain that

$$\begin{aligned} \sum_{\substack{rN \leq n \leq N \\ \|n\alpha - \gamma\| \geq N^{-\nu}}} \frac{1}{\|n\alpha - \gamma\|} &\geq \sum_{k_0 \leq k \leq \log N / \log R - 1} \sum_{n \in A_k} \frac{1}{\|n\alpha - \gamma\|} \\ &\geq \sum_{k_0 \leq k \leq \log N / \log R - 1} R^{k\nu} \cdot \#A_k \\ &\geq \sum_{k_0 \leq k \leq \log N / \log R - 1} \frac{1}{8}N \\ &\geq \frac{1}{8}N(\log N / \log R - k_0 - 1). \end{aligned}$$

This completes the proof of Theorem 1.7 since R and k_0 depend on α and ν only.

8.5 Proof of Theorem 1.8

If $\alpha \notin \mathcal{L} \cup \mathbb{Q}$, then the desired conclusion follows immediately via Theorem 1.7. To prove the other direction, we suppose that $\alpha \in \mathcal{L}$ and explicitly construct a real number γ and a sequence $(N_i)_{i \in \mathbb{N}}$ such that

$$\sum_{1 \leq n \leq N_i} \frac{1}{\|n\alpha - \gamma\|} = o(N_i \log N_i) \quad \text{as } i \rightarrow \infty \quad (8.26)$$

Without loss of generality, we will assume that $\{\alpha\} < \frac{1}{3}$. Indeed, for any integer r we have the inequality $\|n(r\alpha)\| \leq |r|\|n\alpha\|$, which implies that

$$\sum_{1 \leq n \leq N_i} \frac{1}{\|n\alpha\|} \leq |r| \sum_{1 \leq n \leq N_i} \frac{1}{\|n(r\alpha)\|}. \quad (8.27)$$

Now, by Dirichlet's Theorem, we can find $r \in \{\pm 1, \pm 2, \pm 3\}$ such that $\{r\alpha\} < \frac{1}{3}$. The upshot of this and (8.27) is that if $\{\alpha\} \geq \frac{1}{3}$, we simply replace α by $\{r\alpha\}$ in the proof.

Recall that $w(\alpha) = \infty$ for $\alpha \in \mathcal{L}$. Then, by Lemma 1.1, we can find a sequence $\{K_i\}_{i \in \mathbb{N}}$ of positive integers such that

$$\lim_{i \rightarrow \infty} \frac{\log q_{K_{i+1}}}{\log q_{K_i}} = \infty.$$

We will assume that $K_1 > 1$. Since $q_{K_{i+1}} = a_{K_{i+1}}q_{K_i} + q_{K_{i-1}}$ we necessarily have that $a_{K_{i+1}} \rightarrow \infty$ as $i \rightarrow \infty$. Thus, we may also assume that $a_{K_{i+1}} \geq 8$ for all i . Next, we define the sequence $\{b_{k+1}\}_{k \geq 0}$ by setting

$$b_{k+1} := \begin{cases} \lceil \frac{a_{k+1}}{2} \rceil & \text{if } k = K_i \text{ for some } i, \\ 0 & \text{if } a_{k+1} = 1 \text{ or } 2, \\ 1 & \text{otherwise,} \end{cases} \quad (8.28)$$

and we let

$$\gamma := \sum_{k=0}^{\infty} b_{k+1} D_k. \quad (8.29)$$

It is easy to see that this expansion satisfies (3.17) and (3.18). Therefore, the sum (8.29) is absolutely convergent, the real number γ is well defined and the integers b_{k+1} , $k = 0, 1, \dots$ are the Ostrowski coefficients of γ .

Now, for each i let

$$a'_i = \left\lfloor \frac{a_{K_{i+1}}}{4} \right\rfloor \quad \text{and} \quad N_i = a'_i q_{K_i}. \quad (8.30)$$

Since $a_{K_{i+1}} \geq 8$, we have that $2 \leq a'_i < a_{K_{i+1}}$ and so

$$q_{K_i} \leq N_i < q_{K_{i+1}}. \quad (8.31)$$

In order to prove (8.26) we will use the techniques developed in §8.1. First of all we observe that for any positive integer $n \leq N_i$ its Ostrowski coefficient $c_{K_{i+1}}$ satisfies $c_{K_{i+1}} \leq a'_i \leq a_{K_{i+1}}/4$, while the corresponding Ostrowski coefficient of γ is $\geq a_{K_{i+1}}/2$. Hence

$$a_{K_{i+1}}/4 \leq |\delta_{K_{i+1}}| \leq \lceil a_{K_{i+1}}/2 \rceil < a_{K_{i+1}} - 1, \quad (8.32)$$

and the parameter $m = m(n, \alpha, \gamma)$ defined in Lemma 4.3 satisfies

$$m \leq K_i. \quad (8.33)$$

Now, consider the sets $C_{N_i}(u, a)$ and $C_{N_i}^+(m, a, \ell)$ as defined as in §8.1 with $N = N_i$. By definition, these deal with the situation $|\Sigma| \leq 1/2$. By Lemma 5.3 and (8.31), we have that

$$\#C_{N_i}(u, a) \leq \frac{3N_i}{q_{u+1}} + 1 \ll \frac{N_i}{q_{u+1}}. \quad (8.34)$$

In particular, for the case when $u \leq K_i - 1$, this implies that

$$\begin{aligned} & \sum_{u=0}^{K_i-1} \sum_{a=1}^{2a_{u+1}-1} \sum_{n \in C_{N_i}(u, a)} \frac{1}{\|n\alpha - \gamma\|} \\ & \stackrel{(8.3)}{\leq} \sum_{u=0}^{K_i-1} \frac{1}{|D_u|} \sum_{a=1}^{2a_{u+1}-1} \frac{\#C_{N_i}(u, a)}{a} \\ & \ll N_i \sum_{u=0}^{K_i-1} \sum_{a=1}^{2a_{u+1}-1} \frac{1}{a} \ll N_i \sum_{u=0}^{K_i-1} \log(a_{u+1}) \\ & = N_i \log(a_1 \cdots a_{K_i}) \leq N_i \log q_{K_i}. \end{aligned}$$

Next we deal with the set $C_{N_i}(u, a)$ when $u = K_i$. In this case, we have that

$$\#C_{N_i}(K_i, a) \leq 3N_i/q_{K_i+1} + 1 \stackrel{(8.30)}{\leq} 3 \left\lfloor \frac{a_{K_i+1}}{4} \right\rfloor q_{K_i}/q_{K_i+1} + 1 < \frac{3}{4} + 1.$$

Hence

$$\#C_{N_i}(K_i, a) \leq 1, \quad (8.35)$$

for all $a \geq 1$. Furthermore, since $a'_i \leq a_{K_i+1}/4$ and $b_{K_i+1} \geq a_{K_i+1}/2$ the sets $C_{N_i}(K_i, a)$ will be empty whenever $a < a_{K_i+1}/4 - 1$. Indeed, any $n \in C_{N_i}(K_i, a)$ has to lie in one of the four sets given by (8.4) and (8.5). If it lies within those given by (8.4), then

$$n \geq (b_{K_i+1} - (a + 1))q_{K_i} > (a_{K_i+1}/2 - a_{K_i+1}/4)q_{K_i} = (a_{K_i+1}/4)q_{K_i} \geq N_i$$

and this is impossible for $n \in C_{N_i}(K_i, a)$. If n lies within those given by (8.5), then either

$$b_{K_i+1} + (a + 1 - a_{K_i+1}) < 1 + a_{K_i+1}/2 + (a_{K_i+1}/4 - a_{K_i+1}) \leq 0,$$

which is not allowed, or

$$n \geq (b_{K_i+1} - (a + 1 - a_{K_i+1}))q_{K_i} > (a_{K_i+1}/2 - a_{K_i+1}/4 + a_{K_i+1})q_{K_i} > N_i,$$

which is also impossible. Therefore, the contribution from the sets $C_{N_i}(K_i, a)$ to the l.h.s. of (8.26) can be estimated as follows:

$$\begin{aligned} \sum_{a=1}^{2a_{K_i+1}-1} \sum_{n \in C_{N_i}(K_i, a)} \frac{1}{\|n\alpha - \gamma\|} &\stackrel{(8.3)\&(8.35)}{\leq} \sum_{a_{K_i+1}/4 \leq a+1 \leq 2a_{K_i+1}} \frac{1}{a|D_{K_i}|} \\ &\stackrel{(3.5)}{\ll} q_{K_i+1} \stackrel{(3.2)\&(8.30)}{\ll} N_i. \end{aligned} \quad (8.36)$$

Also note that the set $C_{N_i}(u, a)$ in the remaining case of $u = K_i + 1$ is always empty. Indeed, any $n \in C_{N_i}(K_i + 1, a)$ has to lie in one of the sets given by (8.4) and (8.5). In either case, the Ostrowski coefficient of q_{K_i} , which is either b_{K_i+1} or $b_{K_i+1} \pm 1$ is at least $a_{K_i+1}/2 - 1$ and so if $n \in C_{N_i}(K_i + 1, a)$ then

$$n \geq (a_{K_i+1}/2 - 1)q_{K_i} > (a_{K_i+1}/4)q_{K_i} \geq N_i$$

and this is impossible. Therefore, the contribution from the sets $C_{N_i}(K_i + 1, a)$ to the l.h.s. of (8.26) is zero.

Now we analyse the contribution to the l.h.s. of (8.26) arising from the sets $C_{N_i}^+(m, a, \ell)$. Recall that, by definition, we must have that the equalities associated with (8.8) are satisfied. In particular this means that for $k = 0, \dots, m + \ell$ the quantities $|\delta_{k+1}|$ can only be 0, 1, $a_{k+1} - 1$ or a_{k+1} . Then, by (8.32), $t := m + \ell + 1 \leq K_i$. By Lemma 5.3 and (8.30), we have that

$$\#C_{N_i}^+(m, a, \ell) \leq \frac{3N_i}{qt+1} + 1 \ll \frac{N_i}{qt+1}. \quad (8.37)$$

Recall that for a given t there are at most two possible pairs of (m, ℓ) as above for which $C_{N_i}^+(m, a, \ell)$ is non-empty. Furthermore, by (4.13), we obtain that

$$\|n\alpha - \gamma\| \geq a|D_t|.$$

Thus, the contribution from $C_{N_i}^+(m, a, \ell)$ with $t \leq K_i - 1$ is estimated as follows:

$$\sum_{t=2}^{K_i-1} \sum_{m+\ell+1=t} \sum_{a=1}^{2a_{t+1}-1} \sum_{n \in C_{N_i}^+(m, a, \ell)} \frac{1}{\|n\alpha - \gamma\|}$$

$$\begin{aligned}
&\ll \sum_{t=2}^{K_i-1} \sum_{m+\ell+1=t} \sum_{a=1}^{2a_{t+1}-1} \frac{N_i}{a|D_t|q_{t+1}} \\
&\ll N_i \sum_{t=2}^{K_i-1} (1 + \log a_{t+1}) \\
&\ll N_i \max\{K_i, \log(a_1 \dots a_{K_i})\} \ll N_i \log q_{K_i}.
\end{aligned}$$

If $t = K_i$, then

$$0 \leq c_{K_i+1} = b_{K_i+1} \pm (-1)^\ell (a_{K_i+1} - a) \leq N_i/q_{K_i} = a'_i \leq \frac{a_{K_i+1}}{4}.$$

Since $b_{K_i+1} = \lceil \frac{a_{K_i+1}}{2} \rceil$, we then have that $|a_{K_i+1} - a| \leq b_{K_i+1} \leq \frac{1}{2}a_{K_i+1} + \frac{1}{2}$, whence

$$a \geq \frac{1}{2}a_{K_i+1} - \frac{1}{2} \geq \frac{1}{4}a_{K_i+1}.$$

In this case

$$\|n\alpha - \gamma\| \geq a|D_{K_i}| \geq \frac{1}{4}a_{K_i+1}/2q_{K_i+1} \gg 1/q_{K_i} \gg N_i^{-1}.$$

Then, by (8.30) and (8.37), it follows that $\#C_{N_i}^+(m, a, \ell) \ll 1$ and thus

$$\sum_{m+\ell+1=K_i} \sum_{a \geq a_{K_i+1}} \sum_{n \in C_{N_i}^+(m, a, \ell)} \frac{1}{\|n\alpha - \gamma\|} \ll \sum_{a \geq a_{t+1}} \frac{N_i}{q_{K_i+1}a|D_{K_i}|} \ll N_i.$$

Finally, it remains to deal with the case $|\Sigma| > 1/2$. Note that in view of our assumption $0 < \{\alpha\} < \frac{1}{3}$ imposed at the start of the proof, we have that $a_1 \geq 3$. Then, since $K_1 > 1$, we have that $b_1 = 1$ and as is easily seen $\delta_1 = c_1 - b_1 \neq \pm(a_1 - 1)$. This means that the first term of (4.22) is never zero. Hence $1 - |\Sigma| \geq |D_0| = \{\alpha\}$. Then

$$\sum_{\substack{1 \leq n \leq N_i \\ |\Sigma| \geq 1/2}} \frac{1}{\|n\alpha - \gamma\|} \ll N_i. \quad (8.38)$$

Hence every positive integer $n \leq N_i$ has been accounted for and on combining the various estimates above we obtain the upper bound

$$\sum_{1 \leq n \leq N_i} \frac{1}{\|n\alpha - \gamma\|} \ll N_i \log q_{K_i}.$$

By our choice of the sequences $\{K_i\}$ and $\{N_i\}$, this implies (8.26) and thus completes the proof of Theorem 1.8.

9 Proofs of Theorems 2.2, 2.3 and 2.4

9.1 Preliminaries

We begin by describing our strategy and establishing various auxiliary statements. Instead of dealing with the multiplicative problem associated with (2.12), we consider the one-dimensional metrical problem associated with the inequality

$$\|n\beta - \delta\| < \frac{\psi(n)}{\|n\alpha - \gamma\|} := \Psi_\alpha^\gamma(n). \quad (9.1)$$

If $\Psi_\alpha^\gamma(n) \geq \frac{1}{2}$, then (9.1) holds automatically. Hence, to avoid this trivial pathological situation we will assume that

$$\Psi_\alpha^\gamma(n) < \frac{1}{2}. \quad (9.2)$$

Let E_n be the set of $\beta \in [0, 1]$ satisfying (9.1). It is easily verified that

$$|E_n| = 2\Psi_\alpha^\gamma(n).$$

If

$$\sum_{n=1}^{\infty} \Psi_\alpha^\gamma(n) \quad (9.3)$$

converges, the Borel-Cantelli Lemma implies that the set of β 's such that (9.1) (and hence (2.12)) is satisfied infinitely often is of Lebesgue measure zero. Hence, Theorem 2.2 will follow on proving that the conditions of the theorem ensure the convergence of $\sum_{n=1}^{\infty} \Psi_\alpha^\gamma(n)$. Note that in this case (9.2) holds for all sufficiently large n and thus our above argument is justified.

In order to prove Theorem 2.3 we will have to ensure that the sum (9.3) diverges. However, the approximating function Ψ_α^γ is not monotonic so we cannot apply the inhomogeneous version of Khintchine's theorem². Instead we will attempt to make use of known results regarding the Duffin-Schaeffer Conjecture; in particular the Duffin-Schaeffer Theorem. With the above strategy in mind, we now investigate the convergence/divergence behaviour of the sum (9.3).

Lemma 9.1. *Let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be decreasing and $\alpha, \gamma \in \mathbb{R}$ be given. If the sum (2.7) diverges and*

$$R_N(\alpha; \gamma) \gg N \log N \quad \text{for all } N \in \mathbb{N} \quad (9.4)$$

²The inhomogeneous version of Khintchine's theorem states that for any monotonic $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ and $\delta \in \mathbb{R}$, the set $\mathcal{S}^\delta(\psi) := \{\beta \in \mathbb{R} : \|n\beta - \delta\| < \psi(n) \text{ for i.m. } n \in \mathbb{N}\}$ is of full measure if $\sum \psi(n) = \infty$.

then the sum (9.3) diverges. Conversely, if the sum (2.7) converges and (2.11) is satisfied then the sum (9.3) converges.

Proof. By the partial summation formula, for any $N \in \mathbb{N}$ we have that

$$\sum_{n \leq N} \Psi_\alpha^\gamma(n) = \sum_{n \leq N} (\psi(n) - \psi(n+1))R_n(\alpha, \gamma) + \psi(N+1)R_N(\alpha, \gamma). \quad (9.5)$$

Now, if (9.4) holds and (2.7) diverges, then using the monotonicity of ψ and the fact that $\sum_{m \leq n} \log m \asymp n \log n$, we obtain that

$$\begin{aligned} \sum_{n \leq N} \Psi_\alpha^\gamma(n) &\gg \sum_{n \leq N} (\psi(n) - \psi(n+1))n \log n + \psi(N+1)N \log N \\ &\asymp \sum_{n \leq N} (\psi(n) - \psi(n+1)) \sum_{m \leq n} \log m + \psi(N+1) \sum_{m \leq N} \log m \\ &= \sum_{n \leq N} \psi(n) \log n \rightarrow \infty \quad \text{as } N \rightarrow \infty. \end{aligned}$$

The proof of the convergence case of the lemma follows the same line of argument as above but with \gg reversed and makes use of the fact that $\sum_{n \leq N} \psi(n) \log n$ is bounded. \square

Combining Lemma 9.1 with Theorem 1.8 gives the following statement.

Corollary 9.1. *If the sum (2.7) diverges then for all $\alpha \notin \mathfrak{L} \cup \mathbb{Q}$ and for all $\gamma \in \mathbb{R}$, the sum (9.3) diverges.*

Lemma 9.2. *Let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be given and $n \mapsto n\psi(n)$ be decreasing. Furthermore, let $\alpha, \gamma \in \mathbb{R}$. If the sum (2.7) diverges and*

$$S_N(\alpha; \gamma) \gg (\log N)^2 \quad \text{for all } N \in \mathbb{N}, \quad (9.6)$$

then the sum (9.3) diverges. Conversely, if the sum (2.7) converges and (2.10) is satisfied, then the sum (9.3) converges.

Proof. By the partial summation formula, for any $N \in \mathbb{N}$ we have that

$$\begin{aligned} \sum_{n \leq N} \Psi_\alpha^\gamma(n) &= \sum_{n \leq N} (n\psi(n) - (n+1)\psi(n+1))S_n(\alpha, \gamma) \\ &\quad + (N+1)\psi(N+1)S_N(\alpha, \gamma). \end{aligned} \quad (9.7)$$

Now, if (9.6) holds and the sum (2.7) diverges, then using the monotonicity of $n \mapsto n\psi(n)$ and the fact that $\sum_{m \leq n} \frac{\log m}{m} \asymp (\log n)^2$, we obtain that

$$\begin{aligned} \sum_{n \leq N} \Psi_\alpha^\gamma(n) &\gg \sum_{n=1}^N (n\psi(n) - (n+1)\psi(n+1))(\log n)^2 + \\ &\quad + (N+1)\psi(N+1)(\log N)^2 \\ &\gg \sum_{n=1}^N (n\psi(n) - (n+1)\psi(n+1)) \sum_{m=1}^n \frac{\log m}{m} + \\ &\quad + (N+1)\psi(N+1) \sum_{m=1}^N \frac{\log m}{m}. \end{aligned}$$

By the partial summation formula, the r.h.s. is $\sum_{n \leq N} \psi(n) \log n$ and so we have that

$$\sum_{n \leq N} \Psi_\alpha^\gamma(n) \gg \sum_{n \leq N} \psi(n) \log n \rightarrow \infty \quad \text{as } N \rightarrow \infty.$$

The proof of the convergence case of the lemma follows the same line of argument as above but with \gg reversed and makes use of the fact that $\sum_{n \leq N} \psi(n) \log n$ is bounded. \square

9.2 Proof of Theorem 2.2

Combining Lemmas 9.1 and 9.2 together with the observations made at the beginning of §9.1 completes the proof of Theorem 2.2.

9.3 Proof of Theorem 2.3

For $\xi > 0$, define $\mathcal{N} := \mathcal{N}(\xi) \subseteq \mathbb{N}$ by

$$\mathcal{N} = \{n_i\}_{i \in \mathbb{N}} := \{n \in \mathbb{N} : \varphi(n)/n \geq \xi\}. \quad (9.8)$$

By choosing ξ small enough we can guarantee that the set \mathcal{N} has positive lower asymptotic density in \mathbb{N} ; i.e. that

$$\liminf_{N \rightarrow \infty} \#\{n_i \in \mathcal{N} : n_i \leq N\}/N > 0.$$

To see this, first of all note that

$$\begin{aligned} \frac{\varphi(n)}{n} &= \prod_{p|n} \left(1 - \frac{1}{p}\right) = \exp\left(-\sum_{p|n} \frac{1}{p} - \sum_{p|n} \sum_{m=2}^{\infty} \frac{1}{mp^m}\right) \\ &\gg \exp\left(-\sum_{p|n} \frac{1}{p}\right). \end{aligned} \quad (9.9)$$

By [53, Lemma 4], for any $\xi' > 0$ we can find an integer v (depending on ξ') so that for any $x > 0$, the number of integers $1 \leq n \leq x$ which satisfy

$$\sum_{\substack{p|n \\ p \geq v}} \frac{1}{p} \geq \xi'$$

is less than $x/2$. For all other integers we have by (9.9), together with Mertens's Theorem [31, Theorem 429], that

$$\frac{\varphi(n)}{n} \geq c \cdot \exp(-\xi' - \log \log v),$$

for some constant $c > 0$. Therefore choosing

$$\xi = c \cdot \exp(-\xi' - \log \log v)$$

yields a set \mathcal{N} with lower asymptotic density greater than $1/2$, which verifies our assertion. In particular, a consequence of the fact that the density is greater than $1/2$ is that

$$n_i \leq 2i, \quad (9.10)$$

for all $n_i \in \mathcal{N}$ with sufficiently large i . Now let ψ_α^γ denote the approximating function Ψ_α^γ restricted to \mathcal{N} ; i.e. for $n \in \mathbb{N}$

$$\psi_\alpha^\gamma(n) := \begin{cases} \frac{\psi(n)}{\|n\alpha - \gamma\|} & \text{if } n \in \mathcal{N} \\ 0 & \text{otherwise.} \end{cases}$$

By definition, we have that

$$\mathcal{S}^\delta(\psi_\alpha^\gamma) \subset \mathcal{S}^\delta(\Psi_\alpha^\gamma),$$

and that

$$\sum_{n \leq N} \frac{\varphi(n)}{n} \psi_\alpha^\gamma(n) \geq \xi \sum_{n \leq N} \psi_\alpha^\gamma(n). \quad (9.11)$$

Next, it follows from a well know discrepancy result in the theory of uniform distribution [49, Proposition 4], that if n_i is a strictly increasing sequence of positive integers and $\gamma \in \mathbb{R}$, then for almost all $\alpha \in \mathbb{R}$

$$\sum_{i \leq T} \frac{1}{\|n_i \alpha - \gamma\|} \gg T \log T. \quad (9.12)$$

Actually, this statement is explicitly established in the proof of [49, Theorem 2] – see the last displayed inequality on [49, page 516]. This together with (9.10) implies that there is a set $\mathcal{A} \subseteq \mathbb{R}$ of full measure such that for any $\alpha \in \mathcal{A}$

$$\begin{aligned} \sum_{n \leq n_T} \psi_\alpha^\gamma(n) &= \sum_{i \leq T} \psi_\alpha^\gamma(n_i) \\ &= \sum_{i \leq T} (\psi(n_i) - \psi(n_{i+1})) \sum_{j \leq i} \frac{1}{\|n_j \alpha - \gamma\|} \\ &\quad + \psi(n_{T+1}) \sum_{i \leq T} \frac{1}{\|n_i \alpha - \gamma\|} \\ &\gg \sum_{i \leq T} \psi(n_i) \log i \geq \sum_{i \leq T} \psi(2i) \log i \\ &\gg \sum_{n \leq T} \psi(n) \log n. \end{aligned}$$

Thus the divergence of the sum (2.7) implies that

$$\sum_{n=1}^{\infty} \psi_\alpha^\gamma(n) = \infty \quad \forall \alpha \in \mathcal{A} \quad \text{and} \quad \forall \gamma \in \mathbb{R}.$$

The upshot of this together with (9.11) is that for any $\alpha \in \mathcal{A}$ and $\gamma \in \mathbb{R}$ the hypotheses of the Duffin-Schaeffer Theorem, namely (2.3) and (2.4), are satisfied for the function ψ_α^γ . Thus, when $\delta = 0$ the Duffin-Schaeffer Theorem implies that set $\mathcal{S}^\delta(\psi_\alpha^\gamma)$ is of full measure. Hence, it follows that $\mathcal{S}^\delta(\psi_\alpha^\gamma)$ with $\delta = 0$ is of full measure, and this establishes Theorem 2.3.

Remark 9.1. Note that we have proven a little more than what is stated in Conjecture 2.1. Namely, that for any $\gamma \in \mathbb{R}$ and almost all $(\alpha, \beta) \in \mathbb{R}^2$ the inequality

$$\|n\alpha - \gamma\| \|n\beta\|' < \psi(n) \quad (9.13)$$

holds for infinitely $n \in \mathbb{N}$ if the the sum (2.7) diverges, where $\|n\beta\|'$ stands for the distance of $n\beta$ to the nearest integer coprime to n . Observe that the argument

used above would prove Conjecture 2.1 in full (i.e. for any real δ) if we had the inhomogeneous version of the Duffin-Schaeffer Theorem (see Problem 2.2) at hand.

9.4 Proof of Theorem 2.4

Let $\Psi_\alpha^\gamma(n)$ be given by

$$\Psi_\alpha^\gamma(n) := \frac{\psi(n)}{\|n\alpha - \gamma\|} \quad \text{with} \quad \psi(n) := \frac{1}{n(\log n)^2 \log \log \log n}.$$

Recall that

$$\frac{\varphi(n)}{n} \gg \frac{1}{\log \log n},$$

see for example [31, Theorem 328]. It follows, by the partial summation formula together with Theorem 1.8, that for any irrational $\alpha \in \mathbb{R} \setminus \mathcal{L}$,

$$\begin{aligned} \sum_{n \leq N} \frac{\varphi(n)}{n} \Psi_\alpha^\gamma(n) &\gg \sum_{n \leq N} \frac{\psi(n)}{\log \log n} \\ &= \sum_{n \leq N} \left(\frac{\psi(n)}{\log \log n} - \frac{\psi(n+1)}{\log \log(n+1)} \right) \sum_{m=1}^n \frac{1}{\|m\alpha - \gamma\|} \\ &\quad + \frac{\psi(N+1)}{\log \log(N+1)} \sum_{m=1}^N \frac{1}{\|m\alpha - \gamma\|} \\ &\gg \sum_{n \leq N} \frac{\psi(n) \log n}{\log \log n} = \sum_{n \leq N} \frac{1}{n \log n \log \log n \log \log \log n}. \end{aligned}$$

Thus

$$\sum_{n=1}^{\infty} \frac{\varphi(n)}{n} \Psi_\alpha^\gamma(n) = \infty.$$

It now follows, on assuming the truth of the Duffin-Schaeffer Conjecture for functions $\Psi_\alpha^\gamma(n)$, that for almost all $\beta \in \mathbb{R}$ the inequality

$$|n\beta - s| < \Psi_\alpha^\gamma(n)$$

holds for infinitely coprime pairs $(n, s) \in \mathbb{N} \times \mathbb{Z}$. This completes the proof.

10 Proof of Theorem 2.1

The approach we develop in this section inherits some ideas from the theory of regular/ubiquitous systems – see [7, 9]. To begin with we recall some basic statements. Note that the conclusion of Theorem 2.1 is trivial when $\alpha \in \mathbb{Q}$. Hence, throughout the proof we will assume that α is irrational.

10.1 A zero-one law and quasi-independence on average

We begin by recalling the following zero-one law originally discovered by Cassels [19], see also [14].

Lemma 10.1. *Let $\Psi : \mathbb{N} \rightarrow [0, +\infty)$ be any function such that $\Psi(n) \rightarrow 0$ as $n \rightarrow \infty$. Then, the set $\mathcal{W}(\Psi)$ of $x \in [0, 1]$ such that $\|nx\| < \Psi(n)$ for infinitely many $n \in \mathbb{N}$ is either of Lebesgue measure zero or Lebesgue measure one.*

The obvious consequence of this result is that establishing Theorem 2.1 only requires us to show that the set of interest, that is the set of $\beta \in [0, 1]$ such that (2.9) holds infinitely often, is of positive Lebesgue measure. To accomplish this task we will employ the following generalisation of the Borel-Cantelli Lemma from probability theory, see either of [52, Lemma 5], [9, §8] or [12, §2.1].

Lemma 10.2. *Let $E_t \subset [0, 1]$ be a sequence of Lebesgue measurable sets such that*

$$\sum_{t=1}^{\infty} |E_t| = \infty. \quad (10.1)$$

Suppose that there exists a constant $C > 0$ such that

$$\sum_{t,t'=1}^T |E_t \cap E_{t'}| \leq C \left(\sum_{t=1}^T |E_t| \right)^2 \quad (10.2)$$

for infinitely many $T \in \mathbb{N}$. Then

$$|\limsup_{t \rightarrow \infty} E_t| \geq \frac{1}{C}. \quad (10.3)$$

The independence condition (10.2) is often referred to as *quasi-independence on average* and together with the divergent sum condition guarantees that the

associated lim sup set is of positive measure. It does not guarantee full measure; i.e. that $|\limsup_{t \rightarrow \infty} E_t| = 1$. However, this is not an issue if we already know (by some other means) that the lim sup set satisfies a zero-one law.

Remark 10.1. In view of Lemma 10.1, the value of C (as long as it is positive and finite) in Lemma 10.2 is of no interest. The point is that if we can show that $|\mathcal{W}(\Psi)| > 0$, then Lemma 10.1 implies full measure; i.e. $|\mathcal{W}(\Psi)| = 1$.

10.2 Setting up a limsup set

Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $R \in \mathbb{N}$ satisfy $R > 1$. Given $t, k \in \mathbb{N}$, let $\Omega_{t,k}$ be the set of real numbers $\beta \in [0, 1]$ such that there exists a triple $(n, r, s) \in \mathbb{N} \times \mathbb{Z}^2$ of coprime integers such that

$$\begin{cases} R^{-k-1} \leq |n\alpha - r| < R^{-k}, \\ |n\beta - s| < R^{-t+k}, \\ R^{t-1} < n \leq R^t. \end{cases} \quad (10.4)$$

Lemma 10.3. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, let $(q_\ell)_{\ell \geq 0}$ be the sequence of denominators of the principal convergents of α , and let $R \geq 256$. Then for any positive integers t and k such that $2R^{-k} < \|q_2\alpha\|$ and*

$$R^{k+1} \leq q_\ell < R^{t-1} \quad (10.5)$$

for some ℓ , one has the following estimate on the Lebesgue measure of $\Omega_{t,k}$:

$$|\Omega_{t,k}| \geq \frac{1}{2}.$$

Proof. Let $\beta \in [0, 1]$. By Minkowski's Theorem for convex bodies (see [20, p.71]), there are integers n, s, r , not all zero, satisfying the system of inequalities

$$\begin{cases} |n\alpha - r| < R^{-k}, \\ |n\beta - s| < R^{-t+k}, \\ |n| \leq R^t. \end{cases} \quad (10.6)$$

In view of (10.5) we have that $t > 0$ and $t-k > 0$. Hence $R^{-k} < \frac{1}{2}$ and $R^{-t+k} < \frac{1}{2}$, and n cannot be zero as otherwise $n = r = s = 0$. Also, without loss of generality

we can assume that $n > 0$ and n, r, s are coprime. Note that if $\beta \in [0, 1] \setminus \Omega_{t,k}$ then we also necessarily have that either

$$\begin{cases} \|n\alpha\| = |n\alpha - r| < R^{-k-1}, \\ 1 \leq n \leq R^t, \end{cases}$$

or

$$\begin{cases} \|n\alpha\| = |n\alpha - r| < R^{-k}, \\ 1 \leq n \leq R^{t-1}. \end{cases}$$

In view of (10.5), Lemma 6.1 is applicable to either of the above systems and therefore the number of integers n satisfying at least one of these systems is bounded above by $64R^{t-k-1}$. Furthermore, observe that for every n the measure of $\beta \in [0, 1]$ satisfying $|n\beta - s| < R^{-t+k}$ for some $s \in \mathbb{Z}$ is $2R^{-t+k}$ (see, for example, [52, Lemma 8]). Hence, the Lebesgue measure of $[0, 1] \setminus \Omega_{t,k}$ is bounded above by

$$64R^{t-k-1} \times 2R^{-t+k} = 128R^{-1} \leq \frac{1}{2}$$

as, by hypothesis, $R \geq 256$. Hence, $|\Omega_{t,k}| \geq \frac{1}{2}$ as required. \square

Let \mathcal{T}^* be any subset of pairs of positive integers (t, k) satisfying (10.5) for some ℓ . The precise choice of \mathcal{T}^* will be made later. Further, given a positive integer pair $(t, k) \in \mathcal{T}^*$, define

$$N(t, k) := \left\{ (n, r, s) \in \mathbb{N} \times \mathbb{Z}^2 : \begin{cases} R^{t-1} < n \leq R^t, \\ R^{-k-1} < |n\alpha - r| < R^{-k}, \\ 0 \leq s \leq n, \gcd(n, r, s) = 1 \end{cases} \right\}. \quad (10.7)$$

Clearly, for two different pairs (t, k) and (t', k') from \mathcal{T}^*

$$(n, r, s) \in N(t, k) \quad \& \quad (n', r', s') \in N(t', k') \quad \implies \quad n \neq n'. \quad (10.8)$$

By the definition of $\Omega_{t,k}$, we have that

$$\begin{aligned} \Omega_{t,k} &\subset \bigcup_{(n,r,s) \in N(t,k)} \{ \beta \in \mathbb{R} : |n\beta - s| < R^{-t+k} \} \\ &\subset \bigcup_{(n,r,s) \in N(t,k)} \{ \beta \in \mathbb{R} : |\beta - s/n| < R^{-2t+k+1} \}. \end{aligned}$$

Let $Z(t, k)$ be a maximal subcollection of $N(t, k)$ such that

$$\left| \frac{s_1}{n_1} - \frac{s_2}{n_2} \right| > R^{-2t+k} \quad (10.9)$$

for any distinct triples (n_1, r_1, s_1) and (n_2, r_2, s_2) from $Z(t, k)$. By the maximality of $Z(t, k)$, it follows that

$$\Omega_{t,k} \subset \bigcup_{(n,r,s) \in Z(t,k)} \left\{ \beta \in \mathbb{R} : \left| \beta - \frac{s}{n} \right| < (R+1)R^{-2t+k} \right\}.$$

Since, by definition, for any $(t, k) \in \mathcal{T}^*$ condition (10.5) is satisfied for some ℓ , Lemma 10.3 is applicable, and we have that $|\Omega_{t,k}| \geq \frac{1}{2}$. Therefore,

$$\frac{1}{2} \leq |\Omega_{t,k}| \leq \#Z(t, k) \times (2R+2)R^{-2t+k},$$

and so $\#Z(t, k) \geq \frac{1}{4R+4}R^{2t-k}$. By (10.9), we also have that $\#Z(t, k) \leq R^{2t-k}$. Thus, we conclude that

$$C_1 R^{2t-k} \leq \#Z(t, k) \leq R^{2t-k} \quad \text{with} \quad C_1 = \frac{1}{4R+4}. \quad (10.10)$$

Now, given $\xi \in \mathbb{R}$, let

$$E_{t,k}(\xi) = \{ \beta \in \mathbb{R} : |\beta - \xi| < \psi(R^t)R^{-t+k} \}. \quad (10.11)$$

Furthermore, define

$$E_{t,k} := \bigcup_{(n,r,s) \in Z(t,k)} E_{t,k}(s/n) \quad (10.12)$$

and let E be the set of $\beta \in \mathbb{R}$ such that $\beta \in E_{t,k}$ for infinitely many pairs $(t, k) \in \mathcal{T}^*$. The following fairly straightforward statement reveals the role of E in establishing Theorem 2.1.

Lemma 10.4. *Let E be as above. Then $E \subset [0, 1]$ and for every $\beta \in E \setminus \mathbb{Q}$ we have that*

$$\|n\alpha\| \|n\beta\| < \psi(n) \quad (10.13)$$

for infinitely many $n \in \mathbb{N}$.

Proof. Without loss of generality, we can assume that $\psi(n) \rightarrow 0$ as $n \rightarrow \infty$. Let $\beta \in E \setminus \mathbb{Q}$. Then, there exist infinitely many pairs $(t, k) \in \mathcal{T}^*$ such that

$$\begin{cases} n \leq R^t, \\ |n\alpha - r| < R^{-k}, \\ |\beta - s/n| < \psi(R^t)R^{-t+k} \end{cases}$$

for some $(n, r, s) \in \mathbb{N} \times \mathbb{Z}^2$. Since $t > k$ for $(t, k) \in \mathcal{T}^*$, t must take arbitrarily large values. Since ψ is decreasing, we have that

$$\begin{aligned} \|n\alpha\| \|n\beta\| &\leq n |n\alpha - r| \cdot |\beta - s/n| \\ &\leq R^t R^{-k} \psi(R^t) R^{-t+k} \\ &= \psi(R^t) \leq \psi(n). \end{aligned} \tag{10.14}$$

Since $\psi(R^t) \rightarrow 0$ as $t \rightarrow \infty$, if there are only finitely many n arising this way, we would be able to find an $n \in \mathbb{N}$ such that $\|n\alpha\| \|n\beta\| = 0$. Since α is irrational, we would get that $\|n\beta\| = 0$, which means that $\beta \in \mathbb{Q}$ and contradicts the assumption that $\beta \in E \setminus \mathbb{Q}$. Hence, there must be infinitely many $n \in \mathbb{N}$ satisfying (10.14), and hence (10.13). The inclusion $E \subset [0, 1]$ follows from the fact that the approximants s/n to β lie in $[0, 1]$. The proof is thus complete. \square

Remark 10.2. By Lemmas 10.1, 10.2 and 10.4, the proof of Theorem 2.1 would be complete if we found a subset \mathcal{T}^* of suitably ordered pairs (t, k) such that the associated sequence $E_{t,k}$ satisfies conditions (10.1) and (10.2). In the next subsection we present an explicit choice of \mathcal{T}^* . Subsequently, we deal with establishing conditions (10.1) and (10.2) of Lemma 10.2.

10.3 Choosing the indexing set \mathcal{T}^*

The goal of this section is to make a choice of the indexing set \mathcal{T}^* of pairs of positive integers (t, k) introduced in §10.2. Recall, that we are given a monotonically decreasing function $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ such that the sum (2.7) diverges. In what follows we extend ψ to all real numbers $x \geq 1$ by setting $\psi(x) = \psi(\lfloor x \rfloor)$. Clearly, the extended function is decreasing. Without loss of generality, we can assume that

$$n\psi(n) \leq \frac{1}{2} \quad \text{for all } n \in \mathbb{N}. \tag{10.15}$$

If this were not the case, we could replace ψ with $\psi_1(q) = \min\{\psi(q), (2q)^{-1}\}$, which is monotonically decreasing and satisfies the divergence condition – see [5, Lemma 4] for a similar argument.

Next, using the Cauchy condensation test, we obtain that

$$\sum_{t=1}^{\infty} t R^t \psi(R^t) = \infty. \tag{10.16}$$

Explicitly,

$$\begin{aligned}
\infty &= \sum_{n=1}^{\infty} \psi(n) \log n = \sum_{t=1}^{\infty} \sum_{R^{t-1} \leq n < R^t} \psi(n) \log n \\
&\leq \sum_{t=1}^{\infty} \sum_{R^{t-1} \leq n < R^t} \psi(R^{t-1}) \log R^t \\
&= \sum_{t=1}^{\infty} (R^t - R^{t-1}) \psi(R^{t-1}) \log R^t \\
&\ll \sum_{t=1}^{\infty} t R^t \psi(R^t).
\end{aligned}$$

Let $\eta > 0$ be a sufficiently small real parameter and t_0 be a sufficiently large integer. Define

$$\mathcal{T}(t_0, \eta) := \{t \in \mathbb{N} : t \geq t_0, \psi(R^t) \geq R^{-(1+\eta)t}\}.$$

Since the sum $\sum_{t=1}^{\infty} t R^{-\eta t}$ converges for $\eta > 0$, the divergence condition (10.16) implies that

$$\sum_{t \in \mathcal{T}(t_0, \eta)} t R^t \psi(R^t) = \infty. \quad (10.17)$$

Let

$$\nu_1 = \frac{1}{3} + \eta \quad \text{and} \quad \nu_2 = \frac{1}{3} + 2\eta, \quad (10.18)$$

and let \mathcal{T} be the subset of $t \in \mathcal{T}(t_0, \eta)$ such that

$$R^{\nu_2 t} \leq q_\ell < R^{t-1}$$

for some ℓ . We now show that we can assume that

$$\sum_{t \in \mathcal{T}} t R^t \psi(R^t) = \infty. \quad (10.19)$$

Indeed, by (10.17), this is true if the sum over the complement of \mathcal{T} converges. Note that t lies in the complement of \mathcal{T} if

$$q_\ell < R^{\nu_2 t} \quad \text{and} \quad R^{t-1} \leq q_{\ell+1} \quad (10.20)$$

for some ℓ . In particular, this implies that

$$\|q_\ell \alpha\| \stackrel{(3.5)}{<} \frac{1}{q_{\ell+1}} \stackrel{(10.20)}{\leq} R^{-t+1} = R(R^{\nu_2 t})^{-1/\nu_2} \leq R q_\ell^{-1/\nu_2}.$$

Note that the latter holds only finitely often for sufficiently small η if the exponent of approximation of α is less than 3. Alternatively, if we are assuming (2.8), then we have that

$$\psi(q_\ell) \geq Rq_\ell^{-1/\nu_2}$$

for all sufficiently large ℓ . In this case

$$\|q_\ell\alpha\| < Rq_\ell^{-1/\nu_2} \leq \psi(q_\ell)$$

for infinitely many ℓ . Then, for every $\beta \in [0, 1]$ we have that

$$\|q_\ell\alpha\| \cdot \|q_\ell\beta\| \leq \|q_\ell\alpha\| < \psi(q_\ell) \quad (10.21)$$

for infinitely many ℓ . Thus, if (10.19) does not hold the conclusion of Theorem 2.1 holds anyway. Thus, from now on (10.19) will be assumed.

Finally we let

$$\mathcal{T}^* := \{(t, k) : t \in \mathcal{T}, \lceil \nu_1 t \rceil \leq k < \lfloor \nu_2 t \rfloor\}.$$

We now verify that the sum of the measures of the sets $E_{t,k}$ taken over $(t, k) \in \mathcal{T}^*$ is divergent. Moreover, we provide an estimate for the rate of divergence. In what follows

$$S_{\mathcal{T}}(T) := \sum_{t \in \mathcal{T}, t \leq T} tR^t\psi(R^t), \quad (10.22)$$

and \sum^* indicates the summation over $(t, k) \in \mathcal{T}^*$; for example, $\sum_{t \leq T}^*$ means ‘sum over $(t, k) \in \mathcal{T}^*$ with $t \leq T$ ’.

Lemma 10.5. *For any $T > t_0$ we have that*

$$\eta C_1 S_{\mathcal{T}}(T) \leq \sum_{t \leq T}^* |E_{t,k}| \leq 2\eta S_{\mathcal{T}}(T), \quad (10.23)$$

where C_1 is as in (10.10). In particular,

$$\sum_{t \leq T}^* |E_{t,k}| \rightarrow \infty \quad \text{as } T \rightarrow \infty. \quad (10.24)$$

Proof. Let $(t, k) \in \mathcal{T}^*$ with $t \leq T$. Recall that for any distinct triples (n_1, r_1, s_1) and (n_2, r_2, s_2) from $Z(t, k)$ we have that (10.9) is satisfied. Furthermore, by

(10.11), (10.15) and the fact that $R^{-t+k} < 1/2$, the radii of $E_{t,k}(s/n)$ and $E_{t,k}(s'/n')$ are $\leq \frac{1}{2}R^{-2t+k}$. Hence $E_{t,k}(s/n)$ and $E_{t,k}(s'/n')$ are disjoint for distinct triples (n_1, r_1, s_1) and (n_2, r_2, s_2) from $Z(t, k)$. Therefore, on using (10.10), it follows that

$$\begin{aligned} |E_{t,k}| &= \sum_{(n,r,s) \in Z(t,k)} |E_{t,k}(s/n)| \\ &= 2\psi(R^t)R^{-t+k} \cdot \#Z(t, k) \end{aligned} \quad (10.25)$$

$$\stackrel{(10.10)}{\geq} 2\psi(R^t)R^{-t+k} \cdot C_1R^{2t-k} = 2C_1R^t\psi(R^t), \quad (10.26)$$

and

$$\begin{aligned} |E_{t,k}| &= 2\psi(R^t)R^{-t+k} \cdot \#Z(t, k) \\ &\stackrel{(10.10)}{\leq} 2\psi(R^t)R^{-t+k} \cdot R^{2t-k} = 2R^t\psi(R^t). \end{aligned} \quad (10.27)$$

Now, for each fixed $t \in \mathcal{T}$ the number of different k such that $(t, k) \in \mathcal{T}^*$ is $\lfloor \nu_2 t \rfloor - \lceil \nu_1 t \rceil \geq \nu_2 t - \nu_1 t - 2 = \eta t - 2 \geq \eta t/2$, as long as $t_0 \geq 2/\eta$. Also, $\lfloor \nu_2 t \rfloor - \lceil \nu_1 t \rceil \leq \eta t$. Then, (10.23) readily follow from (10.26) and (10.27), while the divergence condition (10.24) follows from (10.23) and (10.17). \square

10.4 Overlaps estimates for $E_{t,k}$

In the previous section, we established the divergent sum condition (10.1) of Lemma 10.2 for the sets $E_{t,k}$ with $(t, k) \in \mathcal{T}^*$. In order to complete the proof of Theorem 2.1 it remains to establish the quasi-independence on average condition (10.2) of Lemma 10.2 for these sets. Observe, that in view of (10.23), this boils down to showing that for T sufficiently large

$$\sum_{t \leq T}^* \sum_{t' \leq T'}^* |E_{t,k} \cap E_{t',k'}| \ll S_{\mathcal{T}}(T)^2. \quad (10.28)$$

10.4.1 Preliminary analysis

Let (t, k) and (t', k') be in \mathcal{T}^* . In particular, we have that $t, t' \geq t_0$,

$$\psi(R^t) \geq R^{-(1+\eta)t} \quad \text{and} \quad \psi(R^{t'}) \geq R^{-(1+\eta)t'}, \quad (10.29)$$

and also that

$$\lceil \nu_1 t \rceil \leq k < \lfloor \nu_2 t \rfloor \quad \text{and} \quad \lceil \nu_1 t' \rceil \leq k' < \lfloor \nu_2 t' \rfloor. \quad (10.30)$$

Our goal is to estimate the measure of $E_{t,k} \cap E_{t',k'}$ for $(t, k) \neq (t', k')$. To begin with, note that, by (10.12)

$$E_{t,k} \cap E_{t',k'} = \bigcup_{\substack{(n,r,s) \in Z(t,k) \\ (n',r',s') \in Z(t',k')}} E_{t,k}(s/n) \cap E_{t',k'}(s'/n'). \quad (10.31)$$

Clearly,

$$|E_{t,k}(s/n) \cap E_{t',k'}(s'/n')| \leq \min \{ |E_{t,k}(s/n)|, |E_{t',k'}(s'/n')| \}.$$

Together with (10.11), this gives that

$$|E_{t,k}(s/n) \cap E_{t',k'}(s'/n')| \leq 2 \min \{ \psi(R^t) R^{-t+k}, \psi(R^{t'}) R^{-t'+k'} \}. \quad (10.32)$$

Using (10.9) and (10.11) we obtain that for any given $(n', r', s') \in Z(t', k')$ the number of triples $(n, r, s) \in Z(t, k)$ such that

$$E_{t,k}(s/n) \cap E_{t',k'}(s'/n') \neq \emptyset \quad (10.33)$$

is at most

$$2 + \frac{|E_{t',k'}(s'/n')|}{R^{-2t+k}} = 2 + \frac{2\psi(R^{t'}) R^{-t'+k'}}{R^{-2t+k}}.$$

By (10.10), we have that $\#Z(t', k') \leq R^{2t'-k'}$. Hence the total number of pairs of triples $(n', r', s') \in Z(t', k')$ and $(n, r, s) \in Z(t, k)$ such that $E_{t,k}(s/n) \cap E_{t',k'}(s'/n') \neq \emptyset$ is at most

$$2 \left(1 + \frac{\psi(R^{t'}) R^{-t'+k'}}{R^{-2t+k}} \right) R^{2t'-k'}.$$

Together with (10.32), this gives that

$$\begin{aligned} |E_{t,k} \cap E_{t',k'}| &\leq 8\psi(R^t) R^t \psi(R^{t'}) R^{t'} \\ &\ll |E_{t,k}| |E_{t',k'}| \end{aligned} \quad (10.34)$$

provided that

$$\frac{\psi(R^{t'}) R^{-t'+k'}}{R^{-2t+k}} \geq 1. \quad (10.35)$$

Since the roles of (t, k) and (t', k') can be reversed in the above argument estimate (10.34) also holds when

$$\frac{\psi(R^t)R^{-t+k}}{R^{-2t'+k'}} \geq 1. \quad (10.36)$$

The upshot is that if either (10.35) or (10.36) holds then we are in good shape. In short, (10.34) together with (10.26) and (10.27) implies that the sets $E_{t,k}$ and $E_{t',k'}$ are pairwise quasi-independent; namely that

$$|E_{t,k} \cap E_{t',k'}| \ll |E_{t,k}| |E_{t',k'}|.$$

10.4.2 Further analysis

We now use a divergent technique to estimate from above the number of pairs of triples $(n, r, s) \in Z(t, k)$ and $(n', r', s') \in Z(t', k')$ such that (10.33) holds. First of all note that (10.33) implies that

$$\left| \beta - \frac{s}{n} \right| < \psi(R^t)R^{-t+k} \quad \text{and} \quad \left| \beta - \frac{s'}{n'} \right| < \psi(R^{t'})R^{-t'+k'}$$

for some $\beta \in [0, 1]$. Hence, by the triangle inequality, we get that

$$\left| \frac{s}{n} - \frac{s'}{n'} \right| \leq 2 \max\{\psi(R^t)R^{-t+k}, \psi(R^{t'})R^{-t'+k'}\}. \quad (10.37)$$

Then, multiplying (10.37) by $n'n$ and using the fact that $n \leq R^t$ and that $n' \leq R^{t'}$, we obtain that

$$|n's - ns'| \leq 2 \max\{\psi(R^t)R^{t'+k}, \psi(R^{t'})R^{t+k}\} =: \Delta. \quad (10.38)$$

Thus the original counting problem related to (10.33) is replaced by the problem of estimating the number of solutions to (10.38). This is typical for the type of problem under consider, see [32] or [52]. However, the available techniques to analyse solutions to (10.38) assume that the pairs (n, s) and (n', s') are coprime and thus for distinct pairs we have that

$$n's - ns' \neq 0. \quad (10.39)$$

Unfortunately, we are not able to impose such an assumption and thus we need to develop a different argument.

In this section, let us continue with the task of counting solutions to (10.38) under the condition that (10.39) holds. With this in mind, first of all observe

that condition (10.39) together with (10.38) implies that $\Delta \geq 1$. Fix $n, n' \in \mathbb{N}$. Clearly, n and n' uniquely define r and r' , since they are the closest integers to $n\alpha$ and $n'\alpha$ respectively. Thus, to fulfill our goal it will be sufficient to count the number of different pairs (s, s') subject to (10.38) and (10.39) simultaneously. We consider two cases: (i) the rank of the \mathbb{Z} -module generated by the collection of vectors (s, s') is 1, and (ii) the rank is 2. Clearly, these cases cover all possible options.

Rank 1 case. In this case we have that all vectors (s, s') in question are collinear. Then there is a fixed non-zero integer vector (s_0, s'_0) such that any other integer vector (s, s') in question has the form $(s, s') = \ell(s_0, s'_0)$ for some $\ell \in \mathbb{Z}$. Since $1 \leq |n's - ns'| \leq \Delta$, we obtain that $1 \leq |\ell| \cdot |n's_0 - ns'_0| \leq \Delta$. Thus, $|\ell| \leq \Delta/|n's_0 - ns'_0| \leq \Delta$. Therefore, the number of pairs (s, s') in the rank one case is no more than 2Δ .

Rank 2 case. In this case there are 2 linearly independent vectors (s, s') . All the vectors (s, s') in question lie in the convex subset of points $(x, y) \in \mathbb{R}^2$ defined by the following system of inequalities

$$|n'x - ny| \leq \Delta, \quad 0 \leq x \leq n.$$

The volume of this set is easily seen to be 2Δ . Hence, by Blichfeld's Theorem³ [18], this body contains at most $4\Delta + 2 \leq 6\Delta$ integer vectors.

The upshot of our discussion is that in either case the number of integer vectors (s, s') in question is $\leq 6\Delta$. Hence, using (10.32) and the definition of Δ , we obtain the following estimate

$$\begin{aligned} \sum_{(s, s')} |E_{t, k}(s/n) \cap E_{t', k'}(s'/n')| &\leq \\ &\leq 24 \max\{\psi(R^t)R^{t+k}, \psi(R^{t'})R^{t+k'}\} \cdot \min\{\psi(R^t)R^{-t+k}, \psi(R^{t'})R^{-t'+k'}\} \\ &= 24R^{-t}R^{-t'} \max\{\psi(R^t)R^{t+k}, \psi(R^{t'})R^{t+k'}\} \cdot \min\{\psi(R^t)R^{t+k}, \psi(R^{t'})R^{t+k'}\} \\ &= 24R^{-t}R^{-t'} \cdot \psi(R^t)R^{t+k} \cdot \psi(R^{t'})R^{t+k'} \\ &= 24\psi(R^t)R^k \psi(R^{t'})R^{k'}. \end{aligned}$$

Recall that, by definition, $Z(t, k) \subset N(t, k)$, and so n satisfies the condition $\|n\alpha\| < R^{-k}$. Since for $(t, k) \in \mathcal{T}^*$ inequalities (10.5) are satisfied for some ℓ ,

³Blichfeld's Theorem states that for any convex bounded body $B \subset \mathbb{R}^n$ and any lattice Λ in \mathbb{R}^n such that $\text{rank}(B \cap \Lambda) = n$ the cardinality of $B \cap \Lambda$ is $\leq n! \frac{\text{vol}_n(B)}{\det \Lambda} + n$.

Lemma 6.1 is applicable. This implies that the number of different integers n in question is $\leq 32R^{t-k}$. Similarly, the number of distinct values which can be realized by the integer n' is $\leq 32R^{t'-k'}$. Therefore,

$$\sum_{n's - ns' \neq 0} |E_{t,k}(s/n) \cap E_{t',k'}(s'/n')| \leq \underbrace{24576}_{24 \cdot 32^2} \psi(R^t)R^t \psi(R^{t'})R^{t'}, \quad (10.40)$$

where the sum is taken over $(n, r, s) \in Z(t, k)$ and $(n', r', s') \in Z(t', k')$ subject to condition (10.39). The upshot is that if (10.39) holds then again we are in good shape; the sets $E_{t,k}$ and $E_{t',k'}$ are pairwise quasi-independent.

10.4.3 The remaining case

In this section we consider the case when none of the conditions (10.35), (10.36) or (10.39) holds. Thus, for the rest of the proof we will assume that

$$\frac{\psi(R^{t'})R^{-t'+k'}}{R^{-2t+k}} \leq 1, \quad \frac{\psi(R^t)R^{-t+k}}{R^{-2t'+k'}} \leq 1 \quad (10.41)$$

and investigate the following subset of the overlap between the sets $E_{t,k}$ and $E_{t',k'}$:

$$\bigcup_{n's - ns' = 0} E_{t,k}(s/n) \cap E_{t',k'}(s'/n'), \quad (10.42)$$

where the union is taken over $(n, r, s) \in Z(t, k)$ and $(n', r', s') \in Z(t', k')$ subject to the condition

$$n's - ns' = 0. \quad (10.43)$$

Before we continue with the analysis of (10.42), we first show that t and t' satisfy the inequalities

$$t \leq \left(1 + \frac{6\eta}{5-6\eta}\right) t' \quad \text{and} \quad t' \leq \left(1 + \frac{6\eta}{5-6\eta}\right) t. \quad (10.44)$$

To see this, first of all note that (10.41) together with (10.29) imply that

$$\frac{R^{-(1+\eta)t'} R^{-t'+k'}}{R^{-2t+k}} \leq 1 \quad \text{and} \quad \frac{R^{-(1+\eta)t} R^{-t+k}}{R^{-2t'+k'}} \leq 1.$$

In view of (10.30), the first inequality above implies that

$$\left(\frac{5}{3} - 2\eta\right)t \leq 2t - k \leq (2 + \eta)t' - k' \leq \frac{5}{3}t'$$

whence the first inequality associated with (10.44) follows. The proof of the second inequality is identical, with the roles of (t, k) and (t', k') interchanged.

Now we return to estimating the measure of (10.42). Let β be any element of (10.42). Then there exist $(n, r, s) \in Z(t, k)$ and $(n', r', s') \in Z(t', k')$ satisfying (10.43), such that

$$\begin{cases} R^{t-1} \leq n \leq R^t, \\ |n\alpha - r| < R^{-k}, \\ |n\beta - s| < R^{-t+k} \end{cases} \quad (10.45)$$

and

$$\begin{cases} R^{t'-1} \leq n' \leq R^{t'}, \\ |n'\alpha - r'| < R^{-k'}, \\ |n'\beta - s'| < R^{-t'+k'}. \end{cases} \quad (10.46)$$

Note that since the vectors (n, r, s) and (n', r', s') are primitive and distinct and that $n > 0$ and $n' > 0$, the vectors (n, r, s) and (n', r', s') are not collinear. Hence, the cross product of (n, r, s) and (n', r', s') ; i.e.

$$(A, B, C) := (n, r, s) \times (n', r', s'),$$

is non-zero integer vector. By (10.43), we have that

$$B = -ns' + n's = 0.$$

Therefore,

$$|A| + |C| > 0. \quad (10.47)$$

Without loss of generality, we can assume that $0 < \alpha < 1$. Then $0 \leq r \leq n$ and $0 \leq r' \leq n'$. Further, observe that

$$C\alpha = (nr' - n'r)\alpha = (n\alpha - r)r' - (n'\alpha - r')r, \quad (10.48)$$

from which it follows that

$$|C| \leq \frac{1}{\alpha} \left(R^{-k+t'} + R^{-k'+t} \right). \quad (10.49)$$

Similarly, since $n's - ns' = 0$, we obtain that

$$A = rs' - r's = (r - n\alpha)s' - (r' - n'\alpha)s.$$

As before, since $0 < \beta < 1$, we obtain that

$$|A| \leq R^{-k+t'} + R^{-k'+t}. \quad (10.50)$$

Assuming that $\eta < 1/9$, by (10.18), (10.45) and the inequalities $\lceil \nu_1 t \rceil \leq k < \lfloor \nu_2 t \rfloor$, we obtain that

$$\begin{aligned} |n\alpha - r| &< R^{-k} \leq R^{-\nu_1 t}, \\ |n\beta - s| &< R^{-t+k} \leq R^{-(1-\nu_2)t} \leq R^{-\nu_1 t}. \end{aligned} \quad (10.51)$$

Observe that

$$\begin{aligned} (1, \alpha, \beta) \times (n, r, s) &= \left(\begin{vmatrix} \alpha & \beta \\ r & s \end{vmatrix}, - \begin{vmatrix} 1 & \beta \\ n & s \end{vmatrix}, \begin{vmatrix} 1 & \alpha \\ n & r \end{vmatrix} \right) \\ &= (-\alpha(n\beta - s) + \beta(n\alpha - r), n\beta - s, -(n\alpha - r)). \end{aligned}$$

Hence, on using (10.51) and the fact that $0 \leq \alpha, \beta \leq 1$, we obtain that

$$|(1, \alpha, \beta) \times (n, r, s)| \leq \sqrt{6} R^{-\nu_1 t}. \quad (10.52)$$

A similar argument shows that

$$|(1, \alpha, \beta) \times (n', r', s')| \leq \sqrt{6} R^{-\nu_1 t'}. \quad (10.53)$$

In particular, since $n \geq R^{t-1}$, (10.52) implies that the (acute) angle θ between $(1, \alpha, \beta)$ and (n, r, s) satisfies

$$\begin{aligned} |\sin \theta| &= \frac{|(1, \alpha, \beta) \times (n, r, s)|}{|(1, \alpha, \beta)| \cdot |(n, r, s)|} \\ &\leq \frac{\sqrt{6} R^{-\nu_1 t}}{n} \\ &\leq \sqrt{6} R R^{-(1+\nu_1)t}. \end{aligned}$$

Similarly, since $n' \geq R^{t'-1}$, (10.53) implies that the (acute) angle θ' between $(1, \alpha, \beta)$ and (n', r', s') satisfies

$$|\sin \theta'| \leq \sqrt{6} R R^{-(1+\nu_1)t'}.$$

Let ϱ be the angle between (n, r, s) and (n', r', s') . Then, by the triangle inequality for the projective distance (see, for example, [6, §3]),

$$|\sin \varrho| \leq |\sin \theta| + |\sin \theta'|$$

$$\leq 2 \max\{|\sin \theta|, |\sin \theta'|\}.$$

On the other hand, the angle $\tilde{\theta}$ between $(1, \alpha, \beta)$ and the vector subspace of \mathbb{R}^3 spanned by (n, r, s) and (n', r', s') is at most $\min\{\theta, \theta'\}$ and thus satisfies the inequality

$$|\sin \tilde{\theta}| \leq \min\{|\sin \theta|, |\sin \theta'|\}.$$

Hence the volume of the parallelepiped generated by $(1, \alpha, \beta)$, (n, r, s) and (n', r', s') is

$$\begin{aligned} & \underbrace{|(1, \alpha, \beta)|}_{\leq \sqrt{3}} \cdot \underbrace{|(n, r, s)|}_{\leq \sqrt{3}n} \cdot \underbrace{|(n', r', s')|}_{\leq \sqrt{3}n'} \cdot |\sin \varrho| \cdot |\sin \tilde{\theta}| \\ & \leq 6\sqrt{3} n n' |\sin \theta| \cdot |\sin \theta'| \\ & \leq 36\sqrt{3} R^2 R^t R^{t'} R^{-(1+\nu_1)t} R^{-(1+\nu_1)t'} \\ & \leq 36\sqrt{3} R^2 R^{-\nu_1 t} R^{-\nu_1 t'}. \end{aligned} \tag{10.54}$$

We also have that this volume is equal to

$$|(1, \alpha, \beta) \cdot ((n, r, s) \times (n', r', s'))| = |(1, \alpha, \beta) \cdot (A, B, C)| = |C\beta + A|.$$

Combining this with (10.54), implies that

$$|C\beta + A| \leq 36\sqrt{3} R^2 R^{-\nu_1 t} R^{-\nu_1 t'}. \tag{10.55}$$

In particular, this together with (10.47) and the facts that $A \in \mathbb{Z}$ and that t can be taken arbitrarily large, implies that $C \neq 0$. Combining (10.49), (10.50) and (10.55), we have proved the following statement.

Lemma 10.6. *Let $\Upsilon = R^{-k+t'} + R^{-k'+t}$ and $\Theta = 36\sqrt{3} R^2 R^{-\nu_1 t} R^{-\nu_1 t'}$. Then (10.42) is a subset of the following*

$$\bigcup_{C=1}^{\lceil \Upsilon/\alpha \rceil} \{\beta \in [0, 1] : \|C\beta\| < \Theta\}.$$

Recall that if $\Theta < 1/2$, then

$$|\{\beta \in [0, 1] : \|C\beta\| < \Theta\}| = 2\Theta.$$

Without loss of generality, the condition $\Theta < 1/2$ can be assumed since t and t' can be taken sufficiently large. Note that

$$t + t' \leq 2 \max\{t, t'\} \stackrel{(10.44)}{\leq} \frac{10}{5 - 6\eta} \min\{t, t'\}. \tag{10.56}$$

Then, using Lemma 10.6 and the inequalities $\lceil \nu_1 t \rceil \leq k < \lfloor \nu_2 t \rfloor$, we obtain that the measure of the set (10.42) is

$$\begin{aligned}
&\leq \frac{72\sqrt{3}R^2}{\alpha} R^{-\nu_1 t} R^{-\nu_1 t'} \cdot (R^{-k+t'} + R^{-k'+t}) \\
&\leq \frac{72\sqrt{3}R^2}{\alpha} \cdot \left(R^{-2\nu_1 t+(1-\nu_1)t'} + R^{-2\nu_1 t'+(1-\nu_1)t} \right) \\
&\stackrel{(10.44)}{\leq} \frac{72\sqrt{3}R^2}{\alpha} \cdot \left(R^{-2\nu_1 t+(1-\nu_1)\left(1+\frac{6\eta}{5-6\eta}\right)t} + R^{-2\nu_1 t'+(1-\nu_1)\left(1+\frac{6\eta}{5-6\eta}\right)t'} \right) \\
&= \frac{72\sqrt{3}R^2}{\alpha} \cdot \left(R^{-\frac{11\eta-12\eta^2}{5-6\eta}t} + R^{-\frac{11\eta-12\eta^2}{5-6\eta}t'} \right) \\
&\leq \frac{144\sqrt{3}R^2}{\alpha} \cdot R^{-\frac{11\eta-12\eta^2}{5-6\eta} \min\{t,t'\}} \\
&\stackrel{(10.56)}{\leq} \frac{144\sqrt{3}R^2}{\alpha} \cdot R^{-\frac{11\eta-12\eta^2}{10}(t+t')} \leq \frac{144\sqrt{3}R^2}{\alpha} \cdot R^{-\eta(t+t')}
\end{aligned}$$

provided that $\eta < 1/12$. Thus,

$$\left| \bigcup_{n's-n s'=0} E_{t,k}(s/n) \cap E_{t',k'}(s'/n') \right| \leq \frac{144\sqrt{3}R^2}{\alpha} \cdot R^{-\eta(t+t')}, \quad (10.57)$$

where, recall, the union is taken over $(n, r, s) \in Z(t, k)$ and $(n', r', s') \in Z(t', k')$ subject to condition (10.43) and assuming that (10.41) holds.

10.5 The finale

Estimates (10.40) and (10.57) combined together show that whenever conditions (10.41) are satisfied, we have that for $(t, k) \neq (t', k')$

$$|E_{t,k} \cap E_{t',k'}| \ll \psi(R^t)R^t \psi(R^{t'})R^{t'} + R^{-\eta(t+t')}. \quad (10.58)$$

This estimate also holds when conditions (10.41) are not satisfied, this time as a consequence of (10.34). Thus, (10.58) together with (10.23) implies that

$$\begin{aligned}
&\sum_{t \leq T}^* \sum_{t' \leq T'}^* |E_{t,k} \cap E_{t',k'}| \\
&\ll \sum_{t \leq T}^* \sum_{t' \leq T'}^* \psi(R^t)R^t \psi(R^{t'})R^{t'} + \sum_{t \leq T}^* \sum_{t' \leq T'}^* R^{-\eta(t+t')} + S_{\mathcal{T}}(T)
\end{aligned}$$

$$\begin{aligned}
&= \left(\sum_{t \in \mathcal{T}, t \leq T} \sum_{\lceil \nu_1 t \rceil \leq k < \lfloor \nu_2 t \rfloor} \psi(R^t) R^t \right)^2 \\
&\quad + \left(\sum_{t \in \mathcal{T}, t \leq T} \sum_{\lceil \nu_1 t \rceil \leq k < \lfloor \nu_2 t \rfloor} R^{-\eta t} \right)^2 + S_{\mathcal{T}}(T) \\
&\leq \left(\sum_{t \in \mathcal{T}, t \leq T} t \psi(R^t) R^t \right)^2 + \left(\sum_{t \in \mathcal{T}, t \leq T} t R^{-\eta t} \right)^2 + S_{\mathcal{T}}(T) \\
&\leq S_{\mathcal{T}}(T)^2 + \left(\sum_{t=1}^{\infty} t R^{-\eta t} \right)^2 + S_{\mathcal{T}}(T) \ll S_{\mathcal{T}}(T)^2
\end{aligned}$$

for sufficiently large T , since $S_{\mathcal{T}}(T) \rightarrow \infty$ as $T \rightarrow \infty$. This establishes (10.28) and thereby completes the proof of Theorem 2.1.

Acknowledgements. From establishing the first results to getting to this stage, this paper has taken over five years to complete. It has been a long but nevertheless interesting journey which has given rise to new and (in our collective judgments) fundamental problems, especially within the classical theory of inhomogeneous Diophantine approximation such as Conjecture 2.1 and Problem 2.2. One consequence of the five year time span is that we have missed Wolfgang Schmidt's eightieth birthday by two years (almost certainly more once the paper comes out in print). Hopefully it has been worth it: over the extra two years the paper, we, and of course Wolfgang, have nicely matured! The mathematical and personal influence of Wolfgang on the three of us has been immense, and we will forever be in his debt. His remarkable creativity and vision are still influencing current research; in particular at York. Thank you!

The authors are grateful to the anonymous reviewer whose comments have been most valuable, especially for drawing our attention to various classical results missing from an earlier version of the paper.

SV would like to thank his wonderful parents, Lalji and Manchaben, for graciously accepting my life, my love for Bridget and also for being fantastic grandparents to the dynamic duo Ayesha and Iona. All of these are a million miles from their norm. In short, they have been and remain to be my greatest teachers.

References

- [1] D. Badziahin, and S. Velani, *Multiplicatively badly approximable numbers and generalised Cantor sets*. Adv. Math., 225 (2011), 2766–2796.
- [2] J. Beck, *Probabilistic Diophantine approximation I: Kronecker sequences*. Ann. of Math. (2) 140 (1994), no. 2, 449–502.
- [3] J. Beck, *Probabilistic Diophantine approximation: Randomness in Lattice Point Counting*. Springer Monographs in Mathematics, 2014.
- [4] H. Behnke, *Zur theorie der diophantischen approximationen*. Abh. Math. Sem. Univ. Hamburg 3 (1924), no. 1, 261–318.
- [5] V. Beresnevich, *On approximation of real numbers by real algebraic numbers*. Acta Arith. 90 (1999), no. 2, 97–112.
- [6] V. Beresnevich, *Rational points near manifolds and metric Diophantine approximation*. Ann. of Math. (2) 175 (2012), no. 1, 187–235.
- [7] V. Beresnevich, V. Bernik, and M. Dodson, *Regular systems, ubiquity and Diophantine approximation*. A panorama of number theory or the view from Baker’s garden (Zrich, 1999), 260–279, Cambridge Univ. Press, Cambridge, 2002.
- [8] V. Beresnevich, V. Bernik, M. Dodson, and S. Velani, *Classical metric diophantine approximation revisited*. In Chen, W.W.L., Gowers, W.T., Halberstam, H., Schmidt, W.M., Vaughan, R.C. (eds.) Analytic Number Theory: Essays in Honour of Klaus Roth. Cambridge University Press, 2009, 38–61.
- [9] V. Beresnevich, D. Dickinson, and S. Velani, *Measure theoretic laws for lim sup sets*. Mem. Amer. Math. Soc. 179 (2006), no. 846, x+91 pp.
- [10] V. Beresnevich, D. Dickinson, and S. Velani, *Diophantine approximation on planar curves and the distribution of rational points*. With an Appendix II by R. C. Vaughan. Ann. of Math. (2) 166 (2007), no. 2, 367–426.
- [11] V. Beresnevich, A. Haynes, and S. Velani, *Multiplicative zero-one laws and metric number theory*. Multiplicative zero-one laws and metric number theory. Acta Arith. 160 (2013), no. 2, 101–114.

- [12] V. Beresnevich, F. Ramírez, and S. Velani, *Metric Diophantine Approximation: aspects of recent work*. in Badziahin, D., Gorodnik, A., and Pey-erimhoff, N. (eds.) Dynamics and Analytic Number Theory:. Cambridge: Cambridge University Press, pp. 1–95.
- [13] V. Beresnevich, R. C. Vaughan, and S. Velani, *Inhomogeneous Diophantine approximation on planar curves*. Math. Ann. 349 (2011), no. 4, 929-942.
- [14] V. Beresnevich, and S. Velani, *A note on zero-one laws in metrical Diophantine approximation*. Acta Arith. 133 (2008), no. 4, 363-374.
- [15] V. Beresnevich, and S. Velani, *A note on simultaneous Diophantine approximation on planar curves*. Math. Ann. 337 (2007), no. 4, 769-796.
- [16] V. Bernik, D. Kleinbock, and G. Margulis, *Khintchine-type theorems on manifolds: the convergence case for standard and multiplicative versions*. Internat. Math. Res. Notices 2001, no. 9, 453-486.
- [17] A. S. Besicovitch, *Sets of fractional dimensions (IV): on rational approximation to real numbers*, J. Lond. Math. Soc. **9** (1934), 126–131.
- [18] H. F. Blichfeldt, *Notes on geometry of numbers*, Bull. Amer. Math. Soc. 27 (1921), 150-153.
- [19] J. W. S. Cassels, *Some metrical theorems in Diophantine approximation. I*, Proc. Cambridge Philos. Soc., 46 (1950), pp. 209-218.
- [20] J. W. S. Cassels, *An introduction to the geometry of numbers*, Corrected reprint of the 1971 edition. Classics in Mathematics. Springer-Verlag, Berlin, 1997.
- [21] S. D. Chowla, *Some problems of diophantine approximation (I)*. *Math. Z.*, 33 (1931), no. 1, 544–563.
- [22] H. G. Diamond, and J. D. Vaaler, *Estimates for partial sums of continued fraction partial quotients*, Pacific. J. Math. 122 (1986), no.1, 73-82.
- [23] M. Drmota, R. F. Tichy, *Sequences, discrepancies and applications*. Lecture Notes in Mathematics, 1651. Springer-Verlag, Berlin, 1997.
- [24] R. J. Duffin, and A. C. Schaeffer, *Khintchine’s problem in metric Diophantine approximation*. Duke Math. J. 8 (1941), 243-255.

- [25] M. Einsiedler, A. Katok, and E. Lindenstrauss, *Invariant measures and the set of exceptions to Littlewood's conjecture*. Ann. of Math. (2) 164 (2006), no. 2, 513-560.
- [26] P. Gallagher, *Metric simultaneous Diophantine approximation*. Journal London Math. Soc. 37 (1962), 387-390.
- [27] A. Gorodnik, P. Vishe, *Simultaneous Diophantine approximation - logarithmic improvements*. Preprint: arXiv:1601.03525
- [28] G. Hardy, and J. Littlewood, *Some problems of diophantine approximation: The lattice-points of a right-angled triangle*. Proceedings of the London Mathematical Society (2) 20 (1922), 15-36.
- [29] G. Hardy, and J. Littlewood, *Some problems of diophantine approximation: The lattice-points of a right-angled triangle (second memoir)*. Abhandlungen aus dem Mathematischen Seminar der Hamburgischen Universität 9 (1922), 212-249.
- [30] G. Hardy, and J. Littlewood, *Some problems of diophantine approximation: a series of cosecants*. Bull. Calcutta Math. Soc. 20 (1930), 251-266.
- [31] G. Hardy, and E. Wright, *An introduction to the theory of numbers*. Sixth edition, Oxford University Press, Oxford, 2008.
- [32] G. Harman, *Metric Number Theory*. LMS Monographs New Series, vol. 18, Clarendon Press, 1998.
- [33] A. Haynes, A. D. Pollington, and S. L. Velani. *The Duffin-Schaeffer Conjecture with extra divergence*. Math Annalen 353 (2012), 259-273.
- [34] Loo Keng Hua, Yuan Wang, *Applications of number theory to numerical analysis*. Springer-Verlag, 1981.
- [35] V. Jarník: *Diophantischen Approximationen und Hausdorffsches Mass*, Mat. Sbornik **36** (1929), 371-382.
- [36] A. Y. Khintchine, *Einige Sätze über Kettenbrüche, mit Anwendungen auf die Theorie der Diophantischen Approximationen*. Math. Ann. 92 (1924), 115-125.
- [37] A. Y. Khintchine, *Continued fractions*. Dover Publications, 1997.

- [38] J. F. Koksma, Diophantische Approximationen. (German) Reprint. Springer-Verlag, Berlin-New York, 1974.
- [39] N. M. Korobov, Number-theoretic methods in approximate analysis. Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow. 1963.
- [40] A. H. Kruse, Estimates of $\sum_{k=1}^N k^{-s} \langle kx \rangle^{-t}$. *Acta Arith.* 12 (1966/1967), 229–261.
- [41] L. Kuipers, and H. Niederreiter, *Uniform distribution of sequences*. Wiley-Interscience, John Wiley & Sons, New York-London-Sydney, 1974.
- [42] S. Lang, *Report on diophantine approximations*. Bull. de la Soc. Math. de France 93 (1965), 117-192.
- [43] S. Lang, *Introduction to Diophantine approximations. New expanded edition*. Springer-Verlag, New York, 1995.
- [44] Thái Hoàng Lê, Jeffrey D. Vaaler, Sums of products of fractional parts. Proc. Lond. Math. Soc. (3) 111 (2015), no. 3, 561–590.
- [45] E. Lindenstrauss, U. Shapira, Homogeneous orbit closures and applications. *Ergodic Theory Dynam. Systems* 32 (2012), no. 2, 785–807.
- [46] A. Pollington, and S. Velani, *On a problem in simultaneous Diophantine approximation: Littlewood’s conjecture*. Acta Math. 185 (2000), 287-306.
- [47] A. Rockett, and P. Szűsz, *Continued Fractions*. World Scientific, Singapore, 1992.
- [48] V. Saligrama, *Deterministic Designs with Deterministic Guarantees: Toeplitz Compressed Sensing Matrices, Sequence Designs and System Identification*. Information Theory, IEEE Transactions on. DOI: 10.1109/TIT.2012.2206951.
- [49] W. M. Schmidt, *Metrical theorems on fractional parts of sequences*. Trans. Amer. Math. Soc. 110 (1964), 493-518.
- [50] Y. Sinai, and C. Ulcigrai, *Estimates from above of certain double trigonometric sums*. J. Fixed Point Theory Appl. 6 (2009), no. 1, 93-113.
- [51] U. Shapira, *A solution to a problem of Cassels and Diophantine properties of cubic numbers*. Ann. of Math. (2) 173 (2011), no. 1, 543-557.

- [52] V. G. Sprindžuk, *Metric Theory of Diophantine Approximation*. Scripta Series in Mathematics. New York: John Wiley & Sons, 1979.
- [53] J. D. Vaaler, *On the metric theory of Diophantine approximation*. Pacific J. Math. **76** (1978) 527-539.
- [54] A. Venkatesh, *The work of Einsiedler, Katok and Lindenstrauss on the Littlewood conjecture*. Bull. Amer. Math. Soc. 45 (2008), 117-134.
- [55] A. Walfisz, Über einige trigonometrische Summen. (German) *Math. Z.* 33 (1931), no. 1, 564–601.
- [56] A. Walfisz, Über einige trigonometrische Summen II. (German) *Math. Z.* 35 (1932), no. 1, 774–788.

VB, SV :

Department of Mathematics, University of York,
Heslington, York, YO10 5DD, England
e-mails: `victor.beresnevich@york.ac.uk`
`sanju.velani@york.ac.uk`

AH :

Department of Mathematics, University of Houston,
Texas, USA
e-mail: `haynes@math.uh.edu`