UNIVERSITY of York

This is a repository copy of *Phytoremediation of metals using lemongrass* (*Cymbopogon citratus (D.C.) Stapf.*) grown under different levels of red mud in soil amended with biowastes.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/114700/</u>

Version: Accepted Version

Article:

Gautam, Meenu, Pandey, Divya and Agrawal, Madhoolika (2016) Phytoremediation of metals using lemongrass (Cymbopogon citratus (D.C.) Stapf.) grown under different levels of red mud in soil amended with biowastes. International journal of phytoremediation. pp. 555-562. ISSN 1522-6514

https://doi.org/10.1080/15226514.2016.1267701

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/



Phytoremediation of metals using lemongrass (Cymbopogon citratus (D.C.) Stapf.) grown under different levels of red mud in soil amended with bio-wastes

Journal:	International Journal of Phytoremediation			
Manuscript ID	BIJP-2016-0266.R2			
Manuscript Type:	Original Article			
Date Submitted by the Author:	06-Oct-2016			
Complete List of Authors:	Gautam, Meenu; Science, Botany Pandey, Divya ; Stockholm Environment Institute Agrawal, Madhoolika; Banaras Hindu University, Ecology Research Laboratory, Department of Botany			
Keywords:	Cymbopogon citratus, Phytoremediation, Red mud			

SCHOLARONE[™] Manuscripts



Phytoremediation of metals using lemongrass (Cymbopogon citratus (D.C.) Stapf.)
grown under different levels of red mud in soil amended with bio-wastes
Author(s): Meenu Gautam ¹ , Divya Pandey ² , Madhoolika Agrawal ³
¹ Meenu Gautam
Research Scholar
Laboratory of Air Pollution and Global Climate Change,
Department of Botany, Institute of Science,
Banaras Hindu University, Varanasi-221005, India
Email ID: meenu400@gmail.com
² Divya Pandey
AXA Junior Research (Post-doctoral) Fellow,
Stockholm Environment Institute at York,
Grimston House, University of York,
Heslington, York, YO10 5DD
Email ID: pandey.divyaa85@gmail.com
³ Corresponding author
Prof. Madhoolika Agrawal
Laboratory of Air Pollution and Global Climate Change,

Department of Botany, Institute of Science,

Banaras Hindu University, Varanasi-221005, India.

Phone no. 0091-542-2368156

Fax no.: 0091-542-2368174

Email ID: madhoo.agrawal@gmail.com

1 Abstract

Due to hostile condition of red mud (RM), its utilization for vegetation is restricted. Therefore, RM with bio-wastes as soil amendment may offer suitable combination to support plant growth with reduced risk of metal toxicity. To evaluate the effects of RM on soil properties, plant growth performance and metal accumulation in lemongrass, a study was conducted using different RM concentrations (0, 5, 10 and 15 % w/w) in soil amended with bio-wastes (cowdung manure (CD) or sewage-sludge (SS)). Application of RM in soil with bio-wastes improved organic matter and nutrient contents, and caused reduction in phytoavailable metal contents. Total plant biomass was increased under all treatments, maximally at 5 % RM in soil with SS (51.7 %) and CD (91.4 %) compared to control (no RM and bio-wastes). Lemongrass acted as a potential metal tolerant plant due to metal tolerance index >100 %. Based on translocation and bioconcentration factors, lemongrass acted as potential phytostabilizer of Fe, Mn and Cu in roots and was found efficient in translocation of Al, Zn, Cd, Pb, Cr, As and Ni from roots to shoot. The study suggests that 5 % RM with bio-wastes preferably SS may be used to enhance phytoremediation potential of lemongrass.

16 Key words: *Cymbopogon citratus*; phytoremediation; red mud

23 1 Introduction

Bauxite residue, commonly referred as red mud (RM) is generated in large quantities during Bayer's process of alumina extraction. It is highly alkaline and saline residue mainly composed of Fe₂O₃, Al₂O₃, CaO, Na₂O, TiO₂, SiO₂, P₂O₅ and VO₅⁻¹. Globally, \approx 150 million tons of RM is generated annually; management of such a quantity of waste is an increasing problem because utilization of only 2-3 % of generated residue has been estimated in a productive way². Nowadays, several applications of RM have been suggested viz. manufacturing of ceramics, building materials, pigments, paints, as adsorbent and catalyst, but challenge remains to find economically viable options to utilize significant amount of residue generated every year². A decade ago, slurry disposal was practiced in adjoining areas, nearby estuaries/lagoons, as filler at depleted mine- and quarry sites or stored in nearby dammed valley³. Improper disposal of poorly treated residue results in several environmental problems including contamination of surface and ground water through leaching, alteration in soil properties and plant community structure, and several other health issues related to human and wildlife³. For an instance, Aika spill in Hungary caused contamination of vast areas of agricultural land with RM that contained elevated levels of toxic metals with consequent impacts on plants ⁴. To date, dry stacking is the most popular choice of RM disposal with relatively lower risk of environmental contamination. However, study by Power et al. "see ref. 5" reported air pollution as major problem associated with its dry disposal practice.

To manage such waste dumps, different phytotechnologies have been developed, which are cost effective, and also offer sustainable and eco-friendly options ⁶. Phytoremediation of RM is, however, a challenging task due to its high alkalinity, salinity, elevated levels of potentially toxic metals, poor water retention and nutrient supplying capacities that limit establishment of plants on RM dumps ⁷. Therefore, prior to establishment of vegetation on such dumps, its unfavorable properties need to be improved to support plant growth coupled with enhanced phytoremediation potential. Studies have been conducted with RM in combination with bio-wastes (sewage-sludge, vegetative dry dust, animal manures, bacteria and mycorrhiza) as soil amendments, which improved soil properties and plant performance with low phytotoxic effects ^{7,8}. These studies have prompted us to assess the utilization of animal manure and sewage-sludge in combination with RM, which may boost the levels of dissolved organic carbon and nutrient availability, and lower phytoavilability of toxic metals. This strategy offers twin benefits in industrial as well as organic waste management ^{9,10}.

Cymbopogon citratus (D.C.) Stapf., commonly known as lemongrass, is a metal tolerant plant that withstand the harsh environmental conditions ¹¹. Israila *et al.* "see ref. 12," have identified lemongrass as a potential metal (Cd, Ni and Pb) accumulator grown on scrap-metal dumpsite at Dakace, Zaria-Nigeria, and suggested its suitability for phytoremediation of metal contaminated sites. Lemongrass cultivation is also widely practiced for stabilization of slopes and restoration of alkaline and saline soils ¹³.

The present study was conducted using lemongrass grown under varying red mud treatments in soil amended with cowdung manure or sewage-sludge (1) to assess the physico-chemical properties of soil under different soil treatments (2) to evaluate the phytoremediation potential of lemongrass, and (3) to assess the influence of metals on plant growth performance under varying soil treatments.

67 2 Materials and methods

The experiment was conducted in Botanical Garden, Banaras Hindu University $(25^{\circ}18' \text{ N } 82^{\circ})$ 01' E and 76.19 m above sea level) from February 03 to August 03, 2013. During the experiment, mean monthly maximum and minimum temperature were 41.4 and 12.5° C,

International Journal of Phytoremediation

respectively. Mean monthly maximum and minimum relative humidity were 89.4 and 29.3
%, respectively, and total rainfall during the period was 110.0 mm.

Red mud was obtained in the form of dry lumps (≈ 70 % solid cake) from the dumping yard, situated nearly 900 m from the premises of HINDALCO Industries Ltd., Renukoot, India. Lemongrass, known for its medicinal value was found growing naturally in the planted areas of RM dumps under plantation project called "Sanjeevani." Sewage-sludge (SS) and cowdung manure (CD) were collected from Dinapur municipal sewage treatment plant, Varanasi and dairy farm, B.H.U., respectively. Garden soil (C) was dug out upto 30 cm depth from Botanical Garden, B.H.U. Lemongrass was incurred from Faculty of Ayurveda, Institute of Medical Sciences, B.H.U. After removing stone and plant materials, RM, SS, CD and C were air dried, crushed, passed through sieve (2 mm mesh size) and mixed in a definite proportion to obtain varying soil treatments (Table S1-1). Prior to the present study, an experiment was conducted using lemongrass grown under varying RM concentrations (0, 10, 20, 30, 40 and 50 % w/w) in soil amended with CD or SS and maximum biomass was obtained under 10 % RM treatment. Therefore, we have selected 5, 10 and 15 % RM w/w with CD or SS for the present study.

Different soil treatments thus obtained were filled into cylindrical plastic pots (diameter, 25 cm; height, 50 cm). In total, there were 90 pots (9 treatments \times 10 replicates); each filled with 10 kg of different soil treatments. Pots were left at experimental site for 14 days for pathogen destruction, physico-chemical stabilization and proper conditioning of treated soil due to bio-waste amendments ¹⁴. On fifteenth day, one plant slip (shoot length: 15 cm, root length: 5 cm) of lemongrass was transplanted into each pot. Amount of water was standardized to avoid the leakage from the pots at different ages. Amounts of watering were 0.4, 0.6, 0.8 and 1.0 L between 0-40, 41-80, 81-120, 121-180 DAT, respectively given in each pot, every alternate day.

For physico-chemical analyses of different soil treatments, samples from three pots per treatment were taken out using soil corer (5 cm diameter and 10 cm depth) just before transplantation. Each sample was air dried, crushed and passed through sieve of 2 mm mesh size. The pH and EC of samples were measured in aqueous suspension of 1:5 (w/v) using pH meter (Model EA940, Orion, U.S.A) and conductivity meter (Model 303, Systronics, India), respectively. Total organic carbon (TOC) and total nitrogen (TN) contents were determined following Walkley and Black's rapid titration method "see ref. 15" and Gerhardt automatic N analyzer (Model KB8S, Germany), respectively. Available phosphorous (AP) was estimated by Olsen's method ¹⁶. Cation exchange capacity (CEC) through exchangeable cations extraction was determined following repeated leaching method ¹⁷. Phytoavailable metals in different soil treatments were extracted using 0.05 M EDTA solution ¹⁸. Exchangeable cations and EDTA extractable metal contents were determined using Atomic Absorption Spectrophotometer (Analyst-800, Perkin Elmer Inc., Norwalk, CT, USA).

Harvesting was done in triplicate by taking out entire plant along with roots and soil from the pots at 180 days after transplantation (DAT). Plants were gently jerked and after separating easily removable soil, roots were washed under running water to remove adhering soil particles. Thereafter, plant parts were washed twice using de-ionized water to avoid metal contamination. Afterwards, roots and shoot were separated and oven dried at 80° C until constant weights were attained. Dry weights of roots and shoot were measured for biomass determination.

Dried samples of roots, shoot and soil in triplicate were homogenized by grinding to fine powder using mortar and pestle. Contents of Al, Fe, Zn, Cu, Mn, Ni, Pb, Cr and Cd were determined on Atomic Absorption Spectrophotometer (AAS) (Analyst-800, Perkin Elmer Inc., Norwalk, CT, USA) after acid digestion (HNO₃ and HClO₄ in 9:4 ratio) following method of Gaidajis "see ref. 19". Moreover, AAS equipped with mercury hydride system

(MHS-15, Perkin Elmer Inc., Shelton, CT, USA) was used to determine As content in acid digested plant- and soil samples following the method by Welz and Sucmanová "see ref. 20". The choice of metals viz. Fe, Al, Ni, Cr, Cd, Ni, As and Pb was based on their high concentrations in RM and potentially phytotoxic effects ²¹. Micronutrients such as Zn, Mn and Cu showed low phytoavailabilities at circum-neutral to alkaline pH 22 .

Precision and accuracy of analysis was assured through repeated analysis of samples against National Institute of Standard and Technology, Standard Reference Material (NBS SRM-1570) for all metals. Blank and drift standards (Sisco Research Laboratories Pvt. Ltd., India) were run after every five sample runs to calibrate the instrument. Results were found within ± 2 % of the certified value. Coefficients of variation of replicate analysis were determined for different determinations and precision of analysis. Variations were found to be less than 10 %.

Metal tolerance ability of the plant was determined through metal tolerance index (MTI) "see ref. 23," whereas its phytoextraction potential was estimated using translocation (TF) and bioconcentration factors (BCF_{plant}), calculated using the formulae by Qihang et al. "see ref. 24". <u>ut</u>

$$MTI (\%) = \frac{Total \ plant \ biomass \ under \ treatment}{Total \ plant \ biomass \ under \ control}$$

$$TF = \frac{Metal \ contents \ in \ shoot \ tissue}{Metal \ contents \ in \ root \ tissue}$$

$$BCF_{plant} = rac{Metal \ contents \ in \ plant \ tissue}{Metal \ contents \ in \ soil}$$

Statistical significance of differences between physico-chemical properties of different soil treatments, total and phytoavailable metal contents, plant biomass, metal contents in plant, TF and BCF values for different treatments were tested by one way analysis of variance

URL: http://mc.manuscriptcentral.com/bijp Email: Jason.White@po.state.ct.us

(ANOVA) followed by Duncan's multiple range test as post hoc. Prior to conducting significance testing, normality and homoscedasticity of data were tested with Kolmogorov-Smirnov and Levene's test, respectively and distribution was found normal based on resulted p values above 0.05 in all cases. Pearson's correlation and linear regression analysis were performed for metal contents in soil and plants grown under different treatments. All the statistical tests were performed using SPSS software, IBM SPSS Statistics 20.0 (IBM, Armonk, NY, USA).

3 Results and discussion

3.1 Soil properties before transplantation

Due to high alkalinity and EC value of RM (Table S1-2), its increasing concentrations in bio-waste amended soil resulted in significant increases in their levels (Table 1). The pH of RM was above, whereas EC value was below threshold levels suggested as remediation target 25 . However, RM treated soils exhibited circum-neutral pH and were in the range of remediation target (pH, 5.5-9.0 and EC, < 4mS cm⁻¹)²⁵. High pH and EC may be ascribed to utilization of caustic soda during Bayer's process, resulting in an increase in soluble and free forms of caustic soda content in RM¹. Insignificant change in CEC values from CD_{RM0} to CD_{RM15} and SS_{RM0} to SS_{RM15} treatments may be due to its low value in RM compared to CD and SS. respectively (Table 1). In RM, more than 50 % of exchangeable cations are dominated by Na^+ , whereas in both CD and SS, the same is dominated by exchangeable Na^+ , Ca^{2+} and Mg^{2+} ions. Bio-wastes in combination with RM are known to ameliorate the low CEC value of RM thereby prompting exchange of other macronutrients (Ca^{2+} , Mg^{2+} and K^{+} ions), essential for better plant growth ²⁶.

162 Notably higher levels of TOC, TN and AP were observed in CD and SS compared to RM
163 (Table S1-2) and C (Table 1). Being low in TOC (0.63 %), TN (trace to 0.02 %) and AP

(0.09 %), RM is limited in major nutrients required for plant's growth and development ²⁷. Organic matter and humus are crucial for soil pedogenesis²⁸. Addition of CD and SS in soil prior to RM addition resulted in significant increases in the levels of TOC, TN and AP in CD_{RM0} and SS_{RM0} treatments, respectively compared to control (Table 1). However, increasing RM treatments in CD or SS amended soil caused gradual decline in TN and AP, whereas TOC showed insignificant change across treatments. However, their levels were significantly higher compared to control. Organic carbon in soil is of paramount importance in determining soil aggregate stability which consequently influences gaseous exchange, phytoavailable nutrients, water storage and transport ²⁹. Jones *et al.* "see ref. 9" found that addition of poultry manure to RM stimulated active microbial biomass, thus enhancing soil aggregate stability. RM was found deficient in TN content; therefore its increasing proportions in bio-waste amended soil may have caused reduction in its value across treatments. A decline in AP may be attributed to appreciable amount of sesquioxides (>50 %) in RM resulting in high P retention capacity due to formation of insoluble metal phosphates

Contents of Al, Fe, Pb, Cd, As, Cr and Ni were higher, whereas Mn, Zn and Cu contents were found low in RM compared to CD and SS (Table S1-2). Xue et al. "see ref. 1" reported nutrient deficiency as a potential limiting factor for vegetative growth on RM. When compared with soil quality guidelines defined by NOAA "see ref. 30", contents of Cd and Cr (in RM and SS), Cu (in RM, SS and CD) and Zn (in SS) were higher than their suggested values. Increasing RM concentrations in bio-waste amended soil increased metal contents, but trends were different under CD and SS amendments (Table 1). Trend of total metal contents was Fe>Al>Mn>Zn>Cu>Cr>Pb>Cd>Ni>As under CD_{RM0} to CD_{RM15} and Fe>Al>Zn>Mn>Cr>Cu>Cd>Pb>Ni>As under SS_{RM0} to SS_{RM15} treatments. Higher contents of Zn, Cr and Cd than Mn, Cu and Pb, respectively in later case may be attributed to many

folds higher Zn, Cr and Cd contents due to SS than CD amendment in soil following RM treatments. Contents of Cu, Zn, Cd and Cr under CD_{RM0} to CD_{RM15} treatments were within, whereas under SS_{RM0} to SS_{RM15} treatments were above the soil quality guidelines of NOAA ³⁰. Moreover, As, Pb and Ni contents in all soil treatments comply with NOAA soil quality guideline. Studied metal contents when compared with their screening levels in soil required for plants, Mn, Zn, Cu, Cd and Cr under all soil treatments, whereas Pb and Ni under SS_{RM5} to SS_{RM15} treatments exceeded their prescribed values by NOAA ³⁰.

Phytoavailable metal contents were maximum in SS followed by CD, C and RM (Table S1-2). Micronutrients such as Fe, Mn, Cu, Zn and Ni, were available in lower concentrations in RM due to formation of immobile metal complexes under alkaline condition and adsorption of metals on the surface of RM²⁷. In soil, CD and SS amendments caused significant increases in phytoavailable metals in CD_{RM0} and SS_{RM0} treatments, respectively compared to C. Furthermore, increasing RM concentrations reduced phytoavailable metal contents significantly compared to CD_{RM0} and SS_{RM0} , respectively (Table 1). Circum-neutral pH induced by RM treatments may favor precipitation of metals followed by increase in metal sorption by charged colloids of RM. Cancrinite and hematite are two principal phases of RM, which provide adsorption capacity to RM³¹. Phytoavailable metal contents under different soil treatments thus showed the trend in the order of Fe>Al>Mn>Zn>Cu>Cr>Ni>Cd>As>Pb. Low phytoavailable Mn is a major problem associated with oxidizing environment which favors oxidation of Mn²⁺ to insoluble Mn⁴⁺ in RM ³². Moreover, increase in Fe-oxide content due to RM addition might have played a significant role in reducing phytoavailable Zn, Ni, Pb and Cd contents across treatments ²². Relatively low phytoavailable Cu content may be ascribed to hydroxyl and carboxyl groups supplied by bio-waste amendments, which lead to formation of insoluble and immobile Cu-complexes, resulting in reduced risk of Cuphytotoxicity³³. However, higher phytoavailable Cu than Ni content may be because of

International Journal of Phytoremediation

increase in aqueous dissolved organic carbon content due to RM addition, which may have increased the mobility of Cu followed by Ni complexed to organic matter ²¹. Reduced phytoavailable Cd and Pb contents may be ascribed to their strong antagonistic relationship with Mn and Cu, respectively. Whereas, appreciable amount of Fe-sulfate in RM at circumneutral to alkaline pH is known to reduce labile As content efficiently in soil ³⁴. Thus, low phytoavailabilities of metals in aged bauxite residues are more pronounced than spiked one, which may be attributed to their complex nature and relative behavior of different metals ³.

3.2 Plant biomass

Root, shoot and total plant biomass were significantly higher under RM treatments compared to C (Fig. 1), indicating that they are efficient enough to tolerate high metal contents in soil and in their tissues ³⁵. Maximum increase in root and shoot biomass was observed under 5 % RM treatment followed by a decline under further treatments. Total plant biomass was significantly increased by 59.7 and 91.4 % under CD_{RM5} and SS_{RM5} treatments, respectively compared to control (Fig. 1). Reduction in plant biomass under 10 and 15 % RM treatments may be attributed to metal toxicity. In a similar study conducted with Festuca rubra, maximum increase in its biomass was found in soil treated with 5 % RM concentration, which was ascribed to decrease in metal contents in grass due to increased soil pH ³⁶. Decrease in biomass of plants growing in metal polluted soil is closely related to growth and development of roots, because roots are first organ exposed to elevated metal contents in soil. Due to metal toxicity, poorly developed roots may lead to decrease in nutrient transport and water uptake by plant, thereby affecting shoot and total plant biomass³⁷.

3.3 Metal contents in lemongrass

Among all metals, content of Fe was found maximum, whereas As content was foundminimum in plant biomass under different RM treatments in soil amended with CD or SS

(Table 2). Low As uptake by the plant under RM treatments may be due to its reduced phytoavailable contents ³⁴. Metal contents in lemongrass when compared to phytotoxic thresholds for medicinal plants showed that all the metals were within the prescribed threshold levels ³⁸. However, Cd content in plant under RM treatments in SS amended soil exceeded its threshold level (5-30 ppm)³⁸. Increase in Cd content in plant may be due to its higher level in SS compared to CD amendments in soil and low adsorption capacity of Cd (1.35 mmol g⁻¹) onto adsorbent surface of RM resulting in its increased mobility and phytoavailability³¹. It should be noteworthy that although Cd content in the plant was higher than its threshold level, no such severe phytotoxic effect of Cd on total plant biomass was reflected which may be due to synthesis of Cd induced phytochelatin complex that masks Cdphytotoxicity ³⁹. Moreover, Fe, Cu, Cr, Cd, Ni, Zn (except under C and CD_{RM0} to CD_{RM15}) treatments) and Pb (except under C and CD_{RM0} treatments) contents in plant exceeded the WHO safe limits (20, 10, 1.5, 0.3, 1.5, 50 and 10 mg kg⁻¹, respectively) for medicinal plants, however, Mn and As were within their safe limits (200 and 5 mg kg⁻¹, respectively) under all treatments ⁴⁰.

Pearson's correlation analysis showed that significant positive correlations exist between metal contents in plant biomass with their levels under different RM treatments in bio-waste amended soil (Table S1-3). Indeed, RM was found to be a predominant source of Al, Fe, Pb, Cd, Ni, Pb, Cr and As, whereas CD and SS acted as sources of Mn, Zn and Cu in different soil treatments (Table 1). Furthermore, linear regression models also confirmed that increasing concentrations of RM in soil amended with CD or SS significantly increased the metal contents in lemongrass (Figs. S1-1 and S1-2). Similar relationships were reported for lemongrass grown in soil treated with increasing contents of Cd, Hg and Pb³⁵. A weak magnitude of relationship for Mn ($R^2 = 0.657$) and As ($R^2 = 0.722$) in plant under different RM treatments in CD amended soil may be attributed to their antagonistic relationship with

Cd ^{14,41}. Moreover, moderate relation for Zn ($R^2=0.776$) and Ni ($R^2=0.757$) may be because of competitive behavior of Fe with Ni followed by Zn for binding sites in plant ⁴². This could also be attributed to relatively low phytoavailable Zn and Ni contents compared to Fe in treated soil.

3.4

3.4 Metal tolerance index of lemongrass

Assessing metal tolerance is of paramount importance while selecting plant for phytoremediation⁴³. To determine metal tolerant behavior of plant, one of the most common parameters used is metal tolerance index (MTI)⁴⁴. Based on total plant biomass, MTI of lemongrass under different soil treatments showed remarkable differences in tolerance to high metal contents compared to C. The MTI under CD_{RM0} CD_{RM5} CD_{RM10} CD_{RM15} SS_{RM0} SS_{RM5} SS_{RM10} and SS_{RM15} treatments were 129.7, 151.7, 126.7, 115.1, 150.1, 191.4, 167.4 and 120.7 %, respectively. Although MTI varied with increasing metal contents due to RM addition in bio-waste amended soil, its value was found >100 % under all treatments, thereby categorizing lemongrass as metal tolerant plant (Table S1-4), capable to grow in metal polluted soil. Metal tolerant plants are known to elicit their ability to tolerate metal induced reactive oxygen species by increasing their enzymatic and non-enzymatic antioxidants, proline accumulation, synthesis of phytochelatins and metallothioneins for detoxification and homeostasis ^{35,45}.

3.5 Translocation and bioconcentration factors for metals

The translocation factor (TF) determines effectiveness in metal movement from roots to shoot, whereas bioconcentration factor (BCF_{plant}) is used in determining the uptake and accumulation of metals from soil into plant biomass ⁴⁶. Plant with BCF_{plant} value >1, indicates its efficiency to uptake and accumulate metals from soil, while with BCF_{plant} value <1 is metal excluder ⁴⁷. Thus by comparing BCF_{plant} and TF values of metals under different soil

treatments, we can compare plant's ability in extracting metals from soil and then translocating them to easily harvestable part of the plant. Plant with both BCF_{plant} and TF values >1 can be used as a suitable candidate for hyper-accumulation of metals, while plants with TF <1 can phytostabilize metals in roots ⁴⁸.

No significant change in TF value for Cu was found under all treatments, whereas Al, Zn, Pb, Cd, Ni, Cr and As showed maximum TF values under CD_{RM0} and SS_{RM0} followed by a gradual decline under increasing RM treatments (Table S1-5). A decline in TF values under RM treatments may be attributed to decrease in available metals absorbed by the roots and their further transport to shoot. The TF values of Fe, Mn and Cu were found < 1 indicating restricted movement of metals from roots to shoot. However, TF values of Al, Zn, Pb, Cd, Ni, Cr and As were found ≥ 1 which illustrates the efficiency of plant in transport of metals from roots to shoot. Effectiveness of lemongrass in metal (Al, Zn, Cd, Pb, Ni, Cr and As) movement from roots to shoot is more likely due to an efficient metal transporter "see ref. 49" and possibly due to metal sequestration in leaf vacuole and apoplast "see ref. 50". TF values of Pb, Cd and Ni were higher, whereas that of Cu and Zn (except in C) were found lower than those reported in lemongrass grown on scrap metal dumpsite ¹². Thus based on TF values, lemongrass was found efficient in translocation of Al, Zn, Cd, Pb, Ni, Cr and As from roots to shoot and acted as a potential phytostabilizer of Fe, Mn and Cu in roots.

The BCF_{plant} values for Mn, Zn, Cu, Ni and Pb increased, whereas for Al, Fe, Cd, Cr and As decreased with increase in RM concentration in bio-waste amended soil, BCF_{plant} values for all metals were < 1 under all soil treatments (Table S1-5). BCF_{plant} values for Zn, Cu, Pb, Cd and Ni in the plant were found less than those reported in lemongrass collected from scrap metal dumpsite ¹². Low BCF_{plant} values for studied metals in the plant under different soil treatments may be attributed to decrease in their phytoavailable contents with increase in soil pH and metal adsorption capacity of RM¹. In the present study, based on BCF_{plant} values,
lemongrass acted as a potential metal excluder.

313 4 Conclusions

The study showed that red mud in combination with sewage-sludge in soil led to more significant improvement in organic matter and nutrient contents compared to cowdung amended soil. Increasing red mud concentrations in bio-waste amended soil increased total metal contents with simultaneous reduction in their phytoavailable contents. Studied metals except Cd in lemongrass were found within the phytotoxic thresholds for medicinal plants. All metals except Mn exceeded the WHO safe limit for consumption in medicinal plants. A significant improvement in plant biomass was observed under red mud treatments compared to control soil with maximum increase under 5 % RM treatment. Lemongrass was found to be a potential metal tolerant plant as metal tolerance index was more than 100 %. Based on translocation factor, lemongrass acted as a potential phytostabilizer of Fe, Mn and Cu in roots, whereas it was found efficient in translocation of Al, Zn, Cd, Pb, Ni, Cr and As from roots to shoot. Moreover, lemongrass was identified as a potential metal excluder due to bioconcentration factors <1 for studied metals. The study suggests that 5 % RM in combination with bio-wastes; preferably SS may be used as a soil quality enhancer by reducing metal toxicity and also make it suitable for lemongrass cultivation coupled with enhanced phytoremediation efficiency. However, a much more detailed investigation over the likely environmental impacts of this strategy under field condition is needed.

331 6 Acknowledgements

Authors are thankful to the Head, Department of Botany, Institute of Science, B.H.U.,
Varanasi and HINDALCO Industries Ltd., Renukoot, U.P. for providing all the necessary
facilities during the research work. We are thankful to AXA Junior Research (Post-doctoral)

International Journal of Phytoremediation

33	5 Fellowship, York, United Kingdom and Council of Scientific and Industrial Research (CSIR),
33	6 New Delhi, India for financial assistance in the form of Research Associateship (Divya
33	Pandey) and Senior Research Fellowship (Meenu Gautam), respectively. We are also grateful
33	to anonymous reviewers for their critical comments and helpful suggestions which helped us
33	9 in improving the manuscript.
34	o a la construcción de l
34	1
34	2
34	3
34	4
34	5
34	5
34	7
34	8
34	9
35	D
35	1
35	2
35	3
35	4
	16

2 3 4	355	References
5 6 7	356	¹ Xue S, Zhu F, Kong X, Wu C, Huang L, Huang N, Hartley W. A review of the
8	357	characterization and revegetation of bauxite residues (Red mud). Environ Sci Pollut
10 11 12	358	Res. 2016; 23:1120-1132.
13 14	359	² Evans K. The History, Challenges, and New Developments in the Management and Use of
15 16 17	360	Bauxite Residue. J Sustain Metall. 2016;1-6.
18 19	361	³ Mayes WM, Burke IT, Gomes HI, Anton AD, Molnár M, Feigl V, Ujaczki E. Advances in
20 21 22	362	Understanding Environmental Risks of Red Mud After the Ajka Spill, Hungary. J
23 24 25	363	Sustain Metall. 2016;13:1-2.
26 27	364	⁴ Ruyters S, Mertens J, Vassilieva E, Dehandschutter B, Poffijn A, Smolders E. The red mud
28 29	365	accident in Ajka (Hungary): plant toxicity and trace metal bioavailability in red mud
30 31 32	366	contaminated soil. Environ Sci Technol. 2011;45:1616-1622.
33 34 25	367	⁵ Power G, Gräfe M, Klauber C. Bauxite residue issues: I. Current management, disposal and
36 37 38	368	storage practices. Hydrometallurgy. 2011;108:33-45.
39 40	369	⁶ Marques APGC, Rangel AOSS Castro PML. Remediation of heavy metal contaminated
41 42	370	soils: phytoremediation as potentially promising clean-up technology. Crit Rev
43 44 45	371	Environ Sci Technol. 2009; 39:622–654.
46 47	372	⁷ Santini TC, Kerr JL, Warren LA. Microbially-driven strategies for bioremediation of bauxite
48 49 50	373	residue. J Hazard Mater. 2015;293:131-57.
51 52 52	374	⁸ Ujaczki É, Feigl V, Molnár M, Vaszita E, Uzinger N, Erdélyi A, Gruiz K. The potential
54 55	375	application of red mud and soil mixture as additive to the surface layer of landfill cover
56 57 58 59	376	system. J Environ Sci. 2016;44:189-96.

3		
4		
5		
6		
7		
/ ^		
8		
9		
1	0	
1	1	
1	ò	
4	ົ	
1	3	
1	4	
1	5	
1	6	
1	7	
1	8	
÷	ă	
י ה	5	
2	J	
2	1	
2	2	
2	3	
2	4	
2	5	
	с Р	
~	2	
2	1	
2	8	
2	9	
3	0	
3	1	
ົ້	່ວ	
0	~	
3	3	
3	4	
3	5	
3	6	
3	7	
2 2	'n	
ົງ	ი ი	
ں م	9	
4	Ů	
4	1	
4	2	
4	3	
4	4	
۵	5	
7	c	
4	0	
4	1	
4	8	
4	9	
5	0	
5	1	
5	י ס	
0 5	~	
о -	3	
5	4	
5	5	
5	6	
5	7	
5	'n	
0 F	0 0	
с С	9	
6	()	

⁹Jones BEH, Haynes RJ, Phillips IR. Influence of organic waste and residue mud additions on
chemical, physical and microbial properties of bauxite residue sand. Environ Sci Pollut
Res. 2011;18:199-211.

¹⁰Courtney R, Timpson J. Reclamation of fine fraction bauxite processing residue (red mud)
 amended with coarse fraction residue and gypsum. Water Air Soil Pollut. 2005;164:91–
 102.

¹¹Das M, Maiti SK. Growth of *Cymbopogon citratus* and *Vetiveria zizanioides* on Cu mine
 tailings amended with chicken manure and manure-soil mixtures: A pot scale study. Int
 J Phytorem. 2009;11:651–663.

- ¹²Israila YZ, Bola AE, Emmanuel GC, Ola IS. The effect of application of EDTA on the
 phytoextraction of heavy metals by *Vetivera zizanioides*, *Cymbopogon citrates* and
 Helianthus annus. Int J Environ Monit Anal. 2015;3:38-43.
- ¹³Bienes R, Jimenez-Ballesta R, Ruiz-Colmenero M, Arevalo D, Alvarez A, Marques MJ.
 Incidence of the growing of aromatics and medicinal plants on the erosion of
 agricultural soils. Span J Rural Develop. 2010;1:19-28.
- ¹⁴Singh RP, Agrawal M. Variations in heavy metal accumulation, growth and yield of rice
 plants grown at different sewage sludge amendment rates. Ecotoxicol Environ Saf.
 2010;73:632-41.
- ¹⁵Allison FE. Soil organic matter and its role in crop production. Amsterdam (Netherlands):
 Elsevier Scientific Publishing Company; 1973.
- ¹⁶Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of Available Phosphorus in Soils
 by Extraction with Sodium Bicarbonate. US: Circular No. 939, Department of
 Agriculture. 1954.p.19.

¹⁷Hesse PR. A textbook of soil chemical analysis. In: Murray J, editor. A textbook of soil
chemical analysis. London; 1971.p.35-88.

¹⁸Quevauviller P, Rauret G, Rubio R, López-Sánchez JF, Ure A, Bacon J, Muntau H.
Certified reference materials for the quality control of EDTA-and acetic acidextractable contents of trace elements in sewage-sludge amended soils (CRMs 483 and
405 484). Fresen J Anal Chem. 1997;357:611-618.

¹⁹Gaidajis G. Ambient concentrations of total suspended particulate matter and its elemental
constituents at the wider area of the mining facilities of TVX Hellas in Chalkidiki,
Greece. J Environ Sci Heal A. 2003;38:2509-2520.

²⁰Welz B, Šucmanová M. L-Cysteine as reducing and releasing agent for determination of
 antimony and arsenic using flow injection hydride generation atomic absorption
 spectrometry—Part 1. Optimization of analytical parameters. Analyst. 1993;118:1417 23.

²¹Lockwood CL, Stewart DI, Mortimer RJ, Mayes WM, Jarvis AP, Gruiz K, Burke IT.
Leaching of copper and nickel in soil-water systems contaminated by bauxite residue
(red mud) from Ajka, Hungary: importance of soil organic matter. Environ Sci Pollut
Res. 2015;22:10800-10.

²²Jones BE, Haynes RJ, Phillips IR. Effect of amendment of bauxite processing sand with
 organic materials on its chemical, physical and microbial properties. J Environ Manag.
 2010;91:2281-8.

²³Wang S, Shi X, Sun H, Chen Y, Pan H, Yang X, Rafiq T. Variations in metal tolerance and
accumulation in three hydroponically cultivated varieties of *Salix integra* treated with
lead. PloS one. 2014; 9:108-568.

423	²⁴ Qihang W, Wang S, Thangavel P, Qingfei L, Zheng H, Jun B, Qui R. Phytostabilization of
424	Jatropha curcas L. in polymetalic acid mine tailings. Int J Phytorem. 2011;13: 788-
425	804.
426	²⁵ Gräfe M, Klauber C. Bauxite residue issues: IV. Old obstacles and new pathways for in situ
427	residue bioremediation. Hydrometallurgy. 2011;108:46-59.
428	²⁶ Tipping E, Smith EJ, Lawlor AJ, Hughes S, Stevens PA. Predicting the release of metals
429	from ombrotrophic peat due to drought-induced acidification. Environ Pollut
430	2003;123:239-53.
431	²⁷ Lacatusu R, Kiselev A, Stroe VM, Rizea N, Lungu M, Lazar R, Stanciuburileanu MM,
432	Calciu I, Popa RG, Filipescu L. Plant Growth Suitable Nutritive Red Mud Composite
433	Materials from Romanian Dry Landfilled Red Mud II. Formulation nutritive composite
434	materials and plant growth tests at laboratory and glasshouse scale. Revista De Chimie.
435	2014;65:1294-1305.
436	²⁸ Filcheva E, Noustorova M, Gentcheva-Kostadinova SV, Haigh MJ. Organic accumulation
437	and microbial action in surface coal-mine spoils, Pernik, Bulgaria. Ecol Eng.
438	2000;15:1-5.
439	²⁹ Zhu F, Xue S, Hartley W, Huang L, Wu C, Li X. Novel predictors of soil genesis following
440	natural weathering processes of bauxite residues. Environ Sci Pollut Res.
441	2016;23:2856-2863.
442	³⁰ Buchmann MF. NOAA Screening Quick Reference Tables, NOAA OR&R Report 08-1.
443	Seattle (WA): Office of Response and Restoration Division, National Oceanic and
444	Atmospheric Administration; 2008-34.
	20

URL: http://mc.manuscriptcentral.com/bijp Email: Jason.White@po.state.ct.us

445	³¹ Santona L, Castaldi P, Melis P. Evaluation of interaction mechanisms between red muds
446	and heavy metals. J Hazard Mater. 2006;136: 324-329.
447	³² Gherardi MJ, Rengel Z. Bauxite residue sand has capacity to rapidly decrease availability of
448	added manganese. Plant and Soil. 2001;234:143-51.
449	³³ Guan TX, He HB, Zhang XD, Bai Z. Cu fractions, mobility and bioavailability in soil-
450	wheat system after Cu-enriched livestock manure applications. Chemosphere.
451	2011;82:215–222.
452	³⁴ Hartley W, Edwards R, Lepp NW. Arsenic and heavy metal mobility in iron oxide-amended
453	contaminated soils as evaluated by short-and long-term leaching tests. Enviro Pollut.
454	2004;131:495-504.
455	³⁵ Handique GK, Handique AK. Proline accumulation in lemongrass (<i>Cymbopogon flexuosus</i>
456	Stapf.) due to heavy metal stress. J Environ Biol. 2009;30:299-302.
457	³⁶ Gray CW, Dunham SJ, Dennis PG, Zhao FJ, McGrath SP. Field evaluation of in situ
458	remediation of heavy metal contaminated soil using lime and red-mud. Environ Pollut.
459	2006;142:530-539.
460	³⁷ Boonyapookana B, Parkplan P, Techapinyawat S, DeLaune RD, Jugsujinda A.
461	Phytoaccumulation of lead by sunflower (Helianthus annuus), tobacco (Nicotiana
462	tabacum), and vetiver (Vetiveria zizanioides). J Environ Sci Health Part A-
463	Toxic/Hazard Subst Environ Engin. 2005;40:117-137.
464	³⁸ Alloway BJ, Ayres DC. Chemical Principles of Environmental Pollution. London (UK):
465	Chapmann and Hall; 1997.

3
4
5
6
7
8
ğ
10
10
12
13
14
15
16
17
18
10
20
20
21
22
23
24
25
26
27
28
20
29
30
31
32
33
34
35
36
37
20
30
39
40
41
42
43
44
45
46
17
-+1 10
40
49
50
51
52
53
54
55
56
50
57
58
59
60

466	³⁹ Domínguez-Solís JR, López-Martín MC, Ager FJ, Ynsa MD, Romero LC, Gotor C.
467	Increased cysteine availability is essential for cadmium tolerance and accumulation in
468	Arabidopsis thaliana. Plant Biotech J. 2004;2:469-476.
469	⁴⁰ World Health Organization. Quality Control Methods for Medicinal Plant Materials.
470	Geneva, Switzerland. 2005.
471	⁴¹ Sun YB, Zhou QX, Liu WT, An J, Xu ZQ, Wang L. Joint effects of arsenic and cadmium
472	on plant growth and metal bioaccumulation: potential Cd-hyperaccumulator and As-
473	excluder Bidens pilosa L. J Hazard Mater. 2009;165:1023-1028.
474	⁴² Wood BW. Iron-induced nickel deficiency in pecan. Hort Sci. 2013;48:1145-1153.
475	⁴³ Zacchini M, Pietrini F, Mugnozza GS, Iori V, Pietrosanti L, Massacci A. Metal tolerance,
476	accumulation and translocation in poplar and willow clones treated with cadmium in
477	hydroponics. Water Air Soil Pollut. 2009;197: 23-34.
478	⁴⁴ Köhl, Lösch R. Experimental characterization of heavy metal tolerance in plants. In: Prasad
479	MNV, Hagemeyer J, editors. Heavy metal stress in plants. Berlin Heidelberg: Springer;
480	1999.p.371-389.
481	⁴⁵ Emamverdian A, Ding Y, Mokhberdoran F, Xie Y. Heavy metal stress and some
482	mechanisms of plant defense response. The Scientific World Journal. 2015.p.26.

⁴⁶Deng H, Ye ZH, Wong MH. Accumulation of lead, zinc, copper and cadmium by 12
wetland plant species thriving in metal-contaminated sites in China. Environ Pollut.
2004;132:29-40.

486	⁴⁷ Yanqun Z, Yuan L, Jianjun C, Haiyan C, Li Q, Schvartz C. Hyperaccumulation of Pb, Zn
487	and Cd in herbaceous grown on lead-zinc mining area in Yunnan, China. Environ Int.
488	2005; 31:755-762.
489	⁴⁸ Yoon J, Cao X, Zhou Q, Ma LQ. Accumulation of Pb, Cu, and Zn in native plants growing
490	on contaminated Florida site. Sci Total. 2006;368: 456-46.
491	⁴⁹ Baker AJM, McGrath SP, Sidoli CMD, Reeves RD. The possibility of in situ heavy metal
492	decontamination of polluted soils using crops of metal-accumulating plants. Resour
493	Conserv Recy. 1994;11:41-49.
494	⁵⁰ Clemens S, Palmgren MG, Krämer U. A long way ahead: understanding and engineering
495	plant metal accumulation. Trends in Plant Sci. 2002;7:309-315.



Fig. 1 Root, shoot and total plant biomass of lemongrass grown under different RM treatments in soil amended with CD or SS. Values are mean \pm SE. Different letters indicate significant differences at p<0.05 according to Duncan's test. RM: red mud; C: control; CD_{RM0}: soil with CD; CD_{RM5}: 5 % RM in CD_{RM0}; CD_{RM10}: 10 % RM in CD_{RM0}; CD_{RM15}: 15 % RM in CD_{RM0}; SS_{RM0}: soil with SS; SS_{RM5}: 5 % RM in SS_{RM0}; SS_{RM10}:10 % RM in SS_{RM0} and SS_{RM15}:15 % RM in SS_{RM0}.

Table 1 Selected physico-chemical properties, total and phytoavailable metal contents in control and RM treatments in soil amended with CD or SS
$(Mean \pm SE)$

Parameters	С	CD _{RM0}	CD _{RM5}	CD _{RM10}	CD _{RM15}	SS _{RM0}	SS _{RM5}	SS _{RM10}	SS _{RM15}
рН	7.87 ± 0.05^{cd}	7.33±0.01 ^f	7.82 ± 0.02^{d}	8.09 ± 0.02^{b}	8.27 ± 0.01^{a}	7.71 ± 0.04^{e}	7.82 ± 0.01^{d}	7.94±0.01°	8.21±0.04 ^a
$EC (mS cm^{-1})$	0.18 ± 0.01^{i}	0.50 ± 0.00^{e}	$0.54{\pm}0.00^{d}$	$0.56 \pm 0.00^{\circ}$	0.86 ± 0.01^{b}	$0.32{\pm}0.00^{h}$	$0.43{\pm}0.00^{g}$	$0.47{\pm}0.00^{\rm f}$	1.15 ± 0.00^{a}
TOC (%)	0.56 ± 0.10^{d}	$4.19 \pm 0.06^{\circ}$	4.23±0.28 ^c	4.56 ± 0.07^{bc}	4.40 ± 0.10^{bc}	4.77±0.21 ^{ab}	5.15 ± 0.06^{a}	5.18±0.11 ^a	5.09 ± 0.06^{a}
TN (%)	0.12 ± 0.01^{f}	$0.54{\pm}0.02^{a}$	$0.48{\pm}0.02^{abc}$	$0.30{\pm}0.01^{d}$	0.21 ± 0.01^{e}	$0.53{\pm}0.001^{ab}$	$0.50{\pm}0.03^{abc}$	0.46 ± 0.03^{bc}	$0.45 \pm 0.04^{\circ}$
AP (mg kg ⁻¹)	31.83 ± 1.64^{g}	443.04±37.98 ^{cd}	416.49±14.97 ^{de}	393.78±6.31 ^{ef}	377.26 ± 12.17^{f}	566.64±18.93 ^a	520.32 ± 5.04^{ab}	485.22±4.54 ^{bc}	463.39±5.74 ^{cd}
CEC (meq100 g^{-1})	4.16 ± 0.11^{bc}	$3.76 \pm 0.06^{\circ}$	$3.78\pm0.03^{\circ}$	3.99±0.37°	4.14 ± 0.02^{bc}	4.50 ± 0.06^{ab}	4.57 ± 0.05^{ab}	4.65±0.03 ^a	4.71 ± 0.02^{a}
Total metal conten	nts (mg kg ⁻¹)								
Al*	$0.08{\pm}0.00^{g}$	0.37±0.01 ^g	8.58 ± 0.21^{e}	19.05±0.22°	24.31 ± 0.19^{b}	3.14 ± 0.24^{f}	16.62 ± 0.13^{d}	24.70 ± 0.20^{b}	32.95±0.23 ^a
Fe*	2.57 ± 0.03^{h}	3.38 ± 0.22^{h}	16.26±0.11 ^g	31.84±0.48 ^e	$48.65 \pm 0.56^{\circ}$	21.81 ± 0.23^{f}	35.80 ± 1.12^{d}	51.91±1.06 ^b	67.39±1.38 ^a
Mn	260.67±1.76 ^e	285.43 ± 1.57^{d}	286.68±0.89 ^d	291.42 ± 4.57^{d}	$303.42 \pm 2.02^{\circ}$	308.39±2.19 ^{bc}	310.27±4.53 ^{abc}	314.97 ± 1.49^{ab}	318.16±1.35 ^a
Zn	134.49±7.49 ^e	146.07 ± 2.16^{e}	147.54±3.21 ^{de}	163.14 ± 2.60^{cd}	$170.61 \pm 4.75^{\circ}$	540.94 ± 7.01^{b}	536.03 ± 5.28^{b}	550.94±7.01 ^{ab}	560.32±5.31 ^a
Cu	70.55 ± 0.79^{f}	76.20±1.38 ^e	79.92±2.08 ^e	88.57±1.93 ^d	116.28±2.07 ^c	177.57±2.27 ^b	179.63±2.55 ^{ab}	181.79±1.30 ^{ab}	184.24 ± 0.62^{a}
Pb	18.27 ± 0.48^{i}	18.79 ± 0.72^{h}	25.53±0.46 ^g	43.26±0.75 ^e	$45.94{\pm}0.43^{d}$	36.49 ± 0.23^{f}	49.03±0.08°	61.80±0.11 ^b	73.31±0.21 ^a
Cd	1.46 ± 0.05^{h}	$2.14{\pm}0.02^{h}$	12.75±0.06 ^g	33.50±0.11 ^f	40.25±0.17 ^e	50.98 ± 0.83^{d}	73.30±1.65°	93.08 ± 2.89^{b}	116.58±3.51 ^a
Ni	11.69 ± 0.81^{f}	12.48 ± 0.34^{f}	13.22 ± 0.25^{f}	16.79±0.38 ^e	18.78±0.75 ^e	26.55 ± 0.25^{d}	35.27±1.24 ^c	39.68 ± 1.04^{b}	49.75±0.27 ^a
Cr	16.94±0.75 ^g	17.97±0.39 ^g	68.31 ± 0.96^{f}	84.07 ± 1.42^{ef}	112.12 ± 3.79^{e}	310.05 ± 5.03^{d}	351.06±3.89 ^c	410.96 ± 0.43^{b}	476.42±37.34 ^a
As	$0.13{\pm}0.01^{g}$	$0.59{\pm}0.01^{fg}$	0.99 ± 0.06^{ef}	1.37±0.06 ^e	1.88 ± 0.06^{d}	3.47±0.24 ^c	$3.80{\pm}0.20^{\circ}$	4.80 ± 0.35^{b}	5.73±0.07 ^a
Phytoavailable me	etal contents (n	ng kg ⁻¹)							
Al	57.86±0.79 ^c	58.21±0.83°	47.00 ± 2.52^{e}	42.33 ± 1.45^{f}	36.00±1.00 ^g	78.51±1.87 ^a	63.67 ± 0.88^{b}	57.89±0.48°	52.36 ± 1.06^{d}
Fe	86.43 ± 0.86^{cd}	103.30±3.43 ^b	82.16 ± 0.90^{de}	72.44±1.27 ^{ef}	66.67 ± 0.33^{f}	135.96±3.59 ^a	111.67±4.41 ^b	92.88±1.28 ^c	81.98±0.56 ^{de}
Mn	35.71±1.17 ^b	35.69±0.24 ^b	32.70±0.36 ^{cd}	31.34 ± 0.47^{de}	28.33 ± 0.88^{f}	41.69±0.24 ^a	37.44 ± 0.08^{b}	34.67±1.02 ^{bc}	29.63±1.93 ^{ef}
Zn	5.07 ± 0.20^{g}	11.19 ± 0.01^{d}	9.98±0.11 ^e	$8.44{\pm}0.03^{f}$	$8.22{\pm}0.40^{ m f}$	18.47 ± 0.21^{a}	16.42 ± 0.24^{b}	15.90 ± 0.34^{b}	14.56±0.73°
Cu	4.24 ± 0.20^{ef}	6.97 ± 0.12^{d}	4.61 ± 0.29^{e}	$3.64{\pm}0.25^{\rm f}$	$2.77 {\pm} 0.08^{g}$	12.65±0.03 ^a	10.67 ± 0.15^{b}	9.80±0.45°	9.49±0.26°
Pb	$0.03{\pm}0.00^{\rm f}$	$0.04{\pm}0.00^{bc}$	$0.03{\pm}0.00^{de}$	$0.03{\pm}0.00^{ef}$	$0.03{\pm}0.00^{g}$	$0.05{\pm}0.00^{a}$	0.04 ± 0.00^{b}	$0.04{\pm}0.00^{bc}$	$0.03{\pm}0.00^{cd}$
Cd	0.27 ± 0.01^{e}	$0.35{\pm}0.00^{b}$	$0.32{\pm}0.00^{d}$	0.27 ± 0.00^{e}	$0.25{\pm}0.00^{\mathrm{f}}$	$0.47{\pm}0.00^{a}$	0.35 ± 0.00^{b}	$0.33 \pm 0.00^{\circ}$	$0.32{\pm}0.00^{d}$
Ni	2.14 ± 0.06^{de}	$2.34{\pm}0.02^{\circ}$	2.16 ± 0.01^{de}	2.07±0.01 ^e	$1.83{\pm}0.04^{\rm f}$	$2.84{\pm}0.01^{a}$	2.43 ± 0.01^{b}	2.23 ± 0.01^{d}	2.17 ± 0.02^{d}
Cr	4.27 ± 0.13^{d}	2.68 ± 0.24^{e}	2.41 ± 0.15^{ef}	$1.89{\pm}0.15^{\rm f}$	$1.07{\pm}0.05^{g}$	6.97 ± 0.15^{a}	6.57 ± 0.28^{ab}	5.96 ± 0.06^{b}	5.23±0.03°
As	$0.01{\pm}0.001^{\rm f}$	$0.08 \pm 0.004^{\circ}$	0.07 ± 0.002^{cd}	0.06 ± 0.004^{d}	0.03 ± 0.001^{e}	0.13 ± 0.01^{a}	$0.10{\pm}0.001^{b}$	$0.07 \pm 0.003^{\circ}$	$0.04{\pm}0.002^{e}$

*g kg⁻¹; EC: electrical conductivity; TOC: total organic carbon; TN: total nitrogen; AP: available phosphorous; CEC: cation exchange capacity; RM: red mud; CD: cowdung manure; SS: sewage sludge; C: control; CD_{RM0}: soil with CD; CD_{RM0}; S $^{\circ}$ RM in CD_{RM0}; CD_{RM10}: 10 % RM in CD_{RM0}; CD_{RM10}: 15 % RM in CD_{RM0}; SS_{RM0}: soil with SS; SS_{RM5}: 5% RM in SS_{RM0}; SS_{RM10}: 10 % RM in SS_{RM0} and SS_{RM10}: 15 % RM in SS_{RM0}. Numbers with different letters in same row differ significantly at p < 0.05 as per the Duncan's test.

Table 2 Metal contents (mg kg⁻¹) in lemongrass grown under control and RM treatments in soil amended with CD or SS (Mean \pm SE).

Metals	С	CD _{RM0}	CD _{RM5}	CD _{RM10}	CD _{RM15}	SS _{RM0}	SS _{RM5}	SS _{RM10}	SS _{RM15}
Al	0.01 ± 0.00^{i}	0.19 ± 0.00^{h}	1.66 ± 0.02^{f}	2.83 ± 0.05^{d}	3.55±0.06 ^b	0.45 ± 0.02^{g}	2.37 ± 0.02^{e}	3.27±0.05 ^c	3.85±0.13 ^a
Fe	$0.63{\pm}0.02^{\rm f}$	0.65 ± 0.01^{f}	2.54 ± 0.07^{e}	$3.18 \pm 0.01^{\circ}$	$3.58 {\pm} 0.03^{b}$	2.76 ± 0.03^{d}	$3.12 \pm 0.10^{\circ}$	3.49 ± 0.06^{b}	4.06 ± 0.11^{a}
Mn	52.03±2.46 ^e	51.23±2.77 ^e	62.54 ± 2.95^{d}	$75.28 \pm 0.26^{\circ}$	84.64 ± 1.77^{b}	74.58±0.96°	88.87 ± 2.36^{b}	97.54±3.11 ^a	102.39±0.23 ^a
Zn	25.35±1.30 ^g	29.03 ± 0.67^{fg}	29.59 ± 1.88^{f}	31.45 ± 0.67^{f}	37.70±0.83 ^e	139.87 ± 1.79^{d}	143.67±1.54°	154.58±0.91 ^b	161.09 ± 1.06^{a}
Cu	41.84 ± 2.77^{f}	45.21±1.49 ^{ef}	$47.14 \pm 0.66^{\text{ef}}$	51.40±2.01 ^e	58.87 ± 0.21^{d}	$69.60 \pm 0.50^{\circ}$	77.85 ± 3.96^{b}	81.68 ± 1.25^{ab}	85.98 ± 2.98^{a}
Pb	$2.16\pm0.01^{\rm h}$	6.00±0.38 ^g	10.18 ± 0.18^{f}	21.96 ± 0.27^{d}	25.27±0.43°	16.52 ± 1.47^{e}	$29.69 \pm 0.10^{\circ}$	43.35±1.87 ^b	52.47 ± 1.22^{a}
Cd	$1.32{\pm}0.04^{g}$	$1.89{\pm}0.03^{g}$	10.33 ± 0.52^{f}	23.93±0.74 ^e	26.99±1.1 ^e	38.16 ± 0.54^{d}	$53.82 \pm 3.32^{\circ}$	68.12 ± 2.45^{b}	79.43 ± 0.48^{a}
Ni	3.78 ± 0.02^{g}	$5.95{\pm}0.09^{f}$	6.44±0.07 ^f	7.69±0.20 ^e	8.18 ± 0.16^{de}	8.69 ± 0.17^{d}	13.31±0.48°	17.59 ± 0.05^{b}	22.56 ± 0.06^{a}
Cr	$8.38{\pm}0.08^{\rm f}$	9.34±0.03 ^e	9.68±0.29 ^e	10.64 ± 0.19^{d}	10.74 ± 0.11^{d}	$37.07 \pm 0.02^{\circ}$	38.82 ± 0.61^{b}	$40.32{\pm}0.02^{a}$	41.03 ± 0.16^{a}
As	$0.10{\pm}0.01^{\rm f}$	0.74 ± 0.04^{e}	0.86 ± 0.03^{de}	0.94 ± 0.03^{d}	$0.98{\pm}0.02^{d}$	2.15±0.01°	2.49 ± 0.11^{b}	2.53±0.01 ^b	2.72 ± 0.01^{a}

RM: red mud; CD: cowdung manure; SS: sewage-sludge; C: control; CD_{RM0}: soil with CD; CD_{RM5}: 5 % RM in CD_{RM0}; CD_{RM1}: 10 % RM in CD_{RM0}; CD_{RM15}: 15 % RM in CD_{RM0}; SS_{RM0}: soil with SS; SS_{RM5}: 5% RM in SS_{RM0}; SS_{RM10}: 10 % RM in SS_{RM0} and SS_{RM1}: 15 % RM in SS_{RM0}. Numbers with different letters in same row differ significantly at p < 0.05 as per the Duncan's test.

1	
2	
3	
4 5	
6	
7	
8	
9	
10	
11	
12	
14	
15	
16	
17	
18	
19 20	
21	
22	
23	
24	
25	
26 27	
28	
29	
30	
31	
32	
33	
35	
36	
37	
38	
39	
40 41	
41 42	
43	
44	
45	
46	
47	
48 ⊿0	
49 50	
51	
52	
53	
54	
55	

Table S1-1 Details of different treatments

C CD _{RM0}	
CD _{RM0}	Unamended soil (Control)
CD	Unamended soil and cowdung manure (2:1 w/w)
CD_{RM5}	5% red mud + 95% CD _{RM0}
CD _{RM10}	10% red mud + 90 % CD _{RM0}
CD _{RM15}	15% red mud + 85% CD _{RM0}
SS_{RM0}	Unamended soil and sewage-sludge (2:1 w/w)
SS _{RM5}	5% red mud + 95% SS _{RM0}
SS _{RM10}	10% red mud + 90 % SS _{RM0}
SS _{RM15}	15% red mud + $85%$ SS _{RM0}

2
5
3
4
5
ĉ
ю
7
8
õ
9
10
11
10
12
13
14
15
15
16
17
10
10
19
20
21
22
23
24
24
25
26
07
21
28
29
20
30
31
32
20
33
34
35
00
30
37
38
00
39
40
41
40
42
43
44
15
45
46
47
10
40
49
50
51 51
51
52
53
51
54
55
56
57
57
58
59
60
1 11 1

Table S1-2 Physico-chemical properties, total and phytoavailable metal contents in red mud,

 sewage-sludge and cowdung manure

Physicochemical properties	Red mud	Sewage-sludge	Cowdung manure
pH	9.21±0.01	7.72±0.02	7.77±0.01
$EC (mS cm^{-1})$	2.96 ± 0.02	1.95 ± 0.01	0.943±0.01
TOC (%)	0.30 ± 0.24	6.58±0.02	5.44±0.02
TN (%)	nd	1.05 ± 0.03	1.09±0.11
AP (mg kg ⁻¹)	6.46 ± 0.02	638.32 ± 8.22	572.24±3.08
CEC (meq 100 ⁻¹ g)	4.39±0.01	6.10±0.06	5.49±0.06
Total metals (mg kg ⁻¹)			
Al*	187.7±4.79	2.42 ± 0.04	0.01 ± 0.00
Fe*	335.60±13.84	22.85 ± 0.84	0.35 ± 0.03
Mn	215.29±6.14	$382.28{\pm}10.48$	355.44±2.99
Zn	179.80±4.69	1087.66 ± 54.44	204.87±7.71
Cu	122.61±2.95	325.01±8.79	188.63±4.34
Pb	162.86±5.93	66.26±3.64	6.91±0.51
Cd	252.18±4.24	169.76 ± 7.56	0.64 ± 0.05
Ni	58.51±0.96	56.90±2.61	4.72±0.16
Cr	418.75±8.32	315.42±8.42	8.04±0.23
As	18.07±1.15	10.67±0.36	$0.04{\pm}0.002$
Phytoavailable metals (mg kg ⁻¹)			
Al	187.7±4.79	2.42 ± 0.04	0.01 ± 0.0001
Fe	83.41±0.52	283.64±5.90	220.12±4.04
Mn	21.91±2.15	54.91±0.11	42.79±0.12
Zn	5.07±0.32	64.81±0.22	16.42±0.02
Cu	4.62±0.12	42.74±1.04	28.74±1.55
Pb	0.03 ± 0.001	0.07±0.001	$0.04{\pm}0.001$
Cd	0.27±0.02	0.71 ± 0.02	0.44 ± 0.01
Ni	1.45 ± 0.04	3.62±0.07	3.24±0.05
Cr	1.04 ± 0.08	5.12±0.48	2.75±0.16
As	0.05 ± 0.002	1.24±0.05	0.02±0.001

*g kg⁻¹; nd: not detected; EC: electrical conductivity; TOC: total organic carbon; TN: total nitrogen; AP: available phosphorous; CEC: cation exchange capacity

Table S1-3 Pearson's correlation coefficient between metal contents in soil and plant biomass under control and different red mud treatments in soil amended with both cowdung manure and sewage-sludge

Metals	R-values			
	Cowdung manure	Sewage-sludge		
Al	0.994***	0.991***		
Fe	0.935***	0.930***		
Mn	0.749^{***}	0.909^{***}		
Zn	0.681^{**}	0.991***		
Cu	0.911****	0.925^{***}		
Pb	0.967***	0.994^{***}		
Cd	0.992***	0.994^{***}		
Ni	0.870****	0.980^{***}		
Cr	0.865***	0.955***		
As	0.849***	0.945***		

S3

2
2
3
4
5
6
7
/
8
9
10
12
13
14
15
15
16
17
18
10
13
20
21
22
22
20
24
25
26
27
21
28
29
30
00
31
32
33
34
05
35
36
37
38
00
39
40
41
12
7 <u>6</u> 40
43
44
45
16
40
4/
48
49
50
50
51
52
53
5/
54
55
56
57
50
20
60

Table S	51-4	Classification	of plant	based	on	metal	toleranc	e

No.MTI1 $0 \le 2$	(0.()	
1 $0 \le 2$	(%) C	lassification of plant
	5 H	ighly sensitive
$2 \qquad 25 \le$	50 S	ensitive
3 $50 \leq$	75 N	loderate
4 $75 \leq$	100 T	olerant
$5 \geq 100$) H	ighly tolerant

Table S1-5 Translocation (TF) and bioconcentration (BCF _{plant}) factors for studied metals in lemongrass grown under control and RM treatments in soil amended with
CD or SS (Mean \pm SE).

Metals	Parameters	С	CD _{RM0}	CD _{RM5}	CD _{RM10}	CD _{RM15}	SS _{RM0}	SS _{RM5}	SS _{RM10}	SS _{RM15}
Al	TF	1.48 ± 0.12^{d}	2.26 ± 0.08^{a}	$2.09{\pm}0.00^{ab}$	2.08 ± 0.07^{ab}	2.08 ± 0.00^{ab}	2.10 ± 0.15^{ab}	1.88 ± 0.07^{bc}	1.85 ± 0.07^{bc}	1.77±0.09°
	BCF _{plant}	$0.08{\pm}0.00^{e}$	$0.52{\pm}0.01^{a}$	$0.19{\pm}0.00^{b}$	$0.15 \pm 0.01^{\circ}$	$0.15 \pm 0.00^{\circ}$	$0.15 \pm 0.01^{\circ}$	$0.14{\pm}0.00^{\circ}$	0.13 ± 0.01^{cd}	$0.12{\pm}0.00^{d}$
Fe	TF	$0.16{\pm}0.00^{d}$	0.17 ± 0.00^{cd}	0.17 ± 0.01^{cd}	0.19 ± 0.01^{bc}	0.23 ± 0.00^{a}	0.12 ± 0.00^{e}	0.15 ± 0.01^{d}	$0.17{\pm}0.00^{d}$	0.21 ± 0.00^{b}
	BCF _{plant}	$0.25{\pm}0.01^{a}$	0.19 ± 0.03^{b}	$0.16 \pm 0.01^{\circ}$	0.10 ± 0.00^{e}	0.07 ± 0.00^{g}	0.13 ± 0.00^{d}	$0.09{\pm}0.00^{ m f}$	$0.07 {\pm} 0.00^{ m gh}$	$0.06{\pm}0.00^{ m h}$
Mn	TF	$0.47{\pm}0.01^{d}$	0.56 ± 0.06^{cd}	$0.75{\pm}0.08^{ab}$	0.80 ± 0.04^{ab}	0.81 ± 0.03^{a}	$0.48{\pm}0.04^{d}$	0.66 ± 0.04^{bc}	$0.79{\pm}0.05^{ab}$	0.77 ± 0.03^{ab}
	BCF _{plant}	$0.20{\pm}0.01^{de}$	0.18 ± 0.01^{e}	0.22 ± 0.01^{d}	$0.25 \pm 0.00^{\circ}$	0.28 ± 0.01^{b}	$0.24 \pm 0.01^{\circ}$	$0.29{\pm}0.00^{b}$	0.31 ± 0.01^{a}	$0.32{\pm}0.01^{a}$
Zn	TF	2.61 ± 0.12^{d}	$1.74{\pm}0.07^{de}$	1.56 ± 0.59^{ef}	1.16 ± 0.09^{ef}	$1.04{\pm}0.04^{\rm f}$	1.52 ± 0.06^{a}	$1.41{\pm}0.07^{ab}$	1.32 ± 0.09^{b}	$1.30\pm0.05^{\circ}$
	BCF _{plant}	0.22 ± 0.02^{cd}	0.17 ± 0.01^{e}	0.20 ± 0.01^{cd}	0.19 ± 0.00^{de}	$0.22 \pm 0.00^{\circ}$	0.26 ± 0.01^{b}	$0.27{\pm}0.01^{ab}$	$0.28{\pm}0.01^{ab}$	$0.29{\pm}0.00^{a}$
Cu	TF	$0.56{\pm}0.03^{a}$	$0.63{\pm}0.01^{a}$	0.66 ± 0.01^{a}	0.66 ± 0.04^{a}	$0.68{\pm}0.05^{a}$	0.66 ± 0.00^{a}	$0.68{\pm}0.00^{a}$	$0.69{\pm}0.00^{a}$	$0.74{\pm}0.14^{a}$
	BCF _{plant}	$0.59{\pm}0.04^{a}$	$0.59{\pm}0.01^{a}$	0.59 ± 0.02^{a}	0.58 ± 0.02^{a}	0.51 ± 0.01^{b}	$0.39{\pm}0.00^{d}$	0.43 ± 0.03^{cd}	0.45 ± 0.01^{bcd}	0.47 ± 0.02^{bc}
Pb	TF	2.24 ± 0.05^{b}	2.34 ± 0.14^{b}	$1.80 \pm 0.08^{\circ}$	1.17 ± 0.04^{de}	1.04 ± 0.05^{e}	2.81 ± 0.27^{a}	$1.80{\pm}0.00^{\circ}$	1.42 ± 0.05^{d}	1.34 ± 0.03^{de}
	BCF _{plant}	$0.12\pm0.00^{\rm f}$	0.33 ± 0.02^{e}	0.40 ± 0.01^{d}	$0.51 \pm 0.01^{\circ}$	0.59 ± 0.02^{b}	0.45 ± 0.04^{cd}	0.61 ± 0.00^{b}	0.70 ± 0.03^{a}	0.72 ± 0.02^{a}
Cd	TF	$1.64{\pm}0.16^{a}$	$1.60{\pm}0.05^{a}$	1.53 ± 0.14^{ab}	1.43 ± 0.05^{ab}	1.29 ± 0.02^{b}	1.59 ± 0.01^{a}	1.51 ± 0.11^{ab}	1.40 ± 0.03^{ab}	1.46 ± 0.09^{ab}
	BCF _{plant}	$0.90{\pm}0.03^{a}$	$0.89{\pm}0.02^{a}$	$0.81{\pm}0.04^{b}$	$0.71 \pm 0.02^{\circ}$	0.63 ± 0.01^{d}	0.75 ± 0.02^{bc}	$0.73 \pm 0.03^{\circ}$	$0.73 \pm 0.03^{\circ}$	0.68 ± 0.02^{cd}
Ni	TF	$2.03{\pm}0.05^{a}$	2.00 ± 0.12^{a}	1.77 ± 0.24^{ab}	1.70 ± 0.18^{ab}	1.68 ± 0.03^{ab}	1.60 ± 0.13^{b}	1.45 ± 0.09^{b}	1.39 ± 0.04^{b}	1.40 ± 0.03^{b}
	BCF _{plant}	$0.33\pm0.02^{\text{d}}$	0.48 ± 0.01^{ab}	$0.49\pm0.01^{\text{a}}$	0.46 ± 0.01^{ab}	0.44 ± 0.02^{b}	0.33 ± 0.00^{d}	$0.38\pm0.02^{\text{c}}$	0.44 ± 0.01^{b}	0.45 ± 0.00^{ab}
Cr	TF	$1.21{\pm}0.00^{a}$	$1.21{\pm}0.00^{a}$	1.13 ± 0.00^{b}	1.08 ± 0.00^{d}	1.04±0.00 ^g	$1.08\pm0.00^{\circ}$	1.05 ± 0.00^{e}	$1.04{\pm}0.00^{\rm f}$	1.03 ± 0.00^{h}
	BCF _{plant}	$0.50{\pm}0.02^{a}$	$0.52{\pm}0.01^{a}$	$0.14{\pm}0.01^{b}$	0.13 ± 0.00^{bc}	0.10 ± 0.00^{de}	0.12 ± 0.00^{bcd}	0.11 ± 0.00^{cde}	0.10 ± 0.00^{cde}	0.09 ± 0.01^{e}
As	TF	$1.18{\pm}0.01^{ab}$	$1.06 \pm 0.00^{\circ}$	$1.05\pm0.00^{\circ}$	$1.04 \pm 0.00^{\circ}$	$1.04 \pm 0.00^{\circ}$	1.20±0.00a	$1.19{\pm}0.01^{ab}$	1.17 ± 0.04^{ab}	1.13 ± 0.03^{b}
	BCF _{plant}	$0.78{\pm}0.09^{bc}$	$1.26{\pm}0.10^{a}$	$0.88{\pm}0.06^{b}$	0.69 ± 0.04^{cd}	0.52 ± 0.01^{de}	0.63 ± 0.05^{cde}	0.66 ± 0.06^{cde}	0.53 ± 0.04^{de}	0.47 ± 0.01^{e}

RM: red mud; CD: cowdung manure; SS: sewage-sludge; C: control; CD_{RM0}: soil with CD; CD_{RM5}: 5 % RM in CD_{RM0}; CD_{RM10}: 10 % RM in CD_{RM0}; CD_{RM15}: 15 % RM in CD_{RM0}, SS_{RM0}: soil with SS; SS_{RM5}: 5% RM in SS_{RM0}; SS_{RM0}: no solution state in same row differ significantly at p < 0.05 as per the Duncan's test.





Fig. S1-1 Linear regression between metal content in soil (independent variable) and entire plant biomass (dependent variable) under control and red mud treatments soil amended with cowdung manure or sewage-sludge (n=3). Levels of significance are indicated: ns, not significant; *, p < 0.05; **, p < 0.01; ***, p < 0.001.

S6





Fig. S1-2 Linear regression between metal content in soil (independent variable) and entire plant biomass (dependent variable) under control and red mud treatments soil amended with cowdung manure or sewage-sludge (n=3). Levels of significance are indicated: ns, not significant; *, p < 0.05; **, p < 0.01; ***, p < 0.001.