



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/116744/>

Version: Accepted Version

Article:

Headlam, J, Ford, AC and Gracie, DJ (2017) Reactive versus proactive therapeutic drug monitoring in IBD patients treated with infliximab: A self-fulfilling prophecy. *Clinical Gastroenterology and Hepatology*, 15 (10). p. 1638. ISSN: 1542-3565

<https://doi.org/10.1016/j.cgh.2017.05.017>

© 2017 by the AGA Institute. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Accepted Manuscript

Reactive versus proactive therapeutic drug monitoring in IBD patients treated with infliximab: A self-fulfilling prophecy

John Headlam, Alexander C. Ford, David J. Gracie



PII: S1542-3565(17)30592-X
DOI: [10.1016/j.cgh.2017.05.017](https://doi.org/10.1016/j.cgh.2017.05.017)
Reference: YJCGH 55245

To appear in: *Clinical Gastroenterology and Hepatology*
Accepted Date: 12 May 2017

Please cite this article as: Headlam J, Ford AC, Gracie DJ, Reactive versus proactive therapeutic drug monitoring in IBD patients treated with infliximab: A self-fulfilling prophecy, *Clinical Gastroenterology and Hepatology* (2017), doi: 10.1016/j.cgh.2017.05.017.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

TITLE PAGE

Title: Reactive versus proactive therapeutic drug monitoring in IBD patients treated with infliximab: A self-fulfilling prophecy.

Short “running” title: Therapeutic drug monitoring in IBD

Authors: John Headlam¹, Alexander C. Ford^{1,2}, David J. Gracie^{1,2}

¹Leeds Gastroenterology Institute, St. James’s University Hospital, Leeds, UK.

²Leeds Institute of Biomedical and Clinical Sciences, University of Leeds, Leeds, UK.

Grant Support: None

Abbreviations: IBD inflammatory bowel disease
TDM therapeutic drug monitoring

Correspondence: Dr. David Gracie

Leeds Gastroenterology Institute

Room 125

4th Floor

Bexley Wing

St. James's University Hospital

Beckett Street

Leeds

United Kingdom

LS9 7TF

Email: djgracie1982@doctors.org.uk

Telephone: +447980765615

Facsimile: +441132429722

Word count: 517

Disclosures: JH: none to declare. ACF: none to declare. DJG: none to declare.

Writing assistance: None

Author contributions: JH, ACF and DJG drafted the letter. All authors contributed to and approved the final draft of the manuscript.

Dear Editor,

We read the article by Papamichael *et al.* with interest.¹ This was a retrospective cohort study comparing the long-term outcomes of inflammatory bowel disease (IBD) patients treated with infliximab. The authors concluded that proactive therapeutic drug monitoring (TDM) was associated with better clinical outcomes including greater drug durability, less need for IBD-related surgery or hospitalization, and a lower risk of developing antibodies to infliximab than those in whom TDM was reactive. However, there are some issues with the study design that we would like to draw attention to.

Reactive TDM was performed only in patients who presented with symptoms consistent with current disease activity, or those who had experienced a prior infusion reaction. Thus, the apparent association of proactive TDM with improved clinical disease outcomes is likely to be largely related to the disparity in disease activity between patients in the proactive and reactive TDM groups at study entry. Although the authors acknowledge this as a limitation, they did not provide data on disease activity at the time of first institution of TDM.

In addition, the proportion of patients receiving escalated dosing of infliximab at the time of inclusion in the study was significantly higher in the reactive TDM group (51% vs. 35%; $P = 0.009$), again suggesting a significant difference in inflammatory activity between the two groups. Furthermore, at the time of allocation to proactive versus reactive TDM, antibodies to infliximab were present in a significantly higher proportion of patients in the reactive versus the proactive groups (28% versus 5%), which is likely to explain the increased risk of subsequent infusion reaction seen in this group of patients, as has been described previously.²

The TAXIT study was a randomized controlled trial assessing the impact of TDM in infliximab treated IBD patients.³ In this study, infliximab responders with optimized

infliximab trough levels were randomized to a dosing regimen based only on physician's assessment of clinical disease activity, or a group where dosing was based on TDM. There was no difference in clinical and biological remission between the two groups after 12 months of therapy, but the proportion of patients flaring during the maintenance phase of the study was lower in those with TDM-based infliximab dosing. These findings indicate that Papamichael *et al.* are likely to have overestimated the superiority of proactive over reactive TDM on clinical outcomes in IBD.

Optimization of anti-TNF therapy may be associated with improved outcomes in IBD,⁴ and further studies investigating how better to personalize these treatments are welcome.⁵ Proactive TDM is the logical approach to the long-term management of patients treated with infliximab but, on the basis of the findings of the current study, its superiority over reactive TDM cannot be ascertained.

REFERENCES

1. Papamichael K, Chachu KA, Vajravelu R, et al. Improved Long-term Outcomes of Patients With Inflammatory Bowel Disease Receiving Proactive Compared With Reactive Monitoring of Serum Concentrations of Infliximab. Clin Gastroenterol Hepatol. 2017. Epub 2017/04/04.

2. Krintel SB, Grunert VP, Hetland ML, et al. The frequency of anti-infliximab antibodies in patients with rheumatoid arthritis treated in routine care and the associations with adverse drug reactions and treatment failure. *Rheumatology (Oxford, England)*. 2013;52(7):1245-53. Epub 2013/03/06.
3. Vande Casteele N, Ferrante M, Van Assche G, et al. Trough concentrations of infliximab guide dosing for patients with inflammatory bowel disease. *Gastroenterology*. 2015;148(7):1320-9.e3. Epub 2015/03/01.
4. Vande Casteele N, Khanna R, Levesque BG, et al. The relationship between infliximab concentrations, antibodies to infliximab and disease activity in Crohn's disease. *Gut*. 2015;64(10):1539-45. Epub 2014/10/23.
5. <https://www.pantsdb.co.uk/#/about>.