

This is a repository copy of *A national-scale assessment of climate change impacts on species: assessing the balance of risks and opportunities for multiple taxa.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/119271/>

Version: Accepted Version

Article:

Pearce-Higgins, James W., Beale, Colin Michael orcid.org/0000-0002-2960-5666, Oliver, Tom et al. (20 more authors) (2017) A national-scale assessment of climate change impacts on species: assessing the balance of risks and opportunities for multiple taxa. *Biological Conservation*. pp. 124-134. ISSN 0006-3207

<https://doi.org/10.1016/j.biocon.2017.06.035>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Manuscript Details

Manuscript number	BIOC_2017_227
Title	A national-scale assessment of climate change impacts on species: assessing the balance of risks and opportunities for multiple taxa
Article type	Full Length Article

Abstract

It is important for conservationists to be able to assess the risks that climate change poses to species, in order to inform decision making. Using standardised and repeatable methods, we present a national-scale assessment of the risks of range loss and opportunities for range expansion that climate change could pose for over 3,000 plants and animals. Species were selected by their occurrence in England, the primary focus of the study, but climate change impacts were assessed across Great Britain, widening their geographical relevance. A basic risk assessment that compared projected future changes in potential range with recently observed changes classified 21% of species as being at high risk and 6% at medium risk of range loss under a B1 climate change scenario. A greater number of species were classified as having a medium (16%) or high (38%) opportunity to potentially expand their distribution. A more comprehensive assessment, incorporating additional ecological information, including potentially confounding and exacerbating factors (e.g. dispersal, habitat availability and other constraints), was applied to 402 species, of which 35 % were at risk of range loss and 42 % may expand their range extent. This study covers a temperate region with a significant proportion of species at their poleward range limit; the balance of risks and opportunities from climate change may be different elsewhere. The outcome of both risk assessments varied between taxonomic groups, with bryophytes and vascular plants containing the greatest proportion of species at risk from climate change. Upland habitats contained more species at risk than other habitats. Whilst the overall pattern was clear, confidence was generally low for individual assessments, with the exception of well-studied taxa such as birds. In response to climate change, nature conservation needs to plan for changing species distributions and increasing uncertainty of the future.

Keywords	adaptation; climate change; climate envelope; Great Britain; risk assessment; vulnerability
Taxonomy	Climate Change Adaptation, Nature Conservation, Global Change Vulnerability
Corresponding Author	James Pearce-Higgins
Order of Authors	James Pearce-Higgins, Colin Beale, Tom Oliver, Tom August, Matthew Carroll, Dario Massimino, Nancy Ockendon, Joanne Savage, Christopher Wheatley, Malcolm Ausden, Richard Bradbury, Simon Duffield, Nicholas Macgregor, Colin McClean, Michael Morecroft, Chris Thomas, Olly Watts, Bjorn Beckmann, Richard Fox, Helen Roy, Peter Sutton, Kevin Walker, Humphrey Crick
Suggested reviewers	Rob Brooker, Wendy Foden, Paul Pearce-Kelly

Submission Files Included in this PDF

File Name [File Type]

Risk and opportunities resubmission letter.docx [Cover Letter]

Comments from the editors and reviewers.docx [Response to Reviewers]

Risk and opportunities resubmission track changes.docx [Revised Manuscript with Changes Marked]

Risk and opportunities resubmission appendices track changes.docx [Revised Manuscript with Changes Marked]

Risk and opportunities resubmission.docx [Manuscript File]

Risk and opportunities resubmission appendices.docx [Supporting File]

Appendix 5. Species outcomes from the simplified risk assessment.xlsx [Supporting File]

Appendix 6. Species outcomes from the full risk assessment.xlsx [Supporting File]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

British Trust for Ornithology (BTO)

The Nunnery
Thetford
Norfolk, IP24 2PU
T: 01842 750050
F: 01842 750030



Re: Revisions to BIOC_2017_227

Dear Vincent,

Many thanks for the invitation to resubmit manuscript BIOC_2017_227, retitled '**A national-scale assessment of climate change impacts on species: assessing the balance of risks and opportunities for multiple taxa**'. We are also grateful to the three reviewers for their comments and requests for clarification, which we believe have significantly improved the manuscript. Their comments are duplicated in the response to reviewers, with our specific responses outlined in italics.

Many thanks in advance for reconsidering this submission, and we look forward to hearing from you regarding this manuscript.

With best wishes,

James Pearce-Higgins

(on behalf of all authors)

Comments from the editors and reviewers:

Reviewer #1

- This is a thorough and well constructed though necessarily complex study, though simplified and nicely understandable in the manuscript (supplementary Appendices appear to provide sufficient additional detail). I have only a few points of clarification that I think would benefit the main manuscript:

Thank you for these comments and points of clarification.

1. It would be useful in the Abstract to indicate that although the species analysed were sampled for their occurrence in England (making the study relevant to a geopolitical entity and its conservation assessments), the analytical framework considers their broader distributions where possible, during the bioclimatic modelling and to inform their risk/opportunity within the focal region; as it stands, the focus on England in the Abstract can lead to the assumption that the distribution modelling is geographically restricted. Indeed, it seemed odd to me that the study considered 'English species' ostensibly to inform conservation decision-making, but then examined their risk/opportunity across Britain; why not just consider British species, since many more arctic/alpine species will occur in Scotland, and this could shift the outcome significantly (the balance of increasing/decreasing species)?

This study was funded by Natural England, the Statutory Nature Conservation Body responsible for nature conservation in England, hence the geographical focus, but considered distributions and future projections within Britain, as the geographical area within the UK most relevant for the conservation of these species. Whilst we can see that this does lead to an apparently 'odd' discrepancy, as indicated in the discussion, in actual fact this difference enabled us to include more northern / upland species than we otherwise would have done had we maintained an England only restriction (L361-366). In response to this comment, we have inserted an additional sentence in the abstract to better explain this separation (L40-42), and then in the introduction (L109-111).

2. It is interesting that the Abstract suggests the study can robustly identify general patterns of biodiversity risk/opportunity despite low confidence for individual species assessments, while the opening of the Introduction states that it is necessary to prioritise species for conservation; to some degree this appears to be a tension throughout the manuscript - i.e. the ability to extract generalities, but with low confidence for any given species (at least for many species) - and I think this conflict could be better expressed through the structure of the Discussion.

This is a useful observation, thank you. Additional text to reflect this tension is inserted in L483-490.

3. It would be helpful to know in the main manuscript which climate data were used at the baseline and for future projections, whether from the UK Met Office, or some other data source (e.g. WorldClim); furthermore, the paper appears to use the SRES emissions scenarios based on storylines (B1, A1F etc.), and it is now important to cross-reference these against the more up-to-date RCP approach, and to indicate where the scenarios used here sit within the RCP framework.

Our projections were based upon the outputs from the UK Met Office (UKCP09), which as identified by the reviewer, probably require a little more detail, particularly for international readers. This is now clarified in the text (apologies for this omission), with further detail provided about the projections and how they relate to the RCPs (L183-193).

4. It was unclear to me how the ecological data, where available (e.g. relating to dispersal), was standardised across such divergent groups. For example, is a dispersal-limited bryophyte considered in the same category as a dispersal-limited bird or butterfly, or is there some across-group calibration (e.g. a dispersal-limited bird = a medium-dispersed bryophyte...). This seems a really

problematic issue when comparing across groups, and it'd be helpful to know some more detail on how this was handled.

More detail about how the exacerbating factors, such as dispersal and habitat availability, were used to moderate the outcome of the risk assessment, is now provided L232-241.

5. It would also be very useful to specify directly what the risks/opportunities consist of (lines 122-124), rather than referencing a previous paper.

Text has been inserted to emphasise that the risks are of species decline in parts of its current range, and opportunities for range expansion into other regions, as a result of climate change, and that by combining observed and modelled responses to climate change, the assessment is appropriate for the long time-frame over which climate change is likely to impact species (L126-132).

6. Line 446 - butterflies

Changed - thanks

7. Figure 1 - second box in second column (Is the decline is linked)

Changed - thanks

-Reviewer #2

- This is an important and timely contribution to climate change assessment efforts and the authors have done a good job in explaining their approach, findings and the associated constraints. Because the constraints are well covered I have no significant critical comments. It would be interesting to have more details on the potential exacerbating factors that were used to assess the degree of confidence but this does not overly detract from the value of the paper. The author's are to be congratulated on realising such an ambitious and important initiative.

Many thanks for these positive comments and for the observation about the value of providing more details about the use of exacerbating actors. We have done this, explaining more clearly what these exacerbating factors are and how they are used to influence both the eventual assessment scores, and their confidence (L232-245).

- Reviewer #3

This paper describes use of two methods for assessing the potential impacts of climate change on the distributions of a broad range of species in England. A basic risk assessment method compared projected range changes with those observed, while a more detailed assessment included additional ecological information in projections, as per a previously published approach. The paper describes and compares the results of both methods and specifically examines differences in risk between taxonomic groups and habitat. A detailed discussion of conservation implications is provided.

The study tackles the important and poorly covered issue of national-scale assessments of climate change impacts on biodiversity. The methods it proposes are valid and valuable for this purpose, and provide a good foundation for further development of this field. While the levels of data available for England may be aspirational for many countries, this demonstration of how they may be used is important. The various challenges and shortcomings of the methods are mostly well acknowledged and discussed and the appendices provide enough information on methods and the English species. The paper is well written and data is suitably presented. My major concern regards the species distribution modeling, including use of outdated projections (discussed below). Overall, however, I think the paper is well worth publishing, subject to some suggested revisions.

Thank you for these comments. Our responses to the detailed criticisms are outlined below.

Main concerns

Species Distribution Modeling concerns:

1. Use of IPCC AR4 emissions scenarios (2007) rather than AR5's RCPs (2013).
2. Use of only one model type (i.e. GAM).
3. Use of a single set of bioclimatic variables for all taxonomic groups considered, from birds to plants. This is widely regarded as inadvisable, for obvious reasons.

Updating these would be first prize. However, understanding the amount of work this would involve, exploring the implications of the above for a subset of species and exploring broader effects on results would probably suffice. A "Ways forward" or "Recommended next steps" section or paragraph suggesting methodological improvements would also be beneficial.

You are correct, that many such SDM studies adopt an ensemble approach covering a range of modelling types, GCMs and emissions scenarios. The resulting spread of projections are regarded as more likely to capture the future than a single output. However, we believe we have a good case for our approach of selecting a single modelling approach and GCM that is most likely to work best for our purposes. However, we had not explained this clearly. This justification is now set-out better in L491-508.

We disagree with the suggestion of using different bioclimatic variables for different taxa – many studies use a set of ecologically relevant variables that are likely to work across taxa (e.g. Huntley et al. 2007 modelled the abundance of European birds from 3 variables based largely on a knowledge of plant biogeography). There is increasing evidence that the relationships between species populations and climate have common patterns across taxa (e.g. Pearce-Higgins et al. 2015 Proc Roy Soc B), probably because most of the climate change impacts operate through altered species' interactions rather than directly through a species' physiology and tolerance to different climatic extremes (Ockendon et al. 2014 GCB).

The omission of rare species is a serious concern in drawing conclusions from both methods used, but on the whole, the authors acknowledge and discuss this issue satisfactorily in the discussion. As part of this, they compare the results using GB + European data with GB only data and find, as expected, that the latter are more pessimistic. What's not acknowledged, however, is that for species that are not European endemics, this pessimism is even more pronounced. If possible, assess the impacts of this, and at least discuss it.

As recognised, the difficulties posed by rare species are already discussed in L355-369. However, the specific omission of potential colonists from Europe was not covered. It is difficult to deal with in practice because it is difficult to know the extent to which any lack of occurrence is down to the sea barrier of the English Channel, or down to climate change. However, the reviewer is correct that by focussing on species that currently occur in England, we will not have allowed for potential colonisation of new species, which we know to be happening, at least in some groups. This limitation is now acknowledged (L381-387).

Why present the results of B1 only in the main paper – current National Commitments reflect a likely change of 3°C (UNEP), and given uncertainties in CO2 emissions politics, this could feasibly lead to anywhere between 2 and 4 degrees. You discuss that there is a strong correlation in results from B1 and A1B, which is expected, but by leaving out the A1B results we don't get a sense of the extent of uncertainty or of which new species would become at risk.

Given the length of the manuscript, we are reluctant to introduce significant additional content into the results section, as suggested by the review. The slightly more pessimistic nature of the A1B scenario under the simplified risk assessment is indicated in L277-279, although the results differed little for the full risk assessment (now added for clarity in L305-307). Hopefully by making the information from the A1B scenario available in Tables A1 and A2, Appendix 1 (which wasn't particularly clear from the previous legend text for these tables), and as individual species-level

assessments in Appendices 5 and 6, that achieves the right balance of content in the paper and appendices, whilst making the differences between these two scenarios available to the reader. However, if the editor would like us to include more information from the A1B scenario in the main paper, we can do.

Detailed comments

Title: "A national-scale assessment of climate risk to species". This should read "climate change". Since it's risks and vulnerabilities, the first component of the title might be "climate change impacts on species:..."

The title has been changed, as suggested, to 'A national-scale assessment of climate change impacts on species: assessing the balance of risks and opportunities for multiple taxa'

168: This isn't an estimate of species' responses, but of shifts in climate space/potential range shifts. Please reword.

Changed to potential range shifts as suggested.

191: should subspecies and varieties be included? If these have reproductive compatibility with others in their species as can be expected then this will lead to overestimation of the species' extinction risk. Results discussed in terms of species so this is misleading.

The approach taken matches that of other previous studies of these data (e.g. Dyer et al. 2016 J App Ecol). There is not necessarily a straight-forward response to the taxonomic challenges of some of these species, and how they are split into different sub-species or distinct varieties, particularly as the taxonomy can change through time. We have therefore maintained the current taxonomic distinctions as represented in the species' database, but in order to assess whether this is likely to bias the results, also present the risk assessment outcomes just for full species. The general similarity of assessment for both 'true' species and all taxonomic concepts suggests that the potential risk of overestimation of species' extinction risk is not a large one.

Ln 446 – typo

Ln 508 – "are necessary"

Dealt with, thank you.

Additional changes

A number of other additional textual changes have been made to the manuscript for clarity, and which are shown by submitting a version of the document showing these using 'track changes'. In particular, we have tightened up the use of some of the terminology. We noted a slight discrepancy between the use of the term 'threat' in the appendix and 'risk' in the main manuscript, which has now been standardised to 'risk' throughout. We have also standardised the use of the term 'opportunity', whereas in the original submission, this term was used interchangeably with 'benefit'.

A national-scale assessment of climate ~~change impacts risk to on~~ species: assessing the balance of risks and opportunities for multiple taxa

James W. Pearce-Higgins^{1,2}, Colin M. Beale³, Tom H. Oliver^{4,5}, Tom A. August⁴, Matthew Carroll^{3, 6,7}, Dario Massimino¹, Nancy Ockendon^{1,2}, Joanne Savage⁴, Christopher J. Wheatley³, Malcolm A. Ausden⁶, Richard B. Bradbury^{2,6,7}, Simon J. Duffield⁸, Nicholas A. Macgregor^{8, 9}, Colin J. McClean¹⁰, Michael D. Morecroft⁸, Chris D. Thomas³, Olly Watts⁶, Björn C. Beckmann^{3,4,11}, Richard Fox¹², Helen E. Roy⁴, Peter G. Sutton¹¹, Kevin J. Walker¹³, Humphrey Q.P. Crick⁸

¹ British Trust for Ornithology, The Nunnery, Thetford, Norfolk, IP24 2PU, UK

² Conservation Science Group, Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

³ Department of Biology, University of York, Wentworth Way, York, YO10 5DD, UK

⁴ Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

⁵ School of Biological Sciences, University of Reading, Whiteknights, Reading, Berkshire RG6 6AS, UK.

⁶ Royal Society for the Protection of Birds, The Lodge, Sandy, Beds., SG19 2DL, UK

⁷ RSPB Centre for Conservation Science, The David Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, UK

⁸ Natural England, Eastbrook, Shaftesbury Road, Cambridge, CB2 8DR, UK

22 ⁹ *Durrell Institute of Conservation and Ecology (DICE), School of Anthropology and*
23 *Conservation, University of Kent, Canterbury, Kent, CT2 7NR, UK*

24 ¹⁰*Environment Department, University of York, Wentworth Way, York, YO10 5NG, UK*

25 ¹¹*Orthoptera and Allied Insects Recording Scheme of Britain and Ireland, c/o Biological*
26 *Records Centre, Centre for Ecology & Hydrology, Wallingford, OX10 8BB, UK*

27 ¹²*Butterfly Conservation, Manor Yard, East Lulworth, Wareham, Dorset, BH20 5QP, UK*

28 ¹³ *Botanical Society of Britain and Ireland (BSBI), Room 14 Bridge House, 1-2 Station*
29 *Bridge, Harrogate, North Yorkshire HG1 1SS, UK*

30

31 Corresponding author: Pearce-Higgins, J.W. Email: james.pearce-higgins@bto.org, Tel: +44
32 (0)1842 750050

33

34 Word count (9,248 including references 2,143, tables and figures 763)

35

36 Abstract

37 It is important for conservationists to be able to assess the risks that climate change poses to
38 species, in order to inform decision making. Using standardised and repeatable methods, we
39 present a national-scale assessment of the risks of range loss and opportunities for range
40 expansion; that climate change could pose for over 3,000 plants and animals. Species were
41 selected by their occurrence in England, the primary focus of the study, but climate change
42 impacts were assessed across Great Britain, widening their geographical relevance. future
43 that occur in England. A basic risk assessment that compared projected future changes in
44 potential range with recently observed changes classified 21% of species as being at high risk
45 and 6% at medium risk of range loss under a B1 climate change scenario. A greater number
46 of species were classified as having a medium (16%) or high (38%) opportunity to potentially
47 expand their distribution. A more comprehensive assessment, incorporating additional
48 ecological information, including potentially confounding and exacerbating factors (e.g
49 .dispersal, habitat availability and other constraints), was applied to 402 species, of which 35
50 % were at risk of range loss and 42 % may expand their range extent. This study covers a
51 temperate region with a significant proportion of species at their poleward range limit; ~~–~~ and
52 The-the balance of risks and opportunities from climate change may be different elsewhere.
53 The outcome of both risk assessments varied between taxonomic groups, with bryophytes
54 and vascular plants containing the greatest proportion of species at risk from climate change.
55 Upland habitats contained more species at risk than other habitats. Whilst the overall pattern
56 was clear, confidence was generally low for individual assessments, with the exception of
57 well-studied taxa such as birds. ~~–~~In response to climate change, nature conservation needs to
58 plan for changing species distributions and increasing uncertainty of the future.

60 Keywords: adaptation; climate change; climate envelope; Great Britain; risk assessment;
61 vulnerability

62 **Introduction**

63 To make the best use of conservation resources, it is necessary to prioritise species for action,
64 for example according to their current status and the threats that they face. Globally, the most
65 widely adopted framework for this is the IUCN Red List which quantifies extinction risk
66 using information on the population size and range extent of a species, and the rate of change
67 in those parameters (Mace et al., 2008, IUCN 2016). Anthropogenic climate change is likely
68 to exacerbate the extinction risk of many species over the course of this century (Thomas et
69 al., 2004, Bellard et al., 2012, Warren et al., 2013, Foden et al., 2013). A number of
70 approaches have been developed to assess the potential impact of climate change on species'
71 future status (Akçakaya et al., 2015). One common approach uses species distribution models
72 (widely termed bioclimatic-envelope or climate-envelope models) to link distribution to
73 climate variables and project the likely future impact of climate change on species'
74 distributions (e.g. Thomas et al., 2004, Huntley et al., 2007, Walmsley et al., 2007, Warren et
75 al., 2013). An alternative approach is to undertake vulnerability assessments which may
76 combine a measure of future projected climate change (exposure) with ecological traits to
77 identify the sorts of species most likely to be both sensitive to and lack the capacity to adapt
78 to climate change (e.g. Gardali et al., 2012, Foden et al., 2013).

79 Vulnerability assessments have often been applied to single taxonomic groups within
80 particular regions or countries (e.g. Heikkinen et al., 2010, Barbet-Massin et al., 2012) or,
81 less commonly across a global scale (Jetz et al., 2007, Foden et al., 2013). Relatively few
82 vulnerability assessments have covered the full range of biodiversity present within a
83 particular geographical area, despite the fact that a comprehensive assessment of as many
84 taxa as possible would assist governments and conservation organisations plan and adapt to
85 climate change. Achieving such wide coverage is challenging because many assessments

require taxon-specific information or use approaches that have limited applicability to other taxa (e.g. Heikkinen et al., 2010, Gardali et al., 2012, Moyle et al., 2013). To date, it has been difficult to develop an approach which works across a range of taxa due to the different nature of ecological traits across contrasting taxonomic groups, and the variable availability of data (e.g. of species distributions, trends and traits). The strong tradition of biological recording in Britain across a wide range of taxa provides a rare opportunity to tackle this challenge.

Thomas et al., (2010¹) developed a framework to assess the threats and potential benefits of climate change that is applicable to a wide range of taxa. It uses bioclimatic-envelope models, combined with information on recent trends and additional ecological information, to identify the likelihood of species' range expansion and contraction, and has so far been applied to UK butterflies and some exemplar species from other taxa (Thomas et al., 2010¹). Here, we use a modification of this approach to undertake a climate change vulnerability assessment of more than 3,000 terrestrial and wetland species, (and in a minority of cases, species aggregates and distinctive subspecies or varieties, hereafter all termed 'species' for brevity; see methods) across 17 taxonomic groups in Britain (Table 1). This provides the first opportunity to examine how an important aspect of vulnerability to climate change varies between taxonomic groups, and between species associated with specific habitat types, for as complete a biological assemblage as currently feasible.

This study was developed as part of a wider initiative of Natural England, the government conservation agency in England, to support decision making on adaptation (Natural England 2014) and inform an adaptation plan (Natural England, 2015). It therefore focuses on species in England, the largest of the component countries within the United Kingdom (UK), but

assesses the vulnerability of those species across Great Britain (GB), ~~as this better represents the unit of contiguous land across which conservation decisions are made within the UK.~~
~~the single land mass within which England is located. This ensures that the outputs are also highly relevant for Wales and Scotland, for UK organisations, and more widely.~~

Materials and Methods

The vulnerability assessment involved a number of steps (Figure 1) outlined below:

1. Distribution data for over 5,000 species were collated for a wide range of taxa that occur in England (Table 1).
2. Statistical models linking species' distributions to climate were used to assess the likely impacts of future climate change upon species' potential distributions.
3. Information from these projections was compared with observed changes in species distribution. By assessing recently observed changes in the context of projected future trends, a *simplified risk assessment* could be undertaken rapidly across all species.
4. For a representative subset of 402 species, additional ecological information enabled the application of the full Thomas et al., (2010) framework. By considering the potential for non-climatic factors and ecological constraints to affect species' responses to climate change, this framework produces a more comprehensive assessment (the *full risk assessment*).

Whilst the term 'risk assessment' can have specific meanings in different contexts, we follow Thomas *et al.* (2010) and use it to describe our methodology for assessing the potential risks of species decline and ~~loss~~extirpation in parts of its current range, and opportunities that the same species may expand its distribution into other regions, both as a result of climate change

~~may pose for species. By using a combination of observed and modelled responses to climate change, the methodology deals with the long time-scales over which species' responses to climate change are likely to occur.~~

Species distribution data

Species distribution data for GB were available from a range of biological recording schemes for a total of seventeen taxonomic groups (Table 1) at a hectad (10 km square) resolution. For inclusion, species had to be present in England and recorded from more than 5 hectads (the minimum required for modelling; Hickling *et al.*, 2006). Even with this threshold the climate envelope models (described below) failed to converge for 10% of the most sparsely distributed species, giving a total of 4,540 species for which modelling was possible.

We used data from 1970-89 to represent baseline distributions prior to recent climate change, in order to minimise the risk of species' distributions being unsynchronised with the climate due to recent range shifts (Mason *et al.* 2015). For plants we used the period 1970-86; the time period (Braithwaite & Walker 2012) that most closely matched the data for other taxa. For birds the period 1988-91 was used, which coincided with a national atlas (Gibbons *et al.*, 1993). Cells for which climate data were not available were excluded from analyses. To aid model convergence, small islands, with little data, were also excluded for all taxa apart from birds, leaving 2,561 hectads, or 2,670 for birds.

Recording effort varies~~d~~ between taxa, with the highest coverage for groups with well-developed and popular volunteer recording schemes such as vascular plants ~~and~~, birds ~~and~~ butterflies. To avoid species' distribution models being biased as a result of limited recording effort, we used the program FRESALO (Hill, 2012) to estimate taxon-specific recorder effort in each 10 km square (see below).

Species distribution modelling

We used the climate envelope modelling approach of Beale et al., (2014) across all taxa (Appendix 1). The approach was devised to address the problem of spatial autocorrelation in large-scale species' distribution data, and applies a Bayesian, spatially explicit (Conditional Autoregressive) Generalised Additive Model (GAM) to species' distribution data in order to separate climatic, spatial and random components in determining the distribution of each species. Four bioclimate variables were used to describe spatial variation in the climate, using 1961-1990 averages:

- mean temperature of the coldest month (MTCO): a measure of winter cold.
- growing degree days above 5°C (GDD5): a measure of biologically useful warmth, calculated by applying a spline to mean monthly temperatures for each cell to convert monthly data to daily estimates.
- the coefficient of variation of temperature (cvTemp): a measure of seasonality
- soil moisture (soilWater): a measure of moisture availability calculated following the bucket model of Prentice et al., (1992), which takes inputs of temperature, rainfall, % sun/cloud and soil water capacities.

For birds and a quarter of vascular plants, we initially constructed 50 km resolution species distribution models across Europe to describe the relationship between occurrence and climate using uninformative priors (i.e. with no prior knowledge of what this relationship should be). Once converged, a second model was fitted to hectad data from GB using informative priors from the European-scale analysis. As a result, any strong climatic signal based on the European distribution would remain essentially unchanged when modelled using GB data only, unless there was strong evidence for a different climatic signal within GB. In

cases where there was high uncertainty in the estimation of ~~species' responses potential range~~
~~shifts~~ at a European level, the GB model would be more heavily informed by outputs from
the British component of the model. We tested for differences between both models for birds
and vascular plants under the A1B scenario. Predicted changes were strongly correlated,
although models based on GB only data tended to result in fewer species showing potential
increases in range (Appendix 1). For species for which data from GB only were available,
only the second model was run using uninformative priors.

Future climate projections for the UK were derived from UKCP09, which use outputs from
an ensemble of variants of the HADSM3 climate model to produce a series of probabilistic
outputs for individual climate variables for three IPCC SRES scenarios (A1F1, A1B and B1).
These are regarded as the most suitable climate change projections for the UK, downscaled to
a 25 km grid (Murphy et al. 2009). We considered two contrasting scenarios, the B1 scenario
which is a low emissions scenario projected to lead to a c. 2°C global temperature increase by
the end of this century, (equivalent to RCP4.5) and the A1B scenario, that represents
vulnerabilities under a medium emissions scenario of c. 4°C global warming by the end of
this century, and is (intermediate between RCP6 and RCP8.5) (Rogelj et al., 2012). As there
was a strong correlation between the results of the ~~two A1B and the B1~~ scenarios, we focus
on the B1 results in this paper, and present the. ~~These represent the species' potential~~
~~vulnerabilities to the magnitude of climate change under a low emissions scenario projected~~
~~to lead to a c. 2°C global temperature increase by the end of this century. R~~results from the
A1B scenario ~~are presented~~ in Appendix 1, ~~representing vulnerabilities under a medium~~
~~emissions scenario of c. 4°C global warming by the end of this century.~~

Simplified risk assessment

Distribution data from national schemes were used to identify post-1989 range changes within the baseline historical distribution (1970-89; or 1970-86 for plants and 1988-91 for birds, as described above), and outside this historic range (newly colonised areas). With the exception of birds, distributional changes required correction to account for variation in observer effort (Appendix 2).

Due to limited data availability across adequately sampled squares, it was not possible to use this method to produce effort-corrected observed trends for 1,492 species, leaving a total of 3,048 to which the risk assessment could be applied. Of these, 50 were species aggregates reflecting taxonomic changes over previous decades (1 bird, 3 carabid beetles, 28 bryophyte and 18 vascular plants), 123 were specific subspecies or varieties (38 bryophytes, 2 spiders and 83 vascular plants), and 80 were infraspecies, whose distribution may have been based on partial information, due to the separate recording of taxonomically distinct subspecies or related species aggregates (31 bryophytes, 1 carabid beetle and 48 vascular plants). The inclusion of this mix of taxonomic resolutions did not bias the risk assessment towards species of particular risk or opportunity categories; in a sensitivity analysis there was no significant difference in the allocation to different risk categories between ‘true’ species and these other taxonomic concepts combined, under either the B1 ($\chi^2_4 = 7.93$, $P = 0.094$) or A1B ($\chi^2_4 = 7.44$, $P = 0.11$) scenarios. We have therefore assessed all taxonomic concepts together, but for completeness also present the results for bryophytes and vascular plant species separately, excluding aggregates, subspecies and infraspecies.

Current contractions within the historical range were compared against the magnitude of projected future contractions to assess risk from climate change, whilst observed range expansion was cross-tabulated with the magnitude of projected future range expansion to assess potential ~~threats~~risks and opportunities from climate change (Appendix 3). The

highest ~~threat-risk~~ or opportunity categories were reserved for those species where projected future changes were consistent with observed changes. As the simplified risk assessment may have inflated the potential risk of climate change for species which have suffered recent declines and range contractions for non-climatic reasons, for a subset of 402 species, we also undertook a full risk assessment following the Thomas et al., (2010¹) framework to account for non-climatic factors and constraints.

Full risk assessment

The 402 species (including 4 subspecies / varieties and 1 infraspecies) for full assessment comprised 155 conservation priority species listed under the Section 41 of the Natural Environment and Rural Communities (NERC) Act 2006 (http://www.legislation.gov.uk/ukpga/2006/16/pdfs/ukpga_20060016_en.pdf), termed NERC species, as well as at least 13 randomly selected species from each taxonomic group. This provided a broad appraisal across taxa, while ensuring as many species of highest conservation concern as possible were included. The full risk assessment ~~used provided a confidence level for each observed and projected population trend using~~ additional ecological information ~~on population size and range extent, the link between linking~~ population and range changes to climate, and on potential exacerbating factors (e.g. range extent and population size, ecological constraints associated with habitat-availability, dispersal and species interactions) ~~to moderate the likely risk and opportunity scores, and the overall assessment of confidence (Thomas et al., 2011). Small and range-restricted populations, or species associated with other constraints, received a higher risk score, whilst the likelihood of range expansion was reduced if habitat availability, dispersal ability and other limiting species were judged as likely to result in species achieving a lower level of range expansion than predicted by the models.~~ This information was gathered from a literature search for each

species using Google Scholar and Web of Science, supplemented by additional information from UK species experts (see Acknowledgements). The confidence associated with ecological information was regarded as good if based upon peer-reviewed literature. If it was based on expert knowledge then the expert was asked to assign the confidence level.

The full risk assessment consisted of four stages (Figure 12, Appendix 4), requiring information on observed changes in occurrence within the current range (Stage I), projected changes within the current range (Stage II), observed changes in occurrence outside the current range (Stage III) and projected changes outside the current range (Stage IV). The results of the four stages were synthesised into a single table (Table A4). The overall confidence for species ‘at risk’ was the confidence associated with the assessment of threat, while for species with an opportunity for expansion, likely to benefit we used the confidence associated with ~~the likely opportunity that~~. For species classed as having ‘risks and opportunities’ or ‘limited impact’, we averaged the two confidence scores.

Statistical analysis

Significant differences in the proportion of species allotted to different risk categories were tested by Chi-square, as were contrasts between taxonomic groups and between NERC and other species. Information on the broad habitat associations of the 155 NERC priority species, summarised into wetland, urban, farmland, upland woodland and coastal categories, was used to test the extent to which species’ vulnerability to climate change, from the full risk assessment, varied between habitats.

Formal differences between the results from the simplified and full risk assessments for each of the 402 species assessed using both risk assessment methods were tested by Chi-square test and by regression. For the latter, we converted the categorical risk assessment into rank scores from high risk (-2) to high opportunity (2), with both ‘risks & opportunities’ and

‘limited impact’ categories scored as 0. Scores were regressed within a generalised linear mixed model, with taxonomic identity as a random effect, using PROC MIXED in SAS v9.2. We used the same scores to test for differences in full risk assessment outcomes between different taxa, and between NERC and other species.

Results

Simplified risk assessment

Of the 3,048 species assessed, 640 were classified as being at high risk of a decline in the area of projected suitable climate under the B1 climate change scenario and 188 at medium risk (a total of 27.2% species at risk). A greater number of species were identified as likely to have a medium (486) or high (1,164) potential opportunity as a result of projected increases in the area of potentially suitable climate (totalling 54.1%; Table 2). For only 6 was limited impact predicted. These estimates of risk were similar under the A1B warming scenario ($\chi^2_5 = 2.96$, $P = 0.71$), although with slightly more species (28.1%) classified as being at risk (Appendix 1 [Table A1](#)).

The outcome of the risk assessment varied significantly between taxonomic groups ($\chi^2_{64} = 475.54$, $P < 0.0001$; excluding the limited impact category due to the small sample size). These differences remained ($\chi^2_{32} = 339.73$, $P < 0.0001$) when simply splitting species into those at risk, likely to have an opportunity, or likely to be unaffected (i.e. risks & opportunities and limited impact categories combined). The proportion of species at risk varied from 6% for wasps to 39% for vascular plants, while the proportion of species with opportunity varied from 37% for bryophytes to 90% for wasps (Figure 3). Repeating this appraisal for bryophytes and vascular plants without subspecies and infraspecies produced equivalent assessments for both (bryophytes: high ~~benefit~~ [opportunity](#) 107 spp (25%),

medium ~~opportunity benefit~~ 48 spp. (11%), risks and ~~opportunity benefits~~ 134 spp. (32%),
medium risk 32 spp. (8%), high risk 102 spp. (24%); vascular plants: high ~~opportunity benefit~~
210 spp. (30%), medium ~~opportunity benefit~~ 103 spp. (15%), risks and ~~opportunity benefits~~
131 spp. (19%), medium risk 59 spp. (8%), high risk 200 spp (28%)). The groups with the
greatest proportion of species at risk from climate change were bryophytes and vascular
plants (> 30 % in both cases), whilst a number of groups were largely (>70 %) comprised of
species for which climate change may present an opportunity for range expansion in GB
(ants, bees, centipedes, coccinellid beetles and wasps).

NERC species contained slightly more ‘high risk’ and ‘medium opportunity’ species and
fewer ‘high opportunity’ species than expected from the pattern across the other species ($\chi^2_4 = 10.30, P = 0.036$), but there was no overall difference between these two species groups
when the categories were simplified to risk, opportunity or unaffected ($\chi^2_2 = 1.07, P = 0.58$).

Full risk assessment

Across all 402 species run through the full framework for the B1 scenario, 141 (35.1 %) were
classified as being at high or medium risk of being negatively affected by climate change,
compared to 168 (41.8 %) which were listed as likely to have a medium or high opportunity
(Table 3). Limited impact was predicted for 19% of species. There was no significant
difference from this classification of species under the A1B scenario ($\chi^2_5 = 0.94, P = 0.92$;
Appendix 1 Table A2). The score attributed to species did not vary between NERC species
and the remainder ($F_{1,384} < 0.01, P = 0.99$), but did vary with taxonomic group ($F_{16,384} = 3.38$,
 $P < 0.0001$). The lowest scores, indicating the greatest proportion of species at risk from
climate change, were for bryophytes (n=14), with the highest scores for ants (n=13) and

wasps (n=13), the majority of which were classed as having a high opportunity from climate change (Figure 4).

There was no significant variation overall between habitats in the frequencies of NERC species allocated to different risk categories ($\chi^2_{25} = 33.86$, $P = 0.11$). However, upland was the only habitat with a majority of species (75 %) regarded as being at risk of a decline in the area of projected suitable climate (Figure 5), which contrasted significantly with average of 40% of species across the remaining habitats when lumped together ($\chi^2_5 = 15.59$, $P = 0.008$).

For the majority (314) of species in the full assessment, confidence was poor, for 86 it was medium and good for only two. Confidence scores differed significantly between taxonomic groups ($\chi^2_{16} = 57.23$, $P < 0.0001$), driven primarily by a greater level of confidence for bird assessments (35% of 82 assessments were accorded medium or good confidence) than for other species, where 18% of 320 assessments were classed as having medium confidence, and none good.

Simplified v Full Risk Assessment

There was a strong association between the scores using the simplified and full approaches for species assessed by both ($F_{1, 398} = 955.56$, $P < 0.0001$; $S_F = -0.33 (\pm 0.089) + 0.91 (\pm 0.029) S_S$, where S_F is the full assessment score and S_S the simplified assessment score). The scores from the two frameworks had a close to 1:1 relationship, but the intercept shows that the full assessment on average produced a lower (higher risk or lower opportunity) score by 0.33 (or one third of a category), ~~a significantly higher threat or lower opportunity category.~~

Discussion

Here we present a national-level assessment of species' vulnerability to climate change, covering 3,048 species across 17 taxonomic groups. Consistently for both B1 and A1B scenarios, we found that there was a greater number of species for which potential range is projected to increase as a result of climate change than it is projected to decrease. This was particularly the case when considering the outputs for the simplified framework for all species, where over 50% were classified with a medium or high opportunity from climate change (Table 2), but also applied to 43 % of the subset of species run through the full risk assessment framework, compared with projected negative range impacts for 35% (Table 3). This also concurs with the previously published results of the full risk assessment methodology for butterflies in GB, which used an A2 climate change scenario intermediate between the B1 and A1B scenarios used here (Thomas et al., 2010¹). Of 58 butterfly species, three were regarded as at high risk from climate change, three at medium risk, 10 likely to have a medium opportunity, 14 a high opportunity and 27 limited impact. If turned into rank scores and added to the results of our study, this would place butterflies intermediate between coccinelid beetles and crane flies, with a mean score of 0.52 (Figure 4). Our findings are also consistent with recently observed trends across multiple taxa in the UK where more species are regarded as being impacted positively by climate change than negatively, at least in the short-term (Burns et al., 2016).

It could be argued that by indicating that a greater number of taxa are likely to have an opportunity for range expansion in response to climate change than be at risk of range contraction, our analysis suggests that climate change will have a positive impact upon UK biodiversity. However, before considering this, it is worth noting how our findings may result from both underlying methodological constraints and inherent biological processes.

It was not possible to undertake assessments for 13% of species because there were insufficient data to generate a bioclimate model, and for a further 29% of remaining species there was insufficient information to produce effort-corrected observed trends. Given latitudinal gradients in observer (recorder) effort within the UK, with more recorders in the south than the north, it is likely that a greater proportion of unassessed species were northerly-distributed and may include species more likely to be at risk of adverse climate change impacts than to benefit. However, by selecting species from England, but using data from across GB for their assessment, this enabled us to include more northern and upland species than we otherwise would have done had we undertaken the assessment with distribution data from England alone. In addition, it is possible that more localised and specialised species, which may be species less likely to benefit from climate change (e.g. Warren et al. 2001), were more likely to be data deficient and excluded. We did observe a significant difference between the scores of conservation priority species (many of which are rare and specialised) and others in the simplified assessment, but there was no such difference in the full assessment.

Apart from birds and vascular plants, the biodiversity data underpinning the assessment were from GB only, and in most cases our models do not capture the full range of climatically-suitable conditions in which the species can occur. A comparison of models based on GB data vs. GB + European data for birds and vascular plants, suggested that GB-only projections tended to be slightly more pessimistic than those that included European data, although the two were strongly correlated. Thus, the use of GB-only projections for most groups may have slightly inflated the projected magnitude of risk for those groups, although the assessment for vascular plants, one of the groups with the greatest proportion of species regarded as being at risk from climate change, included European data in the assessment. It is also worth noting that by considering including only species that currently occur in England,

we did not consider the potential for new species to colonise the UK from mainland Europe as a result of climate change. ~~As a result, our results do not anticipate the colonisation of the UK by new species, which is already happening (e.g. Hiley et al., 2013).~~ Thus which probably means that our results may exclude a number of potential colonists to the UK for which climate change provides an opportunity. In other words, the outcome of the risk assessment may be scale- and context-dependent; a species projected to be at risk from climate change across mainland Europe may undergo a poleward shift and colonise the UK, where it would be regarded as having an opportunity for range expansion. This emphasises the value of undertaking assessments such as this at a range of spatial scales, which has rarely been done.

We assumed that the species distribution models describe the main relationships between species' occurrence and terrestrial climate. As we employed widely-used bioclimatic variables, this is probably reasonable for most terrestrial taxa, but for some coastal bird species which use the marine environment, where spatial patterns of changes in sea temperature and other climate related variables may differ from those on land, projections are likely to be less certain. We also have not considered potentially detrimental impacts of sea-level rise and storm surges upon vulnerable coastal habitats and species (e.g. Gilbert et al., 2010; Ausden 2014).

The full assessment that considered ecological factors known to influence observed changes in populations or distributions, or likely constraints on the impacts of climate change, was applied to 402 species only. By excluding these considerations, the simple assessment applied across all species may have over-attributed observed changes to potential impacts of climate change if they were consistent with future projections (such as for farmland birds, crickets, centipedes and millipedes; Eglington & Pearce-Higgins 2012; Beckmann *et al.* 2015; Lee

2015; Burns et al., 2016), or under-estimated the potential magnitude of future climate change impacts if observed changes were opposite to future projections as a result of non-climatic factors. Although both methodologies delivered broadly comparable results, the full assessment did increase the proportion of species projected to experience only a limited impact of climate change, and included a greater proportion of species projected to be at risk.

Finally, there is considerable uncertainty about the likely pace of any distributional shift in response to climate change. Both bird and butterfly communities appear to be lagging behind the rate of warming observed across Europe (Devictor et al., 2012, Massimino et al., 2015); ~~non~~less-mobile groups, such as many of the vascular plants, may well lag even more. The ability of a species to disperse will be an important constraint on the extent to which some species can occupy any new areas of potential range in the future (Barbet-Massin et al., 2012), as will the availability of areas of potentially suitable habitat for colonisation (Thomas et al., 2012; Hiley et al., 2013), and underlying population dynamics (Mair *et al.* 2014).

Although considerable uncertainty remains about the pace of these responses to climate change, these uncertainties were at least partially captured by the full risk assessment, which reduces the likelihood of opportunity as a result of climate change in species with constrained dispersal ability.

Despite the potential methodological constraints, there are good biological reasons to expect more species to be able to expand their range than be at risk of it contracting in response to climate in GB. This is because there are more southern species with potential for northward range expansion in Britain than there are northern species with southern range margins (e.g. butterflies: Asher et al., 2001; vascular plants: Preston et al., 2002; birds: Balmer et al., 2013), with strong latitudinal gradients in species' richness (e.g. Eglington et al., 2015). In combination with largely polewards shifts that are projected to occur in the distribution of a

range of taxa, and are already being observed (Mason et al., 2015), this would lead to more species being likely to expand their distributions in GB, than to contract. Observations of recent trends suggest that this is already the case (Massimino et al., 2015, Burns et al., 2016). Although we assessed that fewer species would be at risk of range contraction from climate change than have an opportunity, species of certain taxonomic groups and habitats were identified as being more vulnerable than others. In particular, the full risk assessments completed for those species of conservation concern for which the required data is available suggested that species associated with upland habitat-types, where increasing temperatures might be expected to result in northwards and upwards range contraction, would be particularly vulnerable to climate change. This is consistent with the results of other studies suggesting that northern or upland birds (Green et al., 2008, Pearce-Higgins 2010), butterflies (Thomas et al., 2010¹) and plants (Hill & Preson 2015) may be more vulnerable to climate change than other species. Multi-taxa assessments have found similar patterns (Walmsley et al., 2007; Araujo *et al* 2011), and there is already evidence of such impacts being observed (Morecroft & Speakman 2015). While many taxonomic groups contain some species likely to be at risk from climate change and others with the potential to expand their distribution, the balance between these two outcomes will vary with the geographical and habitat bias of that group, as well as the ecological characteristics of the species, such as voltinism, diapause strategy, migratory strategy and growth rate (Bale et al., 2002). Other climate-influenced ecological changes will also affect species abundance and distribution in future through altered species interactions (Ockendon et al., 2014).

Geographical differences may partly account for the apparent high sensitivity to future climate change of bryophytes (Figures 3 and 4), many of which have a northern or north-western distribution, associated with cool and damp conditions. Our analysis suggests that of all the taxonomic groups considered, they are likely to be one of the most at risk from a

reduction in areas of suitable climate, conclusions broadly supported by Ellis (2015), who anticipated detrimental impacts of climate change on northern and upland bryophytes, although potential impacts on species associated with oceanic climates were more uncertain. Even though there is some evidence for recent warming being associated with distribution shifts in some bryophytes (Bates & Preston 2011), there are difficulties in disentangling these changes from decreases in acid and nitrogen deposition from the atmosphere (Roth et al., 2013). The basic assessment also identified vascular plants as containing a high proportion of species at risk from climate change. However climate change may provide more of an opportunity for range expansion in a greater proportion of vascular plants than bryophytes; the full risk assessment suggested 17/51 plants but only 1/14 bryophytes have an opportunity for range expansion from climate change (Figure 4), although it is worth noting that bryophytes probably have greater capacity for colonisation than vascular plants due to their spore-driven dispersal. Conversely the majority of Hymenoptera, particularly ants and wasps, have a southern distribution and were ranked as most likely to experience a high opportunity from climate change. This matches previous studies suggesting that populations of many Hymenoptera increase with warmer temperatures (Pearce-Higgins 2010, Burns et al., 2016), probably because they are thermophilic species largely constrained by temperature.

It is noteworthy that the majority (78%) of full risk assessments had poor confidence. If this is the case in Britain, which is one of the best studied and data rich parts of the world, climate change risk assessments in other parts of the world are likely to be even more uncertain. This emphasises the need for long-term monitoring and research to document and understand the impacts of climate change on biodiversity, particularly outside well-studied parts of Europe and North America (Ockendon et al., 2014). As a result, nature conservation organisations will have to integrate uncertainty and flexibility into their response to climate change. The taxa for which assessments were most robust were ~~butterflies~~¹⁰⁰⁶⁹butterflies, where 46% of

species assessments had medium or good confidence (Thomas et al., 201~~10~~), and birds, for which 35% of assessments were associated with medium or good confidence. These are the two best studied taxonomic groups in Britain with respect to the impacts of climate change on their populations (e.g. Devictor et al., 2012, Morecroft & Speakman 2015), and therefore the groups where observed changes can be more confidently attributed to climate change, where appropriate. They are also much better monitored than the other groups, with robust distribution change and annual population estimates adding to the confidence of the risk assessment. Practically speaking, the low confidence of most of the species' assessments in this study that are not of birds and butterflies, means that caution must be applied in they should only be used to assess the judging the risk that climate change poses to individual species with caution, despite that being the original aim of this work. Whilst we may have more confidence with the overall patterns of change, and how they vary between broad taxonomic groups and habitats, there are many reasons why an individual assessment for a species may not be borne out in reality. In the absence of further monitoring and research, many individual assessments should be used with an understanding of the confidence they are associated with and the uncertainty involved in projecting the future.

The main tool underpinning this assessment was climate envelope modelling. Although the results of some basic models have been criticised in the literature (see Beale et al., 2008), there is increasing evidence linking climate envelope model predictions to observed bird population changes (Stephens et al., 2016). The choice of statistical model, general circulation model (GCM) and emission scenario can have a significant impact upon the results of climate envelope models (Dormann et al., 2008, Diniz-Filho et al., 2009). Whilst we could therefore be criticised for using only one modelling approach (Beale et al., 2014) and one GCM (HADSM3), and therefore not capturing the potential full range of possible futures, we have tried to select approaches that give the most plausible futures. The Bayesian;

spatially-explicit GAM used is a significant advance on other modelling approaches, as it-by
accountings for spatially auto-correlated components of a species' distribution (Beale et al.,
2014). whilstFurthermore, in studies such as this, Baker et al., (2017) advocate using the
most suitable GCM for a particular location, which the-the HADSM3 is for the-UKGB. -and
we used newly developed-modelling techniques designed to overcome many of these
problems.-The use of additional GCMs and modelling approaches could yield alternative
projections and assessments of risk as a potential extension of this work. However, these
additional models would be unlikely to alter Ththe generality of the-conclusions from such
models are likely to be broadly realistic-our conclusions at-thefor high-level taxonomic
groups or habitat-levels, or reduce the uncertainty of the individual species assessments.
Instead, what is required is better validation of climate change risk assessment (Wheatley et
al., in press), even if associated with a high degree of uncertainty for individual species.
-The simplified risk assessment makes use of both observed and projected population and
range changes to assess risks and opportunities, allowing assessments to be moderated by the
extent to which observed and projected trends are in accordance. The full risk assessment
additionally makes use of ecological information on links between population or range
changes and climate and on potential exacerbating factors. This information is used to modify
the final risk assessment for those species, and to moderate -to-assess the degree of
confidence in the assessment. Evidence for a strong statistical link between distribution
and/or abundance and climate, or good evidence that changes are not linked to climate,
increased the confidence of the assessment. The quality of evidence around exacerbating
factors such as range or population size, interacting species, habitat availability and dispersal,
also affected the final assessment of confidence. This combination of climate envelope
modelling with ecological information to assess the degree of constraint which species are
likely to face in responding to climate change, and comparison with observed trends, is a step

forward from the basic climate envelope modelling approach, whilst taking account of some of the potential constraints on a species-by-species basis (Thomas *et al.*, 2011).

Implications for nature conservation

This analysis provides as near comprehensive an overview of how species ranges may change within a country under climate change as is currently possible. It goes beyond general principles of anticipating species range shift and provides an evidence-based assessment of the extent of change that is likely. The risk assessment indicates that, at a national level, the distributions of most species are liable to change. In the basic risk assessment only 6 out of 3048 species were identified as having both low risk and low opportunity, whilst the full assessment classified only 75 of 402 species as having both low opportunity and low risk. This is an important finding for nature conservation planning, suggesting that changing distributions are likely to become the norm, not the exception, in the coming years.

Whilst there are many species that could potentially benefit from an expanding area of potentially suitable climate, these opportunities will not be realised if individuals are ~~not~~ ~~able~~unable to disperse. ~~Natural d~~Dispersal may be limited by several factors including ~~habitat~~ fragmentation, ~~barriers of~~ unsuitable habitats or low populations sizes and other pressures affecting healthy populations. Facilitating species movement is therefore likely to be a major challenge for future species conservation. Although many taxa have shown evidence of poleward shifts in their distribution in ~~Britain~~GB (Mason *et al.* 2015), this has been partly facilitated by a network of protected sites (Thomas *et al.* 2012), whose continued conservation and expansion becomes even more important in a changing climate.

The study also provides a greater clarity on the extent of threat to some species, particularly highlighting the vulnerability of upland taxa where many species are adapted to cool, wet conditions. For those species at risk of losing areas of potentially suitable climate,

conservation actions to increase resilience (Morecroft et al., 2012), including the protection of key sites (Gillingham *et al.* 2015) and *refugia* (Suggitt et al., 2014), the maintenance of large or functional connected areas of semi-natural habitats within landscapes (Newson et al., 2014, Oliver et al., 2015, 2017) and direct management to promote *in-situ* persistence (Greenwood et al., 2015) will be important. An example of the latter is the potential to alter the management of vulnerable peatland habitats by raising water levels, likely to benefit plants, invertebrates and birds (Carroll et al., 2011, Bellamy *et al.* 2012). Reducing other non-climatic pressures on upland species may also increase the ability of their populations to cope with climate change (Pearce-Higgins & Green 2014).

The confidence assessments emphasise that individual species assessments should be treated cautiously and that conservationists need to draw upon the full range of information available before decisions are made about climate change adaptation and conservation management. Nevertheless for many species this assessment provides the main indication of potential climate change risks and opportunities and, accordingly, it can also highlight where further investigation and monitoring ~~is~~are necessary. It also emphasises the importance of planning to accommodate greater uncertainty about where species will survive and thrive in future. For site managers, this includes being aware of where their site is located in the context of the overall distribution of priority species (most simply, core, leading or trailing edges) and being prepared to adjust management priorities as situations change. To achieve this aim, the nature conservation organisations involved in this study are working to integrate these and comparable findings into their conservation practice, and to make this larger, emerging evidence base more accessible to conservation practitioners.

Acknowledgements

We would like to thank the many recorders who have contributed data and the national recording schemes and societies (listed in Table 1) that co-ordinate and collate data. In addition to the scheme organisers listed as authors, specific thanks go to Chris Preston and Oli Pescott (British Bryological Society), Mark Telfer (Ground Beetle Recording Scheme), Tony Barber (British Myriapod & Isopod Group (Centipede Recording Scheme)), John Kramer and Alan Stubbs (Crane-fly Recording Scheme), Stuart Ball, Roger Morris, Joan Childs and Ellie Rotheray (Hoverfly Recording Scheme), Keith Alexander (Soldier Beetles, Jewel Beetles and Glow-worms Recording Scheme) ~~and~~, Peter Harvey (Spider Recording Scheme). Such a project would not be possible without their contributions. We are grateful to JNCC and Defra for their support to the Biological Records Centre, and particularly to species experts who contributed their knowledge to parameterise and give feedback on the climate change risk decision framework: Mike Edwards, Dave Hubble, Paul Lee, Roger Morris and Mark Parsons. We are grateful to three anonymous reviewers for their comments on this manuscript. -This project was funded by Natural England.

References

- Akçakaya, H. R., Butchart, S. H. M., Watson, J. E. M. & Pearson, R. G. 2015. Preventing species extinctions resulting from climate change. *Nat. Clim. Change* 4, 1048-1049.
- Araújo, M.B., Alagador, D., Cabeza, M., Nogués-Bravo, D. & Thuiller, W. 2011. Climate change threatens European conservation areas. *Ecol. Lett.* 14, 484-492.
- Asher, J., Warren, [EM](#), Fox, R., Harding, P., Jeffcoate, G. & Jeffcoate, S. 2001. [The](#) Millennium Atlas of Butterflies in Britain and Ireland. OUP, Oxford.

604 Ausden, M. 2014. Climate change adaptation: putting principles into practice. *Env. Manag.*
605 54, 685-698.

606

607 Baker, D.J., Hartley, A.J., Pearce-Higgins, J.W., Jones, R.G. & Willis, S.G. 2017. Neglected
608 issues in using weather and climate information in ecology and biogeography. *Divers. &*
609 *Distrib.* 23, 329-340

610 Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, M., Brown, V.K.,
611 Butterfield, J., Buse, A., Coulson, J.C., Farrar, J., Good, J.E.G., Harrington, R., Hartley, S.,
612 Hefin Jones, T., Lindroth, R.L., Press, M.C., Symrnioudis, I., Watt, A.D. & Whittaker, J.B.
613 2002. Herbivory in global climate research: direct effects of rising temperature on insect
614 herbivores. *Glob. Ch-[ange](#) Biol.* 8, 1-16.

615 Balmer, D.E., Gillings, S., Caffrey, B.J., Swann, R.L., Downie, I.S. & Fuller, R.J. 2013. *Bird*
616 *Atlas 2007–11: the breeding and wintering birds of Britain and Ireland.* BTO Books,
617 Thetford.

618 Barbet-Massin, M., Thuiller, W. & Jiguet, F. 2012. The fate of European breeding birds
619 under climate, land-use and dispersal scenarios. *Glob. Ch-[ange](#) Biol.* 18, 881-890.

620 Bates, J.W. & Preston, C.D. 2011. Can the effects of climate change on British bryophytes be
621 distinguished from those resulting from other environmental changes? In: eds. Zoltan Tuba,
622 Nancy G. Slack & Lloyd R. Stark. *Bryophyte ecology and climate change.* Cambridge
623 University Press, New York. Pp 371-407.

624 Beale, C.M., Brewer, M.J. & Lennon, J.J. 2014. A new statistical framework for the
625 quantification of covariate associations with species distributions. *Meth. Ecol. Evol.* 5, 421-
626 432.

627 Beckmann, B.C., Purse, B.V., Roy, D.B., Roy, H.E., Sutton, P.G., Thomas, C.D. 2015. Two
628 species with an unusual combination of traits dominate responses of British grasshoppers and
629 crickets to environmental change, *PLoS ONE* 10, e0130488

630 Bellamy, P.E., Stephen, L., Maclean, I.S. & Grant, M.C. 2012. Response of blanket bog
631 vegetation to drain blocking. *Appl. Veg. Sci.* 15: 129-135.

632 Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. 2012. Impacts of
633 climate change on the future of biodiversity. *Ecol. Lett.* 15, 365-377.

634 Braithwaite, M.E. & Walker, K.J. 2012. 50 Years of Mapping the British and Irish Flora
635 1962-2012. Botanical Society of the British Isles, London.

636 Burns, F., Eaton, M.A., Barlow, K.E., Beckmann, B.C., Brereton, T., Brooks, D.R., Brown,
637 P.M.J., Fulaij, N.A., Gent, T., Henderson, I., Noble, D.G., Parsons, M., Powney, G.D., Roy,
638 H.E., Stroh, P., Walker, K., Wilkinson, J.W., Wotton, S.R. & Gregory, R.D. 2016.
639 Agricultural Management and Climatic Change Are the Major Drivers of Biodiversity
640 Change in the UK. *PLoS ONE* 11, e0151595.

641 Carroll M.J., Dennis P., Pearce-Higgins J.W. & Thomas C.D. 2011. Maintaining northern
642 peatland ecosystems in a changing climate: effects of soil moisture, drainage and drain
643 blocking on craneflies. *Glob. Change Biol.* 17, 2991-3001.

644 Devictor, V., van Swaay, C., Brereton, T., Brotons, L., Chamberlain, D., Heliölä, J.,
645 Herrando, S., Julliard, R., Kuussaari, M., Linström, Å., Reif, J., Roy, D.B., Schweiger, O.,

646 Settele, J., Stefanescu, C., Van Strien, A., Van Turnhout, C., Vermouzek, Z., De Vries,
 647 M.W., Wynhoff, I. & Jiguet, F. 2012. Differences in the climate debts of birds and butterflies
 648 at a continental scale. *Nat. Clim. Change* 2, 121-124
 649 Diniz-Filho, J.A.F., Bini, L.M., Rangel, T.F., Loyola, R.D., Hof, C., Nogués-Bravo, D. &
 650 Araújo, M.B. 2009. Partitioning and mapping uncertainties in ensembles of forecasts of
 651 species turnover under climate change. *Ecography* 32, 897-906.
 652 Dormann, C.F., McPherson, J.M., Araújo, M.B., Bivand, R., Bolliger, J., Carl, G., Davies,
 653 R.G., Hirzel, A., Jetz, W., Kissling, W.D., Kühn, I., Ohlemüller, R., Peres-Neto, P.R.,
 654 Reineking, B., Schröder, B., Schurr, F.M. & Wilson, R. 2007. Methods to account for spatial
 655 autocorrelation in the analysis of species distributional data: a review. *Ecography* 30, 609-
 656 628.
 657 Eglington, S.M., Brereton, T.M., Tayleur, C.M., Noble, D., Risely, K., Roy, D.B. & Pearce-
 658 Higgins, J.W. 2015. Patterns and causes of covariation in bird and butterfly community
 659 structure. *Land. Ecol.* 30, 1461-1472
 660 Eglington, S.M & Pearce-Higgins, J.W. 2012. Disentangling the relative importance of
 661 change in climate and land-use intensity in driving recent bird population trends. *PLoS ONE*
 662 7, e30407.
 663 Ellis, C. 2015. Implications of climate change for UK bryophytes and lichens. Technical
 664 biodiversity climate change impacts report card technical paper, 8.
 665 Foden, W.B., Butchart, S.H.B., Stuart, S.N., Vié, J.-C., Akçakaya, H.R., Angulo, A.,
 666 DeVantier, L.M., Gutsche, A., Turak, E., Cao, L., Donner, S.D., Katariya, V., Bernard, R.,
 667 Holland, R.A., Hughes, A.F., O'Hanlon, S.E., Garnett, S.T., Şekercioğlu, Ç. H., Mace, G.M.

668 2013. Identifying the World's Most Climate Change Vulnerble Species: A Systematic Trait-
669 Based Assessment of all Birds, Amphibians and Corals. PLoS ONE 8, e65427.

670 Gardali, T., Seavy, N.E., DiGaudio, R.T. & Comrack, L.A. 2012. A climate change
671 vulnerability assessment of California's at-risk birds. PLoS ONE 7, e29507.

672 Gibbons, D.W., Reid, J.B. & Chapman, R.A. 1993. The New Atlas of Breeding Birds in
673 Britain and Ireland: 1988–1991. London, UK: Poyser.

674 Gilbert, G., Brown, A.F. & Wotton, S.R. 2010. Current dynamics and predicted vulnerability
675 to sea-level rise of a threatened Bittern *Botaurus stellaris* population. Ibis 152, 580-589.

676 Green, R.E., Collingham, Y.C. Willis, S.G., Gregory, R.D. & Smith, K.W. 2008,
677 Performance of climate envelope models in retrodicting recent changes in bird population
678 size from observed climatic change. Biol. Lett. 4, 599-602.

679 Greenwood, O., Mossman, H.L., Suggitt, A.J., Curtis, R.J. & Maclean, I.D. 2016. Using *in*
680 *situ* management to conserve biodiversity under climate change. J. Appl. Ecol. 53, 885-894.

681 Hagemeijer, W.J.M. & Blair, M.J. 1997. The EBCC Atlas of European Breeding Birds.
682 T&AD POYSER.

683 Heikkinen, R. K., Luoto, M., Leikola, N., Pöyry, J., Settele, J., Kudrna, O., Marmion, M.,
684 Fronzek, S. & Thuiller, W. 2010. Assessing the vulnerability of European butterflies to
685 climate change using multiple criteria. Biod. & Conserv. 19, 695-723,

686 Hickling, R., Roy, D.B., Hill, J.K., Fox, R. & Thomas, C.D. 2006. The distributions of a wide
687 range of taxonomic groups are expanding polewards. Glob. Ch-[ange](#) Biol. 12, 450-455.

Hiley, J.R., Bradbury, R.B., Holling, M. & Thomas, C.D. 2013. Protected areas act as establishment centres for species colonizing the UK. *Proc. Roy. Soc. B.* 280, 20122310.

Hill, M. O. 2012. Local frequency as a key to interpreting species occurrence data when recording effort is not known. *Meth. Ecol. Evol.* 3, 195–205.

Hill, M. O. & Preston C. D. 2015. Disappearance of boreal plants in southern Britain: habitat loss or climate change? *Biol. J. Linn. Soc.* 115, 598-610.

~~Hiley, J.R., Bradbury, R.B., Holling, M. & Thomas, C.D. 2013. Protected Areas act as establishment centres for species colonising the United Kingdom. *Proc. Roy. Soc. B Biol. Sci.* 280, 1–8.~~

Huntley, B., Green, R.E., Collingham, Y.C. & Willis, S.G. 2007 A climatic atlas of European breeding birds. Lynx Edicions.

IUCN 2016. The IUCN Red List of Threatened Species. Version 2016-1.
www.iucnredlist.org.

Jetz, W., Wilcove, D.S. & Dobson, A.P. 2007. Projected impacts of climate and land-use change on the global diversity of birds. *PLoS Biology Biol.* 5, e157.

Lee, P. 2015. A review of the millipedes (Diplopoda), centipedes (Chilopoda) and woodlice (Isopoda) of Great Britain. Species Status No.23, Natural England Commissioned Reports, 1-170, <http://publications.naturalengland.org.uk/publication/4924476719366144>

Mace, G. M., Collar, N. J., Gaston, K. J., Hilton-Taylor, C., Akçakaya, H.R., Leader-Williams, N., Milner-Gulland, E.J. & Stuart, S.N. 2008. Quantification of extinction risk: IUCN's system for classifying threatened species. *Conserv. Biol.* 22, 1424–1442.

709 Mair, L., Hill, J.K., Fox, R., Botham, M., Brereton, T. & Thomas, C.D. 2014. Abundance
710 change and habitat availability drive species' responses to climate change. *Nat. Clim. Change*
711 4,127-131

712 Mason, S.C, Palmer, G., Fox, R., Gillings, S., Hill, J.K., Thomas, C.D. & Oliver, T.H. 2015.
713 Geographical range margins of many taxonomic groups continue to shift polewards. *Biol. J.*
714 *Linn. Soc.* 115, 586–597

715 Massimino, D., Johnston, A. & Pearce-Higgins, J.W. 2015. The geographical range of British
716 birds expands during 15 years of warming. *Bird Study* 62, 523-534

717 Mitchell, T.D., Carter, T.R., Jones, P.D., Hulme, M. & New, M. 2004. A comprehensive set
718 of high-resolution grids of monthly climate for Europe and the globe: the observed record
719 (1901-2000) and 16 scenarios (2001-2100). Tyndall Centre Working Paper 55. Pp30.

720 Murphy, J.M., Sexton, D.M.H., Jenkins, G.J., Boorman, P.M., Booth, B.B.B., Brown, C.C.,
721 Clark, R.T., Collins, M., Harris, G.R., Kendon, E.J., Betts, R.A., Brown, S.J., Howard, T. P.,
722 Humphrey, K. A., McCarthy, M. P., McDonald, R. E., Stephens, A., Wallace, C., Warren, R.,
723 Wilby, R., Wood, R. A. 2009. UK Climate Projections Science Report: Climate change
724 projections. Met Office Hadley Centre, Exeter.

725 Morecroft, M.D., Crick, H.Q.P., Duffield, S.J., Macgregor, N.A. 2012. Resilience to climate
726 change: translating principles into practice. *J. Appl. Ecol.* 49, 547-551.

727 Morecroft, M. and Speakman, L (eds.) 2015. Terrestrial Biodiversity Climate Change Impacts
728 Summary Report. Living With Environmental Change.
729 <http://www.nerc.ac.uk/research/partnerships/lwec/products/report-cards/biodiversity/>

730 Moyle, P. B., Kiernan, J. D., Crain, P. K. & Quiñones, R. M. 2013. Climate Change
731 Vulnerability of Native and Alien Freshwater Fishes of California: A Systematic Assessment
732 Approach. PLoS ONE 8, e63883

733 Natural England 2014. Climate Change Adaptation Manual – Evidence to support nature
734 conservation in a changing climate (NE546). Natural England, York.
735 (<http://publications.naturalengland.org.uk/publication/5629923804839936>)

736 Natural England 2015. Natural England’s climate change risk assessment and adaptation plan
737 (2015) (NE612). Natural England, York.
738 <http://publications.naturalengland.org.uk/publication/4599517514039296>

739 Newson, S.E., Oliver, T.H., Gillings, S., Crick, H.Q.P., Morecroft, M.D., Duffield, S.J.,
740 Macgregor, N.A. & Pearce-Higgins, J.W. 2014. Can site and landscape-scale environmental
741 attributes buffer bird populations against weather events? *Ecography* 37, 872-882

742 Ockendon, N., Baker, D.J., Carr, J.A., Almond, R.E.A., Amano, T., Bertram, E., Bradbury,
743 R.B., Bradley, C., Butchart, S.H.M., Doswald, N., Foden, W., Gill, D.J.C., Green, R.E.,
744 Sutherland, W.J., Tanner, E.V.J. & Pearce-Higgins, J.W. 2014. Mechanisms underpinning
745 climatic impacts on natural populations: altered species interactions are more important than
746 direct effects. *Glob. Ch-[ange](#) Biol.* 20, 2221-2229

747 Oliver, T. H., Smithers, R. J., Bailey, S., Walmsley, C. A. & Watts, K. 2012. A decision
748 framework for considering climate change adaptation in biodiversity conservation. *J. Appl.*
749 *Ecol.* 49, 1247-1255.

750 Oliver, T.H., Marshall, H.H., Morecroft, M.D., Brereton, T., Prudhomme, C. & Huntingford,
751 C. 2015. Interacting effects if climate change and habitat fragmentation on drought-sensitive
752 butterflies. *Nat. Clim. Ch-[ange](#)* 5, 941-945

753 Oliver, T.H., Gillings, S., Pearce-Higgins, J.W., Brereton, T., Crick, H.Q.P., Duffield, S.J.,
754 Morecroft, M.D. & Roy, D.B. 2017. Large extents of intensive land use limit community
755 reorganisation during climate warming. *Glob. Ch-[ange Biol.](#)* doi: 10.1111/gcb.13587

756 Pearce-Higgins, J.W. 2010. Using diet to assess the sensitivity of northern and upland birds
757 to climate change. *Clim. Res.* 45, 119-130.

758 Pearce-Higgins, J.W. & Green, R.E. 2014. *Birds and Climate Change: Impacts and*
759 *Conservation Responses*. Cambridge University Press, Cambridge.

760 Prentice, C., Cramer, W., Harrison, S.P., Leemans, R., Monserud, R.A. & Solomon, A.M.
761 1992. A global biome model based on plant physiology and dominance, soil properties and
762 climate. *J. Biogeog.* 19, 117-134.

763 Preston, C. D., Pearman, D. A. & Dines, T. D. 2002. *New Atlas of the British and Irish Flora*.
764 Oxford University Press, Oxford, UK.

765 Rogelj, R., Meinshausen, M. & Knutti, R. 2012. Global warming under old and new
766 scenarios using IPCC climate sensitivity range estimates. *Nat. Clim. Ch-[ange](#) 2*, 248-253.

767 Roth, T., Kohli, L., Rihm, B. & Achermann, B. 2013. Nitrogen deposition is negatively
768 related to species richness and species composition of vascular plants and bryophytes in
769 Swiss mountain grassland. *Agric. Ecosys. & Env.* 178, 121-126.

770 Roy, H. E., Adriaens, T., Isaac, N. J. B., Kenis, M., Martin, G. S., Brown, P. M. J., Hautier,
771 L., et al., 2012. Invasive alien predator causes rapid declines of native European ladybirds.
772 *Divers. & Distrib.* 18, 717-725.

773 Sexton, D.M.H., Harris, G. & Murphy, J. 2010. UKCP09: Spatially Coherent Projections.
 774 Met Office
 775 (<http://ukclimateprojections.metoffice.gov.uk/media.jsp?mediaid=88001&filetype=pdf>).
 776 Stephens, P.A., Mason, L.R., Green, R.E., Gregory, R.D., Sauer, J.R., Alison, J., Aunins, A.,
 777 Brotons, L., Butchart, S.H.M., Campedelli, T., Chodkiewicz, T., Chylarecki, P., Crowe, O.,
 778 Elts, J., Escandell, V., Foppen, R.P.B., Heldbjerg, H., Herrando, S., Husby, M., Jiguet, F.,
 779 Lehtikoinen, A., Lindström, Å., Noble, D.G., Paquet, J., Reif, J., Sattler, T., Szép, T.,
 780 Teufelbauer, N., Trautmann, S., van Strien, A.J., van Turnhout, C.A.M., Vorisek, P. & Willis,
 781 S.G. 2016. Consistent response of bird populations to climate change on two continents.
 782 Science 352, 84-87.
 783 Suggitt, A.J., Wilson, R.J., August, T.A., Beale, C.M., Bennie, J.J., Dordolo, A., Fox, R.,
 784 Hopkins, J.J., Isaac, N.J.B., Jorieux, P., Macgregor, N.A., Marcetteau, J., Massimino, D.,
 785 Morecroft, M.D., Pearce-Higgins, J.W., Walker, K., Maclean, I.M.D. 2014. Climate change
 786 refugia for the flora and fauna of England. Sheffield, Natural England, 210pp.
 787 Sutton, P.G. 2015. A review of the scarce and threatened Orthoptera and allied species of
 788 Great Britain, Orthoptera, Dictyoptera, Dermaptera, Phasmida. Species Status No. 21,
 789 Natural England Commissioned Reports, 1-51,
 790 <http://publications.naturalengland.org.uk/publication/5368778738106368>
 791 Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C.,
 792 Erasmus, B.F.N., de Siqueira, M.F., Grainger, A., Hannah, L., Hughes, L., Huntley, B., van
 793 Jaarsveld, A.S., Midgley, G.F., Miles, L., Ortega-Huerta, M.A., Peterson, A.T., Phillips, O.L.,
 794 Williams, S.E. 2004. Extinction risk from climate change. Nature 427, 145-148.

795 Thomas, C.D., Gillingham, P.K., Bradbury, R.B., Roy, D.B., Anderson, B.J., Baxter, J.M.,
796 Bourn, N.A.D., Crick, H.Q.P., Findon, R.A., Fox, R., Hodgson, J.A., Holt, A.R., Morecroft,
797 M.D., O’Hanlon, N., Oliver, T.H., Pearce-Higgins, J.W., Procter, D.A., Thomas, J.A.,
798 Walker, K.J., Walmsley, C.A., Wilson, R.J. & Hill, J.K. 2012. Protected areas facilitate
799 species’ range expansions. *Proc. Nat. Acad. Sci. USA* 109, 14063-14068.

800 Thomas, C.D., Hill, J.K., Anderson, B.J., Bailey, S., Beale, C.M., Bradbury, R.B., Bulman,
801 C.R., Crick, H.Q.P., Eigenbrod, F., Griffiths, H.M., Kunin, W.E., Oliver, T.H., Walsmley,
802 C.A., Watts, K., Worsfold, N.T. & Yardley, T. 201~~10~~. A framework for assessing threats and
803 benefits to species responding to climate change. *Meth. Ecol. & Evol.* 2, 125–142.

804 Walmsley, C.A., Smithers, R.J., Berry, P.M., Harley, M., Stevenson, M.J. & Catchpole, R.
805 2007. MONARCH – Modelling Natural Resource Responses to Climate Change – a synthesis
806 for biodiversity conservation. UKCIP, Oxford.

807 Warren, M.S., Hill, J.K., Thomas, J.A., Asher, J., Fox, R., Huntley, B., Roy, D.B., Telfer,
808 M.G., Jeffcoate, S., Harding, P., Jeffcoate, G., Willis, S.G., Greated-Davies, J.N., Moss, D.
809 & Thomas, C.D. 2001. Rapid responses of British butterflies to opposing forces of climate
810 and habitat change. *Nature* 414: 65–69.

811 Warren, R., VanDerWal, J., Price, J., Welbergen, J.A., Atkinson, I., Ramirez-Villegas, J.,
812 Osborn, T.J., Jarvis, A., Shoo, L.P., Williams, S.E. & Lowe, J. 2013. Quantifying the benefit
813 of early climate change mitigation in avoiding biodiversity loss. *Nat. Clim. Ch-ange* 3: 678-
814 682.

815 Wheatley, C.J., Beale, C.M., Bradbury, R.B., Pearce-Higgins, J.W., Critchlow, R. & Thomas,
816 C.D. in press. Climate change vulnerability for species – assessing the assessments. Glob.
817 Ch-ange Biol.

818 **Table 1.** Summary of the coverage of different species groups by this risk assessment.

Taxon	Recording Scheme	Link	Total species with distribution data	Species for which climate models converged	Species for which trends could be calculated	Conservation priority species with trends calculated
Ants	Bees, Wasps and Ants Recording Society (BWARS)	www.bwars.com	36	28	13	0
Bees	Bees, Wasps and Ants Recording Society (BWARS)	www.bwars.com	225	187	143	6
Birds	British Trust for Ornithology	www.bto.org	180	180 ¹	180	41

Bryophytes	British Bryological Society	www.britishtbryologicalsociety.org.uk	1,049	850	520	1
Carabid beetles	Ground Beetle Recording Scheme	http://www.brc.ac.uk/scheme/ground-beetle-recording-scheme	317	266	175	3
Centipedes & millipedes	British Myriapod and Isopod Group, Centipede and Millipede Recording Schemes	www.bmig.org.uk	85	66	39	0
Cerambycid Beetles	Cerambycidae Recording Scheme	http://www.coleoptera.org.uk/cerambycidae/home	52	40	0	0
Coccinellid beetles	Ladybird Recording Scheme	www.ladybird-survey.org	44	38	17	0

Craneflies	Dipterists Forum, Cranefly Recording Scheme	www.dipteristsforum.org.uk	78	64	11	0
Crickets & grasshoppers	Orthoptera Recording Scheme	www.orthoptera.org.uk	43	31	23	0
Dragonflies & damselflies	British Dragonfly Society, Dragonfly Recording Network	www.british- dragonflies.org.uk	45	35	26	0
Hoverflies	Dipterists Forum, Hoverfly Recording Scheme	www.hoverfly.org.uk	249	213	175	0
Moths	Butterfly Conservation, National Moth Recording	www.mothscount.org/text/27/ national_moth_recording_sch eme.html	668	622	422	58

	Scheme					
Soldier Beetles and allies	Soldier Beetles, Jewel Beetles and Glow-worms Recording Scheme	http://www.brc.ac.uk/scheme/soldier-beetles-jewel-beetles-and-glow-worms-recording-scheme	53	46	22	0
Spiders	Spider Recording Scheme, British Arachnological Society	www.srs.britishspiders.org.uk , www.BritishSpiders.org.uk	512	374	297	7
Vascular plants	Botanical Society of Britiain and Ireland (BSBI)	www.bsbi.org.uk	1,365	1,339 ²	852	38
Wasps	Bees, Wasps and Ants Recording	www.bwars.com	219	161	133	1

	Society (BWARS)					
TOTAL			5,220	4,540	3,048	155

819 ¹Models for two species failed to converge when built using only GB data.

820 ²For 354 of these, European data were also available.

821

Table 2. Cross-tabulation of the risks and opportunities associated with climate change for all 3048 species run through the *simplified risk assessment*, based upon a low emission B1 projection for 2070-2099 (see Tables A3 and A4 for the derivation and interpretation of each category). Values are the numbers of species in each category.

		RISK				
		VERY HIGH	HIGH	MEDIUM	LOW	TOTALS
OPPORTUNITY	LOW	25	1	7	6	39
	MEDIUM	614	157	481	84	1,336
	HIGH	24	27	358	142	551
	VERY HIGH	56	44	662	360	1,122
	TOTALS	719	229	1,508	592	3,048

Table 3. Cross-tabulation of the risks and opportunities associated with climate change for 402 species from all taxonomic groups run through the *full risk assessment*, based upon a low emission B1 projection for 2070-2099. Values in parentheses are the values for the species of conservation concern only.

		RISK				
		VERY HIGH	HIGH	MEDIUM	LOW	TOTALS
OPPORTUNITY	LOW	67 (34)	37 (11)	21 (7)	75 (27)	200 (79)
	MEDIUM	5 (3)	2 (0)	1 (0)	22 (11)	30 (14)
	HIGH	9 (4)	9 (4)	7 (3)	64 (26)	89 (37)
	VERY HIGH	8 (5)	4 (2)	5 (1)	66 (17)	83 (25)
TOTALS		89 (46)	51 (17)	34 (11)	227 (81)	402 (155)




Figure 1. Summary of the processes involved in the application of the *full risk assessment*

(simplified from Thomas et al., 2011), and how those are represented by the various stages of the process. Black boxes indicate the information required prior to risk assessment. Boxes in grey represent the steps of the *simplified risk assessment*.




Figure 2. The historic (1970-1990) probability of occurrence of an example species, *Bombus rudervarius*, (left) and the projected probability of occurrence under a medium emissions A1B scenario (right). Black crosses show actual records and coloured squares show modelled probability of occurrence.




Figure 3. Proportion of species categorised as likely to be at risk or to benefit have an opportunity for expansion from climate change, based upon a low emission B1 projection for 2070-2099, in different taxonomic groups, as assessed by the *simplified risk assessment*. The sample size of species for each group is given in Table 1.




Figure 4. Modelled *full risk assessment* score for each taxonomic group, from a GLM

containing taxonomic group and conservation status. Presented are least-square means from the model with standard errors. A score of 2 is equivalent to high opportunity, 1, medium opportunity, 0 risk and opportunity or no impact, -1 medium risk and -2 high risk.




Figure 5. Proportion of species categorised as likely to be at risk from climate change, or to face have an opportunity, using the *full risk assessment*, according to the habitat each species is associated with. The sample size for each habitat is shown by the number on each column. About half of species contributed information to more than one habitat. Habitat association information was available for the NERC species of conservation concern only. The results are based upon a low emission B1 projection for 2070-2099.

Appendix 1. Bioclimate modelling

To improve the ability of the models to describe associations with climates that are rare or novel for Britain, following Beale et al. (2014), we incorporated data from Europe. European distribution data were acquired from the European Bird Census Council (Hagemeijer & Blair 1997) and the Atlas Florae Europaeae (<http://www.luomus.fi/en/atlas-florae-europaeae-afe-distribution-vascular-plants-europe>) for birds and plants respectively. Iceland and the Faroe Islands were excluded due to their isolation from the rest of Europe, which aided model convergence. Cells east of longitude 29.99° were also excluded to avoid problems of low observer effort. This yielded 2,644 50 km cells across Europe and we identified species' presence within these from the native portions of each species range (excluding locations where European native species have been introduced).

Observed climate data on a 5 km grid from the period 1961-90 were downloaded for Britain from the UK Meteorological Office web site (<http://www.metoffice.gov.uk/climatechange/science/monitoring/ukcp09/>). These were taken to represent the baseline climate that would be used to describe observed baseline species distributions, and were aggregated to a 10 km grid for analysis. Future projection data were downloaded from the UKCP09 user interface (<http://ukclimateprojections-ui.defra.gov.uk>). To ensure that climate data were consistent across adjacent grid cells and that different climate variables were consistent within the same grid cell, we used the Spatially Coherent Projections (Sexton et al., 2010), rescaled to a 10 km resolution to model change. To represent GB climate under global temperature increases of 2°C and 4°C since pre-industrial times, we used 2070-99 for scenarios B1 and A1B respectively (<http://ukclimateprojections.defra.gov.uk/22614>), as equivalent outputs from the more recent RCP scenarios were not available at the time of this work. Projections were based on data

from 11 Regional Climate Model (RCM) ensemble members. For European-scale models, observed climate data from the period 1961-90 were acquired from the Tyndall Centre for Climate Change Research; dataset CRU TS 1.2 (Mitchell 2004). These data were averaged across the required 50 km UTM grid for Europe, and used to calculate the four bioclimatic variables outlined above. Results for the A1B scenario are presented in Tables A1 and A2 for the simplified and full risk assessments respectively.

To test the effect of incorporating European data upon projections for GB, we repeated the models for birds and vascular plants under the A1B scenario using only data for GB. The predicted changes in extent from this model were strongly correlated with predicted changes from models using the European data to generate informative priors ($r = 0.691$, $n = 532$, $P < 0.0001$). There was no significant difference in the relationship between the two measures of projected change between birds and vascular plants ($F_{1, 528} = 0.052$, $P = 0.82$). However, models based on data from GB only tended to result in fewer species showing a potential increase in range (58% forecast to increase using European data compared to 46% from GB only data) which should be remembered when interpreting the results.

Table A1. Cross-tabulation of the ~~threats~~risks and opportunities associated with the A1B climate change scenario for 2070-2099 for all species based upon the *simplified risk assessment* (see Tables A3 and A4 for the derivation of each category). Values are the numbers of species in each category.

		THREAT				RISK
		VERY HIGH	HIGH	MEDIUM	LOW	TOTALS
OPPORTUNITY	LOW	25	1	7	6	39
	MEDIUM	657	135	475	75	1,342
	HIGH	31	23	343	135	532
	VERY HIGH	44	48	677	366	1,135
	TOTALS	757	207	1502	582	3,048

Table A2. Cross-tabulation of the ~~threats~~ risks and opportunities associated with the A1B climate change scenario for 2070-2099 for all species based upon the *full risk assessment*.

Values in parentheses are the values for the NERC species of conservation concern only.

THREAT <u>RISK</u>						
		VERY HIGH	HIGH	MEDIUM	LOW	TOTAL
OPPORTUNITY	LOW	79 (37)	37 (11)	18 (6)	73 (27)	208 (81)
	MEDIUM	2 (2)	2 (0)	4 (1)	21 (8)	28 (11)
	HIGH	8 (5)	7 (3)	5 (4)	66 (27)	86 (39)
	VERY HIGH	6 (4)	3 (2)	5 (2)	66 (16)	80 (24)
	TOTALS	95 (48)	50 (16)	32 (13)	226 (78)	402 (155)

Appendix 2. Correcting for variation in observer effort.

Mixed-effects models of the probability of occurrence within ‘well-sampled’ 1km squares as a function of time, were used to measure trends in area of occupancy within the baseline historical range, whilst minimising the risk of bias from changing observer effort (Roy et al., 2012). Well-sampled squares were defined as those visited on at least three occasions when at least four species of a particular taxonomic group were recorded. Occurrence was modelled within a generalised linear mixed model with site as a random effect and year as a fixed effect using the function WSS (<https://zenodo.org/record/208752#.WFfNiFOLRQI>). The resulting coefficient of the year term was converted into a percentage decadal change in the estimated probability of occupancy. For poorly-surveyed species, the well-sampled squares we analysed are likely to be a small subset of the true historic range of the species, and so our method assumes that the frequency of species loss from these well surveyed squares accurately represents losses across the true historic range.

More recent data from 1990-2009 were analysed at the hectad resolution to document range change and assess colonisation outside of the historical range. Such analyses controlled for recorder effort, indexed as the proportion of species observed in a hectad relative to the total number of species expected, using the program FRESALO (Hill 2012) implemented in ‘sparta’ (citation here: <https://zenodo.org/record/208752#.WFfNiFOLRQI>). We selected a threshold of recorder effort of 0.25 (25% of likely species being recorded) to define an ‘adequately sampled’ square. The number of colonised hectads was calculated as the number of hectads occupied in the second time period but not in the first time period, considering only hectads that were ‘adequately sampled’ in both time periods. This was then divided by the number of ‘adequately sampled’ hectads within the home range which were occupied in

the first time period. This overall change was then converted to a decadal percentage change value.

Appendix 3. Cross-tabulation of risks and opportunities for the simplified risk assessment

Observed contractions within the historical range were compared against the magnitude of projected future contractions to assess risk from climate change, whilst observed range expansion was cross-tabulated with the magnitude of projected future range expansion to assess potential ~~threats~~risks and opportunities from climate change (Table A3). These outputs were cross-tabulated to provide an overall assessment of risks and opportunities for each species (Figure 1; Table A4).

Table A3. Cross-tabulation of likely threat risks to species (top) and opportunity for species (bottom) from climate change based on observed (rows) and projected (columns) decadal changes in range extent within the current range.

		PROJECTED DECREASE			
		>7.5 %	4.0 – 7.5 %	1.0 – 4.0 %	< 1.0 %
OBSERVED DECREASE	>7.5 %	VERY HIGH	VERY HIGH	HIGH	MEDIUM
	4.0 – 7.5 %	VERY HIGH	HIGH	HIGH	MEDIUM
	1.0 – 4.0 %	HIGH	HIGH	MEDIUM	MEDIUM
	< 1.0 %	MEDIUM	MEDIUM	MEDIUM	LOW
		PROJECTED INCREASE			
		>7.5 %	4.0 – 7.5 %	1.0 – 4.0 %	< 1.0 %
OBSERVED INCREASE	>7.5 %	VERY HIGH	VERY HIGH	HIGH	MEDIUM
	4.0 – 7.5 %	VERY HIGH	HIGH	HIGH	MEDIUM
	1.0 – 4.0 %	HIGH	HIGH	MEDIUM	MEDIUM
	< 1.0 %	MEDIUM	MEDIUM	MEDIUM	LOW

Table A4. Cross-tabulation of the risk and opportunities (Table A3) associated with climate change for each species, in order to summarise the risks (columns) and opportunities (rows) for each species.

		RISK			
		VERY HIGH	HIGH	MEDIUM	LOW
OPPORTUNITY	LOW	HIGH RISK	HIGH RISK	MEDIUM RISK	LIMITED IMPACT
	MEDIUM	HIGH RISK	MEDIUM RISK	RISKS & OPPORTUNITY	MEDIUM OPPORTUNITY
	HIGH	MEDIUM RISK	RISKS & OPPORTUNITY	MEDIUM OPPORTUNITY	HIGH OPPORTUNITY
	VERY HIGH	RISKS & OPPORTUNITY	MEDIUM OPPORTUNITY	HIGH OPPORTUNITY	HIGH OPPORTUNITY

Appendix 4. Detail of the methods and information required for full risk assessment

See Figure 1 for an overview of the risk assessment process.

Stage I.

Distribution change data (Stage I.A) were based on Atlas data (for birds) and modelling of recording scheme data held by Biological Records Centre (BRC) as described above for other taxa. Confidence in all bird trends was assessed as good, based on the high coverage and effort. For other taxa, confidence was assessed as good if the mixed model accounting for recorder effort gave a trend where the upper 80% confidence intervals were in the same impact category as the trend (i.e. we were 80% confident that any observed declines were at least that severe), unless experts highlighted that significant changes in recorder effort, taxonomy or identifiability may have contributed to these trends. The linkage between range decline and climate (Stage I.B) was assessed initially by comparison of the direction of observed and projected declines within the current range. If both were negative then this provided evidence for a link (with poor confidence), if they were contradictory in direction then this provided no evidence for a link and if evidence existed in the published literature for a relationship between climate and population or range change, this was regarded as providing evidence of a link with good confidence. In Stage I.C exacerbating factors and associated confidence were assessed from expert opinion and the scientific literature, with a published study supporting the importance of a particular impact on a species' population or distribution regarded as providing evidence with good confidence.

Stage II.

Projected declines within the current range were estimated using outputs from species distribution modelling. Confidence in these projections was assigned as 'high' where

projected and recently observed trends were consistent and the confidence intervals of bioclimatic models (median confidence interval across squares divided by the variance) were less than a threshold value of 0.02 (selected from a visual assessment of the spread of values). Confidence was assigned as medium if the confidence interval threshold was met but projected and observed trends were in opposing directions, indicating that non-climatic factors had driven recent trends. Confidence was low if the median weighted confidence interval was >0.02 , suggesting that the model projections were uncertain.

Stage III.

Stage III.A and III.B were completed as for Stages I.A and I.B, but using information about range expansion rather than contraction. The only difference was that, as described in Thomas et al., (2010¹), decadal population increases in section III.A were calculated relative to the species' status updated every decade, (as opposed to Stage I.A where changes were calculated relative to the species original status).

Stage IV.

Stage IV.A was based on bioclimatic projections of range expansion outside the current range, calculated as $(\text{newly colonised range}) / (\text{newly colonised range} + \text{current range})$. Confidence was assigned as in Stage II.A. Assessments of exacerbating factors likely to limit range expansion, and our confidence in them (Stage IV.C) were again based on expert knowledge and the literature.

Table A5. Summary of the information required at each stage of the full risk assessment(summarised and adapted from Thomas et al., 2010¹)

Stage	Data sources and criteria used
I.A.impact	<p>For bird species the decadal decline within current range was calculated from Atlas data between 1990-2010.</p> <p>For all other taxa, a mixed effects model on BRC data controlling for recorder effort was used.</p>
I.A.confidence	<p>All bird species trends were assigned good confidence.</p> <p>For other taxa, confidence was based on the C. I. from mixed model: if upper 80% C.I. overlaps the next impact category then confidence is poor, otherwise good.</p>
I.B.impact	<p>If both observed trend (I.A.) and projected trend (II.A.) are negative then linkage="Yes". Supplemented with literature review to assess additional linkages with climate</p>
I.B.confidence	<p>Poor if just assessed by comparison of observed (I.A.) and projected (II.A.) trends.</p> <p>Good if robust evidence identified by literature review.</p>
I.C.i.impact	<p>Is current extent <20 000km²? *</p> <p>Additionally for bird species only: is GB population < 10 000 individuals?</p>
I.C.i.confidence	<p>For bird species generally good.</p> <p>For other taxa: poor if just assessed by using current extent data. Good if robust</p>

evidence identified by literature review or supported by expert opinion.

IC.ii.impact Expert knowledge or evidence from literature review supporting at least one of the factors.

I.Cii.confidence Good if robust evidence from peer-reviewed literature. Poor if based on expert knowledge alone.

For birds, due to generally good understanding of the ecology of these species, experts were asked to assign the confidence level where impact was based on unpublished information.

II.A.impact Bioclimate model projected change in occupancy within current range

II.A.confidence a) Are bioclimate confidence intervals below a threshold value (see main text)?

b) Is direction of projected trends (II.A.) in same direction as observed trend (I.A.)?

For bird species: Yes to a)&b) = good, yes to a) only =medium, no to a) =poor.

For other taxa: Yes to a)&b) = good, yes to a) or b) only =medium, no to a) & b) =poor.

II.B. Not applicable

II.C.i.impact As I.C.i

II.Ci.confidence As I.C.i

II.C.ii.impact As I.C.ii

II.Cii.confidence As I.C.ii

III.A.impact	<p>For bird species: decadal increase outside previous range was calculated from Atlas data between 1990 and 2010.</p> <p>Other taxa: mixed model of BRC data of observed increases beyond species' recent historical range** controlling for recorder effort</p>
III.A.confidence	<p>All bird species trends were assigned with good confidence.</p> <p>For other taxa: the model output was compared across 3 different levels of recorder effort - if the level of recorder effort changes the impact category then confidence is poor, otherwise assigned as good.</p>
III.B.impact	<p>If both observed trend (III.A.) and projected trend (IV.A.) are positive then linkage="Yes". Supplemented with literature review to assess additional linkages with climate.</p>
III.B.confidence	<p>Poor if just assessed by comparing observed (III.A.) and projected trends (IV.A.).</p> <p>Good if robust evidence identified in literature review.</p>
III.C.	Not applicable
IV.A.impact	Bioclimate model projected change in occupancy outside the current range
IV.A.confidence	As II.A.
IV.B.	Not applicable
IV.C.i. impact	As I.C.ii
IV.C.i. confidence	As I.C.ii

IV.C.ii. impact	As I.C.ii
-----------------	-----------

IV.C.ii.confidence	As I.C.ii
--------------------	-----------

IV.C.iii. impact	As I.C.ii
------------------	-----------

IV.C.iii.confidence	As I.C.ii
---------------------	-----------

Note we occasionally changed confidence levels in Stage A (usually 1.A.) if experts highlighted concerns regarding distribution data, e.g. significant changes in recorder effort, recent taxonomic splits, issues regarding taxonomic identification etc.

*Current extent is calculated by bioclimate model: probability of a cell being occupied multiplied by the area of a cell = current extent (possible area occupied)

**Number of newly occupied cells outside the current range as a percentage of cells inside current range.

Appendix 5. Species outcomes from the simplified risk assessment

Appendix 6. Species outcomes from the full risk assessment

1 A national-scale assessment of climate change impacts on species: assessing the balance
2 of risks and opportunities for multiple taxa

3 James W. Pearce-Higgins^{1,2}, Colin M. Beale³, Tom H. Oliver^{4,5}, Tom A. August⁴, Matthew
4 Carroll^{3, 6,7}, Dario Massimino¹, Nancy Ockendon^{1,2}, Joanne Savage⁴, Christopher J.
5 Wheatley³, Malcolm A. Ausden⁶, Richard B. Bradbury^{2,6,7}, Simon J. Duffield⁸, Nicholas A.
6 Macgregor^{8, 9}, Colin J. McClean¹⁰, Michael D. Morecroft⁸, Chris D. Thomas³, Olly Watts⁶,
7 Björn C. Beckmann^{3,4,11}, Richard Fox¹², Helen E. Roy⁴, Peter G. Sutton¹¹, Kevin J. Walker¹³,
8 Humphrey Q.P. Crick⁸

9

10 ¹ *British Trust for Ornithology, The Nunnery, Thetford, Norfolk, IP24 2PU, UK*

11 ² *Conservation Science Group, Department of Zoology, University of Cambridge, Downing*
12 *Street, Cambridge CB2 3EJ, UK*

13 ³ *Department of Biology, University of York, Wentworth Way, York, YO10 5DD, UK*

14 ⁴ *Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford,*
15 *Wallingford, Oxfordshire, OX10 8BB, UK*

16 ⁵ *School of Biological Sciences, University of Reading, Whiteknights, Reading, Berkshire*
17 *RG6 6AS, UK.*

18 ⁶ *Royal Society for the Protection of Birds, The Lodge, Sandy, Beds., SG19 2DL, UK*

19 ⁷ *RSPB Centre for Conservation Science, The David Attenborough Building, Pembroke*
20 *Street, Cambridge CB2 3QZ, UK*

21 ⁸ *Natural England, Eastbrook, Shaftesbury Road, Cambridge, CB2 8DR, UK*

22 ⁹ *Durrell Institute of Conservation and Ecology (DICE), School of Anthropology and*
23 *Conservation, University of Kent, Canterbury, Kent, CT2 7NR, UK*

24 ¹⁰*Environment Department, University of York, Wentworth Way, York, YO10 5NG, UK*

25 ¹¹*Orthoptera and Allied Insects Recording Scheme of Britain and Ireland, c/o Biological*
26 *Records Centre, Centre for Ecology & Hydrology, Wallingford, OX10 8BB, UK*

27 ¹²*Butterfly Conservation, Manor Yard, East Lulworth, Wareham, Dorset, BH20 5QP, UK*

28 ¹³ *Botanical Society of Britain and Ireland (BSBI), Room 14 Bridge House, 1-2 Station*
29 *Bridge, Harrogate, North Yorkshire HG1 1SS, UK*

30

31 Corresponding author: Pearce-Higgins, J.W. Email: james.pearce-higgins@bto.org, Tel: +44
32 (0)1842 750050

33

34 Word count (9,248 including references 2,143, tables and figures 763)

35

36 **Abstract**

37 It is important for conservationists to be able to assess the risks that climate change poses to
38 species, in order to inform decision making. Using standardised and repeatable methods, we
39 present a national-scale assessment of the risks of range loss and opportunities for range
40 expansion that climate change could pose for over 3,000 plants and animals. Species were
41 selected by their occurrence in England, the primary focus of the study, but climate change
42 impacts were assessed across Great Britain, widening their geographical relevance. A basic
43 risk assessment that compared projected future changes in potential range with recently
44 observed changes classified 21% of species as being at high risk and 6% at medium risk of
45 range loss under a B1 climate change scenario. A greater number of species were classified as
46 having a medium (16%) or high (38%) opportunity to potentially expand their distribution. A
47 more comprehensive assessment, incorporating additional ecological information, including
48 potentially confounding and exacerbating factors (e.g. dispersal, habitat availability and other
49 constraints), was applied to 402 species, of which 35 % were at risk of range loss and 42 %
50 may expand their range extent. This study covers a temperate region with a significant
51 proportion of species at their poleward range limit; the balance of risks and opportunities
52 from climate change may be different elsewhere. The outcome of both risk assessments
53 varied between taxonomic groups, with bryophytes and vascular plants containing the
54 greatest proportion of species at risk from climate change. Upland habitats contained more
55 species at risk than other habitats. Whilst the overall pattern was clear, confidence was
56 generally low for individual assessments, with the exception of well-studied taxa such as
57 birds. In response to climate change, nature conservation needs to plan for changing species
58 distributions and increasing uncertainty of the future.

60 Keywords: adaptation; climate change; climate envelope; Great Britain; risk assessment;
61 vulnerability

62 **Introduction**

63 To make the best use of conservation resources, it is necessary to prioritise species for action,
64 for example according to their current status and the threats that they face. Globally, the most
65 widely adopted framework for this is the IUCN Red List which quantifies extinction risk
66 using information on the population size and range extent of a species, and the rate of change
67 in those parameters (Mace et al., 2008, IUCN 2016). Anthropogenic climate change is likely
68 to exacerbate the extinction risk of many species over the course of this century (Thomas et
69 al., 2004, Bellard et al., 2012, Warren et al., 2013, Foden et al., 2013). A number of
70 approaches have been developed to assess the potential impact of climate change on species'
71 future status (Akçakaya et al., 2015). One common approach uses species distribution models
72 (widely termed bioclimatic-envelope or climate-envelope models) to link distribution to
73 climate variables and project the likely future impact of climate change on species'
74 distributions (e.g. Thomas et al., 2004, Huntley et al., 2007, Walmsley et al., 2007, Warren et
75 al., 2013). An alternative approach is to undertake vulnerability assessments which may
76 combine a measure of future projected climate change (exposure) with ecological traits to
77 identify the sorts of species most likely to be both sensitive to and lack the capacity to adapt
78 to climate change (e.g. Gardali et al., 2012, Foden et al., 2013).

79 Vulnerability assessments have often been applied to single taxonomic groups within
80 particular regions or countries (e.g. Heikkinen et al., 2010, Barbet-Massin et al., 2012) or,
81 less commonly across a global scale (Jetz et al., 2007, Foden et al., 2013). Relatively few
82 vulnerability assessments have covered the full range of biodiversity present within a
83 particular geographical area, despite the fact that a comprehensive assessment of as many
84 taxa as possible would assist governments and conservation organisations plan and adapt to
85 climate change. Achieving such wide coverage is challenging because many assessments

require taxon-specific information or use approaches that have limited applicability to other taxa (e.g. Heikkinen et al., 2010, Gardali et al., 2012, Moyle et al., 2013). To date, it has been difficult to develop an approach which works across a range of taxa due to the different nature of ecological traits across contrasting taxonomic groups, and the variable availability of data (e.g. of species distributions, trends and traits). The strong tradition of biological recording in Britain across a wide range of taxa provides a rare opportunity to tackle this challenge.

Thomas et al., (2011) developed a framework to assess the threats and potential benefits of climate change that is applicable to a wide range of taxa. It uses bioclimatic-envelope models, combined with information on recent trends and additional ecological information, to identify the likelihood of species' range expansion and contraction, and has so far been applied to UK butterflies and some exemplar species from other taxa (Thomas et al., 2011). Here, we use a modification of this approach to undertake a climate change vulnerability assessment of more than 3,000 terrestrial and wetland species, (and in a minority of cases, species aggregates and distinctive subspecies or varieties, hereafter all termed 'species' for brevity; see methods) across 17 taxonomic groups in Britain (Table 1). This provides the first opportunity to examine how an important aspect of vulnerability to climate change varies between taxonomic groups, and between species associated with specific habitat types, for as complete a biological assemblage as currently feasible.

This study was developed as part of a wider initiative of Natural England, the government conservation agency in England, to support decision making on adaptation (Natural England 2014) and inform an adaptation plan (Natural England, 2015). It therefore focuses on species in England, the largest of the component countries within the United Kingdom (UK), but assesses the vulnerability of those species across Great Britain (GB), the single land mass

within which England is located. This ensures that the outputs are also highly relevant for Wales and Scotland, for UK organisations, and more widely.

Materials and Methods

The vulnerability assessment involved a number of steps (Figure 1) outlined below:

1. Distribution data for over 5,000 species were collated for a wide range of taxa that occur in England (Table 1).
2. Statistical models linking species' distributions to climate were used to assess the likely impacts of future climate change upon species' potential distributions.
3. Information from these projections was compared with observed changes in species distribution. By assessing recently observed changes in the context of projected future trends, a *simplified risk assessment* could be undertaken rapidly across all species.
4. For a representative subset of 402 species, additional ecological information enabled the application of the full Thomas et al., (2011) framework. By considering the potential for non-climatic factors and ecological constraints to affect species' responses to climate change, this framework produces a more comprehensive assessment (the *full risk assessment*).

Whilst the term 'risk assessment' can have specific meanings in different contexts, we follow Thomas *et al.* (2011) and use it to describe our methodology for assessing the potential risks of species decline and extirpation in parts of its current range, and opportunities that the same species may expand its distribution into other regions, both as a result of climate change. By using a combination of observed and modelled responses to climate change, the methodology

deals with the long time-scales over which species' responses to climate change are likely to occur.

Species distribution data

Species distribution data for GB were available from a range of biological recording schemes for a total of seventeen taxonomic groups (Table 1) at a hectad (10 km square) resolution. For inclusion, species had to be present in England and recorded from more than 5 hectads (the minimum required for modelling; Hickling et al., 2006). Even with this threshold the climate envelope models (described below) failed to converge for 10% of the most sparsely distributed species, giving a total of 4,540 species for which modelling was possible.

We used data from 1970-89 to represent baseline distributions prior to recent climate change, in order to minimise the risk of species' distributions being unsynchronised with the climate due to recent range shifts (Mason *et al.* 2015). For plants we used the period 1970-86; the time period (Braithwaite & Walker 2012) that most closely matched the data for other taxa. For birds the period 1988-91 was used, which coincided with a national atlas (Gibbons et al., 1993). Cells for which climate data were not available were excluded from analyses. To aid model convergence, small islands, with little data, were also excluded for all taxa apart from birds, leaving 2,561 hectads, or 2,670 for birds.

Recording effort varied between taxa, with the highest coverage for groups with well-developed and popular volunteer recording schemes such as vascular plants and birds. To avoid species' distribution models being biased as a result of limited recording effort, we used the program FRESALO (Hill, 2012) to estimate taxon-specific recorder effort in each 10 km square (see below).

Species distribution modelling

154 We used the climate envelope modelling approach of Beale et al., (2014) across all taxa
155 (Appendix 1). The approach was devised to address the problem of spatial autocorrelation in
156 large-scale species' distribution data, and applies a Bayesian, spatially explicit (Conditional
157 Autoregressive) Generalised Additive Model (GAM) to species' distribution data in order to
158 separate climatic, spatial and random components in determining the distribution of each
159 species. Four bioclimate variables were used to describe spatial variation in the climate, using
160 1961-1990 averages:

- 161 • mean temperature of the coldest month (MTCO): a measure of winter cold.
- 162 • growing degree days above 5°C (GDD5): a measure of biologically useful warmth,
163 calculated by applying a spline to mean monthly temperatures for each cell to convert
164 monthly data to daily estimates.
- 165 • the coefficient of variation of temperature (cvTemp): a measure of seasonality
- 166 • soil moisture (soilWater): a measure of moisture availability calculated following the
167 bucket model of Prentice et al., (1992), which takes inputs of temperature, rainfall, %
168 sun/cloud and soil water capacities.

169 For birds and a quarter of vascular plants, we initially constructed 50 km resolution species
170 distribution models across Europe to describe the relationship between occurrence and
171 climate using uninformative priors (i.e. with no prior knowledge of what this relationship
172 should be). Once converged, a second model was fitted to hectad data from GB using
173 informative priors from the European-scale analysis. As a result, any strong climatic signal
174 based on the European distribution would remain essentially unchanged when modelled using
175 GB data only, unless there was strong evidence for a different climatic signal within GB. In
176 cases where there was high uncertainty in the estimation of potential range shifts at a

European level, the GB model would be more heavily informed by outputs from the British component of the model. We tested for differences between both models for birds and vascular plants under the A1B scenario. Predicted changes were strongly correlated, although models based on GB only data tended to result in fewer species showing potential increases in range (Appendix 1). For species for which data from GB only were available, only the second model was run using uninformative priors.

Future climate projections for the UK were derived from UKCP09, which use outputs from an ensemble of variants of the HADSM3 climate model to produce a series of probabilistic outputs for individual climate variables for three IPCC SRES scenarios (A1F1, A1B and B1). These are regarded as the most suitable climate change projections for the UK, downscaled to a 25 km grid (Murphy et al. 2009). We considered two contrasting scenarios, the B1 scenario which is a low emissions scenario projected to lead to a c. 2°C global temperature increase by the end of this century (equivalent to RCP4.5) and the A1B scenario, that represents vulnerabilities under a medium emissions scenario of c. 4°C global warming by the end of this century (intermediate between RCP6 and RCP8.5) (Rogelj et al., 2012). As there was a strong correlation between the results of the two scenarios, we focus on the B1 results in this paper, and present the results from the A1B scenario in Appendix 1.

Simplified risk assessment

Distribution data from national schemes were used to identify post-1989 range changes within the baseline historical distribution (1970-89; or 1970-86 for plants and 1988-91 for birds, as described above), and outside this historic range (newly colonised areas). With the exception of birds, distributional changes required correction to account for variation in observer effort (Appendix 2).

Due to limited data availability across adequately sampled squares, it was not possible to use this method to produce effort-corrected observed trends for 1,492 species, leaving a total of 3,048 to which the risk assessment could be applied. Of these, 50 were species aggregates reflecting taxonomic changes over previous decades (1 bird, 3 carabid beetles, 28 bryophyte and 18 vascular plants), 123 were specific subspecies or varieties (38 bryophytes, 2 spiders and 83 vascular plants), and 80 were infraspecies, whose distribution may have been based on partial information, due to the separate recording of taxonomically distinct subspecies or related species aggregates (31 bryophytes, 1 carabid beetle and 48 vascular plants). The inclusion of this mix of taxonomic resolutions did not bias the risk assessment towards species of particular risk or opportunity categories; in a sensitivity analysis there was no significant difference in the allocation to different risk categories between ‘true’ species and these other taxonomic concepts combined, under either the B1 ($\chi^2_4 = 7.93$, $P = 0.094$) or A1B ($\chi^2_4 = 7.44$, $P = 0.11$) scenarios. We have therefore assessed all taxonomic concepts together, but for completeness also present the results for bryophytes and vascular plant species separately, excluding aggregates, subspecies and infraspecies.

Current contractions within the historical range were compared against the magnitude of projected future contractions to assess risk from climate change, whilst observed range expansion was cross-tabulated with the magnitude of projected future range expansion to assess potential risks and opportunities from climate change (Appendix 3). The highest risk or opportunity categories were reserved for those species where projected future changes were consistent with observed changes. As the simplified risk assessment may have inflated the potential risk of climate change for species which have suffered recent declines and range contractions for non-climatic reasons, for a subset of 402 species, we also undertook a full

risk assessment following the Thomas et al., (2011) framework to account for non-climatic factors and constraints.

Full risk assessment

The 402 species (including 4 subspecies / varieties and 1 infraspecies) for full assessment comprised 155 conservation priority species listed under the Section 41 of the Natural Environment and Rural Communities (NERC) Act 2006 (http://www.legislation.gov.uk/ukpga/2006/16/pdfs/ukpga_20060016_en.pdf), termed NERC species, as well as at least 13 randomly selected species from each taxonomic group. This provided a broad appraisal across taxa, while ensuring as many species of highest conservation concern as possible were included. The full risk assessment used additional ecological information on population size and range extent, the link between population and range changes to climate, and on potential exacerbating factors (e.g. range extent and population size, ecological constraints associated with habitat-availability, dispersal and species interactions) to moderate the likely risk and opportunity scores, and the overall assessment of confidence (Thomas et al., 2011). Small and range-restricted populations, or species associated with other constraints, received a higher risk score, whilst the likelihood of range expansion was reduced if habitat availability, dispersal ability and other limiting species were judged as likely to result in species achieving a lower level of range expansion than predicted by the models. This information was gathered from a literature search for each species using Google Scholar and Web of Science, supplemented by additional information from UK species experts (see Acknowledgements). The confidence associated with ecological information was regarded as good if based upon peer-reviewed literature. If it was based on expert knowledge then the expert was asked to assign the confidence level.

The full risk assessment consisted of four stages (Figure 1, Appendix 4), requiring information on observed changes in occurrence within the current range (Stage I), projected changes within the current range (Stage II), observed changes in occurrence outside the current range (Stage III) and projected changes outside the current range (Stage IV). The results of the four stages were synthesised into a single table (Table A4). The overall confidence for species ‘at risk’ was the confidence associated with the assessment of threat, while for species with an opportunity for expansion, we used the confidence associated with that. For species classed as having ‘risks and opportunities’ or ‘limited impact’, we averaged the two confidence scores.

Statistical analysis

Significant differences in the proportion of species allotted to different risk categories were tested by Chi-square, as were contrasts between taxonomic groups and between NERC and other species. Information on the broad habitat associations of the 155 NERC priority species, summarised into wetland, urban, farmland, upland woodland and coastal categories, was used to test the extent to which species’ vulnerability to climate change, from the full risk assessment, varied between habitats.

Formal differences between the results from the simplified and full risk assessments for each of the 402 species assessed using both risk assessment methods were tested by Chi-square test and by regression. For the latter, we converted the categorical risk assessment into rank scores from high risk (-2) to high opportunity (2), with both ‘risks & opportunities’ and ‘limited impact’ categories scored as 0. Scores were regressed within a generalised linear mixed model, with taxonomic identity as a random effect, using PROC MIXED in SAS v9.2. We used the same scores to test for differences in full risk assessment outcomes between different taxa, and between NERC and other species.

Results

Simplified risk assessment

Of the 3,048 species assessed, 640 were classified as being at high risk of a decline in the area of projected suitable climate under the B1 climate change scenario and 188 at medium risk (a total of 27.2% species at risk). A greater number of species were identified as likely to have a medium (486) or high (1,164) potential opportunity as a result of projected increases in the area of potentially suitable climate (totalling 54.1%; Table 2). For only 6 was limited impact predicted. These estimates of risk were similar under the A1B warming scenario ($\chi^2_3 = 2.96$, $P = 0.71$), although with slightly more species (28.1%) classified as being at risk (Appendix 1 Table A1).

The outcome of the risk assessment varied significantly between taxonomic groups ($\chi^2_{64} = 475.54$, $P < 0.0001$; excluding the limited impact category due to the small sample size). These differences remained ($\chi^2_{32} = 339.73$, $P < 0.0001$) when simply splitting species into those at risk, likely to have an opportunity, or likely to be unaffected (i.e. risks & opportunities and limited impact categories combined). The proportion of species at risk varied from 6% for wasps to 39% for vascular plants, while the proportion of species with opportunity varied from 37% for bryophytes to 90% for wasps (Figure 3). Repeating this appraisal for bryophytes and vascular plants without subspecies and infraspecies produced equivalent assessments for both (bryophytes: high opportunity 107 spp (25%), medium opportunity 48 spp. (11%), risks and opportunity 134 spp. (32%), medium risk 32 spp. (8%), high risk 102 spp. (24%); vascular plants: high opportunity 210 spp. (30%), medium opportunity 103 spp. (15%), risks and opportunity 131 spp. (19%), medium risk 59 spp. (8%), high risk 200 spp (28%)). The groups with the greatest proportion of species at risk from climate change were bryophytes and vascular plants (> 30 % in both cases), whilst a number

of groups were largely (>70 %) comprised of species for which climate change may present an opportunity for range expansion in GB (ants, bees, centipedes, coccinellid beetles and wasps).

NERC species contained slightly more ‘high risk’ and ‘medium opportunity’ species and fewer ‘high opportunity’ species than expected from the pattern across the other species ($\chi^2_4 = 10.30$, $P = 0.036$), but there was no overall difference between these two species groups when the categories were simplified to risk, opportunity or unaffected ($\chi^2_2 = 1.07$, $P = 0.58$).

Full risk assessment

Across all 402 species run through the full framework for the B1 scenario, 141 (35.1 %) were classified as being at high or medium risk of being negatively affected by climate change, compared to 168 (41.8 %) which were listed as likely to have a medium or high opportunity (Table 3). Limited impact was predicted for 19% of species. There was no significant difference from this classification of species under the A1B scenario ($\chi^2_5 = 0.94$, $P = 0.92$; Appendix 1 Table A2). The score attributed to species did not vary between NERC species and the remainder ($F_{1,384} < 0.01$, $P = 0.99$), but did vary with taxonomic group ($F_{16,384} = 3.38$, $P < 0.0001$). The lowest scores, indicating the greatest proportion of species at risk from climate change, were for bryophytes (n=14), with the highest scores for ants (n=13) and wasps (n=13), the majority of which were classed as having a high opportunity from climate change (Figure 4).

There was no significant variation overall between habitats in the frequencies of NERC species allocated to different risk categories ($\chi^2_{25} = 33.86$, $P = 0.11$). However, upland was the only habitat with a majority of species (75 %) regarded as being at risk of a decline in the

area of projected suitable climate (Figure 5), which contrasted significantly with average of 40% of species across the remaining habitats when lumped together ($\chi^2_3 = 15.59$, $P = 0.008$).

For the majority (314) of species in the full assessment, confidence was poor, for 86 it was medium and good for only two. Confidence scores differed significantly between taxonomic groups ($\chi^2_{16} = 57.23$, $P < 0.0001$), driven primarily by a greater level of confidence for bird assessments (35% of 82 assessments were accorded medium or good confidence) than for other species, where 18% of 320 assessments were classed as having medium confidence, and none good.

Simplified v Full Risk Assessment

There was a strong association between the scores using the simplified and full approaches for species assessed by both ($F_{1, 398} = 955.56$, $P < 0.0001$; $S_F = -0.33 (\pm 0.089) + 0.91 (\pm 0.029) S_S$, where S_F is the full assessment score and S_S the simplified assessment score). The scores from the two frameworks had a close to 1:1 relationship, but the intercept shows that the full assessment on average produced a lower (higher risk or lower opportunity) score by 0.33 (or one third of a category),.

Discussion

Here we present a national-level assessment of species' vulnerability to climate change, covering 3,048 species across 17 taxonomic groups. Consistently for both B1 and A1B scenarios, we found that there was a greater number of species for which potential range is projected to increase as a result of climate change than it is projected to decrease. This was particularly the case when considering the outputs for the simplified framework for all species, where over 50% were classified with a medium or high opportunity from climate change (Table 2), but also applied to 43 % of the subset of species run through the full risk

assessment framework, compared with projected negative range impacts for 35% (Table 3). This also concurs with the previously published results of the full risk assessment methodology for butterflies in GB, which used an A2 climate change scenario intermediate between the B1 and A1B scenarios used here (Thomas et al., 2011). Of 58 butterfly species, three were regarded as at high risk from climate change, three at medium risk, 10 likely to have a medium opportunity, 14 a high opportunity and 27 limited impact. If turned into rank scores and added to the results of our study, this would place butterflies intermediate between coccinelid beetles and crane flies, with a mean score of 0.52 (Figure 4). Our findings are also consistent with recently observed trends across multiple taxa in the UK where more species are regarded as being impacted positively by climate change than negatively, at least in the short-term (Burns et al., 2016).

It could be argued that by indicating that a greater number of taxa are likely to have an opportunity for range expansion in response to climate change than be at risk of range contraction, our analysis suggests that climate change will have a positive impact upon UK biodiversity. However, before considering this, it is worth noting how our findings may result from both underlying methodological constraints and inherent biological processes.

It was not possible to undertake assessments for 13% of species because there were insufficient data to generate a bioclimate model, and for a further 29% of remaining species there was insufficient information to produce effort-corrected observed trends. Given latitudinal gradients in observer (recorder) effort within the UK, with more recorders in the south than the north, it is likely that a greater proportion of unassessed species were northerly-distributed and may include species more likely to be at risk of adverse climate change impacts than to benefit. However, by selecting species from England, but using data from across GB for their assessment, this enabled us to include more northern and upland

species than we otherwise would have done had we undertaken the assessment with distribution data from England alone. In addition, it is possible that more localised and specialised species, which may be species less likely to benefit from climate change (e.g. Warren et al. 2001), were more likely to be data deficient and excluded. We did observe a significant difference between the scores of conservation priority species (many of which are rare and specialised) and others in the simplified assessment, but there was no such difference in the full assessment.

Apart from birds and vascular plants, the biodiversity data underpinning the assessment were from GB only, and in most cases our models do not capture the full range of climatically-suitable conditions in which the species can occur. A comparison of models based on GB data vs. GB + European data for birds and vascular plants, suggested that GB-only projections tended to be slightly more pessimistic than those that included European data, although the two were strongly correlated. Thus, the use of GB-only projections for most groups may have slightly inflated the projected magnitude of risk for those groups, although the assessment for vascular plants, one of the groups with the greatest proportion of species regarded as being at risk from climate change, included European data in the assessment. It is also worth noting that by including only species that currently occur in England, we did not consider the potential for new species to colonise the UK from mainland Europe as a result of climate change, which is already happening (e.g. Hiley et al., 2013). Thus our results may exclude a number of potential colonists to the UK for which climate change provides an opportunity. In other words, the outcome of the risk assessment may be scale- and context-dependent; a species projected to be at risk from climate change across mainland Europe may undergo a poleward shift and colonise the UK, where it would be regarded as having an opportunity for range expansion. This emphasises the value of undertaking assessments such as this at a range of spatial scales, which has rarely been done.

We assumed that the species distribution models describe the main relationships between species' occurrence and terrestrial climate. As we employed widely-used bioclimatic variables, this is probably reasonable for most terrestrial taxa, but for some coastal bird species which use the marine environment, where spatial patterns of changes in sea temperature and other climate related variables may differ from those on land, projections are likely to be less certain. We also have not considered potentially detrimental impacts of sea-level rise and storm surges upon vulnerable coastal habitats and species (e.g. Gilbert et al., 2010; Ausden 2014).

The full assessment that considered ecological factors known to influence observed changes in populations or distributions, or likely constraints on the impacts of climate change, was applied to 402 species only. By excluding these considerations, the simple assessment applied across all species may have over-attributed observed changes to potential impacts of climate change if they were consistent with future projections (such as for farmland birds, crickets, centipedes and millipedes; Eglington & Pearce-Higgins 2012; Beckmann *et al.* 2015; Lee 2015; Burns et al., 2016), or under-estimated the potential magnitude of future climate change impacts if observed changes were opposite to future projections as a result of non-climatic factors. Although both methodologies delivered broadly comparable results, the full assessment did increase the proportion of species projected to experience only a limited impact of climate change, and included a greater proportion of species projected to be at risk.

Finally, there is considerable uncertainty about the likely pace of any distributional shift in response to climate change. Both bird and butterfly communities appear to be lagging behind the rate of warming observed across Europe (Devictor et al., 2012, Massimino et al., 2015); less-mobile groups, such as many of the vascular plants, may well lag even more. The ability of a species to disperse will be an important constraint on the extent to which some species

can occupy any new areas of potential range in the future (Barbet-Massin et al., 2012), as will the availability of areas of potentially suitable habitat for colonisation (Thomas et al., 2012; Hiley et al., 2013) and underlying population dynamics (Mair *et al.* 2014). Although considerable uncertainty remains about the pace of these responses to climate change, these uncertainties were at least partially captured by the full risk assessment, which reduces the likelihood of opportunity as a result of climate change in species with constrained dispersal ability.

Despite the potential methodological constraints, there are good biological reasons to expect more species to be able to expand their range than be at risk of it contracting in response to climate in GB. This is because there are more southern species with potential for northward range expansion in Britain than there are northern species with southern range margins (e.g. butterflies: Asher et al., 2001; vascular plants: Preston et al., 2002; birds: Balmer et al., 2013), with strong latitudinal gradients in species' richness (e.g. Eglington et al., 2015). In combination with largely polewards shifts that are projected to occur in the distribution of a range of taxa, and are already being observed (Mason et al., 2015), this would lead to more species being likely to expand their distributions in GB, than to contract. Observations of recent trends suggest that this is already the case (Massimino et al., 2015, Burns et al., 2016). Although we assessed that fewer species would be at risk of range contraction from climate change than have an opportunity, species of certain taxonomic groups and habitats were identified as being more vulnerable than others. In particular, the full risk assessments completed for those species of conservation concern for which the required data is available suggested that species associated with upland habitat-types, where increasing temperatures might be expected to result in northwards and upwards range contraction, would be particularly vulnerable to climate change. This is consistent with the results of other studies suggesting that northern or upland birds (Green et al., 2008, Pearce-Higgins 2010), butterflies

(Thomas et al., 2011) and plants (Hill & Preson 2015) may be more vulnerable to climate change than other species. Multi-taxa assessments have found similar patterns (Walmsley et al., 2007; Araujo *et al* 2011), and there is already evidence of such impacts being observed (Morecroft & Speakman 2015). While many taxonomic groups contain some species likely to be at risk from climate change and others with the potential to expand their distribution, the balance between these two outcomes will vary with the geographical and habitat bias of that group, as well as the ecological characteristics of the species, such as voltinism, diapause strategy, migratory strategy and growth rate (Bale et al., 2002). Other climate-influenced ecological changes will also affect species abundance and distribution in future through altered species interactions (Ockendon et al., 2014).

Geographical differences may partly account for the apparent high sensitivity to future climate change of bryophytes (Figures 3 and 4), many of which have a northern or north-western distribution, associated with cool and damp conditions. Our analysis suggests that of all the taxonomic groups considered, they are likely to be one of the most at risk from a reduction in areas of suitable climate, conclusions broadly supported by Ellis (2015), who anticipated detrimental impacts of climate change on northern and upland bryophytes, although potential impacts on species associated with oceanic climates were more uncertain. Even though there is some evidence for recent warming being associated with distribution shifts in some bryophytes (Bates & Preston 2011), there are difficulties in disentangling these changes from decreases in acid and nitrogen deposition from the atmosphere (Roth et al., 2013). The basic assessment also identified vascular plants as containing a high proportion of species at risk from climate change. However climate change may provide more of an opportunity for range expansion in a greater proportion of vascular plants than bryophytes; the full risk assessment suggested 17/51 plants but only 1/14 bryophytes have an opportunity for range expansion from climate change (Figure 4), although it is worth noting that

bryophytes probably have greater capacity for colonisation than vascular plants due to their spore-driven dispersal. Conversely the majority of Hymenoptera, particularly ants and wasps, have a southern distribution and were ranked as most likely to experience a high opportunity from climate change. This matches previous studies suggesting that populations of many Hymenoptera increase with warmer temperatures (Pearce-Higgins 2010, Burns et al., 2016), probably because they are thermophilic species largely constrained by temperature.

It is noteworthy that the majority (78%) of full risk assessments had poor confidence. If this is the case in Britain, which is one of the best studied and data rich parts of the world, climate change risk assessments in other parts of the world are likely to be even more uncertain. This emphasises the need for long-term monitoring and research to document and understand the impacts of climate change on biodiversity, particularly outside well-studied parts of Europe and North America (Ockendon et al., 2014). As a result, nature conservation organisations will have to integrate uncertainty and flexibility into their response to climate change. The taxa for which assessments were most robust were butterflies, where 46% of species assessments had medium or good confidence (Thomas et al., 2011), and birds, for which 35% of assessments were associated with medium or good confidence. These are the two best studied taxonomic groups in Britain with respect to the impacts of climate change on their populations (e.g. Devictor et al., 2012, Morecroft & Speakman 2015), and therefore the groups where observed changes can be more confidently attributed to climate change, where appropriate. They are also much better monitored than the other groups, with robust distribution change and annual population estimates adding to the confidence of the risk assessment. Practically speaking, the low confidence of most of the species' assessments in this study means that caution must be applied in judging the risk that climate change poses to individual species. Whilst we may have more confidence with the overall patterns of change, and how they vary between broad taxonomic groups and habitats, there are many reasons

487 why an individual assessment for a species may not be borne out in reality. In the absence of
488 further monitoring and research, many individual assessments should be used with an
489 understanding of the confidence they are associated with and the uncertainty involved in
490 projecting the future.

491 The main tool underpinning this assessment was climate envelope modelling. Although the
492 results of some basic models have been criticised in the literature (see Beale et al., 2008),
493 there is increasing evidence linking climate envelope model predictions to observed bird
494 population changes (Stephens et al., 2016). The choice of statistical model, general
495 circulation model (GCM) and emission scenario can have a significant impact upon the
496 results of climate envelope models (Dormann et al., 2008, Diniz-Filho et al., 2009). Whilst
497 we could therefore be criticised for using only one modelling approach (Beale et al., 2014)
498 and one GCM (HADSM3), and therefore not capturing the potential full range of possible
499 futures, we have tried to select approaches that give the most plausible futures. The Bayesian
500 spatially-explicit GAM used is a significant advance on other modelling approaches, as it
501 accounts for spatially auto-correlated components of a species' distribution (Beale et al.,
502 2014). Furthermore, in studies such as this, Baker et al., (2017) advocate using the most
503 suitable GCM for a particular location, which the HADSM3 is for GB. The use of additional
504 GCMs and modelling approaches could yield alternative projections and assessments of risk
505 as a potential extension of this work. However, these additional models would be unlikely to
506 alter the generality of our conclusions for high-level taxonomic groups or habitats, or reduce
507 the uncertainty of the individual species assessments. Instead, what is required is better
508 validation of climate change risk assessment (Wheatley et al., in press).

509 The simplified risk assessment makes use of both observed and projected population and
510 range changes to assess risks and opportunities, allowing assessments to be moderated by the

511 extent to which observed and projected trends are in accordance. The full risk assessment
512 additionally makes use of ecological information on links between population or range
513 changes and climate and on potential exacerbating factors. This information is used to modify
514 the final risk assessment for those species, and to moderate the degree of confidence in the
515 assessment. Evidence for a strong statistical link between distribution and/or abundance and
516 climate, or good evidence that changes are not linked to climate, increased the confidence of
517 the assessment. The quality of evidence around exacerbating factors such as range or
518 population size, interacting species, habitat availability and dispersal, also affected the final
519 assessment of confidence. This combination of climate envelope modelling with ecological
520 information to assess the degree of constraint which species are likely to face in responding
521 to climate change, and comparison with observed trends, is a step forward from the basic
522 climate envelope modelling approach, whilst taking account of some of the potential
523 constraints on a species-by-species basis (Thomas et al., 2011).

524 *Implications for nature conservation*

525 This analysis provides as near comprehensive an overview of how species ranges may change
526 within a country under climate change as is currently possible. It goes beyond general
527 principles of anticipating species range shift and provides an evidence-based assessment of
528 the extent of change that is likely. The risk assessment indicates that, at a national level, the
529 distributions of most species are liable to change. In the basic risk assessment only 6 out of 3048
530 species were identified as having both low risk and low opportunity, whilst the full
531 assessment classified only 75 of 402 species as having both low opportunity and low risk.
532 This is an important finding for nature conservation planning, suggesting that changing
533 distributions are likely to become the norm, not the exception, in the coming years.

534 Whilst there are many species that could potentially benefit from an expanding area of
535 potentially suitable climate, these opportunities will not be realised if individuals are unable
536 to disperse. Natural dispersal may be limited by several factors including habitat
537 fragmentation, barriers of unsuitable habitats or low populations sizes and other pressures
538 affecting healthy populations. Facilitating species movement is therefore likely to be a major
539 challenge for future species conservation. Although many taxa have shown evidence of
540 poleward shifts in their distribution in GB (Mason *et al.* 2015), this has been partly facilitated
541 by a network of protected sites (Thomas *et al.* 2012), whose continued conservation and
542 expansion becomes even more important in a changing climate.

543 The study also provides a greater clarity on the extent of threat to some species, particularly
544 highlighting the vulnerability of upland taxa where many species are adapted to cool, wet
545 conditions. For those species at risk of losing areas of potentially suitable climate,
546 conservation actions to increase resilience (Morecroft *et al.*, 2012), including the protection
547 of key sites (Gillingham *et al.* 2015) and *refugia* (Suggitt *et al.*, 2014), the maintenance of
548 large or functional connected areas of semi-natural habitats within landscapes (Newson *et al.*,
549 2014, Oliver *et al.*, 2015, 2017) and direct management to promote *in-situ* persistence
550 (Greenwood *et al.*, 2015) will be important. An example of the latter is the potential to alter
551 the management of vulnerable peatland habitats by raising water levels, likely to benefit
552 plants, invertebrates and birds (Carroll *et al.*, 2011, Bellamy *et al.* 2012). Reducing other non-
553 climatic pressures on upland species may also increase the ability of their populations to cope
554 with climate change (Pearce-Higgins & Green 2014).

555 The confidence assessments emphasise that individual species assessments should be treated
556 cautiously and that conservationists need to draw upon the full range of information available
557 before decisions are made about climate change adaptation and conservation management.

Nevertheless for many species this assessment provides the main indication of potential climate change risks and opportunities and, accordingly, it can also highlight where further investigation and monitoring are necessary. It also emphasises the importance of planning to accommodate greater uncertainty about where species will survive and thrive in future. For site managers, this includes being aware of where their site is located in the context of the overall distribution of priority species (most simply, core, leading or trailing edges) and being prepared to adjust management priorities as situations change. To achieve this aim, the nature conservation organisations involved in this study are working to integrate these and comparable findings into their conservation practice, and to make this larger, emerging evidence base more accessible to conservation practitioners.

Acknowledgements

We would like to thank the many recorders who have contributed data and the national recording schemes and societies (listed in Table 1) that co-ordinate and collate data. In addition to the scheme organisers listed as authors, specific thanks go to Chris Preston and Oli Pescott (British Bryological Society), Mark Telfer (Ground Beetle Recording Scheme), Tony Barber (British Myriapod & Isopod Group (Centipede Recording Scheme)), John Kramer and Alan Stubbs (Crane-fly Recording Scheme), Stuart Ball, Roger Morris, Joan Childs and Ellie Rotheray (Hoverfly Recording Scheme), Keith Alexander (Soldier Beetles, Jewel Beetles and Glow-worms Recording Scheme) and Peter Harvey (Spider Recording Scheme). Such a project would not be possible without their contributions. We are grateful to JNCC and Defra for their support to the Biological Records Centre, and particularly to species experts who contributed their knowledge to parameterise and give feedback on the climate change risk decision framework: Mike Edwards, Dave Hubble, Paul Lee, Roger

Morris and Mark Parsons. We are grateful to three anonymous reviewers for their comments on this manuscript. This project was funded by Natural England.

References

- Akçakaya, H. R., Butchart, S. H. M., Watson, J. E. M. & Pearson, R. G. 2015. Preventing species extinctions resulting from climate change. *Nat. Clim. Change* 4, 1048-1049.
- Araújo, M.B., Alagador, D., Cabeza, M., Nogués-Bravo, D. & Thuiller, W. 2011. Climate change threatens European conservation areas. *Ecol. Lett.* 14, 484-492.
- Asher, J., Warren, M., Fox, R., Harding, P., Jeffcoate, G. & Jeffcoate, S. 2001. *The Millennium Atlas of Butterflies in Britain and Ireland*. OUP, Oxford.
- Ausden, M. 2014. Climate change adaptation: putting principles into practice. *Env. Manag.* 54, 685-698.
- Baker, D.J., Hartley, A.J., Pearce-Higgins, J.W., Jones, R.G. & Willis, S.G. 2017. Neglected issues in using weather and climate information in ecology and biogeography. *Divers. & Distrib.* 23, 329-340.
- Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, M., Brown, V.K., Butterfield, J., Buse, A., Coulson, J.C., Farrar, J., Good, J.E.G., Harrington, R., Hartley, S., Hefin Jones, T., Lindroth, R.L., Press, M.C., Symrnioudis, I., Watt, A.D. & Whittaker, J.B. 2002. Herbivory in global climate research: direct effects of rising temperature on insect herbivores. *Glob. Change Biol.* 8, 1-16.

602 Balmer, D.E., Gillings, S., Caffrey, B.J., Swann, R.L., Downie, I.S. & Fuller, R.J. 2013. Bird
603 Atlas 2007–11: the breeding and wintering birds of Britain and Ireland. BTO Books,
604 Thetford.

605 Barbet-Massin, M., Thuiller, W. & Jiguet, F. 2012. The fate of European breeding birds
606 under climate, land-use and dispersal scenarios. *Glob. Change Biol.* 18, 881-890.

607 Bates, J.W. & Preston, C.D. 2011. Can the effects of climate change on British bryophytes be
608 distinguished from those resulting from other environmental changes? In: eds. Zoltan Tuba,
609 Nancy G. Slack & Lloyd R. Stark. *Bryophyte ecology and climate change*. Cambridge
610 University Press, New York. Pp 371-407.

611 Beale, C.M., Brewer, M.J. & Lennon, J.J. 2014. A new statistical framework for the
612 quantification of covariate associations with species distributions. *Meth. Ecol. Evol.* 5, 421-
613 432.

614 Beckmann, B.C., Purse, B.V., Roy, D.B., Roy, H.E., Sutton, P.G., Thomas, C.D. 2015. Two
615 species with an unusual combination of traits dominate responses of British grasshoppers and
616 crickets to environmental change, *PLoS ONE* 10, e0130488

617 Bellamy, P.E., Stephen, L., Maclean, I.S. & Grant, M.C. 2012. Response of blanket bog
618 vegetation to drain blocking. *Appl. Veg. Sci.* 15: 129-135.

619 Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. 2012. Impacts of
620 climate change on the future of biodiversity. *Ecol. Lett.* 15, 365-377.

621 Braithwaite, M.E. & Walker, K.J. 2012. 50 Years of Mapping the British and Irish Flora
622 1962-2012. Botanical Society of the British Isles, London.

623 Burns, F., Eaton, M.A., Barlow, K.E., Beckmann, B.C., Brereton, T., Brooks, D.R., Brown,
 624 P.M.J., Fulaij, N.A., Gent, T., Henderson, I., Noble, D.G., Parsons, M., Powney, G.D., Roy,
 625 H.E., Stroh, P., Walker, K., Wilkinson, J.W., Wotton, S.R. & Gregory, R.D. 2016.
 626 Agricultural Management and Climatic Change Are the Major Drivers of Biodiversity
 627 Change in the UK. PLoS ONE 11, e0151595.

628 Carroll M.J., Dennis P., Pearce-Higgins J.W. & Thomas C.D. 2011. Maintaining northern
 629 peatland ecosystems in a changing climate: effects of soil moisture, drainage and drain
 630 blocking on crane flies. Glob. Change Biol. 17, 2991-3001.

631 Devictor, V., van Swaay, C., Brereton, T., Brotons, L., Chamberlain, D., Heliölä, J.,
 632 Herrando, S., Julliard, R., Kuussaari, M., Linström, Å., Reif, J., Roy, D.B., Schweiger, O.,
 633 Settele, J., Stefanescu, C., Van Strien, A., Van Turnhout, C., Vermouzek, Z., De Vries,
 634 M.W., Wynhoff, I. & Jiguet, F. 2012. Differences in the climate debts of birds and butterflies
 635 at a continental scale. Nat. Clim. Change 2, 121-124

636 Diniz-Filho, J.A.F., Bini, L.M., Rangel, T.F., Loyola, R.D., Hof, C., Nogués-Bravo, D. &
 637 Araújo, M.B. 2009. Partitioning and mapping uncertainties in ensembles of forecasts of
 638 species turnover under climate change. Ecography 32, 897-906.

639 Dormann, C.F., McPherson, J.M., Araújo, M.B., Bivand, R., Bolliger, J., Carl, G., Davies,
 640 R.G., Hirzel, A., Jetz, W., Kissling, W.D., Kühn, I., Ohlemüller, R., Peres-Neto, P.R.,
 641 Reineking, B., Schröder, B., Schurr, F.M. & Wilson, R. 2007. Methods to account for spatial
 642 autocorrelation in the analysis of species distributional data: a review. Ecography 30, 609-
 643 628.

644 Eglington, S.M., Brereton, T.M., Tayleur, C.M., Noble, D., Risely, K., Roy, D.B. & Pearce-
 645 Higgins, J.W. 2015. Patterns and causes of covariation in bird and butterfly community
 646 structure. *Land. Ecol.* 30, 1461-1472

647 Eglington, S.M & Pearce-Higgins, J.W. 2012. Disentangling the relative importance of
 648 change in climate and land-use intensity in driving recent bird population trends. *PLoS ONE*
 649 7, e30407.

650 Ellis, C. 2015. Implications of climate change for UK bryophytes and lichens. Technical
 651 biodiversity climate change impacts report card technical paper, 8.

652 Foden, W.B., Butchart, S.H.B., Stuart, S.N., Vié, J.-C., Akçakaya, H.R., Angulo, A.,
 653 DeVantier, L.M., Gutsche, A., Turak, E., Cao, L., Donner, S.D., Katariya, V., Bernard, R.,
 654 Holland, R.A., Hughes, A.F., O'Hanlon, S.E., Garnett, S.T., Şekercioğlu, Ç. H., Mace, G.M.
 655 2013. Identifying the World's Most Climate Change Vulnerable Species: A Systematic Trait-
 656 Based Assessment of all Birds, Amphibians and Corals. *PLoS ONE* 8, e65427.

657 Gardali, T., Seavy, N.E., DiGaudio, R.T. & Comrack, L.A. 2012. A climate change
 658 vulnerability assessment of California's at-risk birds. *PLoS ONE* 7, e29507.

659 Gibbons, D.W., Reid, J.B. & Chapman, R.A. 1993. *The New Atlas of Breeding Birds in*
 660 *Britain and Ireland: 1988–1991*. London, UK: Poyser.

661 Gilbert, G., Brown, A.F. & Wotton, S.R. 2010. Current dynamics and predicted vulnerability
 662 to sea-level rise of a threatened Bittern *Botaurus stellaris* population. *Ibis* 152, 580-589.

663 Green, R.E., Collingham, Y.C., Willis, S.G., Gregory, R.D. & Smith, K.W. 2008,
 664 Performance of climate envelope models in retrodicting recent changes in bird population
 665 size from observed climatic change. *Biol. Lett.* 4, 599-602.

666 Greenwood, O., Mossman, H.L., Suggitt, A.J., Curtis, R.J. & Maclean, I.D. 2016. Using *in*
667 *situ* management to conserve biodiversity under climate change. J. Appl. Ecol. 53, 885-894.

668 Hagemeyer, W.J.M. & Blair, M.J. 1997. The EBCC Atlas of European Breeding Birds.
669 T&AD POYSER.

670 Heikkinen, R. K., Luoto, M., Leikola, N., Pöyry, J., Settele, J., Kudrna, O., Marmion, M.,
671 Fronzek, S. & Thuiller, W. 2010. Assessing the vulnerability of European butterflies to
672 climate change using multiple criteria. Biod. & Conserv. 19, 695-723,

673 Hickling, R., Roy, D.B., Hill, J.K., Fox, R. & Thomas, C.D. 2006. The distributions of a wide
674 range of taxonomic groups are expanding polewards. Glob. Change Biol. 12, 450-455.

675 Hiley, J.R., Bradbury, R.B., Holling, M. & Thomas, C.D. 2013. Protected areas act as
676 establishment centres for species colonizing the UK. Proc. Roy. Soc. B. 280, 20122310.

677 Hill, M. O. 2012. Local frequency as a key to interpreting species occurrence data when
678 recording effort is not known. Meth. Ecol. Evol. 3, 195–205.

679 Hill, M. O. & Preston C. D. 2015. Disappearance of boreal plants in southern Britain: habitat
680 loss or climate change? Biol. J. Linn. Soc. 115, 598-610.

681 Huntley, B., Green, R.E., Collingham, Y.C. & Willis, S.G. 2007 A climatic atlas of European
682 breeding birds. Lynx Edicions.

683 IUCN 2016. The IUCN Red List of Threatened Species. Version 2016-1.
684 www.iucnredlist.org.

685 Jetz, W., Wilcove, D.S. & Dobson, A.P. 2007. Projected impacts of climate and land-use
686 change on the global diversity of birds. PLoS Biol. 5, e157.

687 Lee, P. 2015. A review of the millipedes (Diplopoda), centipedes (Chilopoda) and woodlice
688 (Isopoda) of Great Britain. Species Status No.23, Natural England Commissioned Reports, 1-
689 170, <http://publications.naturalengland.org.uk/publication/4924476719366144>

690 Mace, G. M., Collar, N. J., Gaston, K. J., Hilton-Taylor, C., Akçakaya, H.R., Leader-
691 Williams, N., Milner-Gulland, E.J. & Stuart, S.N. 2008. Quantification of extinction risk:
692 IUCN's system for classifying threatened species. *Conserv. Biol.* 22, 1424–1442.

693 Mair, L., Hill, J.K., Fox, R., Botham, M., Brereton, T. & Thomas, C.D. 2014. Abundance
694 change and habitat availability drive species' responses to climate change. *Nat. Clim. Change*
695 4,127-131

696 Mason, S.C, Palmer, G., Fox, R., Gillings, S., Hill, J.K., Thomas, C.D. & Oliver, T.H. 2015.
697 Geographical range margins of many taxonomic groups continue to shift polewards. *Biol. J.*
698 *Linn. Soc.* 115, 586–597

699 Massimino, D., Johnston, A. & Pearce-Higgins, J.W. 2015. The geographical range of British
700 birds expands during 15 years of warming. *Bird Study* 62, 523-534

701 Mitchell, T.D., Carter, T.R., Jones, P.D., Hulme, M. & New, M. 2004. A comprehensive set
702 of high-resolution grids of monthly climate for Europe and the globe: the observed record
703 (1901-2000) and 16 scenarios (2001-2100). Tyndall Centre Working Paper 55. Pp30.

704 Murphy, J.M., Sexton, D.M.H., Jenkins, G.J., Boorman, P.M., Booth, B.B.B., Brown, C.C.,
705 Clark, R.T., Collins, M., Harris, G.R., Kendon, E.J., Betts, R.A., Brown, S.J., Howard, T. P.,
706 Humphrey, K. A., McCarthy, M. P., McDonald, R. E., Stephens, A., Wallace, C., Warren, R.,
707 Wilby, R., Wood, R. A. 2009. UK Climate Projections Science Report: Climate change
708 projections. Met Office Hadley Centre, Exeter.

709 Morecroft, M.D., Crick, H.Q.P., Duffield, S.J., Macgregor, N.A. 2012. Resilience to climate
710 change: translating principles into practice. *J. Appl. Ecol.* 49, 547-551.

711 Morecroft, M. and Speakman, L (eds.) 2015. Terrestrial Biodiversity Climate Change Impacts
712 Summary Report. Living With Environmental Change.
713 <http://www.nerc.ac.uk/research/partnerships/lwec/products/report-cards/biodiversity/>

714 Moyle, P. B., Kiernan, J. D., Crain, P. K. & Quiñones, R. M. 2013. Climate Change
715 Vulnerability of Native and Alien Freshwater Fishes of California: A Systematic Assessment
716 Approach. *PLoS ONE* 8, e63883

717 Natural England 2014. Climate Change Adaptation Manual – Evidence to support nature
718 conservation in a changing climate (NE546). Natural England, York.
719 (<http://publications.naturalengland.org.uk/publication/5629923804839936>)

720 Natural England 2015. Natural England’s climate change risk assessment and adaptation plan
721 (2015) (NE612). Natural England, York.
722 <http://publications.naturalengland.org.uk/publication/4599517514039296>

723 Newson, S.E., Oliver, T.H., Gillings, S., Crick, H.Q.P., Morecroft, M.D., Duffield, S.J.,
724 Macgregor, N.A. & Pearce-Higgins, J.W. 2014. Can site and landscape-scale environmental
725 attributes buffer bird populations against weather events? *Ecography* 37, 872-882

726 Ockendon, N., Baker, D.J., Carr, J.A., Almond, R.E.A., Amano, T., Bertram, E., Bradbury,
727 R.B., Bradley, C., Butchart, S.H.M., Doswald, N., Foden, W., Gill, D.J.C., Green, R.E.,
728 Sutherland, W.J., Tanner, E.V.J. & Pearce-Higgins, J.W. 2014. Mechanisms underpinning
729 climatic impacts on natural populations: altered species interactions are more important than
730 direct effects. *Glob. Change Biol.* 20, 2221-2229

731 Oliver, T. H., Smithers, R. J., Bailey, S., Walmsley, C. A. & Watts, K. 2012. A decision
732 framework for considering climate change adaptation in biodiversity conservation. *J. Appl.*
733 *Ecol.* 49, 1247-1255.

734 Oliver, T.H., Marshall, H.H., Morecroft, M.D., Brereton, T., Prudhomme, C. & Huntingford,
735 C. 2015. Interacting effects of climate change and habitat fragmentation on drought-sensitive
736 butterflies. *Nat. Clim. Change* 5, 941-945

737 Oliver, T.H., Gillings, S., Pearce-Higgins, J.W., Brereton, T., Crick, H.Q.P., Duffield, S.J.,
738 Morecroft, M.D. & Roy, D.B. 2017. Large extents of intensive land use limit community
739 reorganisation during climate warming. *Glob. Change Biol.* doi: 10.1111/gcb.13587

740 Pearce-Higgins, J.W. 2010. Using diet to assess the sensitivity of northern and upland birds
741 to climate change. *Clim. Res.* 45, 119-130.

742 Pearce-Higgins, J.W. & Green, R.E. 2014. *Birds and Climate Change: Impacts and*
743 *Conservation Responses*. Cambridge University Press, Cambridge.

744 Prentice, C., Cramer, W., Harrison, S.P., Leemans, R., Monserud, R.A. & Solomon, A.M.
745 1992. A global biome model based on plant physiology and dominance, soil properties and
746 climate. *J. Biogeog.* 19, 117-134.

747 Preston, C. D., Pearman, D. A. & Dines, T. D. 2002. *New Atlas of the British and Irish Flora*.
748 Oxford University Press, Oxford, UK.

749 Rogelj, R., Meinshausen, M. & Knutti, R. 2012. Global warming under old and new
750 scenarios using IPCC climate sensitivity range estimates. *Nat. Clim. Change* 2, 248-253.

751 Roth, T., Kohli, L., Rihm, B. & Achermann, B. 2013. Nitrogen deposition is negatively
752 related to species richness and species composition of vascular plants and bryophytes in
753 Swiss mountain grassland. *Agric. Ecosys. & Env.* 178, 121-126.

754 Roy, H. E., Adriaens, T., Isaac, N. J. B., Kenis, M., Martin, G. S., Brown, P. M. J., Hautier,
755 L., et al., 2012. Invasive alien predator causes rapid declines of native European ladybirds.
756 *Divers. & Distrib.* 18, 717-725.

757 Sexton, D.M.H., Harris, G. & Murphy, J. 2010. UKCP09: Spatially Coherent Projections.
758 Met Office
759 (<http://ukclimateprojections.metoffice.gov.uk/media.jsp?mediaid=88001&filetype=pdf>).

760 Stephens, P.A., Mason, L.R., Green, R.E., Gregory, R.D., Sauer, J.R., Alison, J., Aunins, A.,
761 Brotons, L., Butchart, S.H.M., Campedelli, T., Chodkiewicz, T., Chylarecki, P., Crowe, O.,
762 Elts, J., Escandell, V., Foppen, R.P.B., Heldbjerg, H., Herrando, S., Husby, M., Jiguet, F.,
763 Lehtikoinen, A., Lindström, Å., Noble, D.G., Paquet, J., Reif, J., Sattler, T., Szép, T.,
764 Teufelbauer, N., Trautmann, S., van Strien, A.J., van Turnhout, C.A.M., Vorisek, P. & Willis,
765 S.G. 2016. Consistent response of bird populations to climate change on two continents.
766 *Science* 352, 84-87.

767 Suggitt, A.J., Wilson, R.J., August, T.A., Beale, C.M., Bennie, J.J., Dordolo, A., Fox, R.,
768 Hopkins, J.J., Isaac, N.J.B., Jorieux, P., Macgregor, N.A., Marcetteau, J., Massimino, D.,
769 Morecroft, M.D., Pearce-Higgins, J.W., Walker, K., Maclean, I.M.D. 2014. Climate change
770 refugia for the flora and fauna of England. Sheffield, Natural England, 210pp.

771 Sutton, P.G. 2015. A review of the scarce and threatened Orthoptera and allied species of
772 Great Britain, Orthoptera, Dictyoptera, Dermaptera, Phasmida. *Species Status No.* 21,

773 Natural England Commissioned Reports, 1-51,
 774 <http://publications.naturalengland.org.uk/publication/5368778738106368>
 775 Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C.,
 776 Erasmus, B.F.N., de Siqueira, M.F., Grainger, A., Hannah, L., Hughes, L., Huntley, B., van
 777 Jaarsveld, A.S., Midgley, G.F., Miles, L., Ortega-Huerta, M.A., Peterson, A.T., Phillips, O.L.,
 778 Williams, S.E. 2004. Extinction risk from climate change. *Nature* 427, 145-148.
 779 Thomas, C.D., Gillingham, P.K., Bradbury, R.B., Roy, D.B., Anderson, B.J., Baxter, J.M.,
 780 Bourn, N.A.D., Crick, H.Q.P., Findon, R.A., Fox, R., Hodgson, J.A., Holt, A.R., Morecroft,
 781 M.D., O'Hanlon, N., Oliver, T.H., Pearce-Higgins, J.W., Procter, D.A., Thomas, J.A.,
 782 Walker, K.J., Walmsley, C.A., Wilson, R.J. & Hill, J.K. 2012. Protected areas facilitate
 783 species' range expansions. *Proc. Nat. Acad. Sci. USA* 109, 14063-14068.
 784 Thomas, C.D., Hill, J.K., Anderson, B.J., Bailey, S., Beale, C.M., Bradbury, R.B., Bulman,
 785 C.R., Crick, H.Q.P., Eigenbrod, F., Griffiths, H.M., Kunin, W.E., Oliver, T.H., Walsmsley,
 786 C.A., Watts, K., Worsfold, N.T. & Yardley, T. 2011. A framework for assessing threats and
 787 benefits to species responding to climate change. *Meth. Ecol. & Evol.* 2, 125-142.
 788 Walmsley, C.A., Smithers, R.J., Berry, P.M., Harley, M., Stevenson, M.J. & Catchpole, R.
 789 2007. MONARCH – Modelling Natural Resource Responses to Climate Change – a synthesis
 790 for biodiversity conservation. UKCIP, Oxford.
 791 Warren, M.S., Hill, J.K., Thomas, J.A., Asher, J., Fox, R., Huntley, B., Roy, D.B., Telfer,
 792 M.G., Jeffcoate, S., Harding, P., Jeffcoate, G., Willis, S.G., Greatorex-Davies, J.N., Moss, D.
 793 & Thomas, C.D. 2001. Rapid responses of British butterflies to opposing forces of climate
 794 and habitat change. *Nature* 414, 65-69.

795 Warren, R., VanDerWal, J., Price, J., Welbergen, J.A., Atkinson, I., Ramirez-Villegas, J.,
796 Osborn, T.J., Jarvis, A., Shoo, L.P., Williams, S.E. & Lowe, J. 2013. Quantifying the benefit
797 of early climate change mitigation in avoiding biodiversity loss. *Nat. Clim. Change* 3, 678-
798 682.

799 Wheatley, C.J., Beale, C.M., Bradbury, R.B., Pearce-Higgins, J.W., Critchlow, R. & Thomas,
800 C.D. in press. Climate change vulnerability for species – assessing the assessments. *Glob.*
801 *Change Biol.*

802 **Table 1.** Summary of the coverage of different species groups by this risk assessment.

Taxon	Recording Scheme	Link	Total species with distribution data	Species for which climate models converged	Species for which trends could be calculated	Conservation priority species with trends calculated
Ants	Bees, Wasps and Ants Recording Society (BWARS)	www.bwars.com	36	28	13	0
Bees	Bees, Wasps and Ants Recording Society (BWARS)	www.bwars.com	225	187	143	6
Birds	British Trust for Ornithology	www.bto.org	180	180 ¹	180	41

Bryophytes	British Bryological Society	www.britishtobryologicalsociety.org.uk	1,049	850	520	1
Carabid beetles	Ground Beetle Recording Scheme	http://www.brc.ac.uk/scheme/ground-beetle-recording-scheme	317	266	175	3
Centipedes & millipedes	British Myriapod and Isopod Group, Centipede and Millipede Recording Schemes	www.bmig.org.uk	85	66	39	0
Cerambycid Beetles	Cerambycidae Recording Scheme	http://www.coleoptera.org.uk/cerambycidae/home	52	40	0	0
Coccinellid beetles	Ladybird Recording Scheme	www.ladybird-survey.org	44	38	17	0

Craneflies	Dipterists Forum, Cranefly Recording Scheme	www.dipteristsforum.org.uk	78	64	11	0
Crickets & grasshoppers	Orthoptera Recording Scheme	www.orthoptera.org.uk	43	31	23	0
Dragonflies & damselflies	British Dragonfly Society, Dragonfly Recording Network	www.british- dragonflies.org.uk	45	35	26	0
Hoverflies	Dipterists Forum, Hoverfly Recording Scheme	www.hoverfly.org.uk	249	213	175	0
Moths	Butterfly Conservation, National Moth Recording	www.mothscount.org/text/27/ national_moth_recording_sch eme.html	668	622	422	58

	Scheme					
Soldier Beetles and allies	Soldier Beetles, Jewel Beetles and Glow-worms Recording Scheme	http://www.brc.ac.uk/scheme/soldier-beetles-jewel-beetles-and-glow-worms-recording-scheme	53	46	22	0
Spiders	Spider Recording Scheme, British Arachnological Society	www.srs.britishspiders.org.uk , www.BritishSpiders.org.uk	512	374	297	7
Vascular plants	Botanical Society of Britiain and Ireland (BSBI)	www.bsbi.org.uk	1,365	1,339 ²	852	38
Wasps	Bees, Wasps and Ants Recording	www.bwars.com	219	161	133	1

	Society (BWARS)					
TOTAL			5,220	4,540	3,048	155

803 ¹Models for two species failed to converge when built using only GB data.

804 ²For 354 of these, European data were also available.

805

Table 2. Cross-tabulation of the risks and opportunities associated with climate change for all 3048 species run through the *simplified risk assessment*, based upon a low emission B1 projection for 2070-2099 (see Tables A3 and A4 for the derivation and interpretation of each category). Values are the numbers of species in each category.

		RISK				
		VERY HIGH	HIGH	MEDIUM	LOW	TOTALS
OPPORTUNITY	LOW	25	1	7	6	39
	MEDIUM	614	157	481	84	1,336
	HIGH	24	27	358	142	551
	VERY HIGH	56	44	662	360	1,122
	TOTALS	719	229	1,508	592	3,048

Table 3. Cross-tabulation of the risks and opportunities associated with climate change for 402 species from all taxonomic groups run through the *full risk assessment*, based upon a low emission B1 projection for 2070-2099. Values in parentheses are the values for the species of conservation concern only.

		RISK				
		VERY HIGH	HIGH	MEDIUM	LOW	TOTALS
OPPORTUNITY	LOW	67 (34)	37 (11)	21 (7)	75 (27)	200 (79)
	MEDIUM	5 (3)	2 (0)	1 (0)	22 (11)	30 (14)
	HIGH	9 (4)	9 (4)	7 (3)	64 (26)	89 (37)
	VERY HIGH	8 (5)	4 (2)	5 (1)	66 (17)	83 (25)
TOTALS		89 (46)	51 (17)	34 (11)	227 (81)	402 (155)




Figure 1. Summary of the processes involved in the application of the *full risk assessment* (simplified from Thomas et al., 2011), and how those are represented by the various stages of the process. Black boxes indicate the information required prior to risk assessment. Boxes in grey represent the steps of the *simplified risk assessment*.




Figure 2. The historic (1970-1990) probability of occurrence of an example species, *Bombus ruderarius*, (left) and the projected probability of occurrence under a medium emissions A1B scenario (right). Black crosses show actual records and coloured squares show modelled probability of occurrence.




Figure 3. Proportion of species categorised as likely to be at risk or to have an opportunity for expansion from climate change, based upon a low emission B1 projection for 2070-2099, in different taxonomic groups, as assessed by the *simplified risk assessment*. The sample size of species for each group is given in Table 1.




Figure 4. Modelled *full risk assessment* score for each taxonomic group, from a GLM

containing taxonomic group and conservation status. Presented are least-square means from the model with standard errors. A score of 2 is equivalent to high opportunity, 1, medium opportunity, 0 risk and opportunity or no impact, -1 medium risk and -2 high risk.




Figure 5. Proportion of species categorised as likely to be at risk from climate change, or to have an opportunity, using the *full risk assessment*, according to the habitat each species is associated with. The sample size for each habitat is shown by the number on each column. About half of species contributed information to more than one habitat. Habitat association information was available for the NERC species of conservation concern only. The results are based upon a low emission B1 projection for 2070-2099.

Appendix 1. Bioclimate modelling

To improve the ability of the models to describe associations with climates that are rare or novel for Britain, following Beale et al. (2014), we incorporated data from Europe. European distribution data were acquired from the European Bird Census Council (Hagemeijer & Blair 1997) and the Atlas Florae Europaeae (<http://www.luomus.fi/en/atlas-florae-europaeae-afe-distribution-vascular-plants-europe>) for birds and plants respectively. Iceland and the Faroe Islands were excluded due to their isolation from the rest of Europe, which aided model convergence. Cells east of longitude 29.99° were also excluded to avoid problems of low observer effort. This yielded 2,644 50 km cells across Europe and we identified species' presence within these from the native portions of each species range (excluding locations where European native species have been introduced).

Observed climate data on a 5 km grid from the period 1961-90 were downloaded for Britain from the UK Meteorological Office web site (<http://www.metoffice.gov.uk/climatechange/science/monitoring/ukcp09/>). These were taken to represent the baseline climate that would be used to describe observed baseline species distributions, and were aggregated to a 10 km grid for analysis. Future projection data were downloaded from the UKCP09 user interface (<http://ukclimateprojections-ui.defra.gov.uk>). To ensure that climate data were consistent across adjacent grid cells and that different climate variables were consistent within the same grid cell, we used the Spatially Coherent Projections (Sexton et al., 2010), rescaled to a 10 km resolution to model change. To represent GB climate under global temperature increases of 2°C and 4°C since pre-industrial times, we used 2070-99 for scenarios B1 and A1B respectively (<http://ukclimateprojections.defra.gov.uk/22614>), as equivalent outputs from the more recent RCP scenarios were not available at the time of this work. Projections were based on data

from 11 Regional Climate Model (RCM) ensemble members. For European-scale models, observed climate data from the period 1961-90 were acquired from the Tyndall Centre for Climate Change Research; dataset CRU TS 1.2 (Mitchell 2004). These data were averaged across the required 50 km UTM grid for Europe, and used to calculate the four bioclimatic variables outlined above. Results for the A1B scenario are presented in Tables A1 and A2 for the simplified and full risk assessments respectively.

To test the effect of incorporating European data upon projections for GB, we repeated the models for birds and vascular plants under the A1B scenario using only data for GB. The predicted changes in extent from this model were strongly correlated with predicted changes from models using the European data to generate informative priors ($r = 0.691$, $n = 532$, $P < 0.0001$). There was no significant difference in the relationship between the two measures of projected change between birds and vascular plants ($F_{1, 528} = 0.052$, $P = 0.82$). However, models based on data from GB only tended to result in fewer species showing a potential increase in range (58% forecast to increase using European data compared to 46% from GB only data) which should be remembered when interpreting the results.

Table A1. Cross-tabulation of the risks and opportunities associated with the A1B climate change scenario for 2070-2099 for all species based upon the *simplified risk assessment* (see Tables A3 and A4 for the derivation of each category). Values are the numbers of species in each category.

		RISK				
		VERY HIGH	HIGH	MEDIUM	LOW	TOTALS
OPPORTUNITY	LOW	25	1	7	6	39
	MEDIUM	657	135	475	75	1,342
	HIGH	31	23	343	135	532
	VERY HIGH	44	48	677	366	1,135
TOTALS		757	207	1502	582	3,048

Table A2. Cross-tabulation of the risks and opportunities associated with the A1B climate change scenario for 2070-2099 for all species based upon the *full risk assessment*. Values in parentheses are the values for the NERC species of conservation concern only.

		RISK				TOTAL S
		VERY HIGH	HIGH	MEDIUM	LOW	
OPPORTUNITY	LOW	79 (37)	37 (11)	18 (6)	73 (27)	208 (81)
	MEDIUM	2 (2)	2 (0)	4 (1)	21 (8)	28 (11)
	HIGH	8 (5)	7 (3)	5 (4)	66 (27)	86 (39)
	VERY HIGH	6 (4)	3 (2)	5 (2)	66 (16)	80 (24)
TOTALS		95 (48)	50 (16)	32 (13)	226 (78)	402 (155)

Appendix 2. Correcting for variation in observer effort.

Mixed-effects models of the probability of occurrence within ‘well-sampled’ 1km squares as a function of time, were used to measure trends in area of occupancy within the baseline historical range, whilst minimising the risk of bias from changing observer effort (Roy et al., 2012). Well-sampled squares were defined as those visited on at least three occasions when at least four species of a particular taxonomic group were recorded. Occurrence was modelled within a generalised linear mixed model with site as a random effect and year as a fixed effect using the function WSS (<https://zenodo.org/record/208752#.WFfNiFOLRQI>). The resulting coefficient of the year term was converted into a percentage decadal change in the estimated probability of occupancy. For poorly-surveyed species, the well-sampled squares we analysed are likely to be a small subset of the true historic range of the species, and so our method assumes that the frequency of species loss from these well surveyed squares accurately represents losses across the true historic range.

More recent data from 1990-2009 were analysed at the hectad resolution to document range change and assess colonisation outside of the historical range. Such analyses controlled for recorder effort, indexed as the proportion of species observed in a hectad relative to the total number of species expected, using the program FRESALO (Hill 2012) implemented in ‘sparta’ (citation here: <https://zenodo.org/record/208752#.WFfNiFOLRQI>). We selected a threshold of recorder effort of 0.25 (25% of likely species being recorded) to define an ‘adequately sampled’ square. The number of colonised hectads was calculated as the number of hectads occupied in the second time period but not in the first time period, considering only hectads that were ‘adequately sampled’ in both time periods. This was then divided by the number of ‘adequately sampled’ hectads within the home range which were occupied in

the first time period. This overall change was then converted to a decadal percentage change value.

Appendix 3. Cross-tabulation of risks and opportunities for the simplified risk assessment

Observed contractions within the historical range were compared against the magnitude of projected future contractions to assess risk from climate change, whilst observed range expansion was cross-tabulated with the magnitude of projected future range expansion to assess potential risks and opportunities from climate change (Table A3). These outputs were cross-tabulated to provide an overall assessment of risks and opportunities for each species (Figure 1; Table A4).

Table A3. Cross-tabulation of likely risks to species (top) and opportunity for species (bottom) from climate change based on observed (rows) and projected (columns) decadal changes in range extent within the current range.

		PROJECTED DECREASE			
		>7.5 %	4.0 – 7.5 %	1.0 – 4.0 %	< 1.0 %
OBSERVED DECREASE	>7.5 %	VERY HIGH	VERY HIGH	HIGH	MEDIUM
	4.0 – 7.5 %	VERY HIGH	HIGH	HIGH	MEDIUM
	1.0 – 4.0 %	HIGH	HIGH	MEDIUM	MEDIUM
	< 1.0 %	MEDIUM	MEDIUM	MEDIUM	LOW
		PROJECTED INCREASE			
		>7.5 %	4.0 – 7.5 %	1.0 – 4.0 %	< 1.0 %
OBSERVED INCREASE	>7.5 %	VERY HIGH	VERY HIGH	HIGH	MEDIUM
	4.0 – 7.5 %	VERY HIGH	HIGH	HIGH	MEDIUM
	1.0 – 4.0 %	HIGH	HIGH	MEDIUM	MEDIUM
	< 1.0 %	MEDIUM	MEDIUM	MEDIUM	LOW

Table A4. Cross-tabulation of the risk and opportunities (Table A3) associated with climate change for each species, in order to summarise the risks (columns) and opportunities (rows) for each species.

		RISK			
		VERY HIGH	HIGH	MEDIUM	LOW
OPPORTUNITY	LOW	HIGH RISK	HIGH RISK	MEDIUM RISK	LIMITED IMPACT
	MEDIUM	HIGH RISK	MEDIUM RISK	RISKS & OPPORTUNITY	MEDIUM OPPORTUNITY
	HIGH	MEDIUM RISK	RISKS & OPPORTUNITY	MEDIUM OPPORTUNITY	HIGH OPPORTUNITY
	VERY HIGH	RISKS & OPPORTUNITY	MEDIUM OPPORTUNITY	HIGH OPPORTUNITY	HIGH OPPORTUNITY

Appendix 4. Detail of the methods and information required for full risk assessment

See Figure 1 for an overview of the risk assessment process.

Stage I.

Distribution change data (Stage I.A) were based on Atlas data (for birds) and modelling of recording scheme data held by Biological Records Centre (BRC) as described above for other taxa. Confidence in all bird trends was assessed as good, based on the high coverage and effort. For other taxa, confidence was assessed as good if the mixed model accounting for recorder effort gave a trend where the upper 80% confidence intervals were in the same impact category as the trend (i.e. we were 80% confident that any observed declines were at least that severe), unless experts highlighted that significant changes in recorder effort, taxonomy or identifiability may have contributed to these trends. The linkage between range decline and climate (Stage I.B) was assessed initially by comparison of the direction of observed and projected declines within the current range. If both were negative then this provided evidence for a link (with poor confidence), if they were contradictory in direction then this provided no evidence for a link and if evidence existed in the published literature for a relationship between climate and population or range change, this was regarded as providing evidence of a link with good confidence. In Stage I.C exacerbating factors and associated confidence were assessed from expert opinion and the scientific literature, with a published study supporting the importance of a particular impact on a species' population or distribution regarded as providing evidence with good confidence.

Stage II.

Projected declines within the current range were estimated using outputs from species distribution modelling. Confidence in these projections was assigned as 'high' where

projected and recently observed trends were consistent and the confidence intervals of bioclimatic models (median confidence interval across squares divided by the variance) were less than a threshold value of 0.02 (selected from a visual assessment of the spread of values). Confidence was assigned as medium if the confidence interval threshold was met but projected and observed trends were in opposing directions, indicating that non-climatic factors had driven recent trends. Confidence was low if the median weighted confidence interval was >0.02 , suggesting that the model projections were uncertain.

Stage III.

Stage III.A and III.B were completed as for Stages I.A and I.B, but using information about range expansion rather than contraction. The only difference was that, as described in Thomas et al., (2011), decadal population increases in section III.A were calculated relative to the species' status updated every decade, (as opposed to Stage I.A where changes were calculated relative to the species original status).

Stage IV.

Stage IV.A was based on bioclimatic projections of range expansion outside the current range, calculated as $(\text{newly colonised range}) / (\text{newly colonised range} + \text{current range})$. Confidence was assigned as in Stage II.A. Assessments of exacerbating factors likely to limit range expansion, and our confidence in them (Stage IV.C) were again based on expert knowledge and the literature.

Table A5. Summary of the information required at each stage of the full risk assessment
(summarised and adapted from Thomas et al., 2011)

Stage	Data sources and criteria used
I.A.impact	<p>For bird species the decadal decline within current range was calculated from Atlas data between 1990-2010.</p> <p>For all other taxa, a mixed effects model on BRC data controlling for recorder effort was used.</p>
I.A.confidence	<p>All bird species trends were assigned good confidence.</p> <p>For other taxa, confidence was based on the C. I. from mixed model: if upper 80% C.I. overlaps the next impact category then confidence is poor, otherwise good.</p>
I.B.impact	<p>If both observed trend (I.A.) and projected trend (II.A.) are negative then linkage="Yes". Supplemented with literature review to assess additional linkages with climate</p>
I.B.confidence	<p>Poor if just assessed by comparison of observed (I.A.) and projected (II.A.) trends.</p> <p>Good if robust evidence identified by literature review.</p>
I.C.i.impact	<p>Is current extent <20 000km²? *</p> <p>Additionally for bird species only: is GB population < 10 000 individuals?</p>
I.C.i.confidence	<p>For bird species generally good.</p> <p>For other taxa: poor if just assessed by using current extent data. Good if robust</p>

evidence identified by literature review or supported by expert opinion.

IC.ii.impact Expert knowledge or evidence from literature review supporting at least one of the factors.

I.Cii.confidence Good if robust evidence from peer-reviewed literature. Poor if based on expert knowledge alone.

For birds, due to generally good understanding of the ecology of these species, experts were asked to assign the confidence level where impact was based on unpublished information.

II.A.impact Bioclimate model projected change in occupancy within current range

II.A.confidence a) Are bioclimate confidence intervals below a threshold value (see main text)?

b) Is direction of projected trends (II.A.) in same direction as observed trend (I.A.)?

For bird species: Yes to a)&b) = good, yes to a) only =medium, no to a) =poor.

For other taxa: Yes to a)&b) = good, yes to a) or b) only =medium, no to a) & b) =poor.

II.B. Not applicable

II.C.i.impact As I.C.i

II.Ci.confidence As I.C.i

II.C.ii.impact As I.C.ii

II.Cii.confidence As I.C.ii

III.A.impact	<p>For bird species: decadal increase outside previous range was calculated from Atlas data between 1990 and 2010.</p> <p>Other taxa: mixed model of BRC data of observed increases beyond species' recent historical range** controlling for recorder effort</p>
III.A.confidence	<p>All bird species trends were assigned with good confidence.</p> <p>For other taxa: the model output was compared across 3 different levels of recorder effort - if the level of recorder effort changes the impact category then confidence is poor, otherwise assigned as good.</p>
III.B.impact	<p>If both observed trend (III.A.) and projected trend (IV.A.) are positive then linkage="Yes". Supplemented with literature review to assess additional linkages with climate.</p>
III.B.confidence	<p>Poor if just assessed by comparing observed (III.A.) and projected trends (IV.A.).</p> <p>Good if robust evidence identified in literature review.</p>
III.C.	Not applicable
IV.A.impact	Bioclimate model projected change in occupancy outside the current range
IV.A.confidence	As II.A.
IV.B.	Not applicable
IV.C.i. impact	As I.C.ii
IV.C.i. confidence	As I.C.ii

IV.C.ii. impact	As I.C.ii
-----------------	-----------

IV.C.ii.confidence	As I.C.ii
--------------------	-----------

IV.C.iii. impact	As I.C.ii
------------------	-----------

IV.C.iii.confidence	As I.C.ii
---------------------	-----------

Note we occasionally changed confidence levels in Stage A (usually 1.A.) if experts highlighted concerns regarding distribution data, e.g. significant changes in recorder effort, recent taxonomic splits, issues regarding taxonomic identification etc.

*Current extent is calculated by bioclimate model: probability of a cell being occupied multiplied by the area of a cell = current extent (possible area occupied)

**Number of newly occupied cells outside the current range as a percentage of cells inside current range.

Appendix 5. Species outcomes from the simplified risk assessment

Appendix 6. Species outcomes from the full risk assessment

Group	Latin name	English name	NERC species	Observed decline	Projected decline	Risk of decline	Observed expansion
Ants	<i>Formica cu</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Formica fu</i>	Negro Ant		0 > -1%	> -1%	LOW	> +7.5%
Ants	<i>Formica sai</i>	NA		0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Ants	<i>Lasius ali</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Lasius fla</i>	Yellow Meac		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Ants	<i>Lasius mix</i>	NA		0 > -1%	> -1%	LOW	+1 to +4%
Ants	<i>Lasius nig</i>	Small Black		0 > -1%	> -1%	LOW	> +7.5%
Ants	<i>Leptothora</i>	Slender Ant		0 > -1%	-4 to -1%	MODERATE	> +7.5%
Ants	<i>Myrmica ru</i>	Red Ant		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Myrmica ru</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Ants	<i>Myrmica sai</i>	NA		0 > -1%	-4 to -1%	MODERATE	> +7.5%
Ants	<i>Myrmica sci</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Myrmica sci</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena al</i>	NA		0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bees	<i>Andrena an</i>	NA		0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Andrena ap</i>	NA		0 > -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Andrena ar</i>	NA		0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bees	<i>Andrena ba</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena bi</i>	Gwynne's Mi		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena bi</i>	NA		0 > -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Andrena bu</i>	NA		0 > -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Andrena ch</i>	NA		0 -4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Bees	<i>Andrena ci</i>	Grey Mining		0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bees	<i>Andrena co</i>	NA		0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Andrena coi</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena dei</i>	NA		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena do</i>	NA		0 > -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Andrena fl</i>	Yellow Leg		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena fu</i>	NA		0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Andrena fu</i>	Tawny Minir		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena fu</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena fu</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena ha</i>	Early Minir		0 -4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ha</i>	NA		0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Andrena he</i>	NA		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena hu</i>	NA		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena la</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena la</i>	Girdled Mir		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena la</i>	NA		0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bees	<i>Andrena mi</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena mi</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ni</i>	NA		0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ov</i>	NA		0 > -1%	> -1%	LOW	> +7.5%

Bees	<i>Andrena pi</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena pr</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena pr</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena sc</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena su</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Andrena sy</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ta</i> .Tormentil M	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Andrena th</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena ti</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena tr</i> .Trimmer's M	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena va</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena wi</i> .NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Anthidium i</i> .Wool-Carder	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Anthophora</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Anthophora</i> Fork Tailec	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Anthophora</i> Hairy Footc	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Anthophora</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Apis melli</i> .Honey Bee	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Bombus hor</i> .Small Garde	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Bombus hum</i> .NA	1	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Bombus jon</i> .Heath Bumb	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bees	<i>Bombus lap</i> .Large Red 1	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Bombus luc</i> .White-Tailec	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Bombus magi</i> .NA	0	< -7.5%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Bombus mon</i> .Mountain Bu	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Bombus mus</i> .Moss Carder	1	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Bombus pas</i> .Common Carc	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Bombus pra</i> .Early Bumb	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Bombus rud</i> .NA	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Bombus syl</i> .NA	1	> -1%	> -1%	LOW	< +1%
Bees	<i>Bombus ter</i> .Buff-Tailec	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Chelostoma</i> Harebell Ca	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Coelioxys i</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Coelioxys i</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Coelioxys i</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Colletes d</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Colletes h</i> .Sea-aster C	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Colletes m</i> .Margined Co	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Colletes s</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Colletes s</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Bees	<i>Epeolus cr</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Epeolus va</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Halictus c</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Halictus r</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Halictus t</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Hoplitis c</i> .NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Hoplitis s</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%

Bees	<i>Hylaeus an</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Hylaeus br</i> Short Horne	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus co</i> Common Yell	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus co</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus co</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus hy</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus pe</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Hylaeus pi</i> NA	0 -7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Hylaeus si</i> Large Yell	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasioglossi</i> Slender Mir	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasioglossi</i> Least Minir	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasioglossi</i> Brassy Mini	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Lasioglossi</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasioglossi</i> NA	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasioglossi</i> Shaggy Mini	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Macropis e</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Megachile</i> Patchwork I	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Megachile</i> Wood-Carvir	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Megachile</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Megachile</i> Willughby's	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Melecta al</i> NA	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Melitta ha</i> NA	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Melitta le</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada fab</i> Fabricius'	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada fla</i> NA	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bees	<i>Nomada fla</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Nomada fla</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada fuc</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Nomada goo</i> Gooden's N	0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Nomada lat</i> NA	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Nomada leu</i> NA	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bees	<i>Nomada mar</i> Marsham's N	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada pan</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bees	<i>Nomada ruf</i> Red-Horned	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bees	<i>Nomada she</i> Dark Nomad	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada str</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%

Bees	<i>Osmia auru</i>	Gold-Fringe	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Osmia bico</i>	Two Coloured	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Osmia rufa</i>	Red Mason	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Panurgus c</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Sphecodes i</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Bees	<i>Sphecodes i</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Sphecodes i</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes i</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Stelis orn</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Stelis pun</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Accipiter i</i>	Goshawk	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Accipiter i</i>	Sparrowhawk	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Acrocephal</i>	Sedge Warbler	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Acrocephal</i>	Reed Warbler	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Actitis hy</i>	Common Sandpiper	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Aegithalos</i>	Long-tailed Tit	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Aix galeri</i>	Mandarin Duck	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Alauda arv</i>	Skylark	1	-4 to -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Alca torda</i>	Razorbill	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Alcedo att</i>	Kingfisher	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Alectoris i</i>	Red-legged Partridge	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Anas acuta</i>	Pintail	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Anas clype</i>	Shoveler	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Anas crecc</i>	Teal	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Anas penel</i>	Widgeon	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Anas platy</i>	Mallard	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Anas querq</i>	Garganey	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Anas strep</i>	Gadwall	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Anser anse</i>	Greylag Goose	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Anthus pet</i>	Rock Pipit	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Anthus pra</i>	Meadow Pipit	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Anthus tri</i>	Tree Pipit	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Apus apus</i>	Swift	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Aquila chr</i>	Golden Eagle	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Ardea cine</i>	Grey Heron	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Asio flamm</i>	Short-eared Owl	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Asio otus</i>	Long-eared Owl	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Athene noc</i>	Little Owl	0	> -1%	> -1%	LOW	+4 to +7.5%

Birds	<i>Aythya fer.</i> Pochard	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Aythya ful.</i> Tufted Duck	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Botaurus s.</i> Bittern	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Branta can.</i> Canada Goose	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Branta leu.</i> Barnacle Goose	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Bucephala a.</i> Goldeneye	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Burhinus o.</i> Stone-curlew	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Buteo bute.</i> Buzzard	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Calidris a.</i> Dunlin	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Birds	<i>Caprimulgus.</i> Nightjar	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Carduelis a.</i> Lesser Redpoll	1	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Carduelis a.</i> Linnet	1	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Carduelis a.</i> Goldfinch	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Carduelis a.</i> Greenfinch	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Carduelis a.</i> Twite	1	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Carduelis a.</i> Siskin	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Cephus gr.</i> Black Gull	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Birds	<i>Certhia fa.</i> Treecreeper	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	Cettia cetti Cetti's Warbler	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Charadrius</i> Little Ringed Plover	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Charadrius</i> Ringed Plover	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Charadrius</i> Dotterel	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Birds	<i>Chroicocephalus</i> Black-headed Gull	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Chrysolophus</i> Golden Pheasant	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Cinclus ci.</i> Dipper	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	<i>Circus aer.</i> Marsh Harrier	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Circus cya.</i> Hen Harrier	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	Circus pyga Montagu's Harrier	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Coccothraustes</i> Hawfinch	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	Columba liv. Feral Pigeon	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Columba oea.</i> Stock Dove	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Columba pa.</i> Woodpigeon	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Corvus cor.</i> Raven	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Corvus cor.</i> Carrion Crow	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Corvus fru.</i> Rook	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Corvus mon.</i> Jackdaw	0	> -1%	> -1%	LOW	< +1%
Birds	<i>Coturnix c.</i> Quail	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Crex crex</i> Corncrake	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Cuculus ca.</i> Cuckoo	1	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Birds	<i>Cyanistes a.</i> Blue Tit	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Cygnus olo.</i> Mute Swan	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Delichon u.</i> House Martin	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Dendrocopos.</i> Great Spotted Woodpecker	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Dendrocopos.</i> Lesser Spotted Woodpecker	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Emberiza c.</i> Corn Bunting	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Emberiza c.</i> Cirl Bunting	1	-4 to -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Emberiza c.</i> Yellowhammer	1	-7.5 to -4%	> -1%	MODERATE	+1 to +4%

Birds	<i>Emberiza s</i>	Reed Bunting	1	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Erithacus i</i>	Robin	0	> -1%	> -1%	LOW	< +1%
Birds	<i>Falco colu</i>	Merlin	0	> -1%	< -7.5%	MODERATE	> +7.5%
Birds	<i>Falco pere</i>	Peregrine	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Falco subb</i>	Hobby	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Falco tinn</i>	Kestrel	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Ficedula h</i>	Pied Flycat	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Fratercula</i>	Puffin	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Fringilla c</i>	Chaffinch	0	> -1%	> -1%	LOW	< +1%
Birds	<i>Fulica atr</i>	Coot	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Fulmarus g</i>	Fulmar	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Birds	<i>Gallinago i</i>	Snipe	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Gallinula c</i>	Moorhen	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Garrulus g</i>	Jay	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Haematopus</i>	Oystercatch	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Hirundo ru</i>	Swallow	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Hydrobates</i>	Storm Petre	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Lagopus la</i>	Red Grouse	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Birds	<i>Lanius col</i>	Red-backed	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Larus arge</i>	Herring Gul	1	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Larus canu</i>	Common Gull	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Larus fusc</i>	Lesser Blac	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Larus mari</i>	Great Black	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Larus melai</i>	Mediterrane	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Limosa lim</i>	Black-taile	1	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Locustella lt</i>	Savi's Warb	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Locustella</i>	Grasshopper	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Loxia spp.</i>	Crossbill s	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Lullula ar</i>	Woodlark	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Luscinia m</i>	Nightingale	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Mergus mer</i>	Goosander	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Mergus ser</i>	Red-breaste	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Morus bass</i>	Gannet	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Birds	<i>Motacilla all</i>	Pied Wagta	0	< -7.5%	-7.5 to -4%	LOW	+1 to +4%
Birds	<i>Motacilla c</i>	Grey Wagtai	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Motacilla l</i>	Yellow Wagt	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Muscicapa s</i>	Spotted Fly	1	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Birds	<i>Numenius a</i>	Curlew	1	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Oenanthe o</i>	Wheatear	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	<i>Oxyura jam</i>	Ruddy Duck	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Panurus bi</i>	Bearded Tit	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Parus majo</i>	Great Tit	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Passer dom</i>	House Spar	1	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Passer mon</i>	Tree Sparro	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Perdix per</i>	Grey Partri	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Periparus c</i>	Coal Tit	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Phalacroco</i>	Shag	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%

Birds	<i>Phalacrocorax</i> Cormorant	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Phasianus</i> Pheasant	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Philomachus</i> Ruff	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Phoenicurus</i> Black Redstart	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Phoenicurus</i> Redstart	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Phylloscopus</i> Chiffchaff	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Phylloscopus</i> Wood Warbler	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Phylloscopus</i> Willow Warbler	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Pica pica</i> Magpie	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Picus viridis</i> Green Woodpecker	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Pluvialis</i> Golden Plover	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Birds	<i>Podiceps cornutus</i> Great Crested Grebe	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Poecile montanus</i> Willow Tit	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Poecile palustris</i> Marsh Tit	1	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	<i>Porzana porzana</i> Spotted Crail	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Prunella montanorum</i> Dunnock	1	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Psittacula krameri</i> Ring-necked Pheasant	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Puffinus puffinus</i> Manx Shearwater	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Pyrrhula pyrrhula</i> Bullfinch	1	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Rallus aquatilis</i> Water Rail	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Recurvirostra</i> Avocet	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Regulus ignicapilla</i> Firecrest	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Regulus regulus</i> Goldcrest	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Birds	<i>Riparia riparia</i> Sand Martin	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Rissa tridactyla</i> Kittiwake	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Saxicola rubra</i> Whinchat	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Saxicola torquata</i> Stonechat	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Scolopax rusticicola</i> Woodcock	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Sitta europaea</i> Nuthatch	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Somateria mollissima</i> Eider	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Sterna dougalli</i> Roseate Tern	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Sterna hirsuta</i> Common Tern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Sterna parvirostris</i> Arctic Tern	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Sterna sandvicensis</i> Sandwich Tern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Sternula albifrons</i> Little Tern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Streptopelia</i> Collared Dove	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Streptopelia</i> Turtle Dove	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Strix aluco</i> Tawny Owl	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Sturnus vulgaris</i> Starling	1	> -1%	> -1%	LOW	< +1%
Birds	<i>Sylvia atricapilla</i> Blackcap	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Sylvia borin</i> Garden Warbler	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Birds	<i>Sylvia communis</i> Whitethroat	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Sylvia curruca</i> Lesser Whitethroat	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Sylvia undata</i> Dartford Warbler	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Tachybaptus</i> Little Grebe	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Tadorna tadorna</i> Shelduck	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Tetrao tetrix</i> Black Grouse	1	> -1%	< -7.5%	MODERATE	> +7.5%

Birds	<i>Tringa tot.</i>	Redshank	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Troglodyte.</i>	Wren	0	> -1%	> -1%	LOW	< +1%
Birds	<i>Turdus ili.</i>	Redwing	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Turdus meri.</i>	Blackbird	0	> -1%	> -1%	LOW	< +1%
Birds	<i>Turdus phi.</i>	Song Thrush	1	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Turdus pil.</i>	Fieldfare	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Turdus tor.</i>	Ring Ouzel	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Birds	<i>Turdus vis.</i>	Mistle Thrush	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Tyto alba</i>	Barn Owl	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Uria aalge</i>	Guillemot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Vanellus v.</i>	Lapwing	1	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Bryophytes	<i>Abietinella</i>	Prickly Tar	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Adelanthus</i>	Deceptive F	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes	<i>Aloina alo.</i>	Common Aloe	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes	<i>Amblystegi</i>	Creeping Fe	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes	<i>Amblystegi</i>	NA	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes	<i>Amphidium</i>	Lapland Yof	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Amphidium</i>	Mougeot's	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Anastrepta</i>	Orkney Notc	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Anastrophy.</i>	Heller's No	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes	<i>Anastrophy.</i>	Comb Notchv	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Andreaea a.</i>	Alpine Rock	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Andreaea r.</i>	Dusky Rock-	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes	<i>Andreaea r.</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Andreaea r.</i>	Black Rock-	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes	<i>Andreaea r.</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Bryophytes	<i>Andreaea r.</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Aneura pin.</i>	Greasewort	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bryophytes	<i>Anoetangi</i>	Summer-moss	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Anomobryum</i>	NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Anomodon v.</i>	Rambling T&	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes	<i>Anthelia j.</i>	Alpine Silv	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Anthoceros</i>	Dotted Horr	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes	<i>Antitrichia</i>	Pendulous V	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes	<i>Aphanolejei</i>	Long-leaved	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Atrichum c.</i>	Fountain Sm	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes	<i>Atrichum u.</i>	Common Smoc	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes	<i>Atrichum u.</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes	<i>Aulacomniu</i>	Bud-headed	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Bryophytes	<i>Aulacomniu</i>	Bog Groove-	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes	<i>Barbilopho.</i>	Atlantic Pa	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Barbilopho.</i>	Bearded Pav	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes	<i>Barbilopho.</i>	Common Paw	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Barbilopho.</i>	Hatcher's F	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes	<i>Barbula coi.</i>	Lesser Birc	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes	<i>Barbula coi.</i>	NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Bryophytes	<i>Barbula un.</i>	Bird's-clav	0	> -1%	> -1%	LOW	+1 to +4%

Bryophytes <i>Bartramia</i> Haller's Apr	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Bartramia</i> Straight-le	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Bartramia</i> Common Appl	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Bazzania</i> t.Lesser Whip	0	-7.5 to -4%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Bazzania</i> t.Greater Whi	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Blasia</i> pus.Common Kett	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Blepharost</i> Hairy Threa	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Blindia</i> acSharp-leave	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Brachydont</i> Bristle-lea	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Brachythec</i> .Whitish Fea	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Brachythec</i> .Sand Feathe	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Brachythec</i> .River Feath	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Brachythec</i> .Rough-stalk	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Brachythec</i> .Smooth-stal	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Breutel</i> ia Golden-head	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Bryoerythr</i> Rufous Bear	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> alpiAlpine Thre	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Bryum</i> argeiSilver-moss	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Bryum</i> borniPotato Bry	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Bryum</i> caesiTufted Thre	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> caesiNA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Bryum</i> dichBicoloured	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> dichNA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Bryum</i> gemm.Small-bud F	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> moraFlabby Thre	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Bryum</i> pallPale Threa	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Bryum</i> pseuMarsh Bryu	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> pseuNA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Bryum</i> pseuNA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Bryum</i> radiWall Threa	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Bryum</i> rubeiCrimson-tu	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calliergon</i> Heart-leave	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Calliergon</i> Giant Spear	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Calliergon</i> Lindberg's	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Notched Pou	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Common Pou	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Mueller's F	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Nees' Pouch	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Calypogeia</i> Bog Pouchw	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Bryophytes <i>Campyliade</i> .Golden Feat	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes <i>Campylium</i> Yellow Star	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Campylium</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Campylophy</i> .Chalk Feath	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Campylopus</i> Bristly Swa	0	-4 to -1%	< -7.5%	HIGH	+4 to +7.5%
Bryophytes <i>Campylopus</i> Compact Swa	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Campylopus</i> Rusty Swan-	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Campylopus</i> Brittle Swa	0	< -7.5%	-4 to -1%	HIGH	> +7.5%

Bryophytes <i>Campylopus</i> Heath Star	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Campylopus</i> Dwarf Swan-	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Cephalozia</i> Two-horned	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Cephalozia</i> Chain Pince	0 -7.5 to -4%	-7.5 to -4%	HIGH	+1 to +4%
Bryophytes <i>Cephalozia</i> Moon-leaved	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Cephalozia</i> Common Three	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Cephalozia</i> Hampe's Thre	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Ceratodon</i> Redshank [r	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Chiloscyph</i> St Winifric	0 -4 to -1%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Cirriphyll</i> Beech Feath	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Cirriphyll</i> Hair-pointe	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Cladopodie</i> Bog Notchw	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Climacium</i> Tree-moss	0 -4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes <i>Cololejeun</i> Rock Pounce	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Cololejeun</i> Minute Pour	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Cololejeun</i> Rossetti's	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Colura cal</i> Fingered Co	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Conocephal</i> Great Scent	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Conostomum</i> Helmet-moss	0 > -1%	< -7.5%	MODERATE	< +1%
Bryophytes <i>Cratoneuroi</i> Fern-leaved	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Cratoneuroi</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Cryphaea h</i> Lateral Cry	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ctenidium</i> Chalk Comb-	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Ctenidium</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Cynodontiu</i> Brunton's I	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Dialytrich</i> Pointed Lat	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Dichodonti</i> Marsh Forkl	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Dichodonti</i> NA	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Dichodonti</i> Transparent	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Dicranella</i> Rufous Forkl	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Dicranella</i> Field Forkl	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Dicranella</i> Variable Fo	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Dicranowei</i> Common Pinc	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Dicranowei</i> Mountain Pi	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Dicranum f</i> Whip Fork-r	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Dicranum fi</i> Dusky Fork-	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Dicranum fi</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Dicranum m</i> Greater For	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Dicranum s</i> Broom Fork-	0 -4 to -1%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Dicranum s</i> Scott's For	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Dicranum si</i> Rusty Fork-	1 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Dicranum t</i> Fragile For	0 > -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Didymodon</i> Pointed Bea	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Didymodon</i> Fallacious	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> Cylindric F	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Didymodon</i> Dusky Bearc	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> Nicholson's	0 > -1%	> -1%	LOW	> +7.5%

Bryophytes <i>Didymodon</i> Rigid Beard-moss	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> Wavy Beard-moss	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Didymodon</i> Brown Beard-moss	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Didymodon</i> Shady Beard-moss	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> Soft-tufted moss	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Diphyscium</i> Nut-moss	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Diplophyllum</i> White Earwort	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Distichium</i> Fine Distichum	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Ditrichum</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Ditrichum</i> Bendy Ditrichum	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ditrichum</i> Curve-leaved	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Douinia ovata</i> Waxy Earwort	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Drepanocla</i> Fertile Feather-moss	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Drepanolej</i> Toothed Porella	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Encalypta</i> Ribbed Extinct	0 < -7.5%	< -7.5%	VERY HIGH	< +1%
Bryophytes <i>Encalypta</i> Spiral Extinct	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Encalypta</i> Common Extinct	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Entodon complanatus</i> Montagne's moss	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Entosthodon</i> Thin Cord-moss	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Entosthodon</i> Muhlenberg's moss	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Entosthodon</i> Blunt Cord-moss	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Ephemerum</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Ephemerum</i> Strap-leaved	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Ephemerum</i> Serrated Earwort	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Eremonotus</i> Clubwort	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Eurhynchium</i> Common Striped	0 > -1%	> -1%	LOW	< +1%
Bryophytes <i>Fissidens</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> Maidenhair	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Fissidens</i> Lesser Pocket	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Fissidens</i> Curnow's Pocket	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> Welsh Pocket	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Fissidens</i> Fatfoot Pocket	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> Rock Pocket	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> Slender Pocket	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> Narrow-leaved	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Fissidens</i> Short-leaved	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Fissidens</i> Purple-stalked	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> Petty Pocket	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> River Pocket	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> Beck Pocket	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Fissidens</i> Common Pocket	0 -7.5 to -4%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Fissidens</i> NA	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> Green Pocket	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Bryophytes <i>Fontinalis</i> Greater Water-moss	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Fossombron</i> Common Frill	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Fossombron</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%

Bryophytes <i>Frullania</i> Dilated Scallop	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Frullania</i> Spotty Scallop	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Frullania</i> Tamarisk Scallop	0 > -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Bryophytes <i>Frullania</i> Sea Scalewort	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Funaria hygrometrica</i> Common Cornerwort	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Grimmia douglasii</i> Donn's Grimmioid	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia funaria</i> String Grimmioid	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia linaresii</i> NA	0 > -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Grimmia pulvinata</i> Grey-cushionwort	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Grimmia radiata</i> Spreading-liverwort	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia tomentosa</i> Twisted Grimmioid	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia trichomanes</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Gymnocolea inflata</i> Inflated Neckwort	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Gymnomitria</i> Braided Frodo	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Gymnomitria</i> Western Frodo	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Gymnomitria</i> White Frostwort	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Gymnostomum verdigris</i> Verdigris-liverwort	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Gymnostomum</i> Blunt-leafed	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Gymnostomum</i> Luisier's-liverwort	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Gyroweisia</i> Slender Stemwort	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Hamatocaulis</i> Varnished Liverwort	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Harpalejeunea</i> Pointed Pouchwort	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Harpanthus</i> Stipular Flowerwort	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Hedwigia ciliata</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Bryophytes <i>Hedwigia stellata</i> Starry Hoarwort	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Hennediella</i> Stanford Scallopwort	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Herbertus</i> Straw Prongwort	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Heterocladus</i> Wry-leaved	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Heterocladus</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Heterocladus</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Homalia trichomanes</i> Blunt Featherwort	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Homalotheca</i> Yellow Featherwort	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Homalotheca</i> Silky Wallwort	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Hookeria linaresii</i> Shining Hooker	0 > -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Hygroamblystidium</i> Fountain Featwort	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Hygroamblystidium</i> Willow Featwort	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Hygrobiella</i> Lax Notchwort	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Hygrohypnum</i> Claw Brook-liverwort	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Hylocomiastrum</i> Shaded Wood	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Hylocomium</i> Glittering	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Hylocomium</i> Flagellate	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Hypnum androdia</i> Mamillate Liverwort	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Hypnum calopogon</i> Downy Plait	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Hypnum cupressinum</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Hypnum cupressinum</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Hypnum cupressinum</i> Great Plait	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Hypnum cupressinum</i> Supine Plait	0 > -1%	> -1%	LOW	+4 to +7.5%

Bryophytes <i>Hypnum jut.</i> Heath Plain	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Isopterygi</i> Neat Silk-r	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Isothecium</i> Larger Mous	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Isothecium</i> Holt's Mous	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Isothecium</i> Slender Mou	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Isothecium</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Isothecium</i> NA	0 -4 to -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Jamesoni</i> Autumn Flap	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Jungermann</i> .Dark-green	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Jungermann</i> .Dwarf Flap	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Kiaeria bl</i> Blytt's For	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Kiaeria fa</i> Sickle-leav	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Kindbergia</i> Common Feat	0 > -1%	> -1%	LOW	< +1%
Bryophytes <i>Kurzia pau</i> Bristly Fir	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Kurzia syl</i> Wood Finger	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Kurzia tri</i> Heath Finge	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Leiocolea</i> Bantry Notc	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Leiocolea</i> Ragged Notc	0 < -7.5%	< -7.5%	VERY HIGH	< +1%
Bryophytes <i>Lejeunea ca</i> Micheli's I	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Lejeunea la</i> Western Pou	0 > -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Lejeunea pa</i> Pearl Pounc	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Lepidozia</i> Rock Finger	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Lepidozia</i> Pearson's I	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Lepidozia</i> Creeping Fi	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Leptobarbu</i> .Beric Bearc	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Leptodicty</i> Kneiff's Fe	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Leptodon si</i> Prince-of-V	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Leptodonti</i> Bent-leaved	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Leskea pol</i> Many-fruite	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Leucobryum</i> Large White	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Leucobryum</i> Smaller Whi	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Leucodon si</i> Squirrel-ta	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Leucodon si</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Loeskeobryi</i> Short-beake	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Lophocolea</i> Bifid Crest	0 > -1%	> -1%	LOW	< +1%
Bryophytes <i>Lophocolea</i> Fragrant Cr	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Lophocolea</i> Variable-le	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Lophozia e</i> Capitate No	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Lophozia i</i> Jagged Notc	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Lophozia si</i> Hill Notchv	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Lophozia vi</i> Tumid Notch	0 > -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Lunularia</i> Crescent-cr	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Marchantia</i> Common Live	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Marchantia</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Marchesini</i> MacKay's Pc	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Marsupella</i> Scorched Ru	0 > -1%	< -7.5%	MODERATE	< +1%
Bryophytes <i>Marsupella</i> Notched Rus	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%

Bryophytes <i>Marsupella</i> NA	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Marsupella</i> Funck's Rus	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Marsupella</i> Stabler's F	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Metzgeria</i> (Rock Veilw	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Metzgeria</i> (Whiskered V	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Metzgeria</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Metzgeria</i> (Forked Veil	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Metzgeria</i> (Hooked Veil	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Metzgeria</i> (Downy Veilv	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Metzgeria</i> (Blueish Veil	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Microbryum</i> Floerke's F	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Microlejeu</i> Fairy Beads	0 > -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Mnium</i> hornSwan's-neck	0 -7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Bryophytes <i>Mnium</i> marg. NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Molendoa</i> w/Warburg's M	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Mylia</i> tayl(Taylor's F)	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Nardia</i> sca.Ladder Flap	0 -7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Neckera</i> co.Flat Necke	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Neckera</i> cr.Crisped Nec	0 -7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Neckera</i> pu.Dwarf Necke	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Nowellia</i> ci.Wood-rust	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Odontoschi</i> .Bog-m Flapv	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Oedipodium</i> Gouty-moss	0 > -1%	< -7.5%	MODERATE	< +1%
Bryophytes <i>Oligotrich</i> Hercynian F	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Orthodonti</i> Cape Threac	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Orthotheci</i> Fine-leaved	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Orthotheci</i> Red Leskea	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Orthotrich</i> Wood Bristl	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> Anomalous F	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Orthotrich</i> White-tippe	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Orthotrich</i> Lyell's Bri	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> Elegant Bri	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> River Brist	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> Rock Bristl	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Orthotrich</i> Showy Brist	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> Straw Brist	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> Shaw's Bris	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Orthotrich</i> Slender Bri	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Oxyrrhynch</i> .Swartz's Fe	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Oxyrrhynch</i> .Dwarf Feath	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Oxyrrhynch</i> .Twist-tip F	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Oxyrrhynch</i> .Showy Feath	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Oxystegus</i> NA	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Palustriel</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Palustriel</i> .NA	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Pellia</i> epi.Overleaf Pe	0 -7.5 to -4%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Pellia</i> nee.Nees' Pelli	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%

Bryophytes <i>Phascum cuc</i> Cuspidate F	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Phascum cuc</i> .NA	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Philonotis</i> Thick-nerve	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Philonotis</i> Fountain A	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Physcomit</i> .Common Blac	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Plagiobryu</i> Zierian Hun	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> Greater Fea	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Plagiochil</i> Killarney F	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> British Fea	0 -7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> Petty Feath	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Plagiochil</i> Western Fea	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiochil</i> Lesser Feat	0 > -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Plagiochil</i> Spotty Feat	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> Prickly Fea	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Plagiomniu</i> Many-fruite	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiomniu</i> Woodsy Thym	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiomniu</i> Marsh Thyme	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiomniu</i> Long-beake	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiopus</i> Oeder's Apr	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiothec</i> .Curved Silk	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Plagiothec</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Plagiothec</i> .NA	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiothec</i> .Bright Silk	0 -4 to -1%	< -7.5%	HIGH	+1 to +4%
Bryophytes <i>Plagiothec</i> .Alder Silk-	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Plagiothec</i> .Woodsy Silk	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Plagiothec</i> .Juicy Silk-	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiothec</i> .Waved Silk-	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Plasteurhy</i> Lesser Stri	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Platyhypni</i> Portuguese	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Platyhypni</i> Long-beake	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Pleuridium</i> Awl-leaved	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Pleurozium</i> Red-stemmed	0 -7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Bryophytes <i>Pogonatum</i> Aloe Hairca	0 > -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Pogonatum</i> Urn Haircap	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Pohlia ann</i> Pale-fruite	0 > -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Pohlia bul</i> Blunt-bud T	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Pohlia cam</i> Crookneck M	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia dru</i> Drummond's	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia elo</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia lut</i> Yellow Thre	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia mel</i> Pink-fruite	0 -7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Pohlia nut</i> Nodding Thu	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Pohlia wah</i> Pale Glauc	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Polytricha</i> .Alpine Hair	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Polytricha</i> .Bank Hairca	0 -4 to -1%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Polytricha</i> .Slender Hai	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Polytrichu</i> Common Hair	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%

Bryophytes <i>Polytrichum</i> NA	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Polytrichum</i> Juniper Hair	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Polytrichum</i> Bristly Hair	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Polytrichum</i> Strict Hair	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Porella pil</i> Pinnate Scale	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Porella pil</i> Wall Scale	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Preissia quadr</i> Narrow Mush	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Pseudocall</i> Three-ranked	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pseudocroc</i> Hornschuch's	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Pseudotaxi</i> Elegant Sil	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Pterogonium</i> Bird's-foot	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Ptilidium</i> Ciliated Fr	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ptilidium</i> Tree Fringe	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Ptilium cr</i> Ostrich-pl	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Ptychomitri</i> Long-shank	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Racomitrium</i> Yellow Fr	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Racomitrium</i> Narrow-lea	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitrium</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Bryophytes <i>Racomitrium</i> Oval-fru	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitrium</i> Long Fringe	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Racomitrium</i> Dense Fring	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Racomitrium</i> Green Mount	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Racomitrium</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Racomitrium</i> Bristly Fr	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitrium</i> Woolly Fr	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Racomitrium</i> Slender Fr	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Radula aqu</i> Brown Scale	0 -7.5 to -4%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Radula compl</i> Even Scale	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Radula lind</i> Lindenberg's	0 < -7.5%	-4 to -1%	HIGH	< +1%
Bryophytes <i>Reboulia hem</i> Hemisphaeri	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Rhabdoweis</i> Toothed Str	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Rhabdoweis</i> Dwarf Stre	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Rhizomnium</i> Felted Thyn	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Rhizomnium</i> Dotted Thyn	0 -4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Bryophytes <i>Rhynchosteg</i> Tender Feat	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Rhynchosteg</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Rhynchosteg</i> Teesdale Fe	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Rhynchosteg</i> Clustered F	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Rhynchosteg</i> Megapolitar	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Rhynchosteg</i> Wall Feathe	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Rhytidiade</i> Little Shag	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Rhytidiade</i> Big Shaggy-	0 -7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Bryophytes <i>Riccardia</i> Jagged Germ	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Riccardia</i> Delicate Ge	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Riccardia</i> Palmate Ger	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Riccia bey</i> Purple Cry	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Riccia cav</i> Cavernous C	0 > -1%	< -7.5%	MODERATE	> +7.5%

Bryophytes <i>Riccia glauca</i> Glaucous C	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Saccogyna</i> Straggling	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Sanionia</i> Sick-leav	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Sarmentypn</i> Ringless Hc	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sarmentypn</i> Twiggy Spea	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania a</i> Lesser Roug	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania a</i> Rough Earw	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania c</i> Thick-set F	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania c</i> Untidy Earv	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Scapania g</i> Western Ear	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Scapania i</i> Heath Earw	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Scapania n</i> Grove Earw	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Scapania s</i> Norwegian F	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania u</i> Marsh Earw	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Scapania u</i> Shady Earw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Scapania u</i> Water Earw	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Schistidium</i> NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Schistidium</i> Thickpoint	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Schistidium</i> Seaside Gri	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Schistidium</i> Upright Bro	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Schistostea</i> Luminous Mc	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Sciuro-hypn</i> Rusty Feat	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Bryophytes <i>Sciuro-hypn</i> Matted Feat	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Scleropodi</i> Tufted Feat	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Scleropodi</i> Glass-wort	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Scorpidium</i> Intermediate	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Scorpidium</i> Rusty Hook-	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Scorpidium</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Scorpidium</i> Hooked Scor	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Scorpiuriu</i> Curving Fea	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Seligeria</i> Sharp Rock-	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Seligeria</i> Dwarf Rock-	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Solenostom</i> Crenulated	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Solenostom</i> Transparent	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Solenostom</i> Egg Flapwor	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Solenostom</i> Shining Fla	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Solenostom</i> Round-fruit	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Sphagnum c</i> Red Bog-mos	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Sphagnum c</i> Red Bog-mos	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum c</i> Compact Bog	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Sphagnum c</i> Twisted Bog	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum c</i> Feathery Bo	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Sphagnum d</i> Cow-horn Bo	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum f</i> Flat-topped	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum f</i> Fringed Bog	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum f</i> Flexuous Bo	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum g</i> Girgensohn'	0	> -1%	< -7.5%	MODERATE	> +7.5%

Bryophytes <i>Sphagnum</i> <i>il</i> Lesser Cow-	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum</i> <i>pl</i> Blunt-leave	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Sphagnum</i> <i>pl</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Sphagnum</i> <i>pl</i> Papillose I	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Sphagnum</i> <i>pl</i> Golden Bog-	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Sphagnum</i> <i>ql</i> Five-ranked	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum</i> <i>rl</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Bryophytes <i>Sphagnum</i> <i>rl</i> Russow's Bc	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum</i> <i>sl</i> Spiky Bog-n	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum</i> <i>sl</i> Lustrous Bc	0 -4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Sphagnum</i> <i>sl</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum</i> <i>sl</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Bryophytes <i>Sphagnum</i> <i>tl</i> Soft Bog-mc	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum</i> <i>tl</i> Rigid Bog-n	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Splachnum</i> <i>al</i> Round-fruit	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Straminerg</i> <i>al</i> Straw Spear	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Syntrichia</i> <i>al</i> Small Hairy	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Syntrichia</i> <i>al</i> Water Screv	0 -4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes <i>Syntrichia</i> <i>al</i> Intermediat	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Syntrichia</i> <i>al</i> Marble Scre	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Syntrichia</i> <i>al</i> Sand-hill S	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Syntrichia</i> <i>al</i> Great Hairy	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Targionia</i> <i>al</i> Orobus-seec	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Tetraphis</i> <i>al</i> Pellucid Fe	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Tetraplodo</i> <i>al</i> Slender Cru	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Thamnobryu</i> <i>al</i> Fox-tail Fe	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Thuidium</i> <i>al</i> Delicate Ta	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Thuidium</i> <i>al</i> Common Tame	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Tortella</i> <i>al</i> Yellow Cris	0 -7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes <i>Tortella</i> <i>il</i> Sassari Cri	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Tortella</i> <i>nl</i> Neat Crisp-	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Tortella</i> <i>tl</i> Frizzled Cr	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Tortula</i> <i>lal</i> Lance-leave	0 > -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Tortula</i> <i>mal</i> Bordered Sc	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Tortula</i> <i>mol</i> Blunt-fruit	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Tortula</i> <i>mul</i> Wall Screw-	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Tortula</i> <i>pr</i> Tall Pottia	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Tortula</i> <i>su</i> Awl-leaved	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Tortula</i> <i>tri</i> Common Pot	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Trichocole</i> <i>al</i> Handsome Wc	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Trichodon</i> <i>al</i> Cylindric I	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Trichostom</i> <i>al</i> Variable Cr	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Trichostom</i> <i>al</i> Curly Crisp	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Tritomaria</i> <i>al</i> Cut Notchw	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Tritomaria</i> <i>al</i> Larger Cut	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Tritomaria</i> <i>al</i> Lyon's Notc	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Ulota</i> <i>bruci</i> Bruch's Pir	0 > -1%	> -1%	LOW	> +7.5%

Bryophytes <i>Ulota crisp</i> Crisped Pir	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Ulota crisp</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Ulota hutch</i> Hutchins' F	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ulota phyl</i> Frizzled Pi	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Weissia</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Weissia br</i> Small-mouth	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Weissia coi</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Weissia loi</i> Crisp Bearc	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Weissia loi</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Zygodon coi</i> Lesser Yoke	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Zygodon ru</i> Park Yoke-r	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Zygodon vi</i> Green Yoke-	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Zygodon vi</i> NA	0 > -1%	-7.5 to -4%	MODERATE	< +1%
Bryophytes <i>Zygodon vi</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Acupalpus</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Acupalpus</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Acupalpus</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Acupalpus</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Agonum ema</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Agonum ful</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Agonum gra</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Carbid beet <i>Agonum mar</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Agonum mue</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Carbid beet <i>Agonum pic</i> NA	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Carbid beet <i>Agonum tho</i> NA	0 -4 to -1%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Agonum vid</i> NA	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Carbid beet <i>Amara aene</i> Common Sun	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Amara apri</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Amara bifr</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Amara cons</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Amara conv</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Amara eque</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Amara eury</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Amara fami</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Amara luci</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Amara ovat</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Amara pleb</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid beet <i>Amara prae</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Amara simi</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Amara tibi</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Anchomenus</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Anisodacty</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Anthracus</i> NA	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Asaphidion</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Asaphidion</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Carbid beet <i>Asaphidion</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Badister bi</i> NA	0 > -1%	> -1%	LOW	+1 to +4%

Carbid beet <i>Badister d</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Badister s</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Badister u</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	-4 to -1%	HIGH	< +1%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	< +1%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 -4 to -1%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Bembidion i</i> .NA	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Carbid beet <i>Blemus dis</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Blethisa m</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beet <i>Bracteon l</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beet <i>Bradycellu</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beet <i>Bradycellu</i> .NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Bradycellu</i> .NA	0 < -7.5%	> -1%	MODERATE	< +1%
Carbid beet <i>Bradycellu</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beet <i>Bradycellu</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Bradycellu</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Broscus ce</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Calathus c</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Calathus e</i> .NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Carbid beet <i>Calathus f</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Calathus m</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Calathus m</i> .NA	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Carbid beet <i>Calathus m</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Calathus r</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Calodromiu</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%

Carbid beetle <i>Calosoma i.</i> NA	1 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Carbid beetle <i>Carabus ar.</i> NA	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Carbid beetle <i>Carabus gl.</i> NA	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Carbid beetle <i>Carabus gr.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Carbid beetle <i>Carabus mo.</i> Necklace G1	1 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beetle <i>Carabus nei</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beetle <i>Carabus pr.</i> NA	0 -4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Carbid beetle <i>Curtonotus</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beetle <i>Curtonotus</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beetle <i>Cymindis a.</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Carbid beetle <i>Demetrias i.</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beetle <i>Demetrias</i> . NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beetle <i>Dromius ag.</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Carbid beetle <i>Dromius me.</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beetle <i>Dromius qu.</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid beetle <i>Dyschirius</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beetle <i>Dyschirius</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beetle <i>Dyschirius</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid beetle <i>Dyschirius</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beetle <i>Dyschirius</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Carbid beetle <i>Elaphrus u.</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beetle <i>Eurynebria</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beetle <i>Harpalus a.</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beetle <i>Harpalus a.</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beetle <i>Harpalus l.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Carbid beetle <i>Harpalus m.</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	< +1%
Carbid beetle <i>Harpalus r.</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid beetle <i>Harpalus r.</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Carbid beetle <i>Harpalus r.</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid beetle <i>Harpalus s.</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beetle <i>Harpalus s.</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beetle <i>Laemostenu.</i> NA	0 > -1%	> -1%	LOW	< +1%
Carbid beetle <i>Leistus fe.</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid beetle <i>Leistus fu.</i> NA	0 -7.5 to -4%	> -1%	MODERATE	+1 to +4%
Carbid beetle <i>Leistus ru.</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid beetle <i>Leistus te.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beetle <i>Licinus de.</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Carbid beetle <i>Loricera p.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beetle <i>Masoreus w.</i> NA	0 > -1%	> -1%	LOW	< +1%
Carbid beetle <i>Miscodera i.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beetle <i>Nebria bre.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beetle <i>Nebria ruf.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid beetle <i>Nebria sal.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Carbid beetle <i>Notiophilu.</i> NA	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid beetle <i>Notiophilu.</i> NA	0 -7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Carbid beetle <i>Notiophilu.</i> NA	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid beetle <i>Notiophilu.</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%

Carbid beet <i>Notiophilu</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Notiophilu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Ocys harpa</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid beet <i>Odacantha</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Oodes helo</i> .NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	+1 to +4%
Carbid beet <i>Ophonus ar</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Ophonus az</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Ophonus pu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Ophonus sci</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Oxypselaph</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Panagaeus</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Paradromiu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Paradromiu</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Carbid beet <i>Paranchus</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid beet <i>Patrobus a</i> .NA	0	-4 to -1%	< -7.5%	HIGH	+1 to +4%
Carbid beet <i>Patrobus a</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Carbid beet <i>Philorhizu</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Philorhizu</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Philorhizu</i> .NA	1	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Platyderus</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Platynus a</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid beet <i>Poecilus c</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Poecilus v</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Pogonus ch</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Pterostichi</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Carbid beet <i>Pterostichi</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid beet <i>Pterostichi</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Carbid beet <i>Pterostichi</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Pterostichi</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Pterostichi</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid beet <i>Pterostichi</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Pterostichi</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Carbid beet <i>Pterostichi</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid beet <i>Pterostichi</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid beet <i>Pterostichi</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Stenolophu</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Stomis pum</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid beet <i>Syntomus fi</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid beet <i>Syntomus o</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid beet <i>Syntomus t</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Synuchus v</i> .NA	0	< -7.5%	-4 to -1%	HIGH	< +1%
Carbid beet <i>Tachys bis</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Carbid beet <i>Trechoble</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Carbid beet <i>Trechus qu</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid beet <i>Trechus ru</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Carbid beet <i>Trechus se</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid beet <i>Trichocelli</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%

Carbid beetle <i>Trichocellus</i> NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Centipedes <i>Cryptops al</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Centipedes <i>Cryptops h</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Centipedes <i>Geophilus</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Centipedes <i>Geophilus</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Centipedes <i>Geophilus</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Centipedes <i>Geophilus</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Centipedes <i>Henia vesu</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Centipedes <i>Lithobius</i> NA	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Centipedes <i>Lithobius</i> NA	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Centipedes <i>Lithobius</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Centipedes <i>Lithobius</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Centipedes <i>Lithobius</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Centipedes <i>Lithobius</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Centipedes <i>Schendyla</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Centipedes <i>Stigmatoga</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Centipedes <i>Strigamia</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Centipedes <i>Strigamia</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Coccinellid <i>Adalia bip</i> Two-spot La	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Coccinellid <i>Adalia dec</i> Ten-spot La	0 > -1%	> -1%	LOW	+1 to +4%
Coccinellid <i>Anatis oce</i> Eyed Ladybi	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Coccinellid <i>Anisosticta</i> Water Ladyb	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinellid <i>Chilocorus</i> Kidney-spot	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinellid <i>Coccidula</i> NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Coccinellid <i>Coccinella</i> Seven-spot	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Coccinellid <i>Coccinella</i> Eleven-spot	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Coccinellid <i>Exochomus</i> (Pine Ladybi	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Coccinellid <i>Halysia se</i> Orange Lady	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Coccinellid <i>Hippodamia</i> Adonis' Lac	0 < -7.5%	> -1%	MODERATE	> +7.5%
Coccinellid <i>Propylea qi</i> Fourteen-sp	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinellid <i>Psyllobora</i> Twentytwo-s	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinellid <i>Rhyzobius</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Coccinellid <i>Scymnus su</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Coccinellid <i>Subcoccine</i> Twentyfour-	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Coccinellid <i>Tytthaspis</i> Sixteen-sp	0 > -1%	> -1%	LOW	> +7.5%
Craneflies <i>Nephrotoma</i> NA	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Craneflies <i>Ptychopter</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Craneflies <i>Ptychopter</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Craneflies <i>Ptychopter</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Craneflies <i>Tipula ful</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Craneflies <i>Tipula lat</i> NA	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Craneflies <i>Tipula lun</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Craneflies <i>Tipula max</i> NA	0 > -1%	> -1%	LOW	< +1%
Craneflies <i>Tipula ole</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Craneflies <i>Tipula unc</i> NA	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Craneflies <i>Tipula var</i> NA	0 > -1%	-4 to -1%	MODERATE	+1 to +4%
Crickets at <i>Chorthippu</i> Lesser Mars	0 > -1%	> -1%	LOW	> +7.5%

Crickets at <i>Chorthippu</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Chorthippu</i> .Meadow Gras	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Conocephala</i> .NA	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Crickets at <i>Conocephala</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Crickets at <i>Ectobius p</i> .Tawny Cock	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Crickets at <i>Ectobius p</i> .Lesser Cock	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Crickets at <i>Forficula</i> .Common Earv	0 > -1%	> -1%	LOW	+4 to +7.5%
Crickets at <i>Forficula</i> .Lesne's Ear	0 > -1%	> -1%	LOW	> +7.5%
Crickets at <i>Leptophyes</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Crickets at <i>Meconema t</i> .NA	0 > -1%	-4 to -1%	MODERATE	+4 to +7.5%
Crickets at <i>Metrioptera</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Crickets at <i>Metrioptera</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Crickets at <i>Myrmeleote</i> .Mottled Gra	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Crickets at <i>Nemobius s</i> .Wood Cricke	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Crickets at <i>Omocestus</i> .Woodland Gr	0 > -1%	< -7.5%	MODERATE	> +7.5%
Crickets at <i>Omocestus</i> .Common Gree	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Crickets at <i>Pholidoptera</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Platycleis</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Stenobothrus</i> .Stripe-wing	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Crickets at <i>Tetrix cep</i> .Cepero's Gr	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Crickets at <i>Tetrix sub</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Crickets at <i>Tetrix und</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Anasimyia</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Anasimyia</i> .NA	0 < -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Anasimyia</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Arctophila</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Baccha elo</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Brachyopa</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Brachyopa</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Brachyopa</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Brachyopa</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Brachypalpi</i> .NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Hoverflies <i>Brachypalpi</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Callicera</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Chalcosyrph</i> .NA	0 -4 to -1%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 -7.5 to -4%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%

Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> μ NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Chrysogaster</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Chrysogaster</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Chrysotoxus</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Chrysotoxus</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Chrysotoxus</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Chrysotoxus</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Chrysotoxus</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Criorhina</i> μ NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Criorhina</i> μ NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Criorhina</i> μ NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Dasysyrphus</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Dasysyrphus</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Hoverflies <i>Dasysyrphus</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Dasysyrphus</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Dasysyrphus</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Didea fasc.</i> NA	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Hoverflies <i>Didea inter.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Epistrophe</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Epistrophe</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Epistrophe</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Epistrophe</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Episyrphus</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Eriozona e.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Eriozona s.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Eristalis</i> NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Eristalis</i> μ NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eristalis</i> μ NA	0 -7.5 to -4%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Eristalis</i> μ NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Eristalis</i> μ NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Eristalis</i> μ NA	0 > -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Eristalis</i> μ NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Eristalis</i> μ NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Eumerus fulvipes</i> Lesser Bull	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eumerus ornatus</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eumerus saundersi</i> NA	0 > -1%	< -7.5%	MODERATE	+1 to +4%

Hoverflies <i>Eumerus st.</i> Lesser Bull	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Eupeodes b.</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eupeodes c.</i> NA	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eupeodes l.</i> NA	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eupeodes l.</i> NA	0 -7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Eupeodes n.</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Eupeodes n.</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Ferdinandea</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Helophilus</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Helophilus</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Helophilus</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Heringia h.</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Heringia p.</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Heringia v.</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Lejogaster</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Leucozona</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Leucozona</i> .NA	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Hoverflies <i>Melangyna</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Hoverflies <i>Melangyna</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Melangyna</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Melangyna</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Melangyna</i> .NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Melangyna</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Melanogaster</i> NA	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Hoverflies <i>Melanogaster</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Melanostoma</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Hoverflies <i>Meligramma</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Meligramma</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Meliscaeva</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Merodon eq.</i> Greater Bull	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Microdon a.</i> NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Hoverflies <i>Microdon m.</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Myathropa</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Myolepta d.</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Neoascia g.</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Neoascia i.</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Neoascia m.</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Neoascia o.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Neoascia p.</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Orthonevra</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Orthonevra</i> NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Hoverflies <i>Orthonevra</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Paragus ha.</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Paragus ti.</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Parasyrphu.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Parasyrphu.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Parasyrphu.</i> NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%

Hoverflies <i>Parhelophi</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Parhelophi</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Pelecocera</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Pipiza aus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Pipiza bim</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Pipiza fen</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Pipiza lugi</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Pipiza lut</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Pipiza noc</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Pipizella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Pipizella</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Platycheiri</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Platycheiri</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Portevinia</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Psilota an</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Rhingia cai</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Riponnensi</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Scaeva sel</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Sericomyia</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Sericomyia</i> .NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Sphegina e</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Sphegina v</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Syritta pi</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Syrphus ri</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Syrphus to</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Syrphus vi</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Trichopsom</i> .NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Hoverflies <i>Triglyphus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Tropidia s</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Xanthandru</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%

Hoverflies <i>Xanthogram</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Xanthogram</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Xylota abi</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Xylota flo</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Xylota jak</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Hoverflies <i>Xylota segi</i> NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Xylota syl</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Xylota tar</i> NA	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Hoverflies <i>Xylota xan</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Millipedes <i>Archiboreo</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Millipedes <i>Blaniulus</i> Spotted Snake	0	> -1%	> -1%	LOW	+1 to +4%
Millipedes <i>Boreoiulus</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Millipedes <i>Brachydesm</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Millipedes <i>Chordeuma</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Millipedes <i>Cylindroi</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Millipedes <i>Cylindroi</i> NA	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Millipedes <i>Cylindroi</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Millipedes <i>Cylindroi</i> Blunt-tailed	0	< -7.5%	> -1%	MODERATE	< +1%
Millipedes <i>Glomeris m</i> Pill Millip	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Millipedes <i>Julus scan</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Millipedes <i>Macroster</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Millipedes <i>Melogona s</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Millipedes <i>Nanogona p</i> Eyed Flat-l	0	> -1%	> -1%	LOW	+1 to +4%
Millipedes <i>Nemasoma v</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Millipedes <i>Ommatoiulu</i> Striped Mil	0	> -1%	-7.5 to -4%	MODERATE	+1 to +4%
Millipedes <i>Ophiodesmu</i> NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Millipedes <i>Ophiulus</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+1 to +4%
Millipedes <i>Polydesmus</i> Common Flat	0	> -1%	> -1%	LOW	+1 to +4%
Millipedes <i>Polydesmus</i> NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Millipedes <i>Polydesmus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Millipedes <i>Tachypodoi</i> White-legge	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths <i>Abraxas gr</i> The Magpie	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths <i>Acasis vir</i> Yellow-barr	0	> -1%	> -1%	LOW	> +7.5%
Moths <i>Achlya fla</i> Yellow Horr	0	> -1%	-7.5 to -4%	MODERATE	< +1%
Moths <i>Acronicta</i> The Sycamor	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths <i>Acronicta</i> Alder Moth	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths <i>Acronicta</i> Light Knot	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths <i>Acronicta</i> Knot Grass	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths <i>Acronicta</i> Dark Dagger	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths <i>Actebia pr</i> Portland Mc	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths <i>Adscita ge</i> Cistus Fore	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths <i>Adscita st</i> The Foreste	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths <i>Aethalura</i> Grey Birch	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths <i>Agriopis a</i> Scarce Umbe	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths <i>Agriopis l</i> Spring Ushe	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths <i>Agriopis m</i> Dotted Bore	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths <i>Agrochola</i> The Brick	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%

Moths	<i>Agrochola</i> <i>fl</i> Flounced Ch	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Agrochola</i> <i>br</i> Brown-spot	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Agrochola</i> <i>rl</i> Red-line Qu	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Agrochola</i> <i>be</i> Beaded Ches	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Agrochola</i> <i>yl</i> Yellow-line	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Agrotis</i> <i>cil</i> Light Feath	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Agrotis</i> <i>ex</i> Heart & Dar	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Agrotis</i> <i>ri</i> Sand Dart	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Alcis</i> <i>juba</i> Dotted Carp	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Aleucis</i> <i>dis</i> Sloe Carpet	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Allophyes</i> <i>g</i> Green-brinc	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Alsophila</i> <i>m</i> March Moth	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Amphipoea</i> <i>c</i> Crinan Ear	0	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Moths	<i>Amphipoea</i> <i>s</i> Saltern Ear	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Amphipoea</i> <i>l</i> Large Ear	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Angerona</i> <i>p</i> Orange Motl	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Anticlea</i> <i>d</i> The Streame	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Moths	<i>Anticollix</i> <i>d</i> Dentated Pu	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Antitype</i> <i>c</i> Grey Chi	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Apamea</i> <i>anc</i> Large Nutme	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Apamea</i> <i>fur</i> The Confuse	0	< -7.5%	-4 to -1%	HIGH	< +1%
Moths	<i>Apamea</i> <i>lit</i> Light Arche	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Apamea</i> <i>obl</i> Crescent St	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Apamea</i> <i>oph</i> Double Lobe	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Apamea</i> <i>sco</i> Slender Bri	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Apamea</i> <i>sor</i> Rustic Shou	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Apamea</i> <i>sub</i> Reddish Lig	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Apamea</i> <i>unai</i> Small Clouc	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Apeira</i> <i>syr</i> Lilac Beaut	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Apocheima</i> <i>s</i> Small Brinc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Aporophylla</i> <i>f</i> Feathered F	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Aporophylla</i> <i>b</i> Black Rusti	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Archanara</i> <i>t</i> Twin-spotte	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Archanara</i> <i>w</i> Webb's Wair	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Archiearis</i> <i>l</i> Light Orang	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Arctia</i> <i>caj</i> Garden Tig	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Arctia</i> <i>vil</i> Cream-spot	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Arenostola</i> <i>f</i> Fen Wainsco	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Asthena</i> <i>al</i> Small White	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Atethmia</i> <i>c</i> Centre-barr	1	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Atolmis</i> <i>ru</i> Red-necked	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Autographa</i> <i>g</i> Gold Spangl	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Autographa</i> <i>p</i> Plain Golde	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Axylia</i> <i>put</i> The Flame	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Bena</i> <i>bicol</i> Scarce Silv	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Biston</i> <i>str</i> Oak Beauty	0	-7.5 to -4%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Blepharita</i> <i>d</i> Dark Brocac	1	< -7.5%	> -1%	MODERATE	+1 to +4%

Moths	<i>Cabera exa</i>	Common Wave	0 > -1%	-4 to -1%	MODERATE	< +1%
Moths	<i>Callimorpha</i>	Scarlet Tig	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Callistege</i>	Mother Ship	0 < -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Calophasia</i>	Toadflax Br	0 < -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Camptogram</i>	Yellow Shel	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Caradrina</i>	Mottled Rus	1 < -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Carsia sor</i>	Manchester	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Catarhoe ci</i>	Royal Mantl	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Catarhoe ri</i>	Ruddy Carpe	0 > -1%	> -1%	LOW	> +7.5%
Moths	<i>Catocala ni</i>	Red Underwi	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Celaena ha</i>	Haworth's M	1 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Cepphis ad</i>	Little Thor	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Cerastis l</i>	White-marke	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Cerura vim</i>	Puss Moth	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Charanyca</i>	Treble Line	0 > -1%	> -1%	LOW	> +7.5%
Moths	<i>Chesias ru</i>	Broom-tip	1 < -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Chilodes m</i>	Silky Wains	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Chlorissa</i>	Small Grass	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Chloroclys</i>	Dark Marble	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Chloroclys</i>	Arran Carpe	0 < -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Chloroclys</i>	Autumn Gree	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Chloroclys</i>	Red-green C	0 > -1%	> -1%	LOW	> +7.5%
Moths	<i>Chortodes</i>	Mere Wainsc	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cidaria fu</i>	Barred Yell	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Clostera ci</i>	Chocolate-t	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Coenobia ri</i>	Small Rufou	0 > -1%	> -1%	LOW	> +7.5%
Moths	<i>Coenocalpe</i>	Slender-str	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Colocasia</i>	Nut-tree Tu	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Colotois p</i>	Feathered T	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Comibaena</i>	Blotched Er	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Conistra l</i>	Dark Chestr	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Conistra ri</i>	Dotted Ches	0 > -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Coscinia ci</i>	Speckled Fo	0 < -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Cosmia aff</i>	Lesser-spot	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cosmia pyr</i>	Lunar-spot	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cosmia tra</i>	The Dun-bar	0 > -1%	> -1%	LOW	+1 to +4%
Moths	<i>Cossus cos</i>	Goat Moth	1 < -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Craniophor</i>	The Coronet	0 > -1%	> -1%	LOW	> +7.5%
Moths	<i>Crocallis</i>	Scalloped C	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Cryphia mu</i>	Marbled Gre	0 < -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Cucullia a</i>	The Wormwo	0 < -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Cucullia a</i>	Star-wort	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Cucullia ci</i>	Chamomile S	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cucullia u</i>	The Shark	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cybosia me</i>	Four-dotted	0 < -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Cyclophora</i>	The Mocha	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Cyclophora</i>	Clay Triple	0 > -1%	< -7.5%	MODERATE	> +7.5%

Moths	<i>Cyclophora</i>	Dingy Mocha	1	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Cyclophora</i>	False Mocha	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Cyclophora</i>	Maiden's Bl	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Cymatophor</i>	Oak Lutestr	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Dasypolia</i>	Brindled Oc	1	< -7.5%	-4 to -1%	HIGH	< +1%
Moths	<i>Deilephila</i>	Elephant Ha	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Deilephila</i>	Small Eleph	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Deltote ba</i>	Silver Barr	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Deltote un</i>	Silver Hook	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Diacrisia</i>	Clouded But	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Diarsia da</i>	Barred Ches	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Moths	<i>Dicallomer</i>	Dark Tussock	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Dichonia a</i>	Merveille c	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Diloba cae</i>	Figure of I	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Discoloxia</i>	Blomer's Ri	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Moths	<i>Drepana fa</i>	Pebble Hook	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Drymonia d</i>	Marbled Bro	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Moths	<i>Dryobotode</i>	Brindled Gr	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Dypterygia</i>	Bird's Wing	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Dyscia fag</i>	Grey Scallo	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Earias clo</i>	Cream-borde	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Egira cons</i>	Silver Clou	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Eilema can</i>	Hoary Footr	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Eilema com</i>	Scarce Foot	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eilema dep</i>	Buff Footma	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Eilema gri</i>	Dingy Footr	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Eilema lur</i>	Common Foot	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Eilema pygi</i>	Pigmy Footr	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eilema sor</i>	Orange Foot	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Elaphria v</i>	Rosy Marble	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Enargia pa</i>	Angle-strip	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Endromis v</i>	Kentish Glo	0	> -1%	< -7.5%	MODERATE	< +1%
Moths	<i>Ennomos al</i>	Canary-shou	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Ennomos au</i>	Large Thor	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Ennomos er</i>	September J	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Ennomos qu</i>	August Thor	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Entephria</i>	Grey Mounta	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Entephria</i>	Yellow-ring	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Epione rep</i>	Bordered Be	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Epirrhoe r</i>	Wood Carpet	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Epirrhoe t</i>	Small Arger	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Epirrita ai</i>	Autumnal Mo	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Epirrita ci</i>	Pale Novemb	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Epirrita f</i>	Small Autum	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Eremobia o</i>	Dusky Sallc	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Eriogaster</i>	Small Eggar	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Euchoeca n</i>	Dingy Shell	0	> -1%	> -1%	LOW	> +7.5%

Moths	<i>Euclidia g</i>	Burnet Com	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Eugnorisma</i>	Plain Clay	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Eugnorisma</i>	Autumnal R	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Eulithis m</i>	The Spinach	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eulithis p</i>	The Phoenix	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Eulithis p</i>	Barred Stra	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Euphyia bi</i>	Cloaked Car	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Eupithecia</i>	Brindled Pt	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Wormwood Pt	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Currant Pug	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Thyme Pug	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Eupithecia</i>	Oak-tree Pt	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Pauper Pug	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Mottled Pug	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Haworth's F	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Eupithecia</i>	Tawny Speck	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Eupithecia</i>	Pinion-spot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Maple Pug	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Marbled Pug	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Eupithecia</i>	Larch Pug	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Toadflax Pt	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Yarrow Pug	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Narrow-wing	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Pimpinel Pt	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Lead-colour	0	> -1%	> -1%	LOW	< +1%
Moths	<i>Eupithecia</i>	Foxglove Pt	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Eupithecia</i>	Satyr Pug	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Eupithecia</i>	Plain Pug	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Shaded Pug	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Bordered Pt	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	White-spot	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Golden-rod	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Common Pug	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Euproctis</i>	Brown-tail	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupsilia t</i>	The Satellit	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Eurois occi</i>	Great Broc	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Euxoa curs</i>	Coast Dart	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Euxoa trit</i>	White-line	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Furcula bi</i>	Alder Kitten	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Furcula bi</i>	Poplar Kitten	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Furcula fu</i>	Sallow Kitten	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Gnophos ob</i>	Scotch Ann	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Gortyna fl</i>	Frosted Ora	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Graphiphora</i>	Double Dart	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Gymnosceli</i>	Double-stri	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Habrosyne</i>	Buff Arches	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Hadena alb</i>	White Spot	1	< -7.5%	> -1%	MODERATE	> +7.5%

Moths	<i>Hadena com</i>	Varied Corc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Hecatera b</i>	Broad-barre	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Heliophobu</i>	Bordered Go	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Hemistola</i>	Small Emera	1	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Hepialus h</i>	Gold Swift	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Hepialus s</i>	Orange Swif	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Herminia g</i>	Small Fan-f	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Hoplodrina</i>	The Uncerta	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Hoplodrina</i>	The Rustic	1	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Horisme te</i>	The Fern	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Hydrelia f</i>	Small Yello	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Hydriomena</i>	May Highfly	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Moths	<i>Hydriomena</i>	Ruddy Highf	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Hylaea fas</i>	Barred Red	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Hyles gall</i>	Bedstraw Ha	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Hypena pro</i>	The Snout	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Hypenodes</i>	Marsh Oblic	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Hypomecis</i>	Pale Oak Be	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Hyppa rect</i>	The Saxon	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Idaea aver</i>	Riband Wave	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Idaea bise</i>	Small Fan-f	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Idaea dimi</i>	Single-dott	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Idaea emar</i>	Small Scall	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Idaea fusc</i>	Dwarf Crear	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Idaea muri</i>	Purple-borc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Idaea seri</i>	Small Dusty	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Idaea subs</i>	Satin Wave	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Idaea trig</i>	Treble Brov	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Ipimorpha</i>	Double Kidr	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Ipimorpha</i>	The Olive	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Itame brun</i>	Rannoch Loc	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Jodis lact</i>	Little Emer	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Lacanobia</i>	Bright-line	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Lacanobia</i>	Light Broc	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Lampropter</i>	Water Carpe	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Moths	<i>Laothoe po</i>	Poplar Hawk	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Larentia c</i>	The Mallow	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Lasiocampa</i>	Oak Eggar	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Lasiocampa</i>	Grass Eggar	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Laspeyria</i>	Beautiful F	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Leucochlae</i>	Beautiful C	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Leucoma sa</i>	White Satir	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Lithomoia</i>	Golden-rod	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Lithophane</i>	Pale Pinior	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Lithophane</i>	Grey Shoulc	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Lithosia q</i>	Four-spotte	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Lobophora</i>	The Seraphi	0	< -7.5%	> -1%	MODERATE	> +7.5%

Moths	<i>Lomographa</i>	White-pink	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Luperina n.</i>	Sandhill R	0	> -1%	> -1%	LOW	< +1%
Moths	<i>Lycia hirt.</i>	Brindled B	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Lygephila i.</i>	The Blackne	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Macaria al.</i>	Sharp-angle	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Macaria no.</i>	Peacock Mo	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Macaria wai.</i>	The V-Moth	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Macrochilo</i>	Dotted Fan-	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Macroglossi</i>	Hummingbird	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Macrothyla</i>	Fox Moth	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Malacosoma</i>	The Lackey	1	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Mamestra b.</i>	Cabbage Mo	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Meganola s.</i>	Small Black	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Melanchra i.</i>	Dot Moth	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Melanthia i.</i>	Pretty Chal	1	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Menophra al.</i>	Waved Umber	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Mesoligia i.</i>	Cloaked Mir	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Miltochristis</i>	Rosy Footma	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Mimas tili.</i>	Lime Hawk-m	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Moma alpiu.</i>	Scarce Merv	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Mythimna a.</i>	White-point	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Mythimna c.</i>	Shoulder-st	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Mythimna o.</i>	Obscure Wai	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Mythimna p.</i>	Common Wair	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Mythimna p.</i>	Striped Wai	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Mythimna p.</i>	Devonshire	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Mythimna t.</i>	Double Line	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Mythimna u.</i>	White-speck	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Naenia typ.</i>	The Gothic	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Nebula sal.</i>	Striped Twi	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Noctua com.</i>	Lesser Yell	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Noctua fim.</i>	Broad-borde	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Nola confu.</i>	Least Black	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Nonagria t.</i>	Bulrush Wai	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Notodonta i.</i>	Iron Promir	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Notodonta i.</i>	Pebble Prom	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Nudaria mu.</i>	Muslin Foot	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Moths	<i>Odezia atr.</i>	Chimney Swe	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Odontopera</i>	Scalloped F	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Odontosia i.</i>	Scarce Prom	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Oligia fas.</i>	Middle-barr	0	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Oligia lat.</i>	Tawny Marbl	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Oligia str.</i>	Marbled Mir	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Oligia ver.</i>	Rufous Minc	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Omphalosce.</i>	Lunar Under	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Operophter.</i>	Winter Moth	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Operophter.</i>	Northern Wi	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Moths	<i>Opisthogra</i>	Brimstone M	0	> -1%	> -1%	LOW	< +1%
Moths	<i>Oria muscu</i>	Brighton W	1	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Orthosia c</i>	Small Quake	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Orthosia g</i>	Hebrew Char	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Orthosia g</i>	Powdered Qu	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Orthosia i</i>	Clouded Dra	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Orthosia m</i>	Blossom Unc	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Orthosia m</i>	Twin-spotte	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Orthosia o</i>	Northern Di	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Orthosia p</i>	Lead-colour	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Ourapteryx</i>	Swallow-tai	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Pachycnemi</i>	Horse Chest	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Panemeria</i>	Small Yello	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Panolis fl</i>	Pine Beauty	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Papestra b</i>	Glaucous Sh	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Paracolax</i>	Clay Fan-f	1	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Paradarisa</i>	Square Spot	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Paradrina</i>	Pale Mottle	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Parascotia</i>	Waved Black	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Parasemia</i>	Wood Tiger	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Parectropi</i>	Brindled Wh	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Pasiphila</i>	Sloe Pug	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Pasiphila</i>	Bilberry Pu	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Pasiphila</i>	Green Pug	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Pelosia mu</i>	Dotted Foot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Pelurga co</i>	Dark Spinac	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Perconia s</i>	Grass Wave	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Peribatode</i>	Willow Beau	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Peridea an</i>	Great Promi	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Perizoma a</i>	Small Rivul	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Perizoma b</i>	Pretty Pini	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Perizoma d</i>	Twin-spot (0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Perizoma f</i>	Sandy Carpe	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Perizoma s</i>	Marsh Carpe	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Phalera bu</i>	Buff-tip	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Phibalapte</i>	Oblique Str	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Philereme</i>	Dark Umber	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Philereme</i>	Brown Scall	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Phlogophor</i>	Angle Shade	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Photedes c</i>	Least Minor	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Phytometra</i>	Small Purpl	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Plagodis d</i>	Scorched Wi	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Plagodis p</i>	Barred Umbe	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Plemyria r</i>	Blue-border	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Plusia fes</i>	Gold Spot	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Plusia put</i>	Lempke's G	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Poecilocamp</i>	December Mo	0	< -7.5%	> -1%	MODERATE	< +1%

Moths	<i>Polia bomby</i>	Pale Shinier	1	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Moths	<i>Polia nebulosa</i>	Grey Arches	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Polia trimaculata</i>	Silvery Arch	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Polymixis</i>	Large Ranunc	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Polymixis</i>	Feathered I	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Polyplocia</i>	Frosted Gre	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Moths	<i>Pteraphera</i>	Small Sera	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Pterostoma</i>	Pale Promi	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Ptilodon</i>	Maple Promi	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Ptilophora</i>	Plumed Prom	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Moths	<i>Pyrrhia um</i>	Bordered Sa	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Rheumaptera</i>	Scarce Tiss	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Rheumaptera</i>	Argent & Sa	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Rheumaptera</i>	Scallop She	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Rhizedra</i>	Large Wains	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Rhyacia sin</i>	Dotted Rust	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Rivula ser.</i>	Straw Dot	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Saturnia pe</i>	Emperor Mot	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Schrankia</i>	Pinion-stre	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Schrankia</i>	White-line	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Scopula em</i>	Rosy Wave	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Scopula fl</i>	Cream Wave	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Scopula im</i>	Small Bloo	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Moths	<i>Scopula ru</i>	Tawny Wave	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Moths	<i>Scopula te</i>	Smoky Wave	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Moths	<i>Scotoptery</i>	Chalk Carpe	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Scotoptery</i>	July Belle	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Scotoptery</i>	Lead Belle	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Selenia dei</i>	Early Thor	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Selenia lu</i>	Lunar Thor	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Selidosema</i>	Bordered Gr	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Semiaspila</i>	Yellow Bell	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Sesia bemb</i>	Lunar Horne	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Setina irr</i>	Dew Moth	0	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Shargacucu</i>	Striped Lyc	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Shargacucu</i>	The Mulleir	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Simyra alb</i>	Reed Dagger	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Spaelotis</i>	Stout Dart	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Spargania</i>	White-bande	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Spilosoma</i>	Buff Ermine	1	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Spilosoma</i>	Water Ermi	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Stilbia an</i>	The Anomalc	1	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Tethea ocu</i>	Figure of I	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Tetheella</i>	Satin Lutes	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Thalophila</i>	Straw Under	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Thera cogn</i>	Chestnut-co	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Moths	<i>Thera cupr</i>	Cypress Car	0	> -1%	> -1%	LOW	> +7.5%

Moths	<i>Thera firm</i>	Pine Carpet	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Thera juni</i>	Juniper Car	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Theria pri</i>	Early Moth	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Tholera ce</i>	Hedge Rusti	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Tholera de</i>	Feathered (1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Thumatha s</i>	Round-wing	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Trichiura</i>	(Pale Eggar	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Trichopter</i>	Early Tooth	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Trichopter</i>	Barred Toth	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Triphosa d</i>	The Tissue	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Trisateles</i>	Olive Cresc	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Tyta luctu</i>	The Four-sp	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Venusia cai</i>	Welsh Wave	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Moths	<i>Watsonalla</i>	Barred Hook	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Xanthia ci</i>	Orange Sall	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Xanthia gi</i>	Dusky-lemor	1	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Xanthia ic</i>	The Sallow	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Xanthia oc</i>	Pale-lemon	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Xanthorhoe</i>	Balsam Carp	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Xanthorhoe</i>	Red Carpet	1	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Xanthorhoe</i>	Garden Carp	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Xanthorhoe</i>	Large Twin-	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Xestia aga</i>	Heath Rusti	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Xestia tri</i>	Double Squa	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Xylena exs</i>	Sword-grass	0	< -7.5%	-4 to -1%	HIGH	< +1%
Moths	<i>Xylena veti</i>	Red Sword-g	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Xylocampa</i>	(Early Grey	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Zanclognat</i>	The Fan-foc	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Zeuzera py</i>	Leopard Mot	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Zygaena loi</i>	Narrow-bor	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Odonata	<i>Aeshna cae</i>	Azure hawke	0	> -1%	< -7.5%	MODERATE	> +7.5%
Odonata	<i>Aeshna gra</i>	Brown hawke	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Odonata	<i>Aeshna jun</i>	Common haw	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Odonata	<i>Anax imper</i>	Emperor dra	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Brachytron</i>	Hairy drag	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Calopteryx</i>	Banded dem	0	> -1%	< -7.5%	MODERATE	> +7.5%
Odonata	<i>Ceriagrion</i>	Small red c	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Coenagrion</i>	Azure damse	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Odonata	<i>Cordulegas</i>	Golden-ring	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Odonata	<i>Cordulia a</i>	Downy emera	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Odonata	<i>Enallagma</i>	(Common blue	0	< -7.5%	> -1%	MODERATE	< +1%
Odonata	<i>Erythromma</i>	Red-eyed da	0	> -1%	< -7.5%	MODERATE	> +7.5%
Odonata	<i>Gomphus vu</i>	Club-tailed	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Ischnura e</i>	Blue-tailed	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Odonata	<i>Ischnura pi</i>	Scarce blue	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Lestes spo</i>	Emerald dan	0	< -7.5%	> -1%	MODERATE	< +1%
Odonata	<i>Libellula</i>	(Broad-bodie	0	< -7.5%	-4 to -1%	HIGH	> +7.5%

Odonata	<i>Libellula</i>	Four-spotted	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Orthetrum</i>	Black-tailed	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Orthetrum</i>	Keeled skimmer	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Pyrrhosoma</i>	Large red club	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Odonata	<i>Somatochlora</i>	Brilliant club	0	< -7.5%	> -1%	MODERATE	> +7.5%
Odonata	<i>Sympetrum</i>	Black dart	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Odonata	<i>Sympetrum</i>	Yellow-wing	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Odonata	<i>Sympetrum</i>	Ruddy dart	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Sympetrum</i>	Common dart	0	< -7.5%	> -1%	MODERATE	> +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH	< +1%
Soldier beetle	<i>Cantharis</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	-4 to -1%	< -7.5%	HIGH	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Soldier beetle	<i>Malthinus</i>	NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Malthinus</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Soldier beetle	<i>Malthodes</i>	NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Podabrus</i>	a. NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	Common Red	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Spiders	<i>Achaearanea</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Achaearanea</i>	NA	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Spiders	<i>Agalenatea</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Agelena</i>	labyrinth	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Agroeca</i>	br. NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Agroeca</i>	in. NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Agroeca</i>	pr. NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Agyneta</i>	de. NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Agyneta</i>	ol. NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Agyneta</i>	rai. NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Agyneta</i>	su. NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Allomengea</i>	NA	0	-4 to -1%	-4 to -1%	MODERATE	< +1%
Spiders	<i>Allomengea</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Alopecosa</i>	i. NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Alopecosa</i>	i. NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Alopecosa</i>	i. NA	0	> -1%	> -1%	LOW	+4 to +7.5%

Spiders	<i>Amaurobius</i> NA	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Amaurobius</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Amaurobius</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Anelosimus</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Anyphaena</i> Buzzing Spi	0 > -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Aphileta m.</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Araeoncus</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Araneus ma</i> NA	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Spiders	<i>Araneus qu</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Araneus st</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Araneus tr</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Araniella</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Araniella</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Arctosa le</i> NA	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Arctosa pe</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Argenna su</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Argyroneta</i> Water Spide	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Atypus aff.</i> Purse Web S	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Ballus cha</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Baryphyma</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Baryphyma</i> NA	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Spiders	<i>Bathyphant</i> NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Bathyphant</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Bathyphant</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Bathyphant</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Bianor aur</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Bolyphante</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Bolyphante</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Centromeri</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Centromeru</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Centromeru</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Centromeru</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Centromeru</i> NA	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Ceratinell</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Ceratinell</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Ceratinell</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ceratinops</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Cercidia p</i> NA	0 < -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Cheiracanti</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Cheiracanti</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Cicurina c</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Clubiona b</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona c</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona c</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona l</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Clubiona n</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Spiders	<i>Clubiona n</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%

Spiders	<i>Clubiona n</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Clubiona p</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona s</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Clubiona t</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Coelotes a</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Coelotes t</i> NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Spiders	<i>Crustulina</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Crustulina</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Cryphoea</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Diaea dors</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Dictyna ar</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Dictyna la</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Dictyna pu</i> Small Mesh-	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Dictyna un</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Dicymbium</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Diplocentr</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Diploceph</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Diploceph</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Diplostyla</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Dismodicus</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Dolomedes</i> Raft Spider	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Donacochar</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Spiders	<i>Drassodes</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Drassyllus</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Drepanotyl</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Dysdera cr</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Dysdera er</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Enoplognat</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Enoplognat</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Enoplognat</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Episinus a</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Episinus t</i> NA	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Spiders	<i>Erigone ar</i> NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Erigone at</i> NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Erigone loi</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Erigone pr</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Ero cambri</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Ero furcat</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ero tuberc</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Euophrys f</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Evarcha ar</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Evarcha fa</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Spiders	<i>Floronia b</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Gibbaranea</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Gnaphosa l</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Gnathonari</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Gongylidie</i>	NA	0	> -1%	> -1%	LOW	< +1%
Spiders	<i>Gongylidiu</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Hahnia nav</i>	NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Halorates</i>	NA	0	> -1%	> -1%	LOW	< +1%
Spiders	<i>Haplodrass</i>	Heath Gras	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Haplodrass</i>	NA	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Haplodrass</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Harpactea</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Heliophanu</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Heliophanu</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Hilaira ex</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Hilaira fr</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Hilaira pe</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Hylyphante</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Hypomma co</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Hypomma fu</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Hypseliste</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Hypsosinga</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Hypsosinga</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Hypsosinga</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Kaestneria</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Kaestneria</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Labulla th</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Larinioide</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Lathys hum</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Latithorax</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Lepthyphan</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Lepthyphan</i>	NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Lepthyphan</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Lepthyphan</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Lepthyphan</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Lepthyphan</i>	NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Lepthyphan</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Leptothrix</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Spiders	<i>Linyphia h</i>	NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Spiders	<i>Linyphia t</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Lophomma p</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Mangora ac</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Maro minuti</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Maso sunde</i>	NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Mecopisthe</i>	Peus' s Long	1	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Meioneta i</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Meioneta m</i>	Thin Weblet	1	< -7.5%	> -1%	MODERATE	< +1%

Spiders	<i>Meioneta r</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Meioneta s</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Metellina i</i> NA	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Spiders	<i>Metellina i</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Metopobact</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Micrargus i</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Micrargus i</i> NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Spiders	<i>Micrargus i</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Microlinyp</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Microlinyp</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Micrommata</i> Green Spide	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Milleriana</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Minyriolus</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Misumena v</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Moebelia p</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Monocephali</i> Broad Groov	1	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Spiders	<i>Monocephali</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Neoscona a</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Neriene cl</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Neriene fu</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Neriene moi</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Neriene pe</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Nesticus c</i> Comb-footec	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Nuctenea u</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Oedothorax</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Oedothorax</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Oedothorax</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Oreonetide</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Ostearius i</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ozyptila b</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ozyptila p</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Ozyptila s</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Ozyptila s</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Ozyptila t</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Pachygnath</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Pachygnath</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Pachygnath</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Panamomops</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Pardosa ag</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Pardosa am</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Pardosa ho</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Pardosa moi</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Pardosa ni</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Pardosa pa</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pardosa pr</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Pardosa pu</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Pardosa sa</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%

Spiders	<i>Pelecopsis</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Pelecopsis</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Pelecopsis</i> NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Spiders	<i>Pelecopsis</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Philodromu</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Philodromu</i> .NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Spiders	<i>Phrurolithu</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pirata lat</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Pirata pir</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Pirata pis</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Spiders	<i>Pisaura mi</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pityohyphai</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Pocadicnem</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pocadicnem</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Poecilonet</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Porrhomma i</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Porrhomma i</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Porrhomma i</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Porrhomma i</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Porrhomma i</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Robertus ai</i> NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Robertus l</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Saaristoa i</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Saaristoa i</i> Triangle H	1	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Spiders	<i>Saloca dic</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Salticus c</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Salticus s</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Satilatlas</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Scotina gr</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Scotophaeu</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Segestria i</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Silometopu</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Silometopu</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Silometopu</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Singa hama</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Sitticus c</i> Sedge Jump	1	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Sitticus pi</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Sitticus s</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Stemonyphai</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Syedra gra</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Tallusia e</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Tapinocyba</i> NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%

Spiders	<i>Tapinocyba</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Tapinopa</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Taranucnus</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Tegenaria</i>	NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Spiders	<i>Tegenaria</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Tegenaria</i>	House Spider	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Tegenaria</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Tetragnatha</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Tetragnatha</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Tetragnatha</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Textrix</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Thanatus</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Theonoe</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Theridion</i>	NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Spiders	<i>Theridion</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Theridion</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Theridion</i>	NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Theridion</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Theridion</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Theridion</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Theridiosoma</i>	Ray Spider	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Tibellus</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Tmetiscus</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Trichopterus</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Trochosa</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Walckenaera</i>	NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Walckenaera</i>	NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Spiders	<i>Walckenaera</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Spiders	<i>Walckenaera</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Xysticus</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Xysticus</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Xysticus</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Xysticus</i>	NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Xysticus</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Zelotes</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Zilla</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Zora</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Zygiella</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular plant	<i>Acer</i>	Field Maple	0	> -1%	> -1%	LOW	+4 to +7.5%

Vascular plant	<i>Aceras</i>	ant	Man Orchid	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular plant	<i>Achillea</i>	p	Sneezewort	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular plant	<i>Aconitum</i>	n	Monk's-hood	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Actaea</i>	spi	Baneberry	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular plant	<i>Adiantum</i>	c	Maidenhair	0	> -1%	> -1%	LOW	> +7.5%
Vascular plant	<i>Adoxa</i>	mosc	Moschatel	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Aethusa</i>	cy	Fool's Pars	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Aethusa</i>	cy	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular plant	<i>Agrimonia</i>		Agrimony	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular plant	<i>Agrostemma</i>		Corncockle	0	> -1%	> -1%	LOW	> +7.5%
Vascular plant	<i>Agrostis</i>	c	Velvet Bent	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Agrostis</i>	c	Bristle Ber	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular plant	<i>Agrostis</i>	g	Black Bent	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Agrostis</i>	s	Creeping Be	0	> -1%	> -1%	LOW	+1 to +4%
Vascular plant	<i>Agrostis</i>	v	Brown Bent	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Ajuga</i>	pyra	Pyramidal F	1	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular plant	<i>Ajuga</i>	rept	Bugle	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular plant	<i>Alchemilla</i>		Alpine Lady	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular plant	<i>Alchemilla</i>		Hairy Lady'	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular plant	<i>Alchemilla</i>		Slender Lac	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular plant	<i>Alchemilla</i>		Smooth Lady	0	> -1%	-7.5 to -4%	MODERATE	+1 to +4%
Vascular plant	<i>Alchemilla</i>		Pale Lady's	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Alisma</i>	lan	Narrow-leav	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular plant	<i>Alisma</i>	pl	Water-plant	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular plant	<i>Alliaria</i>	p	Garlic Must	0	> -1%	> -1%	LOW	+1 to +4%
Vascular plant	<i>Allium</i>	ole	Field Garli	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Allium</i>	sco	Sand Leek	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Allium</i>	urs	Ramsons	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Alnus</i>	glut	Alder	0	-7.5 to -4%	> -1%	MODERATE	< +1%
Vascular plant	<i>Alopecurus</i>		Bulbous Fox	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Alopecurus</i>		Marsh Foxt	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular plant	<i>Alopecurus</i>		Black-grass	0	> -1%	> -1%	LOW	> +7.5%
Vascular plant	<i>Alopecurus</i>		Meadow Fox	0	-7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Vascular plant	<i>Althaea</i>	of	Marsh-malle	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular plant	<i>Anagallis</i>		Scarlet Pin	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular plant	<i>Anagallis</i>		Scarlet Pin	0	> -1%	> -1%	LOW	> +7.5%
Vascular plant	<i>Anagallis</i>		Chaffweed	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular plant	<i>Andromeda</i>		Bog-rosemar	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular plant	<i>Anemone</i>	nei	Wood Anemor	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular plant	<i>Antennaria</i>		Mountain Ey	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Vascular plant	<i>Anthemis</i>	a	Corn Chamom	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular plant	<i>Anthoxanthi</i>		Sweet Verna	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular plant	<i>Anthriscus</i>		Cow Parsley	0	> -1%	> -1%	LOW	+1 to +4%
Vascular plant	<i>Apera</i>	spic	Loose Silky	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular plant	<i>Aphanes</i>	ar	Parsley-pic	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular plant	<i>Aphanes</i>	au	Slender Par	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular plant	<i>Apium</i>	inun	Lesser Mars	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular pl	<i>Apium nodi</i>	Fool's-water	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Arctium lappula</i>	Greater Burdock	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular pl	<i>Arctium minus</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Arenaria serpyllifolia</i>	Thyme-leaved Cress	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Arenaria serotina</i>	Slender Sandwort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Arenaria serotina</i>	Thyme-leaved Cress	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Armoracia rusticana</i>	Horse-radish	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Arrhenathera elatior</i>	False Oat-grass	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Artemisia vulgaris</i>	Mugwort	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Arum italicum</i>	Italian Loropetalum	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Arum maculatum</i>	Lords-and-Ladies	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Asparagus officinalis</i>	Garden Asparagus	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Asperula cynodactylus</i>	Squinancywort	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Asplenium adnigrum</i>	Black Spleenwort	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Asplenium lanceolatum</i>	Lanceolate Spleenwort	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular pl	<i>Asplenium platyneuron</i>	Maidenhair	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Asplenium septentrionale</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Asplenium septentrionale</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Athyrium filix-femina</i>	Lady-fern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Atriplex barbinervis</i>	Babington's Salt-marsh	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Atriplex patula</i>	Common Orache	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular pl	<i>Atriplex patula</i>	Spear-leaved Orache	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Atropa belladonna</i>	Deadly Nightshade	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular pl	<i>Avena fatua</i>	Wild-oat	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Baldellia hederifolia</i>	Lesser Water-pennywort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Ballota nigra</i>	Black Horehound	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular pl	<i>Barbarea vulgaris</i>	Winter-cress	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Vascular pl	<i>Bellis perennis</i>	Daisy	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Beta vulgaris</i>	Beet	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Betula pendula</i>	Silver Birch	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Betula pubescens</i>	Downy Birch	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Betula pubescens</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Bidens biternata</i>	Nodding Burdock	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Blackstonia perfoliata</i>	Yellow-wort	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Blechnum spicatum</i>	Hard-fern	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular pl	<i>Blysmus cristatus</i>	Flat-sedge	1	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Brachypodium pinnatifidum</i>	Tor-grass	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Brachypodium pinnatifidum</i>	False Bromegrass	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Brassica oleracea</i>	Cabbage	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Brassica rapa</i>	Turnip	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Briza media</i>	Quaking-grass	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Briza minor</i>	Lesser Quaking-grass	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Bromopsis inermis</i>	Hairy-brome	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Bromus horridus</i>	Soft-brome	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular pl	<i>Bromus horridus</i>	Least Soft-brome	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Bromus horridus</i>	Common Soft-brome	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Bromus horridus</i>	Sand Soft-brome	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p <i>Bromus rac</i>	Smooth Bro	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Bryonia di</i>	White Bryo	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p <i>Bupleurum</i>	Slender Ha	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Buxus semp</i>	Box	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Cakile mar</i>	Sea Rocket	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Calamagros</i>	Purple Smal	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Calamagros</i>	Narrow Smal	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Callitrich</i>	Intermediat	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Callitrich</i>	Autumnal We	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Callitrich</i>	Blunt-fruit	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Callitrich</i>	Common Wate	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Callitrich</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Calluna vu</i>	Heather	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Caltha pal</i>	Marsh-mari	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Calystegia</i>	Hedge Bind	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Calystegia</i>	NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Calystegia</i>	Great Bind	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Calystegia</i>	Sea Bindwe	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Campanula</i>	Giant Bell	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Campanula</i>	Harebell	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Capsella bi</i>	Shepherd's-	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Cardamine</i>	Large Bitt	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Cardamine</i>	Coralroot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Cardamine</i>	Wavy Bitter	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Cardamine</i>	Hairy Bitt	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Cardamine</i>	Narrow-leav	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Cardamine</i>	Cuckooflow	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Carduus cr</i>	Wetted Th	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Carex acut</i>	Slender Tus	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Carex appr</i>	Fibrous Tus	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Carex aqua</i>	Water Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Carex atra</i>	Black Alpi	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Carex bige</i>	Stiff Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Carex bine</i>	Green-ribb	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p <i>Carex capi</i>	Hair Sedge	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Carex dian</i>	Lesser Tus	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Carex digi</i>	Fingered S	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Carex dioi</i>	Dioecious S	0	> -1%	-7.5 to -4%	MODERATE	+1 to +4%
Vascular p <i>Carex dist</i>	Distant Sec	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Carex divu</i>	Grey Sedge	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Carex divu</i>	Many-leave	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Carex echi</i>	Star Sedge	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p <i>Carex elon</i>	Elongated S	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Carex eric</i>	Rare Spring	1	< -7.5%	< -7.5%	VERY HIGH	< +1%
Vascular p <i>Carex hirt</i>	Hairy Sedge	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Carex host</i>	Tawny Sedge	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Vascular p <i>Carex humi</i>	Dwarf Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular pl	<i>Carex mages</i>	Tall Bog-sedge	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Carex muric</i>	Prickly Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Carex muric</i>	Small-fruit Sedge	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular pl	<i>Carex nigr</i>	Common Sedge	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Carex otrubae</i>	False Foxglove	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Carex ovalis</i>	Oval Sedge	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Carex panicea</i>	Carnation Sedge	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Vascular pl	<i>Carex panicea</i>	Greater Tussock Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Carex pauciflora</i>	Few-flowered Sedge	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular pl	<i>Carex pilulifera</i>	Pill Sedge	0	-7.5 to -4%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Carex pseudocyperus</i>	Cyperus Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Carex punctata</i>	Dotted Sedge	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Carex remota</i>	Remote Sedge	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular pl	<i>Carex riparia</i>	Greater Portulaca Sedge	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Carex spicata</i>	Spiked Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Carex stricta</i>	Thin-spiked Sedge	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Carex vaginata</i>	Sheathed Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Carex vesicaria</i>	Bladder-sedge	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Carex viridula</i>	Common Yellow Sedge	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular pl	<i>Carpinus betula</i>	Hornbeam	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Castanea sativa</i>	Sweet Chestnut	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Catabrosa aquatica</i>	Whorl-grass	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Catapodium maritimum</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Vascular pl	<i>Centaurea jacobaea</i>	Cornflower	1	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Centaureium montanum</i>	Common Centaury	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Centaureium maritimum</i>	Seaside Centaury	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Cephalanthus occidentalis</i>	White Heliotrope	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Cerastium arvense</i>	Sea Mouse-ear	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Cerastium triviale</i>	Common Mouse-ear	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Cerastium vulgatum</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Cerastium vulgatum</i>	NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular pl	<i>Cerastium vulgatum</i>	Dwarf Mouse-ear	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Ceratocarpus trichospermus</i>	Climbing Cereus	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Ceratophyllum demersum</i>	Rigid Hornwort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Ceratophyllum demersum</i>	Soft Hornwort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Chaenorhizum</i>	Small Toadflax	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Chaerophyllum</i>	Rough Chervil	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Chamaemelum nobile</i>	Chamomile	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Chelidonium majus</i>	Greater Celandine	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Chenopodium album</i>	Fat-hen	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular pl	<i>Chenopodium album</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Chenopodium album</i>	Fig-leaved Goosefoot	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Chenopodium album</i>	Stinking Goosefoot	1	> -1%	> -1%	LOW	+1 to +4%
Vascular pl	<i>Chrysanthemum leucanthemum</i>	Corn Marigold	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Chrysosplenium</i>	Opposite-leaved Starwort	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular pl	<i>Cicendia filifolia</i>	Yellow Centaury	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Cichorium intybus</i>	Chicory	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p <i>Cicuta vir</i>	Cowbane	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p <i>Circaea lu</i>	Enchanter's	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Cirsium er</i>	Woolly Thist	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Cirsium he</i>	Melancholy	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Cochlearia</i>	English Sc	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Cochlearia</i>	Common Scu	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Cochlearia</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Cochlearia</i>	Pyrenean Sc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Colchicum</i>	Meadow Saf	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Conopodium</i>	Pignut	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Corallorhi</i>	Coralroot (0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Cornus san</i>	Dogwood	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Cornus sue</i>	Dwarf Corne	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Coronopus</i>	Swine-cress	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Crataegus</i>	Hawthorn	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Crepis cap</i>	Smooth Hawk	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Crepis pal</i>	Marsh Hawk'	0	-4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Vascular p <i>Crithmum m</i>	Rock Samphi	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Cruciata l</i>	Crosswort	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Cryptogram</i>	Parsley Fer	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Cuscuta ep</i>	Dodder	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Cuscuta eu</i>	Greater Doc	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p <i>Cynosurus</i>	Crested Dog	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Cyperus lo</i>	Galingale	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Cystopteris</i>	Brittle Bla	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p <i>Cytisus sc</i>	Broom	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p <i>Cytisus sc</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Dactylorhi</i>	Common Spot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Dactylorhi</i>	Early Marsh	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Dactylorhi</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Dactylorhi</i>	Northern Ma	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Dactylorhi</i>	Narrow-leav	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Daucus car</i>	Carrot	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Daucus car</i>	Wild Carrot	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Daucus car</i>	Sea Carrot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Deschampsia</i>	Wavy Hair-g	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p <i>Deschampsia</i>	Bog Hair-gr	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Dianthus a</i>	Deptford Pi	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Dianthus d</i>	Maiden Pink	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Digitalis</i>	Foxglove	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Dipsacus fi</i>	Wild Teasel	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Dipsacus p</i>	Small Tease	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Draba inca</i>	Hoary Whit	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Draba mura</i>	Wall Whitl	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Drosera an</i>	Great Sunde	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p <i>Drosera in</i>	Oblong-leav	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Dryopteris</i>	Golden-scal	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%

Vascular p <i>Dryopteris</i> Buckler-Fern	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Dryopteris</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Dryopteris</i> Broad Buckl	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Dryopteris</i> Northern Bu	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Dryopteris</i> Male-fern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Dryopteris</i> Mountain Ma	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Dryopteris</i> Rigid Buckl	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Echium vulg</i> Viper's-bug	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Elatine hy</i> Eight-stame	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Eleocharis</i> Common Spik	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Eleocharis</i> Few-flower	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Elymus can</i> Bearded Cou	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p <i>Elytrigia</i> Common Couc	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Empetrum n</i> Crowberry	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Empetrum n</i> Mountain Cr	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Empetrum n</i> Crowberry	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Epilobium</i> Chickweed V	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Epilobium</i> Alpine Will	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Epilobium</i> Great Willc	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p <i>Epilobium</i> Spear-leave	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Epilobium</i> Short-fruit	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Epilobium</i> Marsh Willc	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Epilobium</i> Hoary Willc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Epilobium</i> Pale Willow	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Epilobium</i> Square-stal	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Epipactis</i> Marsh Helle	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Epipactis</i> Green-flowe	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Epipactis</i> Violet Helle	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Equisetum</i> Rough Horse	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Equisetum</i> Shady Horse	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p <i>Equisetum</i> Wood Horset	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p <i>Equisetum</i> Variegated	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Erica cine</i> Bell Heathe	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Erica tetr</i> Cross-leave	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Vascular p <i>Erica vagai</i> Cornish Hea	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Eriophorum</i> Common Cott	0	-4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Vascular p <i>Eriophorum</i> Hare's-tail	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Vascular p <i>Erodium lei</i> Sticky Stor	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Erodium ma</i> Sea Stork's	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Erodium mo</i> Musk Stork'	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Erophila v</i> Common Whit	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Erysimum c</i> Wallflower	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Euonymus ei</i> Spindle	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p <i>Euphorbia</i> Caper Spurg	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Euphorbia</i> Petty Spurg	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Euphorbia</i> Broad-leave	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Euphrasia</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%

Vascular p <i>Euphrasia</i> Confused Ey	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Euphrasia</i> Slender Eye	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Euphrasia</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Euphrasia</i> Chalk Eyeb	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Euphrasia</i> Scottish Ey	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Euphrasia</i> Western Ey	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Vascular p <i>Euphrasia</i> Cornish Ey	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Fagus sylv</i> Beech	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Fallopia ca</i> Black-bindv	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Festuca ar</i> Rush-leaved	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Festuca ar</i> Tall Fescue	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Festuca fi</i> Fine-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Festuca gi</i> Giant Fescue	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Festuca ov</i> Sheep's-fescue	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Festuca pr</i> Meadow Fescue	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Festuca ru</i> Red Fescue	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Festuca ru</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Festuca ru</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Festuca vi</i> Viviparous	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Foeniculum</i> Fennel	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Frankenia</i> Sea-heath	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Fraxinus e</i> Ash	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Fritillaria</i> Fritillary	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Fumaria ba</i> Tall Ranunculus	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Fumaria de</i> Dense-flowered	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Fumaria mu</i> Common Ranunculus	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Fumaria mu</i> Boreau's Ranunculus	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Fumaria oc</i> Western Ranunculus	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Vascular p <i>Fumaria of</i> Common Fumitory	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Fumaria of</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Fumaria of</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Fumaria pa</i> Fine-leaved	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Fumaria pu</i> Purple Ranunculus	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Fumaria va</i> Few-flowered	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Gagea lute</i> Yellow Star	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Galeopsis</i> Red Hemp-nettle	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Galeopsis</i> Large-flowered	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Galeopsis</i> Common Hemp-nettle	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Galeopsis</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p <i>Galium apa</i> Cleavers	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Galium bor</i> Northern Bedstraw	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Galium mol</i> Hedge Bedstraw	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Galium odo</i> Woodruff	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Galium pal</i> Common Marsh Bedstraw	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Galium pum</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Vascular p <i>Galium sax</i> Heath Bedstraw	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p <i>Galium ste</i> Limestone Bedstraw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p <i>Gastridium</i> Nit-grass	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Gaudinia f</i> French Oat-	0 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Genista ti</i> Dyer's Gree	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Gentianella</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Gentianella</i> Chiltern Ge	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Geranium c</i> Long-stalke	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Geranium d</i> Cut-leaved	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Geranium l</i> Shining Cra	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Geranium m</i> Dove's-foot	0 > -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Geranium p</i> Little-Robi	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Geranium p</i> Small-flowe	0 > -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Geranium r</i> Herb-Robert	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Geranium s</i> Wood Crane'	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Geum rival</i> Water Aven	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Geum urban</i> Wood Aven	0 -7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Glechoma h</i> Ground-ivy	0 > -1%	> -1%	LOW	+1 to +4%
Vascular p <i>Glyceria d</i> Small Sweet	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Glyceria f</i> Floating Sv	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Vascular p <i>Glyceria m</i> Reed Sweet-	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Glyceria n</i> Plicate Swe	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Gnaphalium</i> Heath Cudwe	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Gnaphalium</i> Marsh Cudwe	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Goodyera r</i> Creeping La	0 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Groenlandia</i> Opposite-le	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Gymnadenia</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Gymnadenia</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Gymnocarpi</i> Limestone F	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Hammarbya i</i> Bog Orchid	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Hedera hel</i> Common Ivy	0 < -7.5%	> -1%	MODERATE	< +1%
Vascular p <i>Hedera hel</i> Common Ivy	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Hedera 'Hi</i> Irish Ivy	0 > -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Helianthem</i> Common Rock	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Helictotri</i> Meadow Oat-	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Helleborus</i> Stinking He	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p <i>Helleborus</i> Green Helle	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Herminium i</i> Musk Orchid	1 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Herniaria i</i> Smooth Rupt	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Himantoglo</i> Lizard Orch	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Hippocrepis</i> Horseshoe \	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Hippuris v</i> Mare's-tail	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Holcus mol</i> Creeping Sc	0 -7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p <i>Hordelymus</i> Wood Barley	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Hordeum mu</i> Wall Barley	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Hordeum se</i> Meadow Barl	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Humulus lu</i> Hop	0 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Huperzia s</i> Fir Clubmos	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Hyacinthoi</i> Bluebell	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%

Vascular p <i>Hydrocotyl</i> Marsh Penny	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Hymenophyl</i> Tunbridge F	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Hymenophyl</i> Wilson's F	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Hypericum</i> Tutsan	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Hypericum</i> Pale St Joh	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Hypericum</i> Perforate S	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Hypericum</i> Slender St	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Hypericum</i> Square-stal	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Hypericum</i> Wavy St Joh	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Hypochaeri</i> Smooth Cat'	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Ilex aquif</i> Holly	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Illecebrum</i> Coral-neckl	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Inula cony</i> Ploughman's	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Inula helei</i> Elecampane	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Iris pseud</i> Yellow Iris	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Isoetes eci</i> Spring Quil	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Isolepis c</i> Slender Clu	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Jasione moi</i> Sheep's-bit	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Juncus acu</i> Sharp-flowe	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p <i>Juncus acu</i> Sharp Rush	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Juncus alp</i> Alpine Rush	0	> -1%	< -7.5%	MODERATE	< +1%
Vascular p <i>Juncus amb</i> Frog Rush	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Juncus art</i> Jointed Rus	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Juncus bal</i> Baltic Rush	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Juncus buf</i> Toad Rush	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Juncus bulb</i> Bulbous Rus	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Juncus bulb</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Juncus comp</i> Round-fruit	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Juncus fol</i> Leafy Rush	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Juncus inf</i> Hard Rush	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p <i>Juncus tri</i> Three-flowe	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Juniperus</i> Common Jun	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Juniperus</i> Dwarf Junip	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Knautia ar</i> Field Scabi	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Lactuca se</i> Prickly Let	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Lamiastrum</i> Yellow Arch	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Lamium alb</i> White Dead-	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Lamium amp</i> Henbit Deac	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Lamium con</i> Northern De	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Lamium pur</i> Red Dead-ne	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Lapsana coi</i> Nipplewort	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Lathyrus a</i> Yellow Veto	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Lathyrus j</i> Sea Pea	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Lathyrus l</i> Bitter-veto	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p <i>Lathyrus p</i> Marsh Pea	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Legousia h</i> Venus's-loc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Lemna gibb</i> Fat Duckwee	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p <i>Lemna mino</i> Common Duck	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Lemna tris</i> Ivy-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Leontodon</i> Autumn Hawk	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p <i>Leontodon</i> Rough Hawk	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p <i>Leontodon</i> Lesser Hawk	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Lepidium h</i> Smith's Per	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Lepidium r</i> Narrow-leav	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Leucanthem</i> Oxeye Daisy	0	-7.5 to -4%	> -1%	MODERATE	< +1%
Vascular p <i>Leymus are</i> Lyme-grass	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Limonium h</i> Lax-flowere	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Limosella</i> Mudwort	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Linaria vu</i> Common Toa	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Linum bien</i> Pale Flax	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Linum pere</i> Perennial F	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Listera ov</i> Common Twa	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p <i>Lithosperm</i> Purple Gron	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Lolium per</i> Perennial F	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Lonicera p</i> Honeysuckle	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular p <i>Lotus angu</i> Slender Bir	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Vascular p <i>Lotus pedu</i> Greater Bir	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Lotus subb</i> Hairy Bird'	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Luronium n</i> Floating We	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Luzula cam</i> Field Wood-	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p <i>Luzula for</i> Southern We	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Luzula mul</i> Heath Wood-	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p <i>Luzula mul</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Luzula pil</i> Hairy Wood-	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Luzula syl</i> Great Wood-	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p <i>Lychnis fl</i> Ragged-Robi	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Lycopus eu</i> Gypsywort	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Lysimachia</i> Yellow Pimp	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Lysimachia</i> Tufted Loos	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Malus sylv</i> Crab Apple	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Malva mosc</i> Musk-mallow	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Malva sylv</i> Common Mall	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Marrubium</i> White Horeh	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Medicago a</i> Spotted Mec	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Medicago l</i> Black Medic	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Medicago m</i> Bur Medick	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Medicago p</i> Toothed Mec	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Medicago s</i> Lucerne	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Melampyrum</i> Common Cow-	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Melampyrum</i> Small Cow-v	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Melilotus</i> Tall Melilo	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Melittis m</i> Bastard Bal	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Mentha aqu</i> Water Mint	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Mentha pul</i> Pennyroyal	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p <i>Mentha spi</i>	Spear Mint	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Mentha sua</i>	Round-leave	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Mercuriali</i>	Dog's Mercu	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Minuartia</i>	Fine-leaved	1	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Minuartia</i>	Spring Sand	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Moehringia</i>	Three-nerve	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Moenchia e</i>	Upright Chi	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p <i>Montia fon</i>	Blinks	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Montia fon</i>	NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Montia fon</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Muscari ne</i>	Grape-hyaci	1	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Myosotis a</i>	Field Forge	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p <i>Myosotis d</i>	Changing Fo	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Myosotis l</i>	Tufted Forg	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Myosotis s</i>	Creeping Fo	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p <i>Myosotis s</i>	Pale Forget	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Myosotis s</i>	Wood Forget	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Myosurus m</i>	Mousetail	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Myrica gal</i>	Bog-myrtle	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Myriophyll</i>	Whorled Wat	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Najas flex</i>	Slender Nai	1	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Narcissus</i>	Daffodil	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Nardus str</i>	Mat-grass	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Narthecium</i>	Bog Asphode	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Vascular p <i>Nepeta cat</i>	Cat-mint	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Odontites</i>	Red Bartsia	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Odontites</i>	NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Odontites</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Oenanthe a</i>	Fine-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Oenanthe c</i>	Hemlock Wat	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Oenanthe p</i>	Corky-fruit	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Ononis rep</i>	Common Rest	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Onopordum</i>	Cotton This	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Ophiogloss</i>	Adder's-tor	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Ophrys sph</i>	Early Spide	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Orchis mas</i>	Early-purpl	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Orchis ust</i>	Burnt Orchi	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Oreopteris</i>	Lemon-scent	0	-4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Vascular p <i>Ornithogal</i>	Spiked Star	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Ornithogal</i>	Star-of-Bet	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Orobanche</i>	Thyme Broom	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Orobanche</i>	Knapweed Br	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Orobanche</i>	Common Broc	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Oxalis ace</i>	Wood-sorrel	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Papaver ar</i>	Prickly Pop	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Papaver du</i>	Long-headec	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Papaver du</i>	Yellow-juic	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p <i>Papaver rh</i> Common Poppy	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Parapholis</i> Hard-grass	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Parentucel</i> Yellow Bart	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Pediculari</i> Lousewort	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Persicaria</i> Amphibious	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Persicaria</i> Water-pepper	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Persicaria</i> Pale Persic	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Persicaria</i> Redshank	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Persicaria</i> Small Water	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Persicaria</i> Tasteless V	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Persicaria</i> Alpine Bist	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p <i>Petroselin</i> Garden Pars	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Petroselin</i> Corn Parsle	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Peucedanum</i> Milk-parsle	0 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Phalaris a</i> Reed Canary	0 -7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p <i>Phegopteris</i> Beech Fern	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Phleum alp</i> Alpine Cat'	0 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Phleum ber</i> Smaller Cat	0 > -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Phleum pra</i> Timothy	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Phleum pra</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Phragmites</i> Common Reec	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p <i>Phyllitis</i> Hart's-tong	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Picris ech</i> Bristly Oxt	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Pilularia</i> Pillwort	1 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Pimpinella</i> Greater Bu	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Pimpinella</i> Burnet-saxi	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Pinguicula</i> Common Butt	0 -4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Vascular p <i>Plantago c</i> Buck's-horr	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Plantago l</i> Ribwort Pla	0 -4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Plantago m</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Plantago m</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Plantago m</i> Sea Plantai	0 -7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Poa bulbos</i> Bulbous Mea	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Poa humili</i> Spreading M	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Poa nemora</i> Wood Meadow	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Poa pratens</i> NA	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p <i>Poa pratens</i> Smooth Meac	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Poa trivialis</i> Rough Mead	0 > -1%	> -1%	LOW	< +1%
Vascular p <i>Polemonium</i> Jacob's-lac	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Polygala c</i> Chalk Milkv	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Polygala s</i> Heath Milkv	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Polygala v</i> Common Mill	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Polygonatu</i> Solomon's-s	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Polygonatu</i> Angular Sol	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Polygonum</i> Equal-leave	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Polygonum</i> Knotgrass	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Polygonum</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%

Vascular pl	<i>Polygonum</i>	Ray's Knotg	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Polygonum</i>	Cornfield F	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Polypodium</i>	Southern Po	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular pl	<i>Polypodium</i>	Intermediat	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular pl	<i>Polypodium</i>	Polypody	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Polypodium</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Polystichum</i>	Soft Shield	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Populus nigra</i>	Black-poplar	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Potamogeton</i>	Fen Pondwee	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular pl	<i>Potamogeton</i>	Grass-wrack	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Potamogeton</i>	Flat-stalked	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Potamogeton</i>	Broad-leave	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Potamogeton</i>	Fennel Pond	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Potamogeton</i>	Bog Pondwee	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Potamogeton</i>	Hairlike Po	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Potentilla</i>	Trailing To	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Potentilla</i>	Silverweed	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Potentilla</i>	Alpine Cinc	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular pl	<i>Potentilla</i>	Tormentil	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Potentilla</i>	Shrubby Cir	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Potentilla</i>	Spring Cinc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Potentilla</i>	Creeping Ci	0	> -1%	> -1%	LOW	+1 to +4%
Vascular pl	<i>Potentilla</i>	Barren Stra	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Primula elatior</i>	Oxlip	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Primula veris</i>	Cowslip	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Primula vulgaris</i>	Primrose	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Prunella vulgaris</i>	Selfheal	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Prunus cerasifera</i>	Dwarf Cherry	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Prunus domestica</i>	Wild Plum	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Prunus domestica</i>	Plum	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Prunus domestica</i>	Bullace; Dam	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Prunus padalis</i>	Bird Cherry	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular pl	<i>Prunus spinosa</i>	Blackthorn	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Pteridium aquilinum</i>	Bracken	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular pl	<i>Puccinellia maritima</i>	Stiff Saltm	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Pulicaria vulgaris</i>	Common Flea	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Pulmonaria officinalis</i>	Narrow-leave	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Pulsatilla nuttalliana</i>	Pasqueflowe	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular pl	<i>Pyrola media</i>	Intermediat	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Pyrola rotundifolia</i>	Round-leave	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular pl	<i>Pyrola rotundifolia</i>	Wintergreer	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Quercus pedunculata</i>	Sessile Oak	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Quercus robur</i>	Pedunculate	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Radiola linum</i>	Allseed	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Ranunculus acris</i>	Meadow But	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Ranunculus repens</i>	Common Wate	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Ranunculus repens</i>	Corn Butter	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%

Vascular p <i>Ranunculus</i> Bulbous But	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Ranunculus</i> Lesser Cel	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Ranunculus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Ranunculus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Ranunculus</i> Lesser Spe	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Ranunculus</i> River Water	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Ranunculus</i> Round-leave	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Ranunculus</i> Stream Water	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Ranunculus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Ranunculus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Ranunculus</i> Thread-leav	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Ranunculus</i> Three-lobed	1	> -1%	> -1%	LOW	+1 to +4%
Vascular p <i>Raphanus r.</i> Wild Radish	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Reseda lutea</i> Weld	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Rhinanthus</i> Yellow-rat	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Rhinanthus</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Rhinanthus</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Ribes alpi</i> Mountain Cu	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Ribes rubri</i> Red Currant	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Vascular p <i>Ribes spica</i> Downy Curr	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Rorippa mi</i> Narrow-frui	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rorippa na.</i> Water-cress	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rorippa na.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Rorippa sy.</i> Creeping Ye	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rosa arven.</i> Field-rose	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Rosa caesi.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Rosa caesi.</i> Hairy Dog-1	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Rosa canin.</i> Dog-rose	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Rosa micra</i> Small-flowe	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Rosa molli.</i> Soft Downy-	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Rosa rubig.</i> Sweet-briar	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rosa stylo.</i> Short-style	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Rosa tomen.</i> Harsh Downy	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Rubia pere.</i> Wild Madder	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rubus cham.</i> Cloudberry	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Rubus frut.</i> Bramble	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rubus idae.</i> Raspberry	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Rubus saxa.</i> Stone Bramb	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Rumex acet.</i> Common Sor	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Rumex acet.</i> Sheep's Sor	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rumex acet.</i> Narrow-Leav	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Rumex cong.</i> Clustered I	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rumex cris.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Rumex palu.</i> Marsh Dock	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Rumex rupe.</i> Shore Dock	1	> -1%	> -1%	LOW	+1 to +4%
Vascular p <i>Rumex sang.</i> Wood Dock	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Ruppia cir.</i> Spiral Tass	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Vascular pl	<i>Ruppia mar</i>	Beaked Tass	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Ruscus acu</i>	Butcher's-b	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Sagina ape</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Sagina mar</i>	Sea Pearlwe	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Sagina nod</i>	Knotted Pea	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Sagina pro</i>	Procumbent	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Sagina sub</i>	Heath Pearl	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Sagittaria</i>	Arrowhead	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Salicornia</i>	Long-spike	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Salicornia</i>	Common Glas	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Salicornia</i>	Yellow Glas	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Salicornia</i>	One-flower	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Salicornia</i>	Purple Glas	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Salix alba</i>	White Will	0	> -1%	> -1%	LOW	+1 to +4%
Vascular pl	<i>Salix auri</i>	Eared Will	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular pl	<i>Salix capr</i>	Goat Willow	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Salix capr</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Salix cine</i>	Grey Willow	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Salix cine</i>	Grey Willow	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Salix cine</i>	Rusty Will	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Salix frag</i>	Crack-will	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular pl	<i>Salix herb</i>	Dwarf Will	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular pl	<i>Salix lapp</i>	Downy Will	1	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular pl	<i>Salix myrs</i>	Dark-leaved	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Salix phyl</i>	Tea-leaved	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular pl	<i>Salix purp</i>	Purple Will	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Salix repe</i>	Creeping Wi	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Salsola ka</i>	Prickly Sal	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Sambucus e</i>	Dwarf Elder	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Sambucus n</i>	Elder	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Samolus va</i>	Brookweed	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Sanguisorb</i>	Salad Burn	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Sanguisorb</i>	Great Burn	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Sanicula e</i>	Sanicle	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Sarcocorni</i>	Perennial C	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Saussurea</i>	Alpine Saw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Saxifraga</i>	Mossy Saxi	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Scabiosa c</i>	Small Scabi	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Scandix pe</i>	Shepherd's	1	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular pl	<i>Schoenus n</i>	Black Bog-r	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular pl	<i>Scilla aut</i>	Autumn Squi	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Scilla veri</i>	Spring Squi	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Scirpus sy</i>	Wood Club-r	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Scrophular</i>	Water Figw	0	> -1%	> -1%	LOW	+1 to +4%
Vascular pl	<i>Scrophular</i>	Common Figv	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Scrophular</i>	Green Figw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Sedum angl</i>	English Sto	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular pl	<i>Sedum rosea</i>	Roseroot	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Sedum villi</i>	Hairy Stone	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular pl	<i>Selaginella</i>	Lesser Clu	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Vascular pl	<i>Senecio aqu</i>	Marsh Ragw	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Senecio eri</i>	Hoary Ragw	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Vascular pl	<i>Senecio ja</i>	Common Ragv	0	> -1%	> -1%	LOW	+1 to +4%
Vascular pl	<i>Senecio sy</i>	Heath Grou	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Senecio vu</i>	Groundsel	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Senecio vu</i>	Groundsel	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Seriphidium</i>	Sea Wormwo	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Sherardia</i>	Field Madde	0	> -1%	> -1%	LOW	+1 to +4%
Vascular pl	<i>Sibthorpia</i>	Cornish Mor	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular pl	<i>Silaum sil</i>	Pepper-saxi	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Silene con</i>	Sand Catchf	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Silene dio</i>	Red Campior	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Silene gal</i>	Small-flowe	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Silene lat</i>	White Campi	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular pl	<i>Silene noc</i>	Night-flowe	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular pl	<i>Silene nut</i>	Nottingham	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Silene vul</i>	Bladder Car	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Sinapis ar</i>	Charlock	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Sisymbrium</i>	Hedge Musta	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Sium latif</i>	Greater Wat	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Solanum du</i>	Bittersweet	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Solanum ni</i>	Black Night	0	> -1%	> -1%	LOW	> +7.5%
Vascular pl	<i>Solidago v</i>	Goldenrod	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Sonchus ar</i>	Perennial S	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Sonchus as</i>	Prickly Sov	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular pl	<i>Sonchus ol</i>	Smooth Sowt	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular pl	<i>Sonchus pa</i>	Marsh Sowth	0	> -1%	> -1%	LOW	+1 to +4%
Vascular pl	<i>Sorbus ari</i>	Common Whit	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular pl	<i>Sorbus auc</i>	Rowan	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Sorbus dev</i>	Devon White	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Sparganium</i>	Floating Bu	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular pl	<i>Sparganium</i>	Unbranched	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Sparganium</i>	Branched Bu	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Spartina m</i>	Small Cord-	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Spergula a</i>	Corn Spurre	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Spergulari</i>	Lesser Sea-	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Spergulari</i>	Greater Sea	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Spirodela</i>	Greater Duc	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular pl	<i>Stachys ar</i>	Field Wound	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular pl	<i>Stachys of</i>	Betony	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular pl	<i>Stachys pa</i>	Marsh Wound	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular pl	<i>Stachys sy</i>	Hedge Wound	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular pl	<i>Stellaria</i>	Lesser Stit	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular pl	<i>Stellaria</i>	Greater Sti	0	< -7.5%	> -1%	MODERATE	< +1%

Vascular p <i>Stellaria</i> Lesser Chic	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Stellaria</i> Marsh Stitc	1 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Stellaria</i> Bog Stitchv	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Stratiotes</i> Water-soldi	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Subularia</i> Awlwort	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Succisa pr</i> Devil's-bit	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Symphytum</i> Common Comf	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Tamus comm</i> Black Bryor	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p <i>Tanacetum</i> Feverfew	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Taxus bacc</i> Yew	0 > -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Teucrium s</i> Wood Sage	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Thalictrum</i> Common Meac	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Thelypteris</i> Marsh Fern	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Thesium hu</i> Bastard-toe	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Thlaspi ca</i> Alpine Penr	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p <i>Thymus pol</i> Wild Thyme	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p <i>Tilia plat</i> Large-leave	0 > -1%	> -1%	LOW	> +7.5%
Vascular p <i>Tofieldia</i> Scottish As	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Torilis ar</i> Spreading I	1 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Torilis ja</i> Upright Hec	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Tragopogon</i> Goat's-bear	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Trichophor</i> Northern De	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Trichophor</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Trichophor</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Trientalis</i> Chickweed-v	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Trifolium</i> Lesser Tref	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Trifolium</i> Strawberry	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Trifolium</i> Clustered (0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Trifolium</i> Zigzag Clov	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Trifolium</i> Slender Tre	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Trifolium</i> Bird's-foot	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Trifolium</i> White Clov	0 -4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Trifolium</i> Rough Clov	0 < -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Trifolium</i> Knotted Clo	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Trifolium</i> Subterranea	0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Triglochin</i> Sea Arrowgr	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Triglochin</i> Marsh Arro	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Tripleuros</i> Scentless M	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p <i>Trisetum f</i> Yellow Oat-	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Trollius ei</i> Globeflower	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p <i>Typha lati</i> Bulrush	0 > -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Ulex europ</i> Gorse	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Ulex galli</i> Western Gor	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Ulex minor</i> Dwarf Gorse	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Ulmus glab</i> Wych Elm	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Ulmus mino</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Ulmus proc</i> English Elr	0 < -7.5%	> -1%	MODERATE	+1 to +4%

Vascular p <i>Umbilicus</i> Navelwort	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Urtica urei</i> Small Nettl	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Utricularia</i> Bladderwort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Utricularia</i> :NA	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Vascular p <i>Utricularia</i> :Greater Bla	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Vaccinium</i> Bilberry	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p <i>Vaccinium</i> Bog Bilber	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Vaccinium</i> Cowberry	0	-7.5 to -4%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Valeriana</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Vascular p <i>Valeriana</i> Keeled-frui	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Valeriana</i> Narrow-frui	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Valeriana</i> Hairy-frui	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p <i>Valeriana</i> Broad-frui	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Veronica a</i> Green Field	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Veronica b</i> Brooklime	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p <i>Veronica c</i> Pink Water-	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Veronica c</i> Germander S	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular p <i>Veronica h</i> Ivy-leaved	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Veronica h</i> :NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Veronica h</i> :NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p <i>Veronica m</i> Wood Speedv	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Veronica o</i> Heath Spee	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Veronica s</i> Thyme-leave	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p <i>Veronica s</i> :NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Veronica s</i> :NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p <i>Viburnum o</i> Guelder-ros	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Vicia hirs</i> Hairy Tare	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Vicia lute</i> Yellow-vet	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p <i>Vicia orob</i> Wood Bitter	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Vicia parv</i> Slender Tai	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Vicia sati</i> Narrow-leav	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Vicia sati</i> :NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p <i>Vicia sati</i> Common Veto	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Vicia sylv</i> Wood Vetch	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Vinca mino</i> Lesser Peri	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Viola arve</i> Field Pansy	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p <i>Viola cani</i> Heath Dog-v	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p <i>Viola hirt</i> Hairy Viole	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Viola lact</i> Pale Dog-vi	1	> -1%	-4 to -1%	MODERATE	+1 to +4%
Vascular p <i>Viola lute</i> Mountain Pa	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p <i>Viola odor</i> Sweet Viole	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Viola palu</i> Marsh Viole	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p <i>Viola palu</i> :NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p <i>Viola palu</i> :NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p <i>Viola reici</i> Early Dog-v	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p <i>Viola rivin</i> Common Dog-	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p <i>Viola tric</i> Wild Pansy	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p	<i>Viola tric</i>	Seaside Par	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Viola tric</i>	NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p	<i>Vulpia myu</i>	Rat's-tail	0	> -1%	> -1%	LOW	> +7.5%
Vascular p	<i>Zostera ma</i>	Eelgrass	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Zostera no</i>	Dwarf Eelgr	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Agenioideu</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ammophila</i>	Red Banded	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Wasps	<i>Ancistroce</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ancistroce</i>	Wall Mason	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Anoplius c</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Wasps	<i>Anoplius i</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Anoplius n</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Anoplius v</i>	Black Banded	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Wasps	<i>Arachnosp</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Arachnosp</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Arachnosp</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Arachnosp</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Argogoryte</i>	Field Digge	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Astata boo</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Astata pin</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Auplopus c</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Caliadurgu</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Cerceris a</i>	Sand Taile	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Cerceris r</i>	Ornate Tail	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Chrysis an</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Chrysis igr</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Chrysis im</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Chrysis me</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Chrysis vi</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Cleptes sei</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Crabro cri</i>	Slender Boc	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crabro pel</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crabro scu</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Crossoceru</i>	Blunt Taile	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Crossoceru</i>	Slender Dig	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%

Wasps	<i>Crossocerus</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crossocerus</i> 4-Spotted I	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crossocerus</i> Wesmael's I	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Diodontus</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Diodontus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Diodontus</i> Melancholy	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Dipogon</i> su NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Dolichovesp</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Dolichovesp</i> Tree Wasp	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ectemnius</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Ectemnius</i> NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Wasps	<i>Ectemnius</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Ectemnius</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Elampus</i> pa NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Entomognath</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Episyron</i> ri Red Legged	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Eumenes</i> co Heath Potter	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Evagetus</i> c NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Evagetus</i> di NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Gorytes</i> bi NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Gorytes</i> qu 4-Banded Di	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Gorytes</i> tu NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Gymnomerus</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Hedychridi</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Hedychridi</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Hedychridi</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Lindenius</i> i NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Lindenius</i> j NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Mellinus</i> a Field Digge	0	> -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Microdyner</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Miscophus</i> i NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Mutilla</i> eu Large Velv	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Wasps	<i>Myrmosa</i> at NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Nysson</i> dim Small Spuri	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Nysson</i> spi Large Spuri	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Wasps	<i>Nysson</i> tri NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Odynerus</i> m NA	1	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Odynerus</i> s Spiny Masor	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Omalus</i> aen NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Omalus</i> aur NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Omalus</i> vio NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Wasps	<i>Oxybelus</i> a Silver Spir	0	< -7.5%	> -1%	MODERATE	> +7.5%

Wasps	<i>Oxybelus m</i> Pale Jawed	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Oxybelus u</i> Common Spir	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Passaloecu</i> .Horned Blac	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Passaloecu</i> .NA	0 > -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Passaloecu</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Passaloecu</i> .NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Passaloecu</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Pemphredon</i> Mournful W	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Pemphredon</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Podalonia i</i> Hairy Sand	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Pompilus c</i> .Leaden Spic	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Priocnemis</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Priocnemis</i> NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Psen bruxe</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Psen dahlb</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Psen eques</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Psen lutar</i> .NA	0 -7.5 to -4%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Psenulus c</i> .NA	0 > -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Psenulus p</i> Pale Foote	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Rhopalum c</i> .NA	0 < -7.5%	-4 to -1%	HIGH	+1 to +4%
Wasps	<i>Rhopalum c</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Sapyga cla</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Sapyga qui</i> .NA	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Smicromyrme</i> Small Velve	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Spilomena i</i> .NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Spilomena i</i> .NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Symmorphus</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Tachysphex</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Tiphia fem</i> .NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Tiphia mini</i> Small Tiphi	0 > -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Trichrysis</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> Slender Woc	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> Club Horne	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> NA	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Wasps	<i>Vespa crab</i> .Hornet	0 > -1%	< -7.5%	MODERATE	> +7.5%
Wasps	<i>Vespula ge</i> .German Was	0 > -1%	> -1%	LOW	> +7.5%
Wasps	<i>Vespula ru</i> .Red Wasp	0 < -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Vespula vu</i> .Common Was	0 > -1%	> -1%	LOW	> +7.5%

Projected expansion	Benefit from expansion	Final outcome
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Medium benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	Limited impact
> +7.5%	VERY HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	Medium benefit
< +1%	LOW	Limited impact
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	LOW	Limited impact
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	LOW	Limited impact
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	Risks & benefits
< +1%	LOW	Limited impact
< +1%	MODERATE	High risk
< +1%	LOW	Limited impact
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
+1 to +4%	HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit

< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk

+1 to +4%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	LOW	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Risks & benefits
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	MODERATE	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	LOW	Medium risk
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	LOW	Medium risk
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit

+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit

< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Risks & benefits
+1 to +4%	HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Risks & benefits

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Risks & benefits
+1 to +4%	HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	LOW	High risk
+4 to +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk

< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit

> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk

> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit

> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	LOW	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium risk
+4 to +7.5%	VERY HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	LOW	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit

+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit

< +1%	LOW	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	LOW	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Medium benefit
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
+4 to +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit

+1 to +4%	MODERATE	Medium benefit
< +1%	LOW	High risk
+4 to +7.5%	HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits

< +1%	LOW	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit

+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
> +7.5%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit

+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit

< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Risks & benefits
> +7.5%	MODERATE	Medium benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
< +1%	LOW	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits

> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk

< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Risks & benefits
+1 to +4%	HIGH	Medium risk
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits

< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
+1 to +4%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	High risk

< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	LOW	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit

> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
+4 to +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk

< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk

+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit

< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit

[illegible]

[illegible]

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

Group	Latin name	English name	NERC species	Observed decline	Projected decline	Risk of decline	Observed expansion
Ants	<i>Formica cu</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Formica fu</i>	Negro Ant		0 > -1%	> -1%	LOW	> +7.5%
Ants	<i>Formica sa</i>	NA		0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Ants	<i>Lasius ali</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Lasius fla</i>	Yellow Mead		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Ants	<i>Lasius mix</i>	NA		0 > -1%	> -1%	LOW	+1 to +4%
Ants	<i>Lasius nig</i>	Small Black		0 > -1%	> -1%	LOW	> +7.5%
Ants	<i>Leptothora</i>	Slender Ant		0 > -1%	-4 to -1%	MODERATE	> +7.5%
Ants	<i>Myrmica ru</i>	Red Ant		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Myrmica ru</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Ants	<i>Myrmica sa</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Ants	<i>Myrmica sc</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Ants	<i>Myrmica sc</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena al</i>	NA		0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bees	<i>Andrena an</i>	NA		0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Andrena ap</i>	NA		0 > -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Andrena ar</i>	NA		0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bees	<i>Andrena ba</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena bi</i>	Gwynne's M		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena bi</i>	NA		0 > -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Andrena bu</i>	NA		0 > -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Andrena ch</i>	NA		0 -4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Bees	<i>Andrena ci</i>	Grey Mining		0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bees	<i>Andrena co</i>	NA		0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Andrena co</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena de</i>	NA		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena do</i>	NA		0 > -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Andrena fl</i>	Yellow Leg		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena fu</i>	NA		0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Andrena fu</i>	Tawny Mini		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena fu</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena fu</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena ha</i>	Early Mini		0 -4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ha</i>	NA		0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Andrena he</i>	NA		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena hu</i>	NA		0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena la</i>	NA		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena la</i>	Girdled Mi		0 > -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena la</i>	NA		0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bees	<i>Andrena mi</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena mi</i>	NA		0 < -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ni</i>	NA		0 -7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ov</i>	NA		0 > -1%	> -1%	LOW	> +7.5%

Bees	<i>Andrena pi</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena pr</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena pr</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Andrena sc</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena su</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Andrena sy</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena ta</i> .Tormentil M	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Andrena th</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Andrena ti</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena tr</i> .Trimmer's M	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena va</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Andrena wi</i> .NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Anthidium</i> Wool-Carder	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Anthophora</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Anthophora</i> Fork Tailec	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Anthophora</i> Hairy Footc	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Anthophora</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Apis melli</i> .Honey Bee	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Bombus hor</i> Small Gardc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Bombus hum</i> .NA	1	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Bombus jon</i> Heath Bumb	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Bombus lap</i> .Large Red C	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Bombus luc</i> .White-Tailec	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Bombus mag</i> .NA	0	< -7.5%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Bombus mon</i> Mountain Bu	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Bombus mus</i> .Moss Carder	1	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Bombus pas</i> .Common Carc	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Bombus pra</i> .Early Bumb	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Bombus rud</i> .NA	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Bombus syl</i> .NA	1	> -1%	> -1%	LOW	< +1%
Bees	<i>Bombus ter</i> .Buff-Tailec	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Chelostoma</i> Harebell C	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Coelioxys</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Coelioxys</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Coelioxys</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Colletes d</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Colletes h</i> .Sea-aster C	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Colletes m</i> .Margined Co	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Colletes s</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Colletes s</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Bees	<i>Epeolus cr</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Epeolus va</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Halictus c</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bees	<i>Halictus r</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Halictus t</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Hoplitis c</i> .NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Hoplitis s</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%

Bees	<i>Hylaeus an.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Hylaeus br.</i> Short Horned	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus co.</i> Common Yellow	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus co.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus co.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus hy.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Hylaeus pe.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Hylaeus pi.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Hylaeus si.</i> Large Yellow	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasiogloss.</i> Slender Min	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasiogloss.</i> Least Min	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasiogloss.</i> Brassy Min	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Lasiogloss.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Lasiogloss.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Lasiogloss.</i> Shaggy Min	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Macropis e.</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Bees	<i>Megachile</i> Patchwork	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Megachile</i> Wood-Carvin	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Megachile</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Megachile</i> Willughby's	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Melecta al.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Melitta ha.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Melitta le.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada fab.</i> Fabricius'	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada fla.</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bees	<i>Nomada fla.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Nomada fla.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada fuc.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Nomada goo.</i> Gooden's No	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Nomada lat.</i> NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bees	<i>Nomada leu.</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bees	<i>Nomada mar.</i> Marsham's	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada pan.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bees	<i>Nomada ruf.</i> Red-Horned	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Bees	<i>Nomada she.</i> Dark Nomad	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Nomada str.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%

Bees	<i>Osmia auru</i>	Gold-Fringe	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Osmia bico</i>	Two Coloured	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bees	<i>Osmia rufa</i>	Red Mason	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Panurgus c.</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Sphecodes</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Bees	<i>Sphecodes</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Sphecodes</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Bees	<i>Sphecodes</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bees	<i>Stelis orn.</i>	NA	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bees	<i>Stelis pun.</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Accipiter</i>	Goshawk	0	< -7.5%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Accipiter</i>	Sparrowhawk	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Acrocephal</i>	Sedge Warbler	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Acrocephal</i>	Reed Warbler	0	< -7.5%	> -1%	LOW	> +7.5%
Birds	<i>Actitis hy.</i>	Common Sandpiper	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Aegithalos</i>	Long-tailed Tit	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Aix galeri</i>	Mandarin Duck	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Alauda arv.</i>	Skylark	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Alca torda</i>	Razorbill	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Alcedo att.</i>	Kingfisher	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Alectoris</i>	Red-legged Partridge	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Anas acuta</i>	Pintail	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Anas clype.</i>	Shoveler	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Anas crecc.</i>	Teal	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Anas penel.</i>	Widgeon	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Anas platy.</i>	Mallard	0	-4 to -1%	> -1%	LOW	+1 to +4%
Birds	<i>Anas querq.</i>	Garganey	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Anas strep.</i>	Gadwall	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Anser anse.</i>	Greylag Goose	0	< -7.5%	> -1%	LOW	> +7.5%
Birds	<i>Anthus pet.</i>	Rock Pipit	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Anthus pra.</i>	Meadow Pipit	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Anthus tri.</i>	Tree Pipit	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Apus apus</i>	Swift	0	-4 to -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Aquila chr.</i>	Golden Eagle	0	-7.5 to -4%	< -7.5%	MODERATE	> +7.5%
Birds	<i>Ardea cine.</i>	Grey Heron	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Asio flamm.</i>	Short-eared Owl	0	> -1%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Asio otus</i>	Long-eared Owl	0	-4 to -1%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Athene noc.</i>	Little Owl	0	> -1%	> -1%	LOW	+4 to +7.5%

Birds	<i>Aythya fer.</i> Pochard	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Aythya ful.</i> Tufted Duck	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Botaurus s</i> Bittern	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Branta can.</i> Canada Goose	0	> -1%	< -7.5%	MODERATE	> +7.5%
Birds	<i>Branta leu.</i> Barnacle Goose	0	> -1%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Bucephala</i> Goldeneye	0	> -1%	< -7.5%	MODERATE	> +7.5%
Birds	<i>Burhinus o.</i> Stone-curlew	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Buteo bute.</i> Buzzard	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Calidris a.</i> Dunlin	0	> -1%	< -7.5%	VERY HIGH	+4 to +7.5%
Birds	<i>Caprimulgus.</i> Nightjar	1	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Carduelis</i> Lesser Redpoll	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Birds	<i>Carduelis</i> Linnet	1	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Carduelis</i> Goldfinch	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Carduelis</i> Greenfinch	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Carduelis</i> Twite	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Birds	<i>Carduelis</i> Siskin	0	< -7.5%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Cephus gr.</i> Black Guillemot	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Birds	<i>Certhia fa.</i> Tree creeper	0	< -7.5%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	Cettia cetti Cetti's Warbler	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Charadrius</i> Little Ringed Plover	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Charadrius</i> Ringed Plover	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Charadrius</i> Dotterel	0	> -1%	< -7.5%	VERY HIGH	> +7.5%
Birds	<i>Chroicoceph.</i> Black-headed Gull	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Chrysoloph.</i> Golden Pheasant	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Cinclus ci.</i> Dipper	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	<i>Circus aer.</i> Marsh Harrier	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Circus cya.</i> Hen Harrier	1	> -1%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	Circus pygma Montagu's Harrier	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Coccothraux.</i> Hawfinch	1	> -1%	> -1%	MODERATE	+4 to +7.5%
Birds	Columba liv. Feral Pigeon	0	< -7.5%	> -1%	LOW	> +7.5%
Birds	<i>Columba oe.</i> Stock Dove	0	< -7.5%	> -1%	LOW	+4 to +7.5%
Birds	<i>Columba pa.</i> Woodpigeon	0	< -7.5%	> -1%	LOW	+1 to +4%
Birds	<i>Corvus cor.</i> Raven	0	< -7.5%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Corvus cor.</i> Carrion Crow	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Corvus fru.</i> Rook	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Corvus mon.</i> Jackdaw	0	< -7.5%	> -1%	LOW	< +1%
Birds	<i>Coturnix c.</i> Quail	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Crex crex</i> Corncrake	1	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Cuculus ca.</i> Cuckoo	1	-4 to -1%	-4 to -1%	HIGH	+4 to +7.5%
Birds	<i>Cyanistes</i> Blue Tit	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Cygnus olo.</i> Mute Swan	0	< -7.5%	> -1%	LOW	+4 to +7.5%
Birds	<i>Delichon u.</i> House Martin	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Dendrocopos.</i> Great Spotted Woodpecker	0	-7.5 to -4%	> -1%	LOW	> +7.5%
Birds	<i>Dendrocopos.</i> Lesser Spotted Woodpecker	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Emberiza c.</i> Corn Bunting	1	> -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Emberiza c.</i> Cirl Bunting	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Emberiza c.</i> Yellowhammer	1	> -1%	-4 to -1%	HIGH	+1 to +4%

Birds	<i>Emberiza s.</i> Reed Bunting	1	-7.5 to -4%	> -1%	LOW	+4 to +7.5%
Birds	<i>Erithacus</i> Robin	0	< -7.5%	> -1%	LOW	< +1%
Birds	<i>Falco colu</i> Merlin	0	-7.5 to -4%	< -7.5%	MODERATE	> +7.5%
Birds	<i>Falco pere</i> Peregrine	0	< -7.5%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Falco subb</i> Hobby	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Falco tinn</i> Kestrel	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Ficedula h</i> Pied Flycatcher	0	> -1%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Fratercula</i> Puffin	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Birds	<i>Fringilla</i> Chaffinch	0	-4 to -1%	> -1%	LOW	< +1%
Birds	<i>Fulica atr</i> Coot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Fulmarus g</i> Fulmar	0	> -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Gallinago</i> Snipe	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Gallinula</i> Moorhen	0	> -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Garrulus g</i> Jay	0	-4 to -1%	> -1%	LOW	> +7.5%
Birds	<i>Haematopus</i> Oystercatcher	0	-7.5 to -4%	> -1%	LOW	> +7.5%
Birds	<i>Hirundo ru</i> Swallow	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Hydrobates</i> Storm Petrel	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Lagopus la</i> Red Grouse	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Birds	<i>Lanius col</i> Red-backed	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Larus arge</i> Herring Gull	1	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Larus canu</i> Common Gull	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Larus fusc</i> Lesser Black	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Larus mari</i> Great Black	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Larus mela</i> Mediterranean	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Limosa lim</i> Black-tailed	1	> -1%	> -1%	LOW	> +7.5%
Birds	Locustella l Savi's Warbler	1	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Locustella</i> Grasshopper	1	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Loxia spp.</i> Crossbill	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Lullula ar</i> Woodlark	1	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Luscinia m</i> Nightingale	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Mergus mer</i> Goosander	0	> -1%	< -7.5%	VERY HIGH	> +7.5%
Birds	<i>Mergus ser</i> Red-breasted	0	< -7.5%	-7.5 to -4%	HIGH	> +7.5%
Birds	<i>Morus bass</i> Gannet	0	< -7.5%	-4 to -1%	MODERATE	+4 to +7.5%
Birds	Motacilla al Pied/White	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Motacilla</i> Grey Wagtail	0	-4 to -1%	-7.5 to -4%	MODERATE	> +7.5%
Birds	<i>Motacilla</i> Yellow Wagtail	1	> -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Muscicapa</i> Spotted Flycatcher	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Birds	<i>Numenius a</i> Curlew	1	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Oenanthe o</i> Wheatear	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	<i>Oxyura jam</i> Ruddy Duck	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Birds	<i>Panurus bi</i> Bearded Tit	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Parus majo</i> Great Tit	0	< -7.5%	> -1%	LOW	+1 to +4%
Birds	<i>Passer dom</i> House Sparrow	1	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Passer mon</i> Tree Sparrow	1	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Perdix per</i> Grey Partridge	1	> -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Periparus</i> Coal Tit	0	< -7.5%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Phalacroco</i> Shag	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%

Birds	<i>Phalacrocorax</i> Cormorant	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Phasianus</i> Pheasant	0	< -7.5%	> -1%	LOW	+1 to +4%
Birds	<i>Philomachus</i> Ruff	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Birds	<i>Phoenicurus</i> Black Redstart	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Phoenicurus</i> Redstart	0	> -1%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Phylloscopus</i> Chiffchaff	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Phylloscopus</i> Wood Warbler	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Phylloscopus</i> Willow Warbler	0	< -7.5%	-4 to -1%	MODERATE	+1 to +4%
Birds	<i>Pica pica</i> Magpie	0	< -7.5%	> -1%	LOW	+1 to +4%
Birds	<i>Picus viridis</i> Green Woodpecker	0	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Pluvialis</i> Golden Plover	0	> -1%	< -7.5%	VERY HIGH	+1 to +4%
Birds	<i>Podiceps cornutus</i> Great Crested Grebe	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Birds	<i>Poecile montanus</i> Willow Tit	1	> -1%	< -7.5%	VERY HIGH	+1 to +4%
Birds	<i>Poecile palustris</i> Marsh Tit	1	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	<i>Porzana porzana</i> Spotted Crail	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Prunella montanorum</i> Dunnock	1	-7.5 to -4%	> -1%	LOW	+1 to +4%
Birds	<i>Psittacula</i> Ring-necked Pheasant	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Puffinus puffinus</i> Manx Shearwater	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Pyrrhula pyrrhula</i> Bullfinch	1	> -1%	> -1%	LOW	+4 to +7.5%
Birds	<i>Rallus aquatilis</i> Water Rail	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Recurvirostra</i> Avocet	0	< -7.5%	> -1%	LOW	> +7.5%
Birds	<i>Regulus ignicapilla</i> Firecrest	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Regulus regulus</i> Goldcrest	0	< -7.5%	-7.5 to -4%	MODERATE	+4 to +7.5%
Birds	<i>Riparia riparia</i> Sand Martin	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Rissa tridactyla</i> Kittiwake	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Saxicola rubra</i> Whinchat	0	> -1%	-7.5 to -4%	VERY HIGH	+1 to +4%
Birds	<i>Saxicola torquata</i> Stonechat	0	-7.5 to -4%	> -1%	LOW	> +7.5%
Birds	<i>Scolopax rusticicola</i> Woodcock	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Birds	<i>Sitta europaea</i> Nuthatch	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Birds	<i>Somateria maura</i> Eider	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Sterna dougalli</i> Roseate Tern	1	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Sterna hirsuta</i> Common Tern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Birds	<i>Sterna parvirostris</i> Arctic Tern	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Sterna sandvicensis</i> Sandwich Tern	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Sternula albifrons</i> Little Tern	0	> -1%	> -1%	MODERATE	> +7.5%
Birds	<i>Streptopelia</i> Collared Dove	0	-4 to -1%	> -1%	LOW	> +7.5%
Birds	<i>Streptopelia</i> Turtle Dove	1	> -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Strix aluco</i> Tawny Owl	0	< -7.5%	-4 to -1%	MODERATE	+4 to +7.5%
Birds	<i>Sturnus vulgaris</i> Starling	1	-4 to -1%	> -1%	LOW	< +1%
Birds	<i>Sylvia atricapilla</i> Blackcap	0	< -7.5%	> -1%	LOW	+4 to +7.5%
Birds	<i>Sylvia borin</i> Garden Warbler	0	> -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Birds	<i>Sylvia communis</i> Whitethroat	0	< -7.5%	> -1%	LOW	+1 to +4%
Birds	<i>Sylvia curruca</i> Lesser Whitethroat	0	> -1%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Sylvia undata</i> Dartford Warbler	0	-4 to -1%	> -1%	LOW	> +7.5%
Birds	<i>Tachybaptus</i> Little Grebe	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Tadorna tadorna</i> Shelduck	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Tetrao tetrix</i> Black Grouse	1	> -1%	< -7.5%	MODERATE	> +7.5%

Birds	<i>Tringa tot.</i> Redshank	0	> -1%	> -1%	MODERATE	+4 to +7.5%
Birds	<i>Troglodyte.</i> Wren	0	> -1%	> -1%	LOW	< +1%
Birds	<i>Turdus ili.</i> Redwing	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Birds	<i>Turdus mer.</i> Blackbird	0	< -7.5%	> -1%	LOW	< +1%
Birds	<i>Turdus phi.</i> Song Thrush	1	-7.5 to -4%	> -1%	LOW	+1 to +4%
Birds	<i>Turdus pil.</i> Fieldfare	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Birds	<i>Turdus tor.</i> Ring Ouzel	1	> -1%	< -7.5%	VERY HIGH	+1 to +4%
Birds	<i>Turdus vis.</i> Mistle Thrush	0	> -1%	> -1%	LOW	+1 to +4%
Birds	<i>Tyto alba</i> Barn Owl	0	> -1%	> -1%	LOW	> +7.5%
Birds	<i>Uria aalge</i> Guillemot	0	> -1%	> -1%	MODERATE	+1 to +4%
Birds	<i>Vanellus v.</i> Lapwing	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes	<i>Abietinell.</i> Prickly Tar	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Adelanthus</i> Deceptive I	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes	<i>Aloina alo.</i> Common Aloe	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes	<i>Amblystegi.</i> Creeping Fe	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes	<i>Amblystegi.</i> NA	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes	<i>Amphidium</i> Lapland Yol	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Amphidium</i> Mougeot's	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Anastrepta</i> Orkney Notc	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Anastrophy.</i> Heller's No	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes	<i>Anastrophy.</i> Comb Notch	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Andreaea a.</i> Alpine Rock	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Andreaea r.</i> Dusky Rock	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes	<i>Andreaea r.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Andreaea r.</i> Black Rock	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes	<i>Andreaea r.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Bryophytes	<i>Andreaea r.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Aneura pin.</i> Greasewort	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bryophytes	<i>Anoectangi.</i> Summer-moss	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Anomobryum</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Anomodon v.</i> Rambling T	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes	<i>Anthelia j.</i> Alpine Silv	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes	<i>Anthoceros</i> Dotted Horn	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes	<i>Antitrichi.</i> Pendulous I	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes	<i>Aphanoleje.</i> Long-leaved	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Atrichum c.</i> Fountain S	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes	<i>Atrichum u.</i> Common Smo	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes	<i>Atrichum u.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes	<i>Aulacomniu.</i> Bud-headed	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes	<i>Aulacomniu.</i> Bog Groove	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes	<i>Barbilopho.</i> Atlantic P	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes	<i>Barbilopho.</i> Bearded Pav	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes	<i>Barbilopho.</i> Common Paw	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes	<i>Barbilopho.</i> Hatcher's I	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes	<i>Barbula co.</i> Lesser Bir	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes	<i>Barbula co.</i> NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Bryophytes	<i>Barbula un.</i> Bird's-claw	0	> -1%	> -1%	LOW	+1 to +4%

Bryophytes <i>Bartramia</i> Haller's App.	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Bartramia</i> Straight-leaved	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Bartramia</i> Common App.	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Bazzania</i> t.Lesser Whip	0	-7.5 to -4%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Bazzania</i> t.Greater Whip	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Blasia</i> pus.Common Ket	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Blepharost</i> Hairy Threa	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Blindia</i> ac.Sharp-leaved	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Brachydont</i> Bristle-leaved	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Brachythec</i> .Whitish Feather	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Brachythec</i> .Sand Feather	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Brachythec</i> .River Feather	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Brachythec</i> .Rough-stall	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Brachythec</i> .Smooth-stall	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Breutel</i> ia Golden-headed	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Bryoerythr</i> Rufous Bear	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> alpi.Alpine Threa	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Bryum</i> arge.Silver-moss	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Bryum</i> born.Potato Bryu	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Bryum</i> caes,Tufted Threa	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> caes,NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Bryum</i> dich.Bicoloured	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> dich,NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Bryum</i> gemm.Small-bud l	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> mora.Flabby Threa	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Bryum</i> pall.Pale Threa	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Bryum</i> pseu.Marsh Bryu	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Bryum</i> pseu,NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Bryum</i> pseu,NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Bryum</i> radi.Wall Threa	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Bryum</i> rube.Crimson-tul	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calli</i> ergon Heart-leaved	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Calli</i> ergon Giant Spear	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Calli</i> ergon Lindberg's	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Notched Pou	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Common Pou	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Mueller's l	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Calypogeia</i> Nees' Pouch	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Calypogeia</i> Bog Pouchwe	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Campyl</i> iade Golden Feather	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Campylium</i> .Yellow Star	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Campylium</i> .NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Campylophy</i> .Chalk Feather	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Campylopus</i> Bristly Swa	0	-4 to -1%	< -7.5%	HIGH	+4 to +7.5%
Bryophytes <i>Campylopus</i> Compact Swa	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Campylopus</i> Rusty Swan-	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Campylopus</i> Brittle Swa	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%

Bryophytes <i>Campylopus</i> Heath Star	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Campylopus</i> Dwarf Swan	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Cephalozia</i> Two-horned	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Cephalozia</i> Chain Pince	0	-7.5 to -4%	-7.5 to -4%	HIGH	+1 to +4%
Bryophytes <i>Cephalozia</i> Moon-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Cephalozie</i> Common Thre	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Cephalozie</i> Hampe's Thu	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Ceratodon</i> Redshank [r	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Chiloscyph</i> St Winifric	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Cirriphyll</i> Beech Featl	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Cirriphyll</i> Hair-pointe	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Cladopodie</i> Bog Notchw	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Climacium</i> Tree-moss	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Cololejeun</i> Rock Pounce	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Cololejeun</i> Minute Pou	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Cololejeun</i> Rossetti's	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Colura cal</i> Fingered Co	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Conocephal</i> Great Scen	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Conostomum</i> Helmet-moss	0	> -1%	< -7.5%	MODERATE	< +1%
Bryophytes <i>Cratoneuro</i> Fern-leaved	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Cratoneuro</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Cryphaea h</i> Lateral Cry	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ctenidium</i> Chalk Comb	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Ctenidium</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Cynodontiu</i> Brunton's I	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Dialytrich</i> Pointed Lat	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Dichodonti</i> Marsh Fork	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Dichodonti</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Dichodonti</i> Transparent	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Dicranella</i> Rufous Forl	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Dicranella</i> Field Fork	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Dicranella</i> Variable Fo	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Dicranowei</i> Common Pinc	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Dicranowei</i> Mountain P	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Dicranum f</i> Whip Fork-r	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Dicranum f</i> Dusky Fork-	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Dicranum f</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Dicranum m</i> Greater Fo	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Dicranum s</i> Broom Fork-	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Dicranum s</i> Scott's Fo	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Dicranum s</i> Rusty Fork-	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Dicranum t</i> Fragile Fo	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Didymodon</i> Pointed Bea	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Didymodon</i> Fallacious	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> Cylindric I	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Didymodon</i> Dusky Bear	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> Nicholson's	0	> -1%	> -1%	LOW	> +7.5%

Bryophytes <i>Didymodon</i> .Rigid Beard	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> .Wavy Beard	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Didymodon</i> .Brown Beard	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Didymodon</i> .Shady Beard	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Didymodon</i> .Soft-tufted	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Diphyscium</i> .Nut-moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Diplophyll</i> .White Earwort	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Distichium</i> .Fine Distich	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Ditrichum</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Ditrichum</i> .Bendy Ditr	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ditrichum</i> .Curve-leaved	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Douinia ov</i> .Waxy Earwort	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Drepanocla</i> .Fertile Feather	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Drepanolej</i> .Toothed P	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Encalypta</i> .Ribbed Ext	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Bryophytes <i>Encalypta</i> .Spiral Ext	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Encalypta</i> .Common Ext	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Entodon co</i> .Montagne's	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Entosthodo</i> .Thin Cord-r	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Entosthodo</i> .Muhlenberg'	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Entosthodo</i> .Blunt Cord-	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Ephemerum</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Ephemerum</i> .Strap-leaved	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Ephemerum</i> .Serrated Ea	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Eremonotus</i> .Clubwort	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Eurhynchiu</i> .Common Str	0	> -1%	> -1%	LOW	< +1%
Bryophytes <i>Fissidens</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> .Maidenhair	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Fissidens</i> .Lesser Pock	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Fissidens</i> .Curnow's P	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> .Welsh Pock	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Fissidens</i> .Fatfoot P	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> .Rock Pocke	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> .Slender P	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> .Narrow-lea	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Fissidens</i> .Short-leave	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Fissidens</i> .Purple-sta	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> .Petty Pock	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> .River Pock	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> .Beck Pocke	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Fissidens</i> .Common P	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Fissidens</i> .NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Fissidens</i> .Green Pock	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Fissidens</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Bryophytes <i>Fontinalis</i> .Greater Wa	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Fossombron</i> .Common Fri	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Fossombron</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Bryophytes <i>Frullania</i> Dilated Scale	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Frullania</i> Spotty Scale	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Frullania</i> Tamarisk Scale	0 -7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Bryophytes <i>Frullania</i> Sea Scalewort	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Funaria hygrometrica</i> Common Cornerwort	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Grimmia donniana</i> Donn's Grimmia	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia funaria</i> String Grimmia	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia laevigata</i> NA	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Grimmia pulvinata</i> Grey-cushion	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Grimmia radiata</i> Spreading-leaf	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia tomentosa</i> Twisted Grimmia	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Grimmia trichomanes</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Gymnocolea inflata</i> Inflated Neck	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Gymnomitrium</i> Braided Frog	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Gymnomitrium</i> Western Frog	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Gymnomitrium</i> White Frost	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Gymnostomum</i> Verdigris	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Gymnostomum</i> Blunt-leaf	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Gymnostomum</i> Luisier's	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Gyroweisia</i> Slender Stem	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Hamatocaulis</i> Varnished Leaf	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Harpalejeunea</i> Pointed Pot	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Harpanthus</i> Stipular Frog	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Hedwigia ciliata</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Bryophytes <i>Hedwigia stellata</i> Starry Hoar	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Hennediella</i> Stanford Scale	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Herbertus</i> Straw Prong	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Heterocladia</i> Wry-leaved	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Heterocladia</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Heterocladia</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Homalia trichomanes</i> Blunt Feather	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Homalotheca</i> Yellow Feather	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Homalotheca</i> Silky Wall	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Hookeria</i> l. Shining Hoar	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Hygroamblystidium</i> Fountain Fe	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Hygroamblystidium</i> Willow Fea	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Hygrobiella</i> Lax Notchwo	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Hygrohypnum</i> Claw Brook	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Hylocomia</i> Shaded Wood	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Hylocomium</i> Glittering	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Hylocomium</i> Flagellate	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Hypnum andriacum</i> Mamillate I	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Hypnum calypso</i> Downy Plain	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Hypnum cupressinum</i> NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Hypnum cupressinum</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Hypnum cupressinum</i> Great Plain	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Hypnum cupressinum</i> Supine Pla	0 > -1%	> -1%	LOW	+4 to +7.5%

Bryophytes <i>Hypnum jut.</i> Heath Plain	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Isopterygi</i> Neat Silk-r	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Isothecium</i> Larger Mous	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Isothecium</i> Holt's Mous	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Isothecium</i> Slender Mou	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Isothecium</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Isothecium</i> NA	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes <i>Jamesoni</i> Autumn Flap	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Jungermann</i> .Dark-green	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Jungermann</i> .Dwarf Flap	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Kiaeria bl</i> Blytt's Fo	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Kiaeria fa</i> Sickle-lea	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Kindbergia</i> Common Fea	0	> -1%	> -1%	LOW	< +1%
Bryophytes <i>Kurzia pau</i> Bristly Fir	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Kurzia syl</i> Wood Finge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Kurzia tri</i> Heath Finge	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Leiocolea</i> .Bantry Not	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Leiocolea</i> .Ragged Not	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Bryophytes <i>Lejeunea c</i> Micheli's I	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Lejeunea l</i> .Western Pot	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Lejeunea p</i> .Pearl Pounc	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Lepidozia</i> .Rock Finge	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Lepidozia</i> .Pearson's I	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Lepidozia</i> .Creeping F	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Leptobarbu</i> .Beric Bear	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Leptodicty</i> .Kneiff's Fe	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Leptodon s</i> .Prince-of-W	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Leptodonti</i> .Bent-leaved	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Leskea pol</i> .Many-fruite	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Leucobryum</i> Large White	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Leucobryum</i> Smaller Wh	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Leucodon s</i> .Squirrel-ta	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Leucodon s</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Loeskeobry</i> .Short-beake	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Lophocolea</i> Bifid Cres	0	> -1%	> -1%	LOW	< +1%
Bryophytes <i>Lophocolea</i> Fragrant C	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Lophocolea</i> Variable-le	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Lophozia e</i> .Capitate No	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Lophozia i</i> .Jagged Not	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Lophozia s</i> .Hill Notch	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Lophozia v</i> .Tumid Notch	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Lunularia</i> .Crescent-cr	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Marchantia</i> Common Live	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Marchantia</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Marchesini</i> .MacKay's P	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Marsupella</i> Scorched Ru	0	> -1%	< -7.5%	MODERATE	< +1%
Bryophytes <i>Marsupella</i> Notched Rus	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Bryophytes <i>Marsupella</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Marsupella</i> Funck's Rus	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Marsupella</i> Stabler's I	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Metzgeria</i> Rock Veilw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Metzgeria</i> Whiskered V	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Metzgeria</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Metzgeria</i> Forked Veil	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Metzgeria</i> Hooked Veil	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Metzgeria</i> Downy Veilw	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Metzgeria</i> Blueish Veil	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Microbryum</i> Floerke's I	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Microlejeu</i> Fairy Beads	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Mnium</i> horn.Swan's-neck	0	-7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Bryophytes <i>Mnium</i> marg.NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Molendoa</i> w.Warburg's M	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Mylia</i> tayl.Taylor's F	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Nardia</i> sca.Ladder Flap	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Neckera</i> co.Flat Necke	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Neckera</i> cr.Crisped Ne	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Neckera</i> pu.Dwarf Necke	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Nowellia</i> c.Wood-rust	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Odontoschi</i> .Bog-m Flap	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Oedipodium</i> Gouty-moss	0	> -1%	< -7.5%	MODERATE	< +1%
Bryophytes <i>Oligotrich</i> .Hercynian I	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Orthodonti</i> .Cape Threa	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Orthotheci</i> .Fine-leaved	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Orthotheci</i> .Red Leskea	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Orthotrich</i> .Wood Brist	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> .Anomalous I	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Orthotrich</i> .White-tippe	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Orthotrich</i> .Lydell's Br	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> .Elegant Br	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> .River Brist	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> .Rock Brist	0	> -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Orthotrich</i> .Showy Brist	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> .Straw Brist	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Orthotrich</i> .Shaw's Bris	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Orthotrich</i> .Slender Br	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Oxyrrhynch</i> .Swartz's Fe	0	> -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Oxyrrhynch</i> .Dwarf Featl	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Oxyrrhynch</i> .Twist-tip I	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Oxyrrhynch</i> .Showy Featl	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Oxystegus</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Palustriell</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Palustriell</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Pellia</i> epi.Overleaf Pe	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Pellia</i> nee.Nees' Pell:	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%

Bryophytes <i>Phascum cuc</i> Cuspidate I	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Phascum cuc</i> .NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Philonotis</i> Thick-nerve	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Philonotis</i> Fountain A	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Physcomit</i> Common Bla	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Plagiobryu</i> Zierian Hur	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> Greater Fea	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Plagiochil</i> Killarney I	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> British Fea	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> Petty Featl	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Plagiochil</i> Western Fea	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiochil</i> Lesser Fea	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Plagiochil</i> Spotty Fea	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiochil</i> Prickly Fea	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Plagiomniu</i> Many-fruite	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiomniu</i> Woodsy Thyr	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiomniu</i> Marsh Thyme	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiomniu</i> Long-beake	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiopus</i> Oeder's App	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Plagiothec</i> Curved Sill	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Plagiothec</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Plagiothec</i> .NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiothec</i> Bright Sill	0	-4 to -1%	< -7.5%	HIGH	+1 to +4%
Bryophytes <i>Plagiothec</i> Alder Silk-	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Plagiothec</i> Woodsy Sill	0	< -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Plagiothec</i> Juicy Silk-	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Plagiothec</i> Waved Silk-	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Plasteurhy</i> Lesser Str	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Platyhypni</i> Portuguese	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Platyhypni</i> Long-beake	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Pleuridium</i> Awl-leaved	0	> -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Pleurozium</i> Red-stemme	0	-7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Bryophytes <i>Pogonatum</i> Aloe Hairc	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Pogonatum</i> Urn Hairc	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Pohlia ann</i> Pale-fruite	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Pohlia bul</i> Blunt-bud	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Pohlia cam</i> Crookneck	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia dru</i> Drummond's	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia elo</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia lut</i> Yellow Thre	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pohlia mel</i> Pink-fruite	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Pohlia nut</i> Nodding Th	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Pohlia wah</i> Pale Glauc	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Polytricha</i> Alpine Hair	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Polytricha</i> Bank Hairc	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Polytricha</i> Slender Ha	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Polytrichu</i> Common Hair	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%

Bryophytes <i>Polytrichu</i> NA	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Polytrichu</i> Juniper Hair	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Polytrichu</i> Bristly Hair	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Polytrichu</i> Strict Hair	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Porella pi</i> Pinnate Scale	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Porella pl</i> Wall Scale	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Preissia q</i> Narrow Mush	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Pseudocall</i> Three-rank	0 -7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Pseudocros</i> Hornschuch'	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Pseudotaxi</i> Elegant Sil	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Pterogoni</i> Bird's-foot	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Ptilidium</i> Ciliated Fr	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ptilidium</i> Tree Fringe	0 < -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Ptilium cr</i> Ostrich-pl	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Ptychomit</i> Long-shank	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitriu</i> Yellow Fri	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Racomitriu</i> Narrow-lea	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitriu</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	< +1%
Bryophytes <i>Racomitriu</i> Oval-fruite	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitriu</i> Long Fringe	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Racomitriu</i> Dense Fring	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitriu</i> Green Moun	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Racomitriu</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Racomitriu</i> Bristly Fr	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Racomitriu</i> Woolly Fri	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Racomitriu</i> Slender Fr	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Radula aqu</i> Brown Scale	0 -7.5 to -4%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Radula com</i> Even Scale	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Radula lin</i> Lindenberg'	0 < -7.5%	-4 to -1%	HIGH	< +1%
Bryophytes <i>Reboulia h</i> Hemisphaer	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Rhabdoweis</i> Toothed Str	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Rhabdoweis</i> Dwarf Stre	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Rhizomnium</i> Felted Thy	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Rhizomnium</i> Dotted Thy	0 -4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Bryophytes <i>Rhynchoste</i> Tender Fea	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Rhynchoste</i> NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Rhynchoste</i> Teesdale Fe	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Rhynchoste</i> Clustered I	0 > -1%	> -1%	LOW	+1 to +4%
Bryophytes <i>Rhynchoste</i> Megapolita	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Rhynchoste</i> Wall Feath	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Rhytidiade</i> Little Sha	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Rhytidiade</i> Big Shaggy	0 -7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Bryophytes <i>Riccardia</i> Jagged Ger	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Riccardia</i> Delicate Ge	0 < -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Riccardia</i> Palmate Ge	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Riccia bey</i> Purple Cry	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Riccia cav</i> Cavernous C	0 > -1%	< -7.5%	MODERATE	> +7.5%

Bryophytes <i>Riccia glauca</i> Glaucous Clubmoss	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Saccogyna</i> Straggling Moss	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Sanionia uliginosa</i> Sick-leaved Moss	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Sarmentypnum</i> Ringless Hair-moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sarmentypnum</i> Twiggy Spear-moss	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania acaulis</i> Lesser Rough-earwort	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania acaulis</i> Rough Earwort	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania ciliata</i> Thick-set Earwort	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania ciliata</i> Untidy Earwort	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Scapania gracilis</i> Western Earwort	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Scapania imbricata</i> Heath Earwort	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Scapania nivalis</i> Grove Earwort	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Scapania norvegica</i> Norwegian Earwort	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Scapania uliginosa</i> Marsh Earwort	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Scapania uliginosa</i> Shady Earwort	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Scapania uliginosa</i> Water Earwort	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Schistidium</i> NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Schistidium</i> Thickpoint Moss	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Schistidium</i> Seaside Green Moss	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Schistidium</i> Upright Moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Schistostoma</i> Luminous Moss	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Sciuro-hypnum</i> Rusty Feather-moss	0	-7.5 to -4%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Sciuro-hypnum</i> Matted Feather-moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Scleropodium</i> Tufted Feather-moss	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Scleropodium</i> Glass-wort Moss	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Scorpidium</i> Intermediate Moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Scorpidium</i> Rusty Hook-moss	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Scorpidium</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Bryophytes <i>Scorpidium</i> Hooked Scorpidium	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Scorpiurium</i> Curving Feather-moss	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Seligeria selaginella</i> Sharp Rock-moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Seligeria selaginella</i> Dwarf Rock-moss	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Solenostoma</i> Crenulated Moss	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Solenostoma</i> Transparent Moss	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Solenostoma</i> Egg Flapwort	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Solenostoma</i> Shining Flapwort	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Solenostoma</i> Round-fruited Moss	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Sphagnum ciliatum</i> Red Bog-moss	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Bryophytes <i>Sphagnum ciliatum</i> Red Bog-moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum ciliatum</i> Compact Bog-moss	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Bryophytes <i>Sphagnum ciliatum</i> Twisted Bog-moss	0	> -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum ciliatum</i> Feathery Bog-moss	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Sphagnum denticulatum</i> Cow-horn Bog-moss	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum flexuosum</i> Flat-topped Bog-moss	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum flexuosum</i> Fringed Bog-moss	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum flexuosum</i> Flexuous Bog-moss	0	> -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum girgensohnii</i> Girgensohn's Moss	0	> -1%	< -7.5%	MODERATE	> +7.5%

Bryophytes <i>Sphagnum i.</i> Lesser Cow-	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum p.</i> Blunt-leaved	0 < -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Bryophytes <i>Sphagnum p.</i> NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Sphagnum p.</i> Papillose l	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Sphagnum p.</i> Golden Bog-	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Sphagnum q.</i> Five-ranked	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum r.</i> NA	0 < -7.5%	-4 to -1%	HIGH	< +1%
Bryophytes <i>Sphagnum r.</i> Russow's Bo	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum s.</i> Spiky Bog-r	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum s.</i> Lustrous Bo	0 -4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Sphagnum s.</i> NA	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Sphagnum s.</i> NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Bryophytes <i>Sphagnum t.</i> Soft Bog-mo	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Sphagnum t.</i> Rigid Bog-r	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Splachnum</i> .Round-frui	0 > -1%	< -7.5%	MODERATE	+1 to +4%
Bryophytes <i>Straminerg.</i> Straw Spear	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Syntrichia</i> Small Hair	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Syntrichia</i> Water Scre	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Syntrichia</i> Intermediat	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Syntrichia</i> Marble Scre	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Syntrichia</i> Sand-hill S	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Syntrichia</i> Great Hair	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Targionia</i> .Orobus-seed	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Tetraphis</i> ,Pellucid Fo	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Bryophytes <i>Tetraplodo.</i> Slender Cri	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Thamnobryu.</i> Fox-tail Fo	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Thuidium d.</i> Delicate Ta	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Thuidium t.</i> Common Tama	0 > -1%	-7.5 to -4%	MODERATE	+1 to +4%
Bryophytes <i>Tortella f.</i> Yellow Cris	0 -7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Bryophytes <i>Tortella i.</i> Sassari Cr	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Tortella n.</i> Neat Crisp-	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Tortella t.</i> Frizzled C	0 > -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Bryophytes <i>Tortula la.</i> Lance-leaved	0 > -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Tortula ma.</i> Bordered Sc	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Tortula mo.</i> Blunt-frui	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Tortula mu.</i> Wall Screw-	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Tortula pr.</i> Tall Pottia	0 > -1%	> -1%	LOW	+4 to +7.5%
Bryophytes <i>Tortula su.</i> Awl-leaved	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Tortula tr.</i> Common Pot	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Trichocole.</i> Handsome W	0 -4 to -1%	< -7.5%	HIGH	> +7.5%
Bryophytes <i>Trichodon</i> .Cylindric l	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Trichostom.</i> Variable C	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Trichostom.</i> Curly Crisp	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Tritomaria</i> Cut Notchw	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Bryophytes <i>Tritomaria</i> Larger Cut	0 > -1%	-4 to -1%	MODERATE	+1 to +4%
Bryophytes <i>Tritomaria</i> Lyon's Not	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Bryophytes <i>Ulota bruc.</i> Bruch's Pi	0 > -1%	> -1%	LOW	> +7.5%

Bryophytes <i>Ulota cris</i> ,Crisped Pi	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ulota cris</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Ulota hutc</i> ,Hutchins' I	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Ulota phyl</i> ,Frizzled P:	0 > -1%	-4 to -1%	MODERATE	> +7.5%
Bryophytes <i>Weissia</i> ,NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Weissia br</i> ,Small-moutl	0 < -7.5%	> -1%	MODERATE	> +7.5%
Bryophytes <i>Weissia co</i> ,NA	0 < -7.5%	< -7.5%	VERY HIGH	> +7.5%
Bryophytes <i>Weissia lo</i> ,Crisp Bearc	0 > -1%	> -1%	LOW	> +7.5%
Bryophytes <i>Weissia lo</i> ,NA	0 > -1%	< -7.5%	MODERATE	> +7.5%
Bryophytes <i>Zygodon co</i> ,Lesser Yoke	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Bryophytes <i>Zygodon ru</i> ,Park Yoke-r	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Bryophytes <i>Zygodon vi</i> ,Green Yoke-	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Bryophytes <i>Zygodon vi</i> ,NA	0 > -1%	< -7.5%	MODERATE	< +1%
Bryophytes <i>Zygodon vi</i> ,NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Acupalpus</i> ,NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Acupalpus</i> ,NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Acupalpus</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Acupalpus</i> ,NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Agonum ema</i> ,NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Agonum ful</i> ,NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Agonum gra</i> ,NA	0 < -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Carbid bee <i>Agonum mar</i> ,NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Agonum mue</i> ,NA	0 < -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Agonum pic</i> ,NA	0 > -1%	< -7.5%	MODERATE	+4 to +7.5%
Carbid bee <i>Agonum tho</i> ,NA	0 -4 to -1%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Agonum vid</i> ,NA	0 > -1%	-7.5 to -4%	MODERATE	> +7.5%
Carbid bee <i>Amara aene</i> ,Common Sun	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Amara apri</i> ,NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Amara bifr</i> ,NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Amara cons</i> ,NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Amara conv</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Amara eque</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Amara eury</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Amara fami</i> ,NA	0 < -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Amara luci</i> ,NA	0 < -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Amara ovat</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Amara pleb</i> ,NA	0 < -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Amara prae</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Amara simi</i> ,NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Amara tibi</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Anchomenus</i> ,NA	0 > -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Anisodacty</i> ,NA	0 > -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Anthracus</i> ,NA	0 -4 to -1%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Asaphidion</i> ,NA	0 > -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Asaphidion</i> ,NA	0 < -7.5%	> -1%	MODERATE	< +1%
Carbid bee <i>Asaphidion</i> ,NA	0 < -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Badister b</i> ,NA	0 > -1%	> -1%	LOW	+1 to +4%

Carbid bee <i>Badister d</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Badister s</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Badister u</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	-4 to -1%	HIGH	< +1%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Bembidion</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Carbid bee <i>Blemus dis</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Blethisa m</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid bee <i>Bracteon l</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Bradycellu</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Bradycellu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Bradycellu</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Carbid bee <i>Bradycellu</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Bradycellu</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Bradycellu</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Broscus ce</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Calathus c</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Calathus e</i> .NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Carbid bee <i>Calathus f</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Calathus m</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Calathus m</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Carbid bee <i>Calathus m</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Calathus r</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Calodromiu</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%

Carbid bee <i>Calosoma i.</i> NA	1	> -1%	< -7.5%	MODERATE	+1 to +4%
Carbid bee <i>Carabus ar.</i> NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Carbid bee <i>Carabus gl.</i> NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Carbid bee <i>Carabus gr.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Carabus mo.</i> Necklace G1	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Carabus ne.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Carabus pr.</i> NA	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Curtonotus</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Curtonotus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Cymindis a.</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Carbid bee <i>Demetrias .</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Demetrias .</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Dromius ag.</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Carbid bee <i>Dromius me.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Dromius qu.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Dyschirius</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Dyschirius</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Dyschirius</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Dyschirius</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Dyschirius</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Carbid bee <i>Elaphrus u.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Eurynebria</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Harpalus a.</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Harpalus a.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Harpalus l.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Carbid bee <i>Harpalus n.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Carbid bee <i>Harpalus r.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Harpalus r.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Carbid bee <i>Harpalus r.</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Harpalus s.</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Harpalus s.</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Laemostenu.</i> NA	0	> -1%	> -1%	LOW	< +1%
Carbid bee <i>Leistus fe.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Leistus fu.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Leistus ru.</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Leistus te.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid bee <i>Licinus de.</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Carbid bee <i>Loricera p.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Masoreus w.</i> NA	0	> -1%	> -1%	LOW	< +1%
Carbid bee <i>Miscodera .</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Nebria bre.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid bee <i>Nebria ruf.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Carbid bee <i>Nebria sal.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Notiophilu.</i> NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid bee <i>Notiophilu.</i> NA	0	-7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Carbid bee <i>Notiophilu.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Notiophilu.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%

Carbid bee <i>Notiophilu</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Notiophilu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Ocys harpa</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid bee <i>Odacantha</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Oodes helo</i> .NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Ophonus ar</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Ophonus az</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Ophonus pu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Ophonus sc</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Oxypselaph</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Panagaeus</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Paradromiu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Paradromiu</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Carbid bee <i>Paranchus</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid bee <i>Patrobus a</i> .NA	0	-4 to -1%	< -7.5%	HIGH	+1 to +4%
Carbid bee <i>Patrobus a</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Philorhizu</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Philorhizu</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Philorhizu</i> .NA	1	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Platyderus</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Platynus a</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Carbid bee <i>Poecilus c</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Poecilus v</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Pogonus ch</i> .NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Pterostich</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Carbid bee <i>Pterostich</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Carbid bee <i>Pterostich</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Carbid bee <i>Pterostich</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Pterostich</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Pterostich</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid bee <i>Pterostich</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Pterostich</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Carbid bee <i>Pterostich</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Carbid bee <i>Pterostich</i> .NA	0	> -1%	> -1%	LOW	< +1%
Carbid bee <i>Pterostich</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Stenolophu</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Stomis pum</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Carbid bee <i>Syntomus f</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Carbid bee <i>Syntomus o</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Carbid bee <i>Syntomus t</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Synuchus v</i> .NA	0	< -7.5%	-4 to -1%	HIGH	< +1%
Carbid bee <i>Tachys bis</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Carbid bee <i>Trechoblem</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Carbid bee <i>Trechus qu</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Carbid bee <i>Trechus ru</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Carbid bee <i>Trechus se</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Carbid bee <i>Trichocell</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%

Carbid bee <i>Trichocell</i> .NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Centipedes <i>Cryptops a</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Centipedes <i>Cryptops h</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Centipedes <i>Geophilus</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Centipedes <i>Geophilus</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Centipedes <i>Geophilus</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Centipedes <i>Geophilus</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Centipedes <i>Henia vesu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Centipedes <i>Lithobius</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Centipedes <i>Lithobius</i> .NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Centipedes <i>Lithobius</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Centipedes <i>Lithobius</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Centipedes <i>Lithobius</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Centipedes <i>Lithobius</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Centipedes <i>Schendyla</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Centipedes <i>Stigmatoga</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Centipedes <i>Strigamia</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Centipedes <i>Strigamia</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Coccinelid <i>Adalia bip</i> .Two-spot La	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Coccinelid <i>Adalia dec</i> .Ten-spot La	0	> -1%	> -1%	LOW	+1 to +4%
Coccinelid <i>Anatis oce</i> .Eyed Ladyb	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Coccinelid <i>Anisostict</i> .Water Ladyb	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinelid <i>Chilocorus</i> .Kidney-spot	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinelid <i>Coccidula</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Coccinelid <i>Coccinella</i> .Seven-spot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Coccinelid <i>Coccinella</i> .Eleven-spot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Coccinelid <i>Exochomus</i> .Pine Ladyb	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Coccinelid <i>Halysia se</i> .Orange Lady	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Coccinelid <i>Hippodamia</i> .Adonis' Lac	0	< -7.5%	> -1%	MODERATE	> +7.5%
Coccinelid <i>Propylea q</i> .Fourteen-sp	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinelid <i>Psyllobora</i> .Twentytwo-s	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Coccinelid <i>Rhyzobius</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Coccinelid <i>Scymnus su</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Coccinelid <i>Subcoccine</i> .Twentyfour-	0	> -1%	> -1%	LOW	> +7.5%
Coccinelid <i>Tytthaspis</i> .Sixteen-sp	0	> -1%	> -1%	LOW	> +7.5%
Craneflies <i>Nephrotoma</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Craneflies <i>Ptychopter</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Craneflies <i>Ptychopter</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Craneflies <i>Ptychopter</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Craneflies <i>Tipula ful</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Craneflies <i>Tipula lat</i> .NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Craneflies <i>Tipula lun</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Craneflies <i>Tipula max</i> .NA	0	> -1%	> -1%	LOW	< +1%
Craneflies <i>Tipula ole</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Craneflies <i>Tipula unc</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Craneflies <i>Tipula var</i> .NA	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Crickets at <i>Chorthippu</i> .Lesser Mars	0	> -1%	> -1%	LOW	> +7.5%

Crickets at <i>Chorthippu</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Chorthippu</i> .Meadow Grass	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Conocephala</i> .NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Crickets at <i>Conocephala</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Crickets at <i>Ectobius p.</i> Tawny Cock	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Crickets at <i>Ectobius p.</i> Lesser Cock	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Crickets at <i>Forficula</i> .Common Ear	0	> -1%	> -1%	LOW	+4 to +7.5%
Crickets at <i>Forficula</i> .Lesne's Ear	0	> -1%	> -1%	LOW	> +7.5%
Crickets at <i>Leptophyes</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Crickets at <i>Meconema t.</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Crickets at <i>Metrioptera</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Crickets at <i>Metrioptera</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Crickets at <i>Myrmeleote</i> .Mottled Grass	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Crickets at <i>Nemobius s.</i> Wood Cricket	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Crickets at <i>Omocestus</i> .Woodland Grass	0	> -1%	< -7.5%	MODERATE	> +7.5%
Crickets at <i>Omocestus</i> .Common Green	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Crickets at <i>Pholidoptera</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Platycleis</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Crickets at <i>Stenobothrus</i> .Stripe-wing	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Crickets at <i>Tetrix cep.</i> Cepero's Grass	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Crickets at <i>Tetrix sub.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Crickets at <i>Tetrix und.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Anasimyia</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Anasimyia</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Anasimyia</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Arctophila</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Baccha elo.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Brachyopa</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Brachyopa</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Brachyopa</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Brachyopa</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Brachypalpa</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Brachypalpa</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Callicera</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Chalcosyrphus</i> .NA	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%

Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Cheilosia</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Chrysogaster</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Chrysogaster</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Chrysotoxus</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Chrysotoxus</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Chrysotoxus</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Chrysotoxus</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Chrysotoxus</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Criorhina</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Criorhina</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Criorhina</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Dasysyrphus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Dasysyrphus</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Hoverflies <i>Dasysyrphus</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Dasysyrphus</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Dasysyrphus</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Hoverflies <i>Didea fasciata</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Didea intermedia</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Epistrophe</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Epistrophe</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Epistrophe</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Epistrophe</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Episyrphus</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Eriozona e.</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Eriozona s.</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Eristalis</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Eristalis</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eristalis</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Eristalis</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Eristalis</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Eristalis</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Eristalis</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Eristalis</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Eumerus fusca</i> Lesser Bull	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eumerus ornatus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eumerus sabula</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%

Hoverflies <i>Eumerus st.</i> Lesser Bull	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Eupeodes b.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eupeodes c.</i> NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eupeodes l.</i> NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Eupeodes l.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Eupeodes n.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Eupeodes n.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Ferdinande.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Helophilus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Helophilus</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Helophilus</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Heringia h.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Heringia p.</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Heringia v.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Lejogaster</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Leucozona .</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Leucozona .</i> NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Hoverflies <i>Melangyna .</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Hoverflies <i>Melangyna .</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Melangyna .</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Melangyna .</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Melangyna .</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Melangyna .</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Melanogast.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Melanogast.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Melanostom.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Hoverflies <i>Meligramma</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Meligramma</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Meliscaeva</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Merodon eq.</i> Greater Bu.	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Microdon a.</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Hoverflies <i>Microdon m.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Myathropa .</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Myolepta d.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Neoascia g.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Neoascia i.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Neoascia m.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Neoascia o.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Neoascia p.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Orthonevra</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Orthonevra</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Hoverflies <i>Orthonevra</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Paragus ha.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Paragus ti.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Parasyrphu.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Parasyrphu.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Parasyrphu.</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%

Hoverflies <i>Parhelophi</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Parhelophi</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Pelecocera</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Pipiza aus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Pipiza bim</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Pipiza fen</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Hoverflies <i>Pipiza lug</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Pipiza lut</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Pipiza noc</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Pipizella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Pipizella</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Hoverflies <i>Platycheir</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Platycheir</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Platycheir</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Platycheir</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Portevinia</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Hoverflies <i>Psilota an</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Rhingia ca</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Riponnensi</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Scaeva sel</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Sericomyia</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Hoverflies <i>Sericomyia</i> .NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Sphaeropho</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Sphegina e</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Sphegina v</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Syritta pi</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Hoverflies <i>Syrphus ri</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Syrphus to</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Syrphus vi</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Trichopsom</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Triglyphus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Tropidia s</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Volucella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Xanthandru</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%

Hoverflies <i>Xanthogram</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Hoverflies <i>Xanthogram</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Hoverflies <i>Xylota abi</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Xylota flo</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Hoverflies <i>Xylota jak</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Hoverflies <i>Xylota seg</i> NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Xylota syl</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Hoverflies <i>Xylota tar</i> NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Hoverflies <i>Xylota xan</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Millipedes <i>Archiboreo</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Millipedes <i>Bianiulus</i> Spotted Snake	0	> -1%	> -1%	LOW	+1 to +4%
Millipedes <i>Boreoiulus</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Millipedes <i>Brachydesm</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Millipedes <i>Chordeuma</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Millipedes <i>Cylindroi</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Millipedes <i>Cylindroi</i> NA	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Millipedes <i>Cylindroi</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Millipedes <i>Cylindroi</i> Blunt-tailed	0	< -7.5%	> -1%	MODERATE	< +1%
Millipedes <i>Glomeris m</i> Pill Millip	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Millipedes <i>Julus scan</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Millipedes <i>Macrostern</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Millipedes <i>Melogona s</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Millipedes <i>Nanogona p</i> Eyed Flat-b	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Millipedes <i>Nemasoma v</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Millipedes <i>Ommatoiulu</i> Striped Mi	0	> -1%	-7.5 to -4%	MODERATE	+1 to +4%
Millipedes <i>Ophiodesmu</i> NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Millipedes <i>Ophiulus</i> NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Millipedes <i>Polydesmus</i> Common Flat	0	> -1%	> -1%	LOW	+1 to +4%
Millipedes <i>Polydesmus</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Millipedes <i>Polydesmus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Millipedes <i>Tachypodoi</i> White-legge	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths <i>Abraxas gr</i> The Magpie	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths <i>Acasis vir</i> Yellow-bar	0	> -1%	> -1%	LOW	> +7.5%
Moths <i>Achlya fla</i> Yellow Horn	0	> -1%	< -7.5%	MODERATE	< +1%
Moths <i>Acronicta</i> The Sycamo	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths <i>Acronicta</i> Alder Moth	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths <i>Acronicta</i> Light Knot	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths <i>Acronicta</i> Knot Grass	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths <i>Acronicta</i> Dark Dagge	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths <i>Actebia pr</i> Portland Mo	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths <i>Adscita ge</i> Cistus For	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths <i>Adscita st</i> The Forest	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths <i>Aethalura</i> Grey Birch	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths <i>Agriopis a</i> Scarce Umb	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths <i>Agriopis l</i> Spring Ush	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths <i>Agriopis m</i> Dotted Bor	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths <i>Agrochola</i> The Brick	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%

Moths	<i>Agrochola</i> .Flounced Cl	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Agrochola</i> .Brown-spot	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Agrochola</i> .Red-line Qu	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Agrochola</i> .Beaded Ches	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Agrochola</i> .Yellow-line	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Agrotis ci</i> .Light Feat	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Agrotis ex</i> Heart & Da	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Agrotis ri</i> .Sand Dart	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Alcis juba</i> Dotted Carp	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Aleucis di</i> .Sloe Carpet	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Allophyes</i> .Green-brinc	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Alsophila</i> .March Moth	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Amphipoea</i> .Crinan Ear	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Amphipoea</i> .Saltern Ear	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Amphipoea</i> .Large Ear	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Angerona p</i> .Orange Mot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Anticlea d</i> .The Stream	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Moths	<i>Anticollix</i> Dentated Pu	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Antitype c</i> .Grey Chi	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Apamea anc</i> .Large Nutm	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Apamea fur</i> The Confus	0	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Apamea lit</i> .Light Arche	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Apamea obl</i> .Crescent S	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Apamea oph</i> .Double Lob	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Apamea sco</i> .Slender Br	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Apamea sor</i> .Rustic Sho	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Apamea sub</i> .Reddish Lig	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Apamea una</i> .Small Clou	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Apeira syr</i> .Lilac Beau	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Apocheima</i> .Small Brinc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Aporophyla</i> Feathered I	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Aporophyla</i> Black Rust	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Archanara</i> .Twin-spotte	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Archanara</i> .Webb's Wain	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Archiearis</i> Light Orang	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Arctia caj</i> .Garden Tig	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Arctia vil</i> .Cream-spot	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Arenostola</i> Fen Wainsc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Asthen a</i> .Small White	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Atethmia c</i> .Centre-barr	1	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Atolmis ru</i> .Red-necked	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Autographa</i> Gold Spang	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Autographa</i> Plain Golde	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Axylia put</i> .The Flame	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Bena bicol</i> .Scarce Silv	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Biston str</i> .Oak Beauty	0	-7.5 to -4%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Blepharita</i> Dark Brocac	1	< -7.5%	> -1%	MODERATE	+1 to +4%

Moths	<i>Cabera exa</i>	Common Wave	0	> -1%	-4 to -1%	MODERATE	< +1%
Moths	<i>Callimorph</i>	Scarlet Tig	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Callistege</i>	Mother Ship	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Calophasia</i>	Toadflax B	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Camptogram</i>	Yellow She	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Caradrina</i>	Mottled Rus	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Carsia sor</i>	Manchester	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Catarhoe c</i>	Royal Mant	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Catarhoe r</i>	Ruddy Carpe	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Catocala n</i>	Red Underw	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Celaena ha</i>	Haworth's M	1	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Cepphis ad</i>	Little Thor	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Cerastis l</i>	White-marke	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Cerura vin</i>	Puss Moth	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Charanyca</i>	Treble Line	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Chesias ru</i>	Broom-tip	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Chilodes m</i>	Silky Wains	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Chlorissa</i>	Small Grass	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Chloroclys</i>	Dark Marble	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Chloroclys</i>	Arran Carpe	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Chloroclys</i>	Autumn Gree	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Chloroclys</i>	Red-green C	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Chortodes</i>	Mere Wainsc	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cidaria fu</i>	Barred Yell	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Clostera c</i>	Chocolate-t	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Coenobia r</i>	Small Rufou	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Coenocalpe</i>	Slender-st	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Colocasia</i>	Nut-tree T	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Colotois p</i>	Feathered	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Comibaena</i>	Blotched Er	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Conistra l</i>	Dark Chest	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Conistra r</i>	Dotted Ches	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Coscinia c</i>	Speckled Fo	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Cosmia aff</i>	Lesser-spot	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cosmia pyr</i>	Lunar-spot	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cosmia tra</i>	The Dun-ba	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Cossus cos</i>	Goat Moth	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Craniophor</i>	The Corone	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Crocallis</i>	Scalloped C	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Cryphia mu</i>	Marbled Gre	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Cucullia a</i>	The Wormwo	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Cucullia a</i>	Star-wort	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Cucullia c</i>	Chamomile S	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cucullia u</i>	The Shark	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Cybosia me</i>	Four-dotte	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Cyclophora</i>	The Mocha	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Cyclophora</i>	Clay Triple	0	> -1%	< -7.5%	MODERATE	> +7.5%

Moths	<i>Cyclophora</i> Dingy Mocha	1	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Cyclophora</i> False Mocha	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Cyclophora</i> Maiden's Bl	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Cymatophor</i> Oak Lutest	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Dasypolia</i> Brindled Oc	1	< -7.5%	-4 to -1%	HIGH	< +1%
Moths	<i>Deilephila</i> Elephant H	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Deilephila</i> Small Eleph	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Deltote ba</i> Silver Bar	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Deltote un</i> Silver Hool	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Diacrisia</i> Clouded Bu	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Diarsia da</i> Barred Ches	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Moths	<i>Dicallomer</i> Dark Tussock	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Dichonia a</i> Merveille o	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Diloba cae</i> Figure of I	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Discoloxia</i> Blomer's R	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Moths	<i>Drepana fa</i> Pebble Hool	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Drymonia d</i> Marbled Bro	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Moths	<i>Dryobotode</i> Brindled G	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Dypterygia</i> Bird's Wing	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Dyscia fag</i> Grey Scallop	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Earias clo</i> Cream-borde	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Egira cons</i> Silver Clou	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Eilema can</i> Hoary Foot	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Eilema com</i> Scarce Foot	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eilema dep</i> Buff Footm	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Eilema gri</i> Dingy Foot	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Eilema lur</i> Common Foot	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Eilema pyg</i> Pigmy Foot	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eilema sor</i> Orange Foot	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Elaphria v</i> Rosy Marble	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Enargia pa</i> Angle-strip	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Endromis v</i> Kentish Glo	0	> -1%	< -7.5%	MODERATE	< +1%
Moths	<i>Ennomos al</i> Canary-sho	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Ennomos au</i> Large Thorn	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Ennomos er</i> September	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Ennomos qu</i> August Tho	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Entephria</i> Grey Mount	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Entephria</i> Yellow-ring	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Epione rep</i> Bordered Be	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Epirrhoe r</i> Wood Carpe	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Epirrhoe t</i> Small Arge	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Epirrita a</i> Autumnal Mo	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Epirrita c</i> Pale Novemb	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Epirrita f</i> Small Autu	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Eremobia o</i> Dusky Sall	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Eriogaster</i> Small Egg	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Euchoeca n</i> Dingy Shell	0	> -1%	> -1%	LOW	> +7.5%

Moths	<i>Euclidia g</i>	Burnet Com	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Eugnorisma</i>	Plain Clay	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Eugnorisma</i>	Autumnal R	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Eulithis m</i>	The Spinach	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eulithis p</i>	The Phoenix	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Eulithis p</i>	Barred Str	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Euphyia bi</i>	Cloaked Ca	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Eupithecia</i>	Brindled P	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Wormwood P	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Currant Pug	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Thyme Pug	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Eupithecia</i>	Oak-tree P	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Pauper Pug	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Mottled Pug	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Haworth's I	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Eupithecia</i>	Tawny Speck	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Eupithecia</i>	Pinion-spot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Maple Pug	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Marbled Pug	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Eupithecia</i>	Larch Pug	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Toadflax P	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Yarrow Pug	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Narrow-wing	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Pimpinel P	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	Lead-colour	0	> -1%	> -1%	LOW	< +1%
Moths	<i>Eupithecia</i>	Foxglove P	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Eupithecia</i>	Satyr Pug	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Eupithecia</i>	Plain Pug	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Shaded Pug	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Bordered P	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Eupithecia</i>	White-spot	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupithecia</i>	Golden-rod	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Eupithecia</i>	Common Pug	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Euproctis</i>	Brown-tail	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Eupsilia t</i>	The Satell	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Eurois occ</i>	Great Broc	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Euxoa curs</i>	Coast Dart	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Euxoa trit</i>	White-line	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Furcula bi</i>	Alder Kite	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Furcula bi</i>	Poplar Kite	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Furcula fu</i>	Sallow Kite	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Gnophos ob</i>	Scotch Ann	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Gortyna fl</i>	Frosted Ora	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Graphiphora</i>	Double Dart	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Gymnosceli</i>	Double-str	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Habrosyne</i>	Buff Arches	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Hadena alb</i>	White Spot	1	< -7.5%	> -1%	MODERATE	> +7.5%

Moths	<i>Hadena com</i>	Varied Corn	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Hecatera b</i>	Broad-barred	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Heliophobu</i>	Bordered Ge	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Hemistola</i>	Small Emera	1	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Hepialus h</i>	Gold Swift	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Hepialus s</i>	Orange Swif	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Herminia g</i>	Small Fan-t	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Hoplodrina</i>	The Uncerta	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Hoplodrina</i>	The Rustic	1	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Horisme te</i>	The Fern	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Hydrelia f</i>	Small Yello	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Hydriomena</i>	May Highfly	0	-4 to -1%	-4 to -1%	MODERATE	+1 to +4%
Moths	<i>Hydriomena</i>	Ruddy High	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Hylaea fas</i>	Barred Red	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Hyles gall</i>	Bedstraw Ha	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Hypena pro</i>	The Snout	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Hypenodes</i>	Marsh Oblic	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Hypomecis</i>	Pale Oak Be	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Hyppa rect</i>	The Saxon	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Idaea aver</i>	Riband Wave	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Idaea bise</i>	Small Fan-t	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Idaea dimi</i>	Single-dot	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Idaea emar</i>	Small Scal	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Idaea fusc</i>	Dwarf Crea	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Idaea muri</i>	Purple-bor	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Idaea seri</i>	Small Dusty	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Idaea subs</i>	Satin Wave	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Idaea trig</i>	Treble Bro	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Ipimorpha</i>	Double Kid	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Ipimorpha</i>	The Olive	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Itame brun</i>	Rannoch Loc	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Jodis lact</i>	Little Emer	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Lacanobia</i>	Bright-line	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Lacanobia</i>	Light Broc	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Lampropter</i>	Water Carpe	0	-4 to -1%	< -7.5%	HIGH	+4 to +7.5%
Moths	<i>Laothoe po</i>	Poplar Hawl	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Larentia c</i>	The Mallow	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Lasiocampa</i>	Oak Eggar	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Lasiocampa</i>	Grass Eggar	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Laspeyria</i>	Beautiful I	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Leucochlae</i>	Beautiful C	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Leucoma sa</i>	White Satin	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Lithomoia</i>	Golden-rod	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Lithophane</i>	Pale Pinion	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Lithophane</i>	Grey Shoul	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Lithosia q</i>	Four-spott	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Lobophora</i>	The Seraph	0	< -7.5%	> -1%	MODERATE	> +7.5%

Moths	<i>Lomographa</i> White-pink	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Luperina n.</i> Sandhill R	0	> -1%	> -1%	LOW	< +1%
Moths	<i>Lycia hirt.</i> Brindled B	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Lygephila</i> ,The Blackne	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Macaria al</i> Sharp-angle	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Macaria no</i> Peacock Mo	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Macaria wa.</i> The V-Moth	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Macrochilo</i> Dotted Fan	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Macrogloss.</i> Hummingbird	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Macrothyla.</i> Fox Moth	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Malacosoma</i> The Lackey	1	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Mamestra b.</i> Cabbage Mo	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Meganola s.</i> Small Black	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Melanchra</i> ,Dot Moth	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Melanthia</i> ,Pretty Cha	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Menophra a.</i> Waved Umber	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Mesoligia</i> .Cloaked Mir	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Miltochris</i> Rosy Footma	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Mimas tili.</i> Lime Hawk-r	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Moma alpiu.</i> Scarce Merv	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Mythimna a.</i> White-point	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Mythimna c.</i> Shoulder-s	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Mythimna o.</i> Obscure Wa	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Mythimna p.</i> Common Wair	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Mythimna p.</i> Striped Wa	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Mythimna p.</i> Devonshire	0	-4 to -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Mythimna t.</i> Double Line	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Mythimna u.</i> White-speck	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Naenia typ.</i> The Gothic	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Nebula sal.</i> Striped Tw	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Noctua com.</i> Lesser Yel	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Noctua fim.</i> Broad-borde	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Nola confu.</i> Least Black	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Nonagria t.</i> Bulrush Wa	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Notodonta</i> ,Iron Promi	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Notodonta</i> .Pebble Pro	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Nudaria mu.</i> Muslin Foo	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Moths	<i>Odezia atr.</i> Chimney Swe	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Odontopera</i> Scalloped I	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Odontosia</i> ,Scarce Pro	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Oligia fas.</i> Middle-bar	0	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Oligia lat.</i> Tawny Marb	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Oligia str.</i> Marbled Mir	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Oligia ver.</i> Rufous Min	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Omphalosce.</i> Lunar Under	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Operophter.</i> Winter Mot	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Operophter.</i> Northern W	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Moths	<i>Opisthogra</i>	Brimstone M	0	> -1%	> -1%	LOW	< +1%
Moths	<i>Oria muscu</i>	Brighton W	1	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Orthosia c.</i>	Small Quake	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Orthosia g.</i>	Hebrew Char	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Orthosia g.</i>	Powdered Qu	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Orthosia i.</i>	Clouded Dra	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Orthosia m.</i>	Blossom Unc	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Orthosia m.</i>	Twin-spotte	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Orthosia o.</i>	Northern D	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Orthosia p.</i>	Lead-colour	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Ourapteryx</i>	Swallow-ta	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Pachycnemi</i>	Horse Chest	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Panemeria</i>	Small Yello	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Panolis fl.</i>	Pine Beaut	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Papestra b.</i>	Glaucous Sl	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Paracolax</i>	Clay Fan-f	1	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Paradarisa</i>	Square Spot	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Paradrina</i>	Pale Mottl	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Parascotia</i>	Waved Black	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Parasemia</i>	Wood Tiger	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Parectropi</i>	Brindled W	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Pasiphila</i>	Sloe Pug	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Pasiphila</i>	Bilberry P	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Pasiphila</i>	Green Pug	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Pelosia mu.</i>	Dotted Foot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Pelurga co.</i>	Dark Spinac	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Perconia s.</i>	Grass Wave	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Peribatode</i>	Willow Bea	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Peridea an</i>	Great Prom	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Perizoma a.</i>	Small Rivu	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Perizoma b.</i>	Pretty Pin	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Perizoma d.</i>	Twin-spot	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Perizoma f.</i>	Sandy Carpe	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Perizoma s.</i>	Marsh Carpe	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Phalera bu.</i>	Buff-tip	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Phibalapte</i>	Oblique Str	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Philereme</i>	Dark Umber	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Philereme</i>	Brown Scal	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Phlogophor</i>	Angle Shade	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Photedes c.</i>	Least Minor	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Phytometra</i>	Small Purp	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Plagodis d.</i>	Scorched W	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Plagodis p.</i>	Barred Umbe	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Moths	<i>Plemyria r.</i>	Blue-border	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Plusia fes</i>	Gold Spot	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Plusia put.</i>	Lempke's G	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Poecilocam</i>	December Mo	0	< -7.5%	> -1%	MODERATE	< +1%

Moths	<i>Polia bomb</i>	Pale Shinin	1	< -7.5%	< -7.5%	VERY HIGH	< +1%
Moths	<i>Polia nebu</i>	Grey Arches	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Polia trim</i>	Silvery Arc	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Moths	<i>Polymixis</i>	Large Ranu	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Polymixis</i>	Feathered I	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Moths	<i>Polyploca</i>	Frosted Gre	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Moths	<i>Pteraphera</i>	Small Sera	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Pterostoma</i>	Pale Promi	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Ptilodon c</i>	Maple Prom	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Ptilophora</i>	Plumed Pro	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Moths	<i>Pyrrhia um</i>	Bordered S	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Rheumapter</i>	Scarce Tiss	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Moths	<i>Rheumapter</i>	Argent & S	1	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Moths	<i>Rheumapter</i>	Scallop She	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Rhizedra l</i>	Large Wain	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Rhyacia si</i>	Dotted Rus	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Rivula ser</i>	Straw Dot	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Saturnia p</i>	Emperor Mo	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Schrankia</i>	Pinion-str	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Schrankia</i>	White-line	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Scopula em</i>	Rosy Wave	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Scopula fl</i>	Cream Wave	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Scopula im</i>	Small Bloo	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Moths	<i>Scopula ru</i>	Tawny Wave	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Moths	<i>Scopula te</i>	Smoky Wave	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Moths	<i>Scotoptery</i>	Chalk Carpe	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Scotoptery</i>	July Belle	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Moths	<i>Scotoptery</i>	Lead Belle	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Selenia de</i>	Early Thorn	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Selenia lu</i>	Lunar Thorn	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Selidosema</i>	Bordered G	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Semiaspila</i>	Yellow Bel	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Moths	<i>Sesia bemb</i>	Lunar Horne	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Setina irr</i>	Dew Moth	0	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Shargacucu</i>	Striped Lye	1	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Shargacucu</i>	The Mullein	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Simyra alb</i>	Reed Dagger	0	> -1%	> -1%	LOW	+4 to +7.5%
Moths	<i>Spaelotis</i>	Stout Dart	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Spargania</i>	White-bande	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Spilosoma</i>	Buff Ermin	1	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Spilosoma</i>	Water Ermi	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Moths	<i>Stilbia an</i>	The Anomalo	1	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Moths	<i>Tethea ocu</i>	Figure of I	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Moths	<i>Tetheella</i>	Satin Lutes	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Thalpophil</i>	Straw Under	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Thera cogn</i>	Chestnut-co	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Moths	<i>Thera cupr</i>	Cypress Can	0	> -1%	> -1%	LOW	> +7.5%

Moths	<i>Thera firm</i>	Pine Carpet	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Moths	<i>Thera juni</i>	Juniper Carpet	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Theria pri</i>	Early Moth	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Tholera ce</i>	Hedge Rust	1	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Tholera de</i>	Feathered (1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Thumatha s</i>	Round-winged	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Trichiura</i>	Pale Egg	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Moths	<i>Trichopter</i>	Early Tooth	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Trichopter</i>	Barred Tooth	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Triphosa d</i>	The Tissue	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Trisateles</i>	Olive Cresc	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Tyta luctu</i>	The Four-sp	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Venusia ca</i>	Welsh Wave	0	-4 to -1%	< -7.5%	HIGH	+4 to +7.5%
Moths	<i>Watsonalla</i>	Barred Hool	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Moths	<i>Xanthia ci</i>	Orange Salt	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Xanthia gi</i>	Dusky-lemon	1	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Xanthia ic</i>	The Sallow	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Xanthia oc</i>	Pale-lemon	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Xanthorhoe</i>	Balsam Carp	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Moths	<i>Xanthorhoe</i>	Red Carpet	1	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Moths	<i>Xanthorhoe</i>	Garden Carp	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Moths	<i>Xanthorhoe</i>	Large Twin-	0	> -1%	< -7.5%	MODERATE	> +7.5%
Moths	<i>Xestia aga</i>	Heath Rust	1	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Xestia tri</i>	Double Squa	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Xylena exs</i>	Sword-grass	0	< -7.5%	> -1%	MODERATE	< +1%
Moths	<i>Xylena vet</i>	Red Sword-g	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Moths	<i>Xylocampa</i>	Early Grey	0	> -1%	> -1%	LOW	> +7.5%
Moths	<i>Zanclognat</i>	The Fan-fo	0	> -1%	> -1%	LOW	+1 to +4%
Moths	<i>Zeuzera py</i>	Leopard Mo	0	< -7.5%	> -1%	MODERATE	> +7.5%
Moths	<i>Zygaena lo</i>	Narrow-bor	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Odonata	<i>Aeshna cae</i>	Azure hawk	0	> -1%	< -7.5%	MODERATE	> +7.5%
Odonata	<i>Aeshna gra</i>	Brown hawk	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Odonata	<i>Aeshna jun</i>	Common haw	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Odonata	<i>Anax imper</i>	Emperor dra	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Brachytron</i>	Hairy drag	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Calopteryx</i>	Banded dem	0	> -1%	< -7.5%	MODERATE	> +7.5%
Odonata	<i>Ceriagrion</i>	Small red	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Coenagrion</i>	Azure damse	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Odonata	<i>Cordulegas</i>	Golden-ring	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Odonata	<i>Cordulia a</i>	Downy eme	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Odonata	<i>Enallagma</i>	Common blu	0	< -7.5%	-4 to -1%	HIGH	< +1%
Odonata	<i>Erythromma</i>	Red-eyed da	0	> -1%	< -7.5%	MODERATE	> +7.5%
Odonata	<i>Gomphus vu</i>	Club-tailed	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Ischnura e</i>	Blue-tailed	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Odonata	<i>Ischnura p</i>	Scarce blue	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Lestes spo</i>	Emerald dar	0	< -7.5%	> -1%	MODERATE	< +1%
Odonata	<i>Libellula</i>	Broad-bodied	0	< -7.5%	-4 to -1%	HIGH	> +7.5%

Odonata	<i>Libellula</i>	Four-spotted	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Orthetrum</i>	Black-tailed	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Orthetrum</i>	Keeled skimmer	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Odonata	<i>Pyrrhosoma</i>	Large red	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Odonata	<i>Somatochlora</i>	Brilliant	0	< -7.5%	> -1%	MODERATE	> +7.5%
Odonata	<i>Sympetrum</i>	Black darter	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Odonata	<i>Sympetrum</i>	Yellow-winged	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Odonata	<i>Sympetrum</i>	Ruddy darter	0	> -1%	> -1%	LOW	> +7.5%
Odonata	<i>Sympetrum</i>	Common darter	0	< -7.5%	> -1%	MODERATE	> +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	-4 to -1%	< -7.5%	HIGH	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH	< +1%
Soldier beetle	<i>Cantharis</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	-4 to -1%	< -7.5%	HIGH	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Soldier beetle	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Cantharis</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Soldier beetle	<i>Malthinus</i>	NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Malthinus</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Soldier beetle	<i>Malthodes</i>	NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Podabrus</i>	NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	Common Red	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Soldier beetle	<i>Rhagonycha</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Spiders	<i>Achaearanea</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Achaearanea</i>	NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Spiders	<i>Agalenatea</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Agelena</i>	labyrinthica	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Agroeca</i>	brachyura	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Agroeca</i>	pinch	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Agroeca</i>	pinch	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Agyneta</i>	depressa	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Agyneta</i>	oligoloba	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Agyneta</i>	raissa	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Agyneta</i>	subulata	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Allomengea</i>	NA	0	-4 to -1%	-4 to -1%	MODERATE	< +1%
Spiders	<i>Allomengea</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Alopecosa</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Alopecosa</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Alopecosa</i>	NA	0	> -1%	> -1%	LOW	+4 to +7.5%

Spiders	<i>Amaurobius</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Amaurobius</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Amaurobius</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Anelosimus</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Anyphaena</i> Buzzing Sp:	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Aphileta m</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Araeoncus</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Araneus ma</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Spiders	<i>Araneus qu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Araneus st</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Araneus tr</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Araniella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Araniella</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Arctosa le</i> .NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Arctosa pe</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Argenna su</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Argyroneta</i> Water Spide	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Atypus aff</i> Purse Web :	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Ballus cha</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Baryphyma</i> ,NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Baryphyma</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Spiders	<i>Bathyphant</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Bathyphant</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Bathyphant</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Bathyphant</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Bianor aur</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Bolyphante</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Bolyphante</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Centromeri</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Centromeru</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Centromeru</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Centromeru</i> .NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Centromeru</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Ceratinell</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Ceratinell</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Ceratinell</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ceratinops</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Cercidia p</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Cheiracant</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Cheiracant</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Cicurina c</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Clubiona b</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona c</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona c</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona l</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Clubiona n</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Clubiona n</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%

Spiders	<i>Clubiona n.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Clubiona p.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Clubiona s.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Clubiona t.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Coelotes a.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Coelotes t.</i> NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Spiders	<i>Crustulina</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Crustulina</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Cryphoea .</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Diaea dors.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Dictyna ar.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Dictyna la</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Dictyna pu.</i> Small Mesh	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Dictyna un.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Dicymbium</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Diplocentr.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Diploceph.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Diploceph.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Diplostyla</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Dismodicus</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Dolomedes .</i> Raft Spider	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Donacochar.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Spiders	<i>Drassodes .</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Drassyllus</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Drepanotyl.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Dysdera cr.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Dysdera er.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Enoplognat.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Enoplognat.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Enoplognat.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Entelecara</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Episinus a.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Episinus t.</i> NA	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Spiders	<i>Erigone ar.</i> NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Erigone at.</i> NA	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Erigone lo.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Erigone pr.</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Ero cambri.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Ero furcat.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ero tuberc.</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Euophrys f.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Evarcha ar.</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Evarcha fa.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Spiders	<i>Floronia b.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Gibbaranea</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Gnaphosa l.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Gnathonari.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Gongylidie.</i> NA	0	> -1%	> -1%	LOW	< +1%
Spiders	<i>Gongylidiu.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Hahnia nav.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Halorates .</i> NA	0	> -1%	> -1%	LOW	< +1%
Spiders	<i>Haplodrass.</i> Heath Gras	1	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Haplodrass.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Haplodrass.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Harpactea .</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Heliophanu.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Heliophanu.</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Hilaira ex.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Hilaira fr.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Hilaira pe.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Hylyphante.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Hypomma co.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Hypomma fu.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Hypseliste.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Hypsosinga</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Hypsosinga</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Hypsosinga</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Kaestneria</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Kaestneria</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Labulla th.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Larinioide.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Lathys hum.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Latithorax</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Lepthyphan</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Lepthyphan</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Lepthyphan</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Lepthyphan</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Lepthyphan</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Lepthyphan</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Lepthyphan</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Leptothrix</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%
Spiders	<i>Linyphia h.</i> NA	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Spiders	<i>Linyphia t.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Lophomma p.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Mangora ac.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Maro minut.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Maso sunde</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Mecopisthe.</i> Peus' s Long	1	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Meioneta i.</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Meioneta m.</i> Thin Weble	1	< -7.5%	> -1%	MODERATE	< +1%

Spiders	<i>Meioneta r.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Meioneta s.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Metellina .</i> NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Metellina .</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Metopobact.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Micrargus .</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Micrargus .</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Micrargus .</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Microlinyp.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Microlinyp.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Micrommata</i> Green Spide	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Milleriana</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Minyriolus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Misumena v.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Moebelia p.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Monocephal.</i> Broad Groov	1	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Monocephal.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Neoscona a.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Neriene cl.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Neriene fu.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Neriene mo.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Neriene pe.</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Nesticus c.</i> Comb-footed	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Nuctenea u.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Oedothorax</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Oedothorax</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Oedothorax</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Oreonetide.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Ostearius .</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ozyptila b.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Ozyptila p.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Ozyptila s.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Ozyptila s.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Ozyptila t.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Pachygnath.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Pachygnath.</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Pachygnath.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Panamomops</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Pardosa ag.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Pardosa am.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Pardosa ho.</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Pardosa mo.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Pardosa ni.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Pardosa pa.</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pardosa pr.</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Pardosa pu.</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Pardosa sa.</i> NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%

Spiders	<i>Pelecopsis</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Pelecopsis</i> NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Pelecopsis</i> NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Spiders	<i>Pelecopsis</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Philodromu</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Philodromu</i> .NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Philodromu</i> .NA	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Spiders	<i>Phrurolith</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pirata lat</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Pirata pir</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Pirata pis</i> .NA	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Spiders	<i>Pisaura mi</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pityohypha</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Pocadicnem</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Pocadicnem</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Poecilonet</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Porrhomma</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Porrhomma</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Porrhomma</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Porrhomma</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Porrhomma</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Robertus a</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Robertus l</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Saaristoa</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Saaristoa</i> .Triangle H	1	< -7.5%	< -7.5%	VERY HIGH	< +1%
Spiders	<i>Saloca dic</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Salticus c</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Salticus s</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Satilatlas</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Scotina gr</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Scotophaeu</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Segestria</i> .NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Silometopu</i> .NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Silometopu</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Spiders	<i>Silometopu</i> .NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Singa hama</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Sitticus c</i> .Sedge Jump	1	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Sitticus p</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Sitticus s</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Stemonypha</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Syedra gr</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Tallusia e</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Tapinocyba</i> NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%

Spiders	<i>Tapinocyba</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Tapinopa</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Taranucnus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Spiders	<i>Tegenaria</i> NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Spiders	<i>Tegenaria</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Tegenaria</i> House Spider	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Spiders	<i>Tegenaria</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Tetragnath</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Tetragnath</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Tetragnath</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Textrix</i> de NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Spiders	<i>Thanatus</i> s NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Theonoe</i> mi NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Theridion</i> NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Spiders	<i>Theridion</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Theridion</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Theridion</i> NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Theridion</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Theridion</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Theridion</i> NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Theridios</i> Ray Spider	0	> -1%	> -1%	LOW	+1 to +4%
Spiders	<i>Tibellus</i> o NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Tmeticus</i> a NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Trichopter</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Trochosa</i> r NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Spiders	<i>Walckenaer</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Spiders	<i>Walckenaer</i> NA	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Walckenaer</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Spiders	<i>Walckenaer</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Spiders	<i>Xysticus</i> a NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Spiders	<i>Xysticus</i> c NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Spiders	<i>Xysticus</i> k NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Xysticus</i> l NA	0	< -7.5%	> -1%	MODERATE	< +1%
Spiders	<i>Xysticus</i> u NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Spiders	<i>Zelotes</i> la NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Zilla</i> diod NA	0	> -1%	> -1%	LOW	> +7.5%
Spiders	<i>Zora</i> spini NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Spiders	<i>Zygiella</i> x NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p	<i>Acer campe</i> Field Maple	0	> -1%	> -1%	LOW	+4 to +7.5%

Vascular p.	<i>Aceras ant.</i>	Man Orchid	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Achillea p.</i>	Sneezewort	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p.	<i>Aconitum n.</i>	Monk's-hood	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Actaea spi.</i>	Baneberry	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Adiantum c.</i>	Maidenhair	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p.	<i>Adoxa mosc.</i>	Moschatel	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Aethusa cy.</i>	Fool's Pars	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Aethusa cy.</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Agrimonia</i>	Agrimony	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Agrostemma</i>	Corncockle	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Agrostis c.</i>	Velvet Bent	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Agrostis c.</i>	Bristle Bent	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Agrostis g.</i>	Black Bent	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Agrostis s.</i>	Creeping Be	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p.	<i>Agrostis v.</i>	Brown Bent	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Ajuga pyra.</i>	Pyramidal I	1	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Ajuga rept.</i>	Bugle	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular p.	<i>Alchemilla</i>	Alpine Lady	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p.	<i>Alchemilla</i>	Hairy Lady'	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p.	<i>Alchemilla</i>	Slender Lac	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Alchemilla</i>	Smooth Lady	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Vascular p.	<i>Alchemilla</i>	Pale Lady's	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Alisma lan.</i>	Narrow-leav	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Alisma pla.</i>	Water-plant	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p.	<i>Alliaria p.</i>	Garlic Must	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p.	<i>Allium ole.</i>	Field Garli	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Allium sco.</i>	Sand Leek	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Allium urs.</i>	Ramsons	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Alnus glut.</i>	Alder	0	-7.5 to -4%	> -1%	MODERATE	< +1%
Vascular p.	<i>Alopecurus</i>	Bulbous Fox	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Alopecurus</i>	Marsh Foxta	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular p.	<i>Alopecurus</i>	Black-grass	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Alopecurus</i>	Meadow Fox	0	-7.5 to -4%	-4 to -1%	HIGH	+1 to +4%
Vascular p.	<i>Althaea of.</i>	Marsh-malle	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p.	<i>Anagallis</i>	Scarlet Pir	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Anagallis</i>	Scarlet Pir	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Anagallis</i>	Chaffweed	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Andromeda</i>	Bog-rosema	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p.	<i>Anemone ne.</i>	Wood Anemor	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p.	<i>Antennaria</i>	Mountain E	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Anthemis a.</i>	Corn Chamom	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p.	<i>Anthoxanth.</i>	Sweet Verna	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Anthriscus</i>	Cow Parsley	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p.	<i>Apera spic.</i>	Loose Silky	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Aphanes ar.</i>	Parsley-pic	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Aphanes au.</i>	Slender Par	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Apium inun.</i>	Lesser Mars	0	< -7.5%	-4 to -1%	HIGH	> +7.5%

Vascular p. <i>Apium nodi</i> Fool's-water	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Arctium l</i> Greater Burdock	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Arctium mi</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Arenaria s</i> Thyme-leaved	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Arenaria s</i> Slender Samolys	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Arenaria s</i> Thyme-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Armoracia</i> Horse-radish	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Arrhenathe</i> False Oat-grass	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Artemisia</i> Mugwort	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Arum itali</i> Italian Lords-and-Ladies	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Arum macul</i> Lords-and-Ladies	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Asparagus</i> Garden Asparagus	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Asperula c</i> Squinancywort	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Asplenium</i> Black Spleenwort	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Asplenium</i> Lanceolate	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Asplenium</i> Maidenhair	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Asplenium</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Asplenium</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Athyrium f</i> Lady-fern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Atriplex g</i> Babington's	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Atriplex p</i> Common Orach	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Atriplex p</i> Spear-leaved	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Atropa bel</i> Deadly Nightshade	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p. <i>Avena fatu</i> Wild-oat	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Baldellia</i> Lesser Water-pennywort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ballota ni</i> Black Horehound	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Barbarea v</i> Winter-cress	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Vascular p. <i>Bellis per</i> Daisy	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Beta vulga</i> Beet	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Betula pen</i> Silver Birch	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Betula pub</i> Downy Birch	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Betula pub</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Bidens cer</i> Nodding Burdock	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Blackstoni</i> Yellow-wort	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Blechnum s</i> Hard-fern	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p. <i>Blysmus co</i> Flat-sedge	1	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Brachypodi</i> Tor-grass	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Brachypodi</i> False Bromegrass	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Brassica o</i> Cabbage	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Brassica r</i> Turnip	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Briza medi</i> Quaking-grass	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Briza mino</i> Lesser Quaking-grass	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Bromopsis</i> Hairy-bromegrass	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Bromus hor</i> Soft-bromegrass	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Bromus hor</i> Least Soft-bromegrass	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Bromus hor</i> Common Soft-bromegrass	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Bromus hor</i> Sand Soft-bromegrass	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p. <i>Bromus rac.</i> Smooth Bro	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Bryonia di</i> White Bryon	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Bupleurum</i> Slender Ha	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Buxus semp</i> Box	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Cakile mar</i> Sea Rocket	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Calamagros</i> Purple Sma	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Calamagros</i> Narrow Sma	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Callitrich</i> Intermediat	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Callitrich</i> Autumnal Wa	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Callitrich</i> Blunt-frui	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Callitrich</i> Common Wate	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Callitrich</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Calluna vu</i> Heather	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Caltha pal</i> Marsh-mari	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Calystegia</i> Hedge Bind	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Calystegia</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Calystegia</i> Great Bind	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Calystegia</i> Sea Bindwe	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Campanula</i> Giant Bell	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Campanula</i> Harebell	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Capsella b</i> Shepherd's-	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Cardamine</i> Large Bitt	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Cardamine</i> Coralroot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cardamine</i> Wavy Bitt	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Cardamine</i> Hairy Bitt	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Cardamine</i> Narrow-lea	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Cardamine</i> Cuckooflow	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Carduus cr</i> Wetted Thi	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Carex acut</i> Slender Tus	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex appr</i> Fibrous Tus	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Carex aqua</i> Water Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex atra</i> Black Alpi	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Carex bige</i> Stiff Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex bine</i> Green-ribb	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Carex capi</i> Hair Sedge	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Carex dian</i> Lesser Tus	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Carex digi</i> Fingered S	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Carex dioi</i> Dioecious S	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Vascular p. <i>Carex dist</i> Distant Sec	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Carex divu</i> Grey Sedge	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Carex divu</i> Many-leave	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex echi</i> Star Sedge	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p. <i>Carex elon</i> Elongated S	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Carex eric</i> Rare Spring	1	< -7.5%	< -7.5%	VERY HIGH	< +1%
Vascular p. <i>Carex hirt</i> Hairy Sedge	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Carex host</i> Tawny Sedge	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Carex humi</i> Dwarf Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p. <i>Carex mages</i> Tall Bog-sedge	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Carex muric</i> Prickly Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex muric</i> Small-fruited	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Carex nigr</i> Common Sedge	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Carex otru</i> False Foxglove	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Carex oval</i> Oval Sedge	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Carex pani</i> Carnation Sedge	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p. <i>Carex pani</i> Greater Tussock	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex pauc</i> Few-flowered	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Carex pilu</i> Pill Sedge	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Carex pseu</i> Cyperus Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex punc</i> Dotted Sedge	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Carex remo</i> Remote Sedge	0	-4 to -1%	< -7.5%	HIGH	+4 to +7.5%
Vascular p. <i>Carex ripa</i> Greater Portulaca	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Carex spic</i> Spiked Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex stri</i> Thin-spiked	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Carex vagi</i> Sheathed Sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex vesi</i> Bladder-sedge	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Carex viri</i> Common Yellow	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Carpinus b</i> Hornbeam	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Castanea s</i> Sweet Chestnut	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Catabrosa</i> Whorl-grass	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Catapodium</i> NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Vascular p. <i>Centaurea</i> Cornflower	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Centaureum</i> Common Centaury	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Centaureum</i> Seaside Centaury	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cephalanth</i> White Heliotrope	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cerastium</i> Sea Mouse-ear	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Cerastium</i> Common Mouse-ear	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Cerastium</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Cerastium</i> NA	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Cerastium</i> Dwarf Mouse-ear	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Ceratocapn</i> Climbing Cuscuta	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Ceratophyl</i> Rigid Hornwort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ceratophyl</i> Soft Hornwort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Chaenorhin</i> Small Toadflax	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Chaerophyl</i> Rough Chervil	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Chamaemel</i> Chamomile	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Chelidoni</i> Greater Celandine	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Chenopodiu</i> Fat-hen	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Chenopodiu</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Chenopodiu</i> Fig-leaved	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Chenopodiu</i> Stinking Goosefoot	1	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Chrysanth</i> Corn Marigold	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Chrysosple</i> Opposite-leaved	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p. <i>Cicendia f</i> Yellow Centaury	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cichorium</i> Chicory	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p. <i>Cicuta vir.</i> Cowbane	0	-4 to -1%	< -7.5%	HIGH	+4 to +7.5%
Vascular p. <i>Circaea lu</i> Enchanter's	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cirsium er</i> Woolly Thistle	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cirsium he</i> Melancholy	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Cochlearia</i> English Sc	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Cochlearia</i> Common Sc	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Cochlearia</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Cochlearia</i> Pyrenean Sc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Colchicum</i> Meadow Saf	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Conopodium</i> Pignut	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Corallorhi</i> Coralroot (0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cornus san</i> Dogwood	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Cornus sue</i> Dwarf Corne	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Coronopus</i> Swine-cress	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Crataegus</i> Hawthorn	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Crepis cap</i> Smooth Hawk	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Crepis pal</i> Marsh Hawk'	0	-4 to -1%	< -7.5%	HIGH	+1 to +4%
Vascular p. <i>Crithmum m</i> Rock Samph	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Cruciata l</i> Crosswort	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Cryptogram</i> Parsley Fe	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cuscuta ep</i> Dodder	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Cuscuta eu</i> Greater Doc	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Cynosurus</i> Crested Dog	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Cyperus lo</i> Galingale	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Cystopteris</i> Brittle Bla	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Cytisus sc</i> Broom	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p. <i>Cytisus sc</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Dactylorhi</i> Common Spo	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Dactylorhi</i> Early Marsh	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Dactylorhi</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Dactylorhi</i> Northern Ma	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Dactylorhi</i> Narrow-lea	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Daucus car</i> Carrot	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Daucus car</i> Wild Carro	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Daucus car</i> Sea Carrot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Deschampsia</i> Wavy Hair-g	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p. <i>Deschampsia</i> Bog Hair-g	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Dianthus a</i> Deptford P	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Dianthus d</i> Maiden Pinl	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Vascular p. <i>Digitalis</i> Foxglove	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Dipsacus f</i> Wild Tease	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Dipsacus p</i> Small Tease	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Draba inca</i> Hoary Whit	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Draba mura</i> Wall Whitl	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Drosera an</i> Great Sunde	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Drosera in</i> Oblong-lea	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Dryopteris</i> Golden-sca	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%

Vascular p. <i>Dryopteris</i> Buckler-Fern	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Dryopteris</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Dryopteris</i> Broad Buckl	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Dryopteris</i> Northern Bu	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Dryopteris</i> Male-fern	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Dryopteris</i> Mountain Ma	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Dryopteris</i> Rigid Buckl	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Echium vul</i> Viper's-bug	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Elatine hy</i> Eight-stame	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Eleocharis</i> Common Spil	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Eleocharis</i> Few-flower	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Elymus can</i> Bearded Cou	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Elytrigia</i> Common Cou	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Empetrum n</i> Crowberry	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Empetrum n</i> Mountain C	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Empetrum n</i> Crowberry	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Epilobium</i> Chickweed W	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Epilobium</i> Alpine Will	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Epilobium</i> Great Will	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Epilobium</i> Spear-leave	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Epilobium</i> Short-frui	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Epilobium</i> Marsh Will	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Epilobium</i> Hoary Will	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Epilobium</i> Pale Willow	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Epilobium</i> Square-sta	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Epipactis</i> Marsh Helle	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Epipactis</i> Green-flow	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Epipactis</i> Violet Hel	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Equisetum</i> Rough Horse	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Equisetum</i> Shady Horse	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p. <i>Equisetum</i> Wood Horse	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p. <i>Equisetum</i> Variegated	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Erica cine</i> Bell Heath	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Erica tetr</i> Cross-leave	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Vascular p. <i>Erica vaga</i> Cornish Hea	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Eriophorum</i> Common Cot	0	-4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Vascular p. <i>Eriophorum</i> Hare's-tail	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Vascular p. <i>Erodium le</i> Sticky Sto	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Erodium ma</i> Sea Stork's	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Erodium mo</i> Musk Stork'	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Erophila v</i> Common Whit	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Erysimum c</i> Wallflower	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Euonymus e</i> Spindle	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p. <i>Euphorbia</i> Caper Spurg	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Euphorbia</i> Petty Spurg	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Euphorbia</i> Broad-leave	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Euphrasia</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%

Vascular p. <i>Euphrasia</i> Confused Eyeb	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Euphrasia</i> Slender Eyeb	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Euphrasia</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Euphrasia</i> Chalk Eyeb	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Euphrasia</i> Scottish Eyeb	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Euphrasia</i> Western Eyeb	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p. <i>Euphrasia</i> Cornish Eyeb	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Fagus sylv.</i> Beech	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Fallopia c.</i> Black-bindw	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Festuca ar.</i> Rush-leaved	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Festuca ar.</i> Tall Fescue	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Festuca fi.</i> Fine-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Festuca gi.</i> Giant Fescue	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Festuca ov.</i> Sheep's-fescue	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Festuca pr.</i> Meadow Fescue	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Festuca ru.</i> Red Fescue	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Festuca ru.</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Festuca ru.</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Festuca vi.</i> Viviparous	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Foeniculum</i> Fennel	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Frankenia</i> Sea-heath	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Fraxinus e.</i> Ash	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Fritillari.</i> Fritillary	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Fumaria ba.</i> Tall Ranunc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Fumaria de.</i> Dense-flower	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Fumaria mu.</i> Common Ranunc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Fumaria mu.</i> Boreau's Ranunc	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Fumaria oc.</i> Western Ranunc	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Vascular p. <i>Fumaria of.</i> Common Fum	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Fumaria of.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Fumaria of.</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Fumaria pa.</i> Fine-leaved	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Fumaria pu.</i> Purple Ranunc	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Fumaria va.</i> Few-flowered	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Gagea lute.</i> Yellow Star	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Galeopsis</i> Red Hemp-net	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Galeopsis</i> Large-flower	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Galeopsis</i> Common Hemp	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Galeopsis</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Galium apa.</i> Cleavers	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Galium bor.</i> Northern Bedstraw	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Galium mol.</i> Hedge Bedstraw	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Galium odo.</i> Woodruff	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Galium pal.</i> Common Marsh	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Galium pum.</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Vascular p. <i>Galium sax.</i> Heath Bedstraw	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p. <i>Galium ste.</i> Limestone I	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p. <i>Gastridium</i> Nit-grass	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Gaudinia</i> f.French Oat	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Genista</i> ti.Dyer's Gre	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Gentianell</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Gentianell</i> Chiltern Ge	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Geranium</i> c.Long-stalk	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Geranium</i> d.Cut-leaved	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Geranium</i> l.Shining Cra	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Geranium</i> m.Dove's-foot	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Geranium</i> p.Little-Rob	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Geranium</i> p.Small-flowe	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Geranium</i> r.Herb-Rober	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p. <i>Geranium</i> s.Wood Crane'	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Geum</i> rival Water Aven	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Geum</i> urban.Wood Aven	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Glechoma</i> h.Ground-ivy	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Glyceria</i> d.Small Sweet	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Glyceria</i> f.Floating Sw	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p. <i>Glyceria</i> m.Reed Sweet-	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Glyceria</i> n.Plicate Sw	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Gnaphalium</i> Heath Cudwe	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Gnaphalium</i> Marsh Cudwe	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Goodyera</i> r.Creeping La	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Groenlandi</i> Opposite-le	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Gymnadenia</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Gymnadenia</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Gymnocarpi</i> Limestone I	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Hammarbya</i> ,Bog Orchid	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Hedera</i> hel.Common Ivy	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular p. <i>Hedera</i> hel.Common Ivy	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Hedera</i> 'Hi.Irish Ivy	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Helianthem</i> Common Rock	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Helictotri</i> Meadow Oat-	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Helleborus</i> Stinking He	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p. <i>Helleborus</i> Green Helle	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Herminium</i> Musk Orchid	1	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Herniaria</i> ,Smooth Rup	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Himantoglo</i> Lizard Orcl	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Hippocrepis</i> Horseshoe V	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Hippuris</i> v.Mare's-tail	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Holcus</i> mol.Creeping Sc	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p. <i>Hordeum</i> mu.Wood Barley	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Hordeum</i> mu.Wall Barley	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Hordeum</i> se.Meadow Bar	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Humulus</i> lu.Hop	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Huperzia</i> s.Fir Clubmos	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Hyacinthoi</i> Bluebell	0	< -7.5%	-7.5 to -4%	VERY HIGH	< +1%

Vascular p. <i>Hydrocotyl</i> Marsh Penny	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Hymenophyl</i> Tunbridge I	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Hymenophyl</i> Wilson's F	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Hypericum</i> Tutsan	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Hypericum</i> Pale St Jol	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Hypericum</i> Perforate S	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Hypericum</i> Slender St	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Hypericum</i> Square-sta	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Hypericum</i> Wavy St Jol	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Hypochaeri</i> Smooth Cat'	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Ilex aquif</i> Holly	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Illecebrum</i> Coral-neck	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Inula cony</i> Ploughman's	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Inula hele</i> Elecampane	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Iris pseud</i> Yellow Iris	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Isoetes ec</i> Spring Qui	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Isolepis c</i> Slender Clu	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Jasione mo</i> Sheep's-bit	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Juncus acu</i> Sharp-flowe	0	-4 to -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Juncus acu</i> Sharp Rush	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Juncus alp</i> Alpine Rus	0	> -1%	< -7.5%	MODERATE	< +1%
Vascular p. <i>Juncus amb</i> Frog Rush	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Juncus art</i> Jointed Rus	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Juncus bal</i> Baltic Rus	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Juncus buf</i> Toad Rush	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Juncus bul</i> Bulbous Rus	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Juncus bul</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Juncus com</i> Round-frui	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Juncus fol</i> Leafy Rush	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Juncus inf</i> Hard Rush	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Juncus tri</i> Three-flowe	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Juniperus</i> Common Jun	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Juniperus</i> Dwarf Junip	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Knautia ar</i> Field Scab	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Lactuca se</i> Prickly Le	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Lamiastrum</i> Yellow Arch	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Lamium alb</i> White Dead-	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Lamium amp</i> Henbit Dea	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Lamium con</i> Northern De	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Lamium pur</i> Red Dead-ne	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Lapsana co</i> Nipplewort	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Lathyrus a</i> Yellow Veto	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Lathyrus j</i> Sea Pea	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Lathyrus l</i> Bitter-veto	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p. <i>Lathyrus p</i> Marsh Pea	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Legousia h</i> Venus's-loe	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Lemna gibb</i> Fat Duckwee	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p	<i>Lemna mino</i>	Common Duckweed	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p	<i>Lemna tris</i>	Ivy-leaved Duckweed	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Leontodon</i>	Autumn Hawkbit	0	-7.5 to -4%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p	<i>Leontodon</i>	Rough Hawkbit	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p	<i>Leontodon</i>	Lesser Hawkbit	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Lepidium h</i>	Smith's Peppercorn	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Lepidium r</i>	Narrow-leaved Peppercorn	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Leucanthem</i>	Oxeye Daisy	0	-7.5 to -4%	> -1%	MODERATE	< +1%
Vascular p	<i>Leymus are</i>	Lyme-grass	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p	<i>Limonium h</i>	Lax-flowered Sea Purslane	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p	<i>Limosella</i>	Mudwort	0	> -1%	> -1%	LOW	> +7.5%
Vascular p	<i>Linaria vu</i>	Common Toadflax	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p	<i>Linum bien</i>	Pale Flax	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Linum pere</i>	Perennial Flax	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p	<i>Listera ov</i>	Common Twyford	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p	<i>Lithosperm</i>	Purple Gromwell	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Lolium per</i>	Perennial Ryegrass	0	> -1%	> -1%	LOW	> +7.5%
Vascular p	<i>Lonicera p</i>	Honeysuckle	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular p	<i>Lotus angu</i>	Slender Bird's-foot	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Vascular p	<i>Lotus pedu</i>	Greater Bird's-foot	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p	<i>Lotus subb</i>	Hairy Bird's-foot	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p	<i>Luronium n</i>	Floating Waterweed	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p	<i>Luzula cam</i>	Field Wood-rush	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p	<i>Luzula for</i>	Southern Wood-rush	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Luzula mul</i>	Heath Wood-rush	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p	<i>Luzula mul</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p	<i>Luzula pil</i>	Hairy Wood-rush	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p	<i>Luzula syl</i>	Great Wood-rush	0	-7.5 to -4%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p	<i>Lychnis fl</i>	Ragged-Robin	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p	<i>Lycopus eu</i>	Gypsywort	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p	<i>Lysimachia</i>	Yellow Pimpernel	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p	<i>Lysimachia</i>	Tufted Loosestrife	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p	<i>Malus sylv</i>	Crab Apple	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p	<i>Malva mosc</i>	Musk-mallow	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p	<i>Malva sylv</i>	Common Mallow	0	> -1%	> -1%	LOW	> +7.5%
Vascular p	<i>Marrubium</i>	White Horehound	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p	<i>Medicago a</i>	Spotted Medick	0	> -1%	> -1%	LOW	> +7.5%
Vascular p	<i>Medicago l</i>	Black Medick	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p	<i>Medicago m</i>	Bur Medick	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p	<i>Medicago p</i>	Toothed Medick	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p	<i>Medicago s</i>	Lucerne	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p	<i>Melampyrum</i>	Common Cow-wort	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p	<i>Melampyrum</i>	Small Cow-wort	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p	<i>Melilotus</i>	Tall Melilot	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p	<i>Melittis m</i>	Bastard Balm	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p	<i>Mentha aqu</i>	Water Mint	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p	<i>Mentha pul</i>	Pennyroyal	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p. <i>Mentha spi.</i> Spear Mint	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Mentha sua</i> Round-leaved	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Mercurialis</i> .Dog's Merc	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Minuartia</i> .Fine-leaved	1	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Minuartia</i> Spring Sand	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Moehringia</i> Three-nerve	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Moenchia e</i> Upright Cha	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Montia fon</i> Blinks	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Montia fon</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Montia fon</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Muscari ne</i> Grape-hyac	1	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Myosotis a</i> Field Forge	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Myosotis d</i> Changing Fo	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Myosotis l</i> .Tufted For	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Myosotis s</i> Creeping Fo	0	-4 to -1%	-7.5 to -4%	HIGH	+4 to +7.5%
Vascular p. <i>Myosotis s</i> Pale Forge	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Myosotis s</i> Wood Forge	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p. <i>Myosurus m</i> Mousetail	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Myrica gal</i> Bog-myrtle	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Myriophyll</i> Whorled Wat	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Najas flex</i> Slender Na	1	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Narcissus</i> ,Daffodil	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Nardus str</i> .Mat-grass	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Narthecium</i> Bog Asphode	0	> -1%	-4 to -1%	MODERATE	+1 to +4%
Vascular p. <i>Nepeta cat</i> .Cat-mint	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Odontites</i> Red Bartsia	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Odontites</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Odontites</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Oenanthe a</i> Fine-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Oenanthe c</i> .Hemlock Wat	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Oenanthe p</i> Corky-fruit	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Ononis rep</i> Common Res	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Onopordum</i> .Cotton This	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Ophiogloss</i> Adder's-ton	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ophrys sph</i> Early Spide	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Orchis mas</i> Early-purp	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Orchis ust</i> Burnt Orch	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Oreopteris</i> Lemon-scent	0	-4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Vascular p. <i>Ornithogal</i> Spiked Star	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Ornithogal</i> Star-of-Ber	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Orobanche</i> .Thyme Broo	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Orobanche</i> .Knapweed Br	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Orobanche</i> .Common Bro	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Oxalis ace</i> Wood-sorre	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Papaver ar</i> Prickly Pop	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Papaver du</i> Long-head	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Papaver du</i> Yellow-juic	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p. <i>Papaver rh.</i> Common Poppy	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Parapholis</i> Hard-grass	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Parentucel</i> Yellow Bar	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Pediculari</i> Lousewort	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Persicaria</i> Amphibious	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Persicaria</i> Water-pepper	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Persicaria</i> Pale Persic	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Persicaria</i> Redshank	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Persicaria</i> Small Water	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Persicaria</i> Tasteless	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Persicaria</i> Alpine Bist	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p. <i>Petroselin</i> Garden Pars	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Petroselin</i> Corn Parsle	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Peucedanum</i> Milk-parsle	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Phalaris a.</i> Reed Canary	0	-7.5 to -4%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Phegopteri</i> Beech Fern	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Phleum alp</i> Alpine Cat'	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Phleum ber</i> Smaller Cat	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Phleum pra</i> Timothy	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Phleum pra</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Phragmites</i> Common Reec	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Phyllitis</i> Hart's-tong	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Picris ech</i> Bristly Ox	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Pilularia</i> Pillwort	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Pimpinella</i> Greater Bu	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Pimpinella</i> Burnet-sax	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Pinguicula</i> Common But	0	-4 to -1%	-7.5 to -4%	HIGH	+1 to +4%
Vascular p. <i>Plantago c</i> Buck's-horn	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Plantago l</i> Ribwort Pla	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Plantago m</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Plantago m</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Plantago m</i> Sea Planta	0	-7.5 to -4%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Poa bulbos</i> Bulbous Mea	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Poa humili</i> Spreading M	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p. <i>Poa nemora</i> Wood Meadow	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Poa pratens</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Poa pratens</i> Smooth Mea	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Poa trivia</i> Rough Mead	0	> -1%	> -1%	LOW	< +1%
Vascular p. <i>Polemonium</i> Jacob's-lac	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Polygala c</i> Chalk Milk	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Polygala s</i> Heath Milk	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Polygala v</i> Common Mill	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Polygonatu</i> Solomon's-s	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Polygonatu</i> Angular Sol	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Polygonum</i> Equal-leave	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p. <i>Polygonum</i> Knotgrass	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Polygonum</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%

Vascular p. <i>Polygonum</i> Ray's Knot	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Polygonum</i> Cornfield I	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Polypodium</i> Southern Po	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Polypodium</i> Intermediate	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Polypodium</i> Polypody	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Polypodium</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Polystichu</i> Soft Shield	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Populus ni</i> Black-popla	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Potamogeto</i> Fen Pondwee	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Potamogeto</i> Grass-wrack	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Potamogeto</i> Flat-stalk	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Potamogeto</i> Broad-leav	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Potamogeto</i> Fennel Pond	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Potamogeto</i> Bog Pondwee	0	> -1%	-4 to -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Potamogeto</i> Hairlike Po	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Potentilla</i> Trailing To	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Potentilla</i> Silverweed	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Potentilla</i> Alpine Cinc	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Potentilla</i> Tormentil	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Potentilla</i> Shrubby Cin	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Potentilla</i> Spring Cinc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Potentilla</i> Creeping C	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Potentilla</i> Barren Stra	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Primula el</i> Oxlip	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Primula ve</i> Cowslip	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Primula vu</i> Primrose	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Prunella v</i> Selfheal	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Prunus cer</i> Dwarf Cher	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Prunus dom</i> Wild Plum	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Prunus dom</i> Plum	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Prunus dom</i> Bullace; Dar	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Prunus pad</i> Bird Cherry	0	-7.5 to -4%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Prunus spi</i> Blackthorn	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Pteridium</i> Bracken	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p. <i>Puccinelli</i> Stiff Salt	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Pulicaria</i> Common Flea	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Pulmonaria</i> Narrow-leav	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p. <i>Pulsatilla</i> Pasqueflowe	1	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p. <i>Pyrola med</i> Intermediate	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Pyrola rot</i> Round-leav	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Pyrola rot</i> Wintergreen	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Quercus pe</i> Sessile Oal	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Quercus ro</i> Pedunculate	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Radiola li</i> Allseed	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Ranunculus</i> Meadow But	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Ranunculus</i> Common Wate	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ranunculus</i> Corn Butter	1	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%

Vascular p. <i>Ranunculus</i> Bulbous Buttercup	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Ranunculus</i> Lesser Celandine	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Ranunculus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ranunculus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ranunculus</i> Lesser Speedwell	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Ranunculus</i> River Watercress	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Ranunculus</i> Round-leaved	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Ranunculus</i> Stream Watercress	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Ranunculus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Ranunculus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Ranunculus</i> Thread-leaved	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ranunculus</i> Three-lobed	1	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Raphanus</i> r.Wild Radish	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Reseda</i> l.ut.Weld	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Rhinanthus</i> Yellow-rattle	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Rhinanthus</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Rhinanthus</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Ribes</i> alpi.Mountain Currant	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Ribes</i> rubr.Red Currant	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Vascular p. <i>Ribes</i> spic.Downy Currant	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rorippa</i> mi.Narrow-fruited	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rorippa</i> na.Water-cress	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rorippa</i> na.NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rorippa</i> sy.Creeping Yellowcress	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Rosa</i> arven.Field-rose	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rosa</i> caesi.NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rosa</i> caesi.Hairy Dog-rose	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Rosa</i> canin.Dog-rose	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Rosa</i> micra.Small-flowered	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rosa</i> molli.Soft Downy	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rosa</i> rubig.Sweet-briar	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rosa</i> stylo.Short-styled	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Rosa</i> tomen.Harsh Downy	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rubia</i> pere.Wild Madder	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rubus</i> cham.Cloudberry	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Rubus</i> frut.Bramble	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rubus</i> idae.Raspberry	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Rubus</i> saxa.Stone Bramble	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Rumex</i> acet.Common Sorrel	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Rumex</i> acet.Sheep's Sorrel	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rumex</i> acet.Narrow-Leaved	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Rumex</i> cong.Clustered I	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rumex</i> cris,NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Rumex</i> palu.Marsh Dock	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Rumex</i> rupe.Shore Dock	1	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Rumex</i> sang.Wood Dock	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Ruppia</i> cir.Spiral Tassell	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Vascular p. <i>Ruppia mar</i> Beaked Tass	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Ruscus acu</i> Butcher's-s	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Sagina ape</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Sagina mar</i> Sea Pearlwe	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Sagina nod</i> Knotted Pea	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Sagina pro</i> Procumbent	0	-7.5 to -4%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Sagina sub</i> Heath Pear	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Sagittaria</i> Arrowhead	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Salicornia</i> Long-spike	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Salicornia</i> Common Glas	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Salicornia</i> Yellow Glas	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Salicornia</i> One-flower	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Salicornia</i> Purple Glas	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Salix alba</i> White Will	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Salix auri</i> Eared Will	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Salix capr</i> Goat Willow	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Salix capr</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Salix cine</i> Grey Willow	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Salix cine</i> Grey Willow	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Salix cine</i> Rusty Will	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Salix frag</i> Crack-will	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Salix herb</i> Dwarf Will	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Salix lapp</i> Downy Will	1	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p. <i>Salix myrs</i> Dark-leaved	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Salix phyl</i> Tea-leaved	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Salix purp</i> Purple Will	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Salix repe</i> Creeping W	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Salsola ka</i> Prickly Sa	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Sambucus e</i> Dwarf Elder	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Sambucus n</i> Elder	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Samolus va</i> Brookweed	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Sanguisorb</i> Salad Burne	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Sanguisorb</i> Great Burne	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Sanicula e</i> Sanicle	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Sarcocorni</i> Perennial C	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Saussurea</i> Alpine Saw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Saxifraga</i> Mossy Saxi	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Scabiosa c</i> Small Scab	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Scandix pe</i> Shepherd's	1	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p. <i>Schoenus n</i> Black Bog-1	0	> -1%	-7.5 to -4%	MODERATE	> +7.5%
Vascular p. <i>Scilla aut</i> Autumn Squ	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Scilla ver</i> Spring Squ	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Scirpus sy</i> Wood Club-1	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Scrophular</i> Water Figw	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p. <i>Scrophular</i> Common Fig	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Scrophular</i> Green Figw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Sedum angl</i> English Sto	0	< -7.5%	-4 to -1%	HIGH	> +7.5%

Vascular p.	<i>Sedum rosea</i>	Roseroot	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Sedum vill</i>	Hairy Stone	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p.	<i>Selaginell</i>	Lesser Clu	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p.	<i>Senecio aq</i>	Marsh Ragw	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Senecio er</i>	Hoary Ragw	0	> -1%	< -7.5%	MODERATE	+1 to +4%
Vascular p.	<i>Senecio ja</i>	Common Ragv	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p.	<i>Senecio sy</i>	Heath Grou	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Senecio vu</i>	Groundsel	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Senecio vu</i>	Groundsel	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Seriphidiu</i>	Sea Wormwo	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Sherardia</i>	Field Madde	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p.	<i>Sibthorpia</i>	Cornish Mor	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p.	<i>Silaum sil</i>	Pepper-sax	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p.	<i>Silene con</i>	Sand Catch	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Silene dio</i>	Red Campio	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Silene gal</i>	Small-flowe	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Silene lat</i>	White Camp	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p.	<i>Silene noc</i>	Night-flowe	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p.	<i>Silene nut</i>	Nottingham	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Silene vul</i>	Bladder Car	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Sinapis ar</i>	Charlock	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p.	<i>Sisymbrium</i>	Hedge Musta	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Sium latif</i>	Greater War	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Solanum du</i>	Bittersweet	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Vascular p.	<i>Solanum ni</i>	Black Nigh	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Solidago v</i>	Goldenrod	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p.	<i>Sonchus ar</i>	Perennial S	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Sonchus as</i>	Prickly Sov	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p.	<i>Sonchus ol</i>	Smooth Sow	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p.	<i>Sonchus pa</i>	Marsh Sowth	0	> -1%	> -1%	LOW	+1 to +4%
Vascular p.	<i>Sorbus ari</i>	Common Whit	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p.	<i>Sorbus auc</i>	Rowan	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p.	<i>Sorbus dev</i>	Devon White	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Sparganium</i>	Floating Bu	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Sparganium</i>	Unbranched	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Sparganium</i>	Branched Bu	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Spartina m</i>	Small Cord	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Spergula a</i>	Corn Spurre	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Spergulari</i>	Lesser Sea	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Spergulari</i>	Greater Sea	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Spirodela</i>	Greater Duc	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Stachys ar</i>	Field Wound	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p.	<i>Stachys of</i>	Betony	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Stachys pa</i>	Marsh Wound	0	-4 to -1%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Stachys sy</i>	Hedge Wound	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Stellaria</i>	Lesser Sti	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p.	<i>Stellaria</i>	Greater Sti	0	< -7.5%	> -1%	MODERATE	< +1%

Vascular p.	<i>Stellaria</i>	Lesser Chickweed	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Stellaria</i>	Marsh Stitchwort	1	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Stellaria</i>	Bog Stitchwort	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Stratiotes</i>	Water-soldanel	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Subularia</i>	Awlwort	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Succisa pr.</i>	Devil's-bit	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p.	<i>Symphytum</i>	Common Comfrey	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Tamus comm.</i>	Black Bryony	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p.	<i>Tanacetum</i>	Feverfew	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Taxus bacc.</i>	Yew	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p.	<i>Teucrium s.</i>	Wood Sage	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Thalictrum</i>	Common Meadow	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Thelypteris</i>	Marsh Fern	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Thesium hu.</i>	Bastard-toadflax	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Thlaspi ca.</i>	Alpine Pennycress	0	> -1%	< -7.5%	MODERATE	+4 to +7.5%
Vascular p.	<i>Thymus pol.</i>	Wild Thyme	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Vascular p.	<i>Tilia plat.</i>	Large-leaved Lime	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Tofieldia</i>	Scottish Asphodel	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p.	<i>Torilis ar.</i>	Spreading Horsetail	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Torilis ja.</i>	Upright Horsetail	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p.	<i>Tragopogon</i>	Goat's-beard	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Trichophor.</i>	Northern Duckweed	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p.	<i>Trichophor.</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p.	<i>Trichophor.</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Trientalis</i>	Chickweed-flower	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Trifolium</i>	Lesser Trefoil	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Trifolium</i>	Strawberry	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Trifolium</i>	Clustered	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Trifolium</i>	Zigzag Clover	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Trifolium</i>	Slender Trefoil	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Trifolium</i>	Bird's-foot	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Trifolium</i>	White Clover	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Vascular p.	<i>Trifolium</i>	Rough Clover	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Trifolium</i>	Knotted Clover	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p.	<i>Trifolium</i>	Subterranean	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Triglochin</i>	Sea Arrowgrass	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Triglochin</i>	Marsh Arrowgrass	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Tripleuros</i>	Scentless May	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Trisetum f.</i>	Yellow Oatgrass	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p.	<i>Trollius e.</i>	Globeflower	0	< -7.5%	< -7.5%	VERY HIGH	+1 to +4%
Vascular p.	<i>Typha lati.</i>	Bulrush	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p.	<i>Ulex europ.</i>	Gorse	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Ulex galli.</i>	Western Gorse	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Ulex minor</i>	Dwarf Gorse	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p.	<i>Ulmus glab.</i>	Wych Elm	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p.	<i>Ulmus mino.</i>	NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p.	<i>Ulmus proc.</i>	English Elm	0	< -7.5%	> -1%	MODERATE	+1 to +4%

Vascular p. <i>Umbilicus</i> Navelwort	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Urtica ure</i> Small Nettl	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Utriculari</i> Bladderwort	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Utriculari</i> NA	0	-4 to -1%	-7.5 to -4%	HIGH	> +7.5%
Vascular p. <i>Utriculari</i> Greater Bla	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Vaccinium</i> Bilberry	0	< -7.5%	-7.5 to -4%	VERY HIGH	+1 to +4%
Vascular p. <i>Vaccinium</i> Bog Bilber	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Vaccinium</i> Cowberry	0	-7.5 to -4%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Valeriana</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	< +1%
Vascular p. <i>Valeriane</i> Keeled-fru	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Valeriane</i> Narrow-fru	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Valeriane</i> Hairy-frui	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Valeriane</i> Broad-frui	1	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Veronica a</i> Green Field	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Veronica b</i> Brooklime	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Vascular p. <i>Veronica c</i> Pink Water-	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Veronica c</i> Germander S	0	< -7.5%	> -1%	MODERATE	< +1%
Vascular p. <i>Veronica h</i> Ivy-leaved	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Veronica h</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Veronica h</i> NA	0	-4 to -1%	< -7.5%	HIGH	> +7.5%
Vascular p. <i>Veronica m</i> Wood Speed	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Veronica o</i> Heath Spee	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Veronica s</i> Thyme-leave	0	-7.5 to -4%	-7.5 to -4%	HIGH	> +7.5%
Vascular p. <i>Veronica s</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Veronica s</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Vascular p. <i>Viburnum o</i> Guelder-ros	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Vicia hirs</i> Hairy Tare	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Vicia lute</i> Yellow-veto	0	< -7.5%	-7.5 to -4%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Vicia orob</i> Wood Bitter	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Vicia parv</i> Slender Tai	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Vicia sati</i> Narrow-lea	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Vicia sati</i> NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p. <i>Vicia sati</i> Common Veto	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Vicia sylv</i> Wood Vetch	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Vinca mino</i> Lesser Per	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Viola arve</i> Field Pansy	0	> -1%	> -1%	LOW	+4 to +7.5%
Vascular p. <i>Viola cani</i> Heath Dog-v	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Vascular p. <i>Viola hirt</i> Hairy Viole	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Viola lact</i> Pale Dog-v	1	> -1%	-4 to -1%	MODERATE	+1 to +4%
Vascular p. <i>Viola lute</i> Mountain Pa	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Vascular p. <i>Viola odor</i> Sweet Viole	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Viola palu</i> Marsh Viole	0	> -1%	-4 to -1%	MODERATE	> +7.5%
Vascular p. <i>Viola palu</i> NA	0	> -1%	< -7.5%	MODERATE	> +7.5%
Vascular p. <i>Viola palu</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Vascular p. <i>Viola reic</i> Early Dog-v	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p. <i>Viola rivi</i> Common Dog-	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Vascular p. <i>Viola tric</i> Wild Pansy	0	< -7.5%	> -1%	MODERATE	> +7.5%

Vascular p.	<i>Viola tric</i>	Seaside Pai	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Viola tric</i>	NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Vascular p.	<i>Vulpia myu</i>	Rat's-tail	0	> -1%	> -1%	LOW	> +7.5%
Vascular p.	<i>Zostera ma</i>	Eelgrass	0	< -7.5%	> -1%	MODERATE	> +7.5%
Vascular p.	<i>Zostera no</i>	Dwarf Eelg	0	-4 to -1%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Agenioideu</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ammophila</i>	Red Banded	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	> -1%	-7.5 to -4%	MODERATE	+4 to +7.5%
Wasps	<i>Ancistroce</i>	NA	0	< -7.5%	-4 to -1%	HIGH	> +7.5%
Wasps	<i>Ancistroce</i>	Wall Mason	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ancistroce</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Anoplius c</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Wasps	<i>Anoplius i</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Anoplius n</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Anoplius v</i>	Black Banded	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Wasps	<i>Arachnospi</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Arachnospi</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Arachnospi</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Arachnospi</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Argogoryte</i>	Field Digge	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Astata boo</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Astata pin</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Auplopus c</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Caliadurgu</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Cerceris a</i>	Sand Taile	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Cerceris r</i>	Ornate Tail	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Chrysis an</i>	NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Chrysis ig</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Chrysis im</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Chrysis me</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Chrysis vi</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Cleptes se</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Crabro cri</i>	Slender Bo	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crabro pel</i>	NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crabro scu</i>	NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Crossoceru</i>	Blunt Tail	0	< -7.5%	< -7.5%	VERY HIGH	> +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Crossoceru</i>	Slender Dig	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	-4 to -1%	HIGH	+4 to +7.5%
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	MODERATE	> +7.5%

Wasps	<i>Crossocerus</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crossocerus</i> .4-Spotted I	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Crossocerus</i> .Wesmael's I	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Diodontus</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Diodontus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Diodontus</i> .Melancholy	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Dipogon</i> .su.NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Dolichoves</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Dolichoves</i> .Tree Wasp	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Ectemnius</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Ectemnius</i> .NA	0	-7.5 to -4%	< -7.5%	VERY HIGH	> +7.5%
Wasps	<i>Ectemnius</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Ectemnius</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Ectemnius</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Elampus</i> .pa.NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Entomognathus</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Episyron</i> .r.Red Legged	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Eumenes</i> .co.Heath Potter	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Evagetes</i> .c.NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Evagetes</i> .d.NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Gorytes</i> .bi.NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Gorytes</i> .qu.4-Banded D	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Gorytes</i> .tu.NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Gymnomerus</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Hedychridi</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Hedychridi</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Hedychridi</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Lindenius</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Lindenius</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Mellinus</i> .a.Field Digger	0	> -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Microdynerus</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Miscophus</i> .NA	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Mutilla</i> .eu.Large Velvet	0	< -7.5%	-7.5 to -4%	VERY HIGH	> +7.5%
Wasps	<i>Myrmosa</i> .at.NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Nysson</i> .dim.Small Spuri	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Nysson</i> .spi.Large Spuri	0	< -7.5%	< -7.5%	VERY HIGH	+4 to +7.5%
Wasps	<i>Nysson</i> .tri.NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Odynerus</i> .m.NA	1	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Odynerus</i> .s.Spiny Mason	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Omalus</i> .aen.NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Omalus</i> .aur.NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Omalus</i> .vio.NA	0	-7.5 to -4%	> -1%	MODERATE	> +7.5%
Wasps	<i>Oxybelus</i> .a.Silver Spiri	0	< -7.5%	> -1%	MODERATE	> +7.5%

Wasps	<i>Oxybelus m</i> Pale Jawed	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Oxybelus u</i> Common Spic	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Passaloecu</i> .Horned Blac	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Passaloecu</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Passaloecu</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Passaloecu</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Wasps	<i>Passaloecu</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Pemphredon</i> Mournful Wa	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Pemphredon</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Podalonia</i> .Hairy Sand	0	< -7.5%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Pompilus c</i> .Leaden Spic	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Priocnemis</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Priocnemis</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Priocnemis</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Psen bruxe</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Psen dahlb</i> .NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Psen eques</i> NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Psen lutar</i> .NA	0	-7.5 to -4%	> -1%	MODERATE	+1 to +4%
Wasps	<i>Psenulus c</i> .NA	0	> -1%	> -1%	LOW	+1 to +4%
Wasps	<i>Psenulus p</i> .Pale Footed	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Rhopalum c</i> .NA	0	< -7.5%	-4 to -1%	HIGH	+1 to +4%
Wasps	<i>Rhopalum c</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Sapyga cla</i> NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Sapyga qui</i> .NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Smicromyr</i> m.Small Velv	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Spilomena</i> .NA	0	< -7.5%	> -1%	MODERATE	+4 to +7.5%
Wasps	<i>Spilomena</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Symmorphus</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Tachysphex</i> NA	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Tiphia fem</i> .NA	0	> -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Tiphia min</i> .Small Tiph	0	> -1%	> -1%	LOW	+4 to +7.5%
Wasps	<i>Trichrysis</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> Slender Woc	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> Club Horne	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> NA	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Trypoxylon</i> NA	0	-4 to -1%	> -1%	MODERATE	> +7.5%
Wasps	<i>Vespa crab</i> .Hornet	0	> -1%	< -7.5%	MODERATE	> +7.5%
Wasps	<i>Vespula ge</i> .German Was	0	> -1%	> -1%	LOW	> +7.5%
Wasps	<i>Vespula ru</i> .Red Wasp	0	< -7.5%	> -1%	MODERATE	> +7.5%
Wasps	<i>Vespula vu</i> .Common Was	0	> -1%	> -1%	LOW	> +7.5%

Projected expansion	Benefit from expansion	Final outcome
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	Limited impact
+4 to +7.5%	VERY HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk

< +1%	MODERATE	Medium benefit
< +1%	LOW	Limited impact
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	LOW	Limited impact
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	LOW	Limited impact
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	Risks & benefits
< +1%	LOW	Limited impact
< +1%	MODERATE	High risk
< +1%	LOW	Limited impact
< +1%	MODERATE	Medium benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit

< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk

+1 to +4%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	LOW	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Risks & benefits
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	MODERATE	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	LOW	Medium risk
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	LOW	Medium risk
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
> +7.5%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit

+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium benefit

< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+4 to +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	LOW	High risk
+4 to +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk

< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit

> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	Risks & benefits
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk

> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium risk
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit

> +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	LOW	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	LOW	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	LOW	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit

+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	Medium benefit

< +1%	LOW	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
< +1%	LOW	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Medium benefit
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit

+1 to +4%	MODERATE	Medium benefit
< +1%	LOW	High risk
+4 to +7.5%	HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits

< +1%	LOW	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit

+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Medium benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit

+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit

< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Medium benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	High benefit
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
> +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits

> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	LOW	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk

< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Medium benefit
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	LOW	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit

< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits

< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit

< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	LOW	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	High risk

< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
< +1%	LOW	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit

> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium benefit
+4 to +7.5%	MODERATE	Risks & benefits
> +7.5%	HIGH	Risks & benefits
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk

< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
+4 to +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	MODERATE	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
+1 to +4%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Medium benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk

+4 to +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	Medium benefit

< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
+4 to +7.5%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	Risks & benefits
+1 to +4%	HIGH	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
+1 to +4%	MODERATE	Risks & benefits
+4 to +7.5%	MODERATE	Risks & benefits

> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
+1 to +4%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	HIGH	Medium benefit

< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	LOW	High risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	High risk
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
< +1%	MODERATE	High risk
+1 to +4%	HIGH	Medium benefit
+1 to +4%	HIGH	Medium benefit
+4 to +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+1 to +4%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
+4 to +7.5%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	High risk
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	HIGH	Medium benefit
> +7.5%	HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Medium risk
> +7.5%	VERY HIGH	High benefit
+4 to +7.5%	HIGH	Medium benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
< +1%	MODERATE	Risks & benefits
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit
> +7.5%	VERY HIGH	High benefit

Group	Latin Name	English name	NERC na species	Observed decline	Projected decline	Risk of decline
Ants	<i>Formica c</i>	NA	0	< -7.5%	> -1%	LOW
Ants	<i>Formica fu</i>	Negro Ant	0	> -1%	> -1%	LOW
Ants	<i>Formica sa</i>	NA	0	-4 to -1%	< -7.5%	HIGH
Ants	<i>Lasius ali</i>	NA	0	< -7.5%	> -1%	LOW
Ants	<i>Lasius fla</i>	Yellow Mea	0	< -7.5%	> -1%	LOW
Ants	<i>Lasius mix</i>	NA	0	> -1%	> -1%	LOW
Ants	<i>Lasius nig</i>	Small Blac	0	> -1%	> -1%	LOW
Ants	<i>Leptothora</i>	Slender An	0	> -1%	-4 to -1%	MODERATE
Ants	<i>Myrmica ru</i>	Red Ant	0	< -7.5%	> -1%	LOW
Ants	<i>Myrmica ru</i>	NA	0	> -1%	> -1%	LOW
Ants	<i>Myrmica sa</i>	NA	0	> -1%	-4 to -1%	MODERATE
Ants	<i>Myrmica sc</i>	NA	0	-7.5 to -4%	> -1%	LOW
Ants	<i>Myrmica sc</i>	NA	0	> -1%	> -1%	LOW
Bees	<i>Andrena ha</i>	Early Mini	0	-4 to -1%	> -1%	LOW
Bees	<i>Andrena ta</i>	Tormentil	1	< -7.5%	> -1%	LOW
Bees	<i>Andrena va</i>	NA	0	< -7.5%	> -1%	LOW
Bees	<i>Bombus hum</i>	NA	1	> -1%	< -7.5%	HIGH
Bees	<i>Bombus mon</i>	Mountain B	0	> -1%	< -7.5%	HIGH
Bees	<i>Bombus mus</i>	Moss Carde	1	< -7.5%	> -1%	MODERATE
Bees	<i>Bombus rua</i>	NA	1	< -7.5%	> -1%	LOW
Bees	<i>Bombus syl</i>	NA	1	> -1%	> -1%	LOW
Bees	<i>Colletes h</i>	Sea-aster	1	< -7.5%	> -1%	LOW
Bees	<i>Halictus c</i>	NA	0	< -7.5%	> -1%	LOW
Bees	<i>Lasiogloss</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH
Bees	<i>Lasiogloss</i>	NA	0	< -7.5%	-4 to -1%	HIGH
Bees	<i>Lasiogloss</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH
Bees	<i>Melecta al</i>	NA	0	-4 to -1%	> -1%	LOW
Bees	<i>Melitta ha</i>	NA	0	-7.5 to -4%	> -1%	LOW
Bees	<i>Nomada fla</i>	NA	0	> -1%	> -1%	LOW
Bees	<i>Sphecodes</i>	NA	0	< -7.5%	> -1%	LOW
Bees	<i>Sphecodes</i>	NA	0	-7.5 to -4%	> -1%	LOW
Bees	<i>Sphecodes</i>	NA	0	-7.5 to -4%	> -1%	LOW
Birds	<i>Accipiter</i>	Goshawk	0	> -1%	-7.5 to -4%	MODERATE
Birds	<i>Alauda arv</i>	Skylark	1	-4 to -1%	> -1%	LOW
Birds	<i>Anas platy</i>	Mallard	0	> -1%	> -1%	LOW
Birds	<i>Anthus tri</i>	Tree Pipit	1	< -7.5%	-7.5 to -4%	VERY HIGH
Birds	<i>Asio flamm</i>	Short-eare	0	< -7.5%	-7.5 to -4%	VERY HIGH
Birds	<i>Botaurus s</i>	Bittern	1	< -7.5%	> -1%	LOW
Birds	<i>Branta can</i>	Canada Goo	0	> -1%	-7.5 to -4%	MODERATE
Birds	<i>Branta leu</i>	Barnacle G	0	< -7.5%	-7.5 to -4%	VERY HIGH
Birds	<i>Burhinus o</i>	Stone-curl	1	< -7.5%	-4 to -1%	VERY HIGH
Birds	<i>Buteo bute</i>	Buzzard	0	> -1%	-4 to -1%	LOW
Birds	<i>Caprimulgu</i>	Nightjar	1	< -7.5%	> -1%	LOW
Birds	<i>Carduelis</i>	Lesser Red	1	< -7.5%	-7.5 to -4%	VERY HIGH

Birds	<i>Carduelis</i> Linnet	1	> -1%	> -1%	LOW
Birds	<i>Carduelis</i> Twite	1	< -7.5%	-7.5 to -4%	VERY HIGH
Birds	<i>Certhia fa</i> Treecreeper	0	< -7.5%	-7.5 to -4%	MODERATE
Birds	<i>Cettia cet</i> Cetti's Warbler	0	> -1%	> -1%	LOW
Birds	<i>Cinclus ci</i> Dipper	0	-7.5 to -4%	-7.5 to -4%	HIGH
Birds	<i>Circus cya</i> Hen Harrier	1	< -7.5%	-4 to -1%	HIGH
Birds	<i>Coccothrau</i> Hawfinch	1	< -7.5%	> -1%	MODERATE
Birds	<i>Columba li</i> Rock Dove	0	> -1%	> -1%	LOW
Birds	<i>Columba oe</i> Stock Dove	0	> -1%	> -1%	LOW
Birds	<i>Corvus fru</i> Rook	0	> -1%	-4 to -1%	LOW
Birds	<i>Crex crex</i> Corncrake	1	< -7.5%	> -1%	LOW
Birds	<i>Cuculus ca</i> Cuckoo	1	< -7.5%	-4 to -1%	VERY HIGH
Birds	<i>Delichon u</i> House Martin	0	> -1%	> -1%	LOW
Birds	<i>Dendrocopo</i> Great Spot	0	> -1%	> -1%	LOW
Birds	<i>Dendrocopo</i> Lesser Spot	1	< -7.5%	> -1%	LOW
Birds	<i>Emberiza c</i> Corn Bunting	1	< -7.5%	> -1%	LOW
Birds	<i>Emberiza c</i> Cirl Bunting	1	-4 to -1%	> -1%	LOW
Birds	<i>Emberiza c</i> Yellowhammer	1	-7.5 to -4%	> -1%	MODERATE
Birds	<i>Emberiza s</i> Reed Bunting	1	> -1%	> -1%	LOW
Birds	<i>Falco colu</i> Merlin	0	> -1%	< -7.5%	MODERATE
Birds	<i>Falco pere</i> Peregrine	0	-7.5 to -4%	-7.5 to -4%	VERY HIGH
Birds	<i>Falco tinn</i> Kestrel	0	-4 to -1%	> -1%	LOW
Birds	<i>Fulica atr</i> Coot	0	-4 to -1%	> -1%	LOW
Birds	<i>Haematopus</i> Oystercatcher	0	-4 to -1%	> -1%	LOW
Birds	<i>Lagopus la</i> Red Grouse	1	< -7.5%	< -7.5%	VERY HIGH
Birds	<i>Lanius col</i> Red-backed	0	< -7.5%	> -1%	LOW
Birds	<i>Larus arge</i> Herring Gull	1	> -1%	> -1%	LOW
Birds	<i>Larus fusc</i> Lesser Black	0	< -7.5%	> -1%	LOW
Birds	<i>Limosa lim</i> Black-tail	1	> -1%	> -1%	LOW
Birds	<i>Locustella</i> Savi's Warbler	1	< -7.5%	> -1%	LOW
Birds	<i>Locustella</i> Grasshopper	1	< -7.5%	> -1%	LOW
Birds	<i>Lullula ar</i> Woodlark	1	< -7.5%	> -1%	LOW
Birds	<i>Morus bass</i> Gannet	0	> -1%	-4 to -1%	MODERATE
Birds	<i>Motacilla</i> Yellow Wagtail	1	< -7.5%	> -1%	LOW
Birds	<i>Muscicapa</i> Spotted Flycatcher	1	-7.5 to -4%	> -1%	MODERATE
Birds	<i>Numenius a</i> Curlew	1	< -7.5%	-7.5 to -4%	VERY HIGH
Birds	<i>Passer dom</i> House Sparrow	1	> -1%	> -1%	LOW
Birds	<i>Passer mon</i> Tree Sparrow	1	< -7.5%	> -1%	LOW
Birds	<i>Perdix per</i> Grey Partridge	1	< -7.5%	> -1%	HIGH
Birds	<i>Phalacroco</i> Cormorant	0	< -7.5%	> -1%	LOW
Birds	<i>Phoenicuru</i> Black Redstart	0	< -7.5%	> -1%	LOW
Birds	<i>Phoenicuru</i> Redstart	0	< -7.5%	> -1%	VERY HIGH
Birds	<i>Phylloscop</i> Wood Warbler	1	< -7.5%	-7.5 to -4%	VERY HIGH
Birds	<i>Picus viri</i> Green Woodpecker	0	> -1%	> -1%	LOW
Birds	<i>Podiceps c</i> Great Crested	0	-4 to -1%	> -1%	LOW
Birds	<i>Poecile mo</i> Willow Tit	1	-4 to -1%	> -1%	VERY HIGH

Birds	<i>Poecile pa</i> Marsh Tit	1	-7.5 to -4%	-7.5 to -4%	HIGH
Birds	<i>Porzana po</i> Spotted Cr	0	< -7.5%	> -1%	LOW
Birds	<i>Prunella m</i> Dunnock	1	> -1%	> -1%	LOW
Birds	<i>Puffinus p</i> Manx Shear	0	< -7.5%	> -1%	LOW
Birds	<i>Pyrrhula p</i> Bullfinch	1	> -1%	> -1%	LOW
Birds	<i>Rallus aqu</i> Water Rail	0	< -7.5%	> -1%	LOW
Birds	<i>Saxicola r</i> Whinchat	0	< -7.5%	-7.5 to -4%	VERY HIGH
Birds	<i>Saxicola t</i> Stonechat	0	< -7.5%	-7.5 to -4%	LOW
Birds	<i>Sitta euro</i> Nuthatch	0	> -1%	> -1%	LOW
Birds	<i>Sterna dou</i> Roseate Te	1	< -7.5%	> -1%	LOW
Birds	<i>Streptopel</i> Collared D	0	> -1%	> -1%	LOW
Birds	<i>Streptopel</i> Turtle Dov	1	< -7.5%	> -1%	LOW
Birds	<i>Sturnus vu</i> Starling	1	> -1%	> -1%	LOW
Birds	<i>Sylvia bor</i> Garden War	0	-7.5 to -4%	-4 to -1%	VERY HIGH
Birds	<i>Sylvia com</i> Whitethroa	0	> -1%	> -1%	LOW
Birds	<i>Tachybaptu</i> Little Gre	0	> -1%	> -1%	LOW
Birds	<i>Tetrao tet</i> Black Grou	1	> -1%	< -7.5%	MODERATE
Birds	<i>Tringa tot</i> Redshank	0	< -7.5%	> -1%	LOW
Birds	<i>Troglodyte</i> Wren	0	> -1%	> -1%	LOW
Birds	<i>Turdus mer</i> Blackbird	0	> -1%	> -1%	LOW
Birds	<i>Turdus phi</i> Song Thrus	1	> -1%	> -1%	LOW
Birds	<i>Turdus tor</i> Ring Ouzel	1	< -7.5%	< -7.5%	VERY HIGH
Birds	<i>Tyto alba</i> Barn Owl	0	> -1%	> -1%	LOW
Birds	<i>Vanellus v</i> Lapwing	1	-7.5 to -4%	> -1%	HIGH
Bryophytes	<i>Calypogeia</i> Bog Pouchw	0	< -7.5%	-4 to -1%	HIGH
Bryophytes	<i>Dicranum s</i> Rusty Fork	1	< -7.5%	> -1%	LOW
Bryophytes	<i>Didymodon</i> Cylindric	0	> -1%	> -1%	LOW
Bryophytes	<i>Didymodon</i> Brown Bear	0	> -1%	-7.5 to -4%	HIGH
Bryophytes	<i>Leucobryum</i> Large Whit	0	< -7.5%	-7.5 to -4%	VERY HIGH
Bryophytes	<i>Plagiochil</i> Western Fe	0	> -1%	< -7.5%	HIGH
Bryophytes	<i>Plagiomniu</i> Many-fruit	0	< -7.5%	> -1%	LOW
Bryophytes	<i>Plagiothec</i> Bright Sil	0	-4 to -1%	< -7.5%	VERY HIGH
Bryophytes	<i>Porella pl</i> Wall Scale	0	< -7.5%	< -7.5%	VERY HIGH
Bryophytes	<i>Racomitriu</i> Slender Fr	0	> -1%	< -7.5%	HIGH
Bryophytes	<i>Radula aqu</i> Brown Scal	0	-7.5 to -4%	< -7.5%	VERY HIGH
Bryophytes	<i>Rhytidiade</i> Little Sha	0	< -7.5%	-7.5 to -4%	VERY HIGH
Bryophytes	<i>Sphagnum p</i> Golden Bog	0	< -7.5%	< -7.5%	VERY HIGH
Bryophytes	<i>Zygodon vi</i> NA	0	> -1%	-7.5 to -4%	HIGH
Carbid be	<i>Amara ovat</i> NA	0	< -7.5%	> -1%	LOW
Carbid be	<i>Amara tibi</i> NA	0	< -7.5%	> -1%	LOW
Carbid be	<i>Anatis oce</i> Eyed Ladyb	0	< -7.5%	> -1%	LOW
Carbid be	<i>Anthracus</i> NA	0	-4 to -1%	> -1%	LOW
Carbid be	<i>Bembidion</i> NA	0	> -1%	> -1%	LOW
Carbid be	<i>Bembidion</i> NA	0	> -1%	> -1%	LOW
Carbid be	<i>Bradycellu</i> NA	0	< -7.5%	> -1%	LOW
Carbid be	<i>Calathus m</i> NA	0	< -7.5%	> -1%	LOW

Carbid be: <i>Calosoma i</i> NA	1	> -1%	-7.5 to -4%	HIGH
Carbid be: <i>Carabus mo</i> Necklace G	1	< -7.5%	> -1%	LOW
Carbid be: <i>Carabus pr</i> NA	0	-4 to -1%	-4 to -1%	HIGH
Carbid be: <i>Curtonotus</i> NA	0	> -1%	> -1%	LOW
Carbid be: <i>Laemostenu</i> NA	0	> -1%	> -1%	LOW
Carbid be: <i>Ocys harpa</i> NA	0	> -1%	> -1%	LOW
Carbid be: <i>Philorhizu</i> NA	1	< -7.5%	> -1%	LOW
Carbid be: <i>Pterostich</i> NA	0	> -1%	< -7.5%	HIGH
Carbid be: <i>Syntomus f</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Cryptops h</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Geophilus</i> NA	0	< -7.5%	> -1%	MODERATE
Centipede: <i>Geophilus</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Geophilus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Centipede: <i>Geophilus</i> NA	0	< -7.5%	> -1%	MODERATE
Centipede: <i>Henia vesu</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Lithobius</i> NA	0	< -7.5%	-4 to -1%	HIGH
Centipede: <i>Lithobius</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Lithobius</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Lithobius</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Schendyla</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Stigmatoga</i> NA	0	> -1%	> -1%	LOW
Centipede: <i>Strigamia</i> NA	0	< -7.5%	> -1%	LOW
Coccinelid <i>Adalia bip</i> Two-spot L	0	< -7.5%	> -1%	LOW
Coccinelid <i>Adalia dec</i> Ten-spot L	0	> -1%	> -1%	LOW
Coccinelid <i>Anisostict</i> Water Lady	0	< -7.5%	> -1%	LOW
Coccinelid <i>Coccidula</i> NA	0	< -7.5%	-4 to -1%	HIGH
Coccinelid <i>Coccinella</i> Seven-spot	0	-4 to -1%	> -1%	LOW
Coccinelid <i>Coccinella</i> Eleven-spo	0	< -7.5%	> -1%	LOW
Coccinelid <i>Halyzia se</i> Orange Lad	0	< -7.5%	-4 to -1%	HIGH
Coccinelid <i>Hippodamia</i> Adonis' La	0	< -7.5%	> -1%	LOW
Coccinelid <i>Propylea q</i> Fourteen-s	0	< -7.5%	> -1%	LOW
Coccinelid <i>Psyllobora</i> Twentytwo-	0	< -7.5%	> -1%	LOW
Coccinelid <i>Scymnus su</i> NA	0	< -7.5%	> -1%	LOW
Coccinelid <i>Subcoccine</i> Twentyfour	0	> -1%	-4 to -1%	MODERATE
Craneflies <i>Nephrotoma</i> NA	0	-4 to -1%	> -1%	LOW
Craneflies <i>Ptychopter</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Craneflies <i>Ptychopter</i> NA	0	< -7.5%	> -1%	LOW
Craneflies <i>Ptychopter</i> NA	0	< -7.5%	> -1%	LOW
Craneflies <i>Tipula ful</i> NA	0	< -7.5%	> -1%	LOW
Craneflies <i>Tipula lat</i> NA	0	> -1%	-7.5 to -4%	HIGH
Craneflies <i>Tipula lun</i> NA	0	< -7.5%	> -1%	LOW
Craneflies <i>Tipula max</i> NA	0	> -1%	> -1%	LOW
Craneflies <i>Tipula ole</i> NA	0	< -7.5%	> -1%	LOW
Craneflies <i>Tipula unc</i> NA	0	> -1%	-7.5 to -4%	HIGH
Craneflies <i>Tipula var</i> NA	0	> -1%	-4 to -1%	MODERATE

Crickets : <i>Chorthippu</i> NA	0	-7.5 to -4%	> -1%	LOW
Crickets : <i>Conocephal</i> NA	0	> -1%	-4 to -1%	MODERATE
Crickets : <i>Conocephal</i> NA	0	> -1%	> -1%	LOW
Crickets : <i>Ectobius p</i> Tawny Cock	0	< -7.5%	< -7.5%	VERY HIGH
Crickets : <i>Ectobius p</i> Lesser Coc	0	< -7.5%	-7.5 to -4%	VERY HIGH
Crickets : <i>Forficula</i> Common Ear	0	> -1%	> -1%	LOW
Crickets : <i>Forficula</i> Lesne's Ea	0	> -1%	> -1%	LOW
Crickets : <i>Meconema t</i> NA	0	> -1%	-4 to -1%	MODERATE
Crickets : <i>Omocestus</i> Woodland G	0	> -1%	< -7.5%	HIGH
Crickets : <i>Omocestus</i> Common Gre	0	< -7.5%	-7.5 to -4%	VERY HIGH
Crickets : <i>Platycleis</i> NA	0	< -7.5%	> -1%	LOW
Crickets : <i>Tetrix cep</i> Cepero's G	0	< -7.5%	> -1%	LOW
Crickets : <i>Tetrix sub</i> NA	0	> -1%	> -1%	LOW
Hoverflies <i>Anasimyia</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Hoverflies <i>Cheilosia</i> NA	0	< -7.5%	> -1%	LOW
Hoverflies <i>Cheilosia</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Hoverflies <i>Cheilosia</i> NA	0	> -1%	> -1%	LOW
Hoverflies <i>Cheilosia</i> NA	0	-4 to -1%	< -7.5%	HIGH
Hoverflies <i>Eristalis</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Hoverflies <i>Eupeodes b</i> NA	0	< -7.5%	> -1%	MODERATE
Hoverflies <i>Eupeodes n</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Hoverflies <i>Neoascia m</i> NA	0	< -7.5%	> -1%	LOW
Hoverflies <i>Platycheir</i> NA	0	> -1%	-7.5 to -4%	HIGH
Hoverflies <i>Sphaeropho</i> NA	0	> -1%	> -1%	LOW
Hoverflies <i>Sphegina e</i> NA	0	< -7.5%	> -1%	MODERATE
Hoverflies <i>Xylota seg</i> NA	0	-4 to -1%	> -1%	LOW
Millipede: <i>Blaniulus</i> Spotted Sn	0	> -1%	> -1%	LOW
Millipede: <i>Brachydesm</i> NA	0	> -1%	> -1%	LOW
Millipede: <i>Chordeuma</i> NA	0	> -1%	< -7.5%	HIGH
Millipede: <i>Cylindroi</i> Blunt-tail	0	-4 to -1%	> -1%	LOW
Millipede: <i>Glomeris m</i> Pill Milli	0	< -7.5%	< -7.5%	VERY HIGH
Millipede: <i>Melogona s</i> NA	0	> -1%	-7.5 to -4%	HIGH
Millipede: <i>Nanogona p</i> Eyed Flat-	0	> -1%	> -1%	LOW
Millipede: <i>Nemasoma v</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Millipede: <i>Ommatoiulu</i> Striped Mi	0	> -1%	-7.5 to -4%	HIGH
Millipede: <i>Ophiulus</i> NA	0	> -1%	-7.5 to -4%	HIGH
Millipede: <i>Polydesmus</i> Common Fla	0	> -1%	> -1%	LOW
Millipede: <i>Polydesmus</i> NA	0	> -1%	-4 to -1%	MODERATE
Millipede: <i>Tachypodoi</i> White-legg	0	-4 to -1%	< -7.5%	HIGH
Moths <i>Acronicta</i> Knot Grass	1	< -7.5%	> -1%	LOW
Moths <i>Adscita st</i> The Forest	1	< -7.5%	< -7.5%	VERY HIGH
Moths <i>Agrochola</i> Flounced C.	1	< -7.5%	> -1%	LOW
Moths <i>Agrochola</i> Brown-spot	1	< -7.5%	-7.5 to -4%	VERY HIGH
Moths <i>Agrochola</i> Beaded Che	1	< -7.5%	> -1%	LOW
Moths <i>Alcis juba</i> Dotted Car	0	> -1%	< -7.5%	HIGH

Moths	<i>Aleucis di</i> Sloe Carpe	1	< -7.5%	-4 to -1%	HIGH
Moths	<i>Allophyes</i> Green-brin	1	< -7.5%	-7.5 to -4%	VERY HIGH
Moths	<i>Alsophila</i> March Moth	0	< -7.5%	> -1%	LOW
Moths	<i>Apamea anc</i> Large Nutm	1	< -7.5%	> -1%	LOW
Moths	<i>Arctia caj</i> Garden Tig	1	< -7.5%	> -1%	LOW
Moths	<i>Atethmia c</i> Centre-bar	1	-4 to -1%	> -1%	LOW
Moths	<i>Blepharita</i> Dark Broca	1	< -7.5%	> -1%	LOW
Moths	<i>Cabera exa</i> Common Wav	0	> -1%	-4 to -1%	MODERATE
Moths	<i>Caradrina</i> Mottled Ru	1	< -7.5%	> -1%	LOW
Moths	<i>Celaena ha</i> Haworth's	1	< -7.5%	-7.5 to -4%	VERY HIGH
Moths	<i>Chesias ru</i> Broom-tip	1	< -7.5%	> -1%	LOW
Moths	<i>Colotois p</i> Feathered	0	< -7.5%	> -1%	LOW
Moths	<i>Cosmia aff</i> Lesser-spo	0	< -7.5%	> -1%	LOW
Moths	<i>Cossus cos</i> Goat Moth	1	< -7.5%	> -1%	LOW
Moths	<i>Cyclophora</i> Dingy Moch	1	> -1%	> -1%	LOW
Moths	<i>Cyclophora</i> False Moch	1	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Cymatophor</i> Oak Lutest	1	< -7.5%	> -1%	LOW
Moths	<i>Dasypolia</i> Brindled O	1	< -7.5%	-4 to -1%	HIGH
Moths	<i>Diloba cae</i> Figure of	1	< -7.5%	> -1%	LOW
Moths	<i>Eilema sor</i> Orange Foo	0	> -1%	< -7.5%	HIGH
Moths	<i>Ennomos er</i> September	1	< -7.5%	> -1%	LOW
Moths	<i>Ennomos qu</i> August Tho	1	< -7.5%	> -1%	LOW
Moths	<i>Entephria</i> Grey Mount	1	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Eugnorisma</i> Autumnal R	1	< -7.5%	> -1%	LOW
Moths	<i>Eulithis m</i> The Spinac	1	< -7.5%	> -1%	LOW
Moths	<i>Euxoa trit</i> White-line	1	< -7.5%	> -1%	LOW
Moths	<i>Graphiphor</i> Double Dar	1	< -7.5%	> -1%	MODERATE
Moths	<i>Hadena alb</i> White Spot	1	< -7.5%	> -1%	LOW
Moths	<i>Heliophobu</i> Bordered G	1	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Hemistola</i> Small Emer	1	-7.5 to -4%	> -1%	LOW
Moths	<i>Hoplodrina</i> The Rustic	1	> -1%	> -1%	LOW
Moths	<i>Idaea muri</i> Purple-bor	0	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Jodis lact</i> Little Eme	0	< -7.5%	> -1%	LOW
Moths	<i>Lycia hirt</i> Brindled B	1	< -7.5%	> -1%	MODERATE
Moths	<i>Macaria wa</i> The V-Moth	1	< -7.5%	> -1%	MODERATE
Moths	<i>Malacosoma</i> The Lackey	1	< -7.5%	> -1%	LOW
Moths	<i>Melanchra</i> Dot Moth	1	< -7.5%	> -1%	LOW
Moths	<i>Melanthia</i> Pretty Cha	1	< -7.5%	-7.5 to -4%	VERY HIGH
Moths	<i>Mythimna c</i> Shoulder-s	1	< -7.5%	> -1%	LOW
Moths	<i>Operophter</i> Northern W	0	< -7.5%	> -1%	MODERATE
Moths	<i>Oria muscu</i> Brighton W	1	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Orthosia g</i> Powdered Q	1	< -7.5%	> -1%	LOW
Moths	<i>Paracolax</i> Clay Fan-f	1	< -7.5%	> -1%	LOW
Moths	<i>Pelurga co</i> Dark Spina	1	< -7.5%	> -1%	LOW
Moths	<i>Photedes c</i> Least Mino	0	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Polia bomb</i> Pale Shini	1	< -7.5%	-7.5 to -4%	VERY HIGH
Moths	<i>Rheumapter</i> Argent & S	1	< -7.5%	-4 to -1%	HIGH
Moths	<i>Rhizedra l</i> Large Wain	1	-7.5 to -4%	> -1%	LOW
Moths	<i>Scotoptery</i> Chalk Carp	1	< -7.5%	< -7.5%	VERY HIGH

Moths	<i>Selidosema</i>	Bordered G	0	< -7.5%	> -1%	LOW
Moths	<i>Shargacucu</i>	Striped Ly	1	< -7.5%	> -1%	LOW
Moths	<i>Spilosoma</i>	Buff Ermin	1	> -1%	> -1%	LOW
Moths	<i>Stilbia an</i>	The Anomal	1	< -7.5%	-4 to -1%	HIGH
Moths	<i>Tholera ce</i>	Hedge Rust	1	< -7.5%	> -1%	LOW
Moths	<i>Tholera de</i>	Feathered	1	< -7.5%	> -1%	LOW
Moths	<i>Trichiura</i>	Pale Eggar	1	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Trichopter</i>	Early Toot	0	> -1%	-4 to -1%	MODERATE
Moths	<i>Trichopter</i>	Barred Too	1	< -7.5%	> -1%	MODERATE
Moths	<i>Trisateles</i>	Olive Cres	1	< -7.5%	> -1%	LOW
Moths	<i>Tyta luctu</i>	The Four-s	1	< -7.5%	> -1%	LOW
Moths	<i>Watsonalla</i>	Barred Hoo	0	< -7.5%	< -7.5%	VERY HIGH
Moths	<i>Xanthia gi</i>	Dusky-lemo	1	< -7.5%	-7.5 to -4%	VERY HIGH
Moths	<i>Xanthia ic</i>	The Sallow	1	< -7.5%	> -1%	LOW
Moths	<i>Xanthorhoe</i>	Red Carpet	1	< -7.5%	-7.5 to -4%	VERY HIGH
Moths	<i>Xestia aga</i>	Heath Rust	1	< -7.5%	> -1%	MODERATE
Odonata	<i>Aeshna cae</i>	Azure hawk	0	> -1%	< -7.5%	HIGH
Odonata	<i>Aeshna gra</i>	Brown hawk	0	< -7.5%	-4 to -1%	HIGH
Odonata	<i>Aeshna jun</i>	Common hawk	0	< -7.5%	> -1%	LOW
Odonata	<i>Anax imper</i>	Emperor dr	0	> -1%	> -1%	LOW
Odonata	<i>Ceriagrion</i>	Small red	0	< -7.5%	< -7.5%	VERY HIGH
Odonata	<i>Enallagma</i>	Common blu	0	-7.5 to -4%	> -1%	MODERATE
Odonata	<i>Erythromma</i>	Red-eyed d	0	> -1%	< -7.5%	HIGH
Odonata	<i>Ischnura e</i>	Blue-taile	0	-7.5 to -4%	> -1%	LOW
Odonata	<i>Orthetrum</i>	Black-tail	0	> -1%	> -1%	LOW
Odonata	<i>Orthetrum</i>	Keeled ski	0	-7.5 to -4%	< -7.5%	VERY HIGH
Odonata	<i>Pyrrhosoma</i>	Large red	0	< -7.5%	> -1%	LOW
Odonata	<i>Sympetrum</i>	Ruddy dart	0	> -1%	> -1%	LOW
Odonata	<i>Sympetrum</i>	Common dar	0	-7.5 to -4%	> -1%	LOW
Soldier b	<i>Cantharis</i>	NA	0	-4 to -1%	-7.5 to -4%	VERY HIGH
Soldier b	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH
Soldier b	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	LOW
Soldier b	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	LOW
Soldier b	<i>Cantharis</i>	NA	0	-7.5 to -4%	-4 to -1%	HIGH
Soldier b	<i>Malthinus</i>	NA	0	> -1%	-4 to -1%	MODERATE
Soldier b	<i>Malthinus</i>	NA	0	> -1%	> -1%	LOW
Soldier b	<i>Podabrus a</i>	NA	0	> -1%	< -7.5%	HIGH
Soldier b	<i>Rhagonycha</i>	Common Red	0	-7.5 to -4%	> -1%	LOW
Soldier b	<i>Rhagonycha</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH
Soldier b	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	LOW
Soldier b	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	LOW
Soldier b	<i>Rhagonycha</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH
Spiders	<i>Anelosimus</i>	NA	0	> -1%	> -1%	LOW
Spiders	<i>Araneus ma</i>	NA	0	> -1%	< -7.5%	HIGH
Spiders	<i>Araneus qu</i>	NA	0	> -1%	> -1%	LOW
Spiders	<i>Bathypphant</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH

Spiders	<i>Ceratinella</i> NA	0	< -7.5%	< -7.5%	VERY HIGH
Spiders	<i>Clubiona</i> n NA	0	< -7.5%	> -1%	LOW
Spiders	<i>Dictyna</i> pu Small Mesh	1	< -7.5%	-7.5 to -4%	VERY HIGH
Spiders	<i>Diplocephala</i> NA	0	< -7.5%	> -1%	LOW
Spiders	<i>Haplodrass</i> Heath Gras	1	< -7.5%	> -1%	LOW
Spiders	<i>Mecopisthes</i> Peus' s Lon	1	< -7.5%	> -1%	LOW
Spiders	<i>Meioneta</i> m Thin Weble	1	< -7.5%	> -1%	LOW
Spiders	<i>Monocephala</i> Broad Groo	1	< -7.5%	-4 to -1%	HIGH
Spiders	<i>Porrhomma</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH
Spiders	<i>Porrhomma</i> NA	0	< -7.5%	> -1%	MODERATE
Spiders	<i>Saaristoia</i> Triangle H	1	< -7.5%	-7.5 to -4%	VERY HIGH
Spiders	<i>Salticus</i> s NA	0	> -1%	> -1%	LOW
Spiders	<i>Sitticus</i> c Sedge Jump	1	< -7.5%	> -1%	LOW
Spiders	<i>Tapinocyba</i> NA	0	-4 to -1%	< -7.5%	VERY HIGH
Spiders	<i>Trichoptera</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH
Spiders	<i>Walckenaera</i> NA	0	> -1%	< -7.5%	HIGH
Vascular	<i>Acera</i> ant Man Orchid	0	> -1%	< -7.5%	HIGH
Vascular	<i>Ajuga</i> pyra Pyramidal	1	> -1%	-7.5 to -4%	HIGH
Vascular	<i>Blysmus</i> co Flat-sedge	1	< -7.5%	-7.5 to -4%	VERY HIGH
Vascular	<i>Bupleurum</i> Slender Ha	1	< -7.5%	> -1%	LOW
Vascular	<i>Calamagrostis</i> Purple Sma	0	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Calamagrostis</i> Narrow Sma	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Calystegia</i> Great Bind	0	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Carex</i> eric Rare Sprin	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Carex</i> viri Common Yel	0	> -1%	> -1%	LOW
Vascular	<i>Centaurea</i> Cornflower	1	< -7.5%	-7.5 to -4%	VERY HIGH
Vascular	<i>Cephalanthus</i> White Hell	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Chamaemelum</i> Chamomile	1	< -7.5%	> -1%	LOW
Vascular	<i>Chenopodium</i> Fig-leaved	0	> -1%	> -1%	LOW
Vascular	<i>Chenopodium</i> Stinking G	1	> -1%	> -1%	LOW
Vascular	<i>Cicendia</i> f Yellow Cen	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Dactylorhiza</i> Common Spo	0	< -7.5%	> -1%	LOW
Vascular	<i>Dactylorhiza</i> Narrow-lea	0	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Dianthus</i> a Deptford P	1	< -7.5%	> -1%	LOW
Vascular	<i>Euphrasia</i> Chalk Eyeb	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Euphrasia</i> Cornish Ey	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Fumaria</i> pu Purple Ram	1	-7.5 to -4%	> -1%	LOW
Vascular	<i>Galeopsis</i> Red Hemp-n	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Herminium</i> Musk Orchi	1	> -1%	< -7.5%	HIGH
Vascular	<i>Illecebrum</i> Coral-neck	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Luronium</i> n Floating W	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Melittis</i> m Bastard Ba	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Mentha</i> pul Pennyroyal	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Minuartia</i> Fine-leave	1	> -1%	-7.5 to -4%	HIGH
Vascular	<i>Muscari</i> ne Grape-hyac	1	> -1%	> -1%	LOW
Vascular	<i>Najas</i> flex Slender Na	1	> -1%	< -7.5%	HIGH
Vascular	<i>Phyllitis</i> Hart' s-ton	0	< -7.5%	> -1%	LOW

Vascular	<i>Pilularia</i>	Pillwort	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Potamogeton</i>	Grass-wrack	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Pulsatilla</i>	Pasqueflower	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Quercus</i>	rope	0	-4 to -1%	> -1%	LOW
Vascular	<i>Ranunculus</i>	Corn Buttercup	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Ranunculus</i>	Three-lobed	1	> -1%	> -1%	LOW
Vascular	<i>Rubus</i>	saxatilis	0	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Rumex</i>	rupestris	1	> -1%	> -1%	LOW
Vascular	<i>Salix</i>	lappacea	1	> -1%	< -7.5%	HIGH
Vascular	<i>Scandix</i>	peucedana	1	> -1%	-7.5 to -4%	HIGH
Vascular	<i>Scrophularia</i>	Water Figwort	0	> -1%	> -1%	LOW
Vascular	<i>Silene</i>	gallica	1	< -7.5%	> -1%	LOW
Vascular	<i>Sium</i>	latifolium	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Sparganium</i>	Branched Bur-reed	0	< -7.5%	> -1%	LOW
Vascular	<i>Spartina</i>	maritima	1	< -7.5%	> -1%	LOW
Vascular	<i>Stellaria</i>	Marshall's	1	< -7.5%	> -1%	LOW
Vascular	<i>Torilis</i>	arvensis	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Valeriana</i>	lanceolata	1	< -7.5%	< -7.5%	VERY HIGH
Vascular	<i>Veronica</i>	campanulata	0	-7.5 to -4%	> -1%	MODERATE
Vascular	<i>Viola</i>	lactuca	1	> -1%	-4 to -1%	MODERATE
Wasps	<i>Agenioidea</i>	NA	0	< -7.5%	> -1%	LOW
Wasps	<i>Ancistrocheilus</i>	Wall Mason	0	< -7.5%	> -1%	LOW
Wasps	<i>Cerceris</i>	ornata	0	> -1%	> -1%	LOW
Wasps	<i>Crabro</i>	pelus	0	< -7.5%	> -1%	LOW
Wasps	<i>Crossocerus</i>	NA	0	< -7.5%	> -1%	LOW
Wasps	<i>Entomognathus</i>	NA	0	< -7.5%	> -1%	LOW
Wasps	<i>Hedychridi</i>	NA	0	> -1%	> -1%	LOW
Wasps	<i>Nysson</i>	spilargus	0	< -7.5%	< -7.5%	VERY HIGH
Wasps	<i>Odynerus</i>	maritimus	1	< -7.5%	> -1%	LOW
Wasps	<i>Oxybelus</i>	argenteus	0	< -7.5%	> -1%	LOW
Wasps	<i>Pemphredon</i>	mournful	0	< -7.5%	> -1%	LOW
Wasps	<i>Priocnemis</i>	NA	0	< -7.5%	> -1%	LOW
Wasps	<i>Priocnemis</i>	NA	0	> -1%	> -1%	LOW
Wasps	<i>Smicromyrme</i>	Small Velvet	0	> -1%	> -1%	LOW

Associated confidence	Observed expansion	Projected expansion	Benefit from expansion	Associated confidence	Final outcome
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
MEDIUM	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	+1 to +4%	HIGH	POOR	Risks & benefits
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	+1 to +4%	HIGH	POOR	High benefit
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
MEDIUM	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	+1 to +4%	+4 to +7.5%	MODERATE	MEDIUM	Medium benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	HIGH	POOR	Risks & benefits
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	Medium benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	< +1%	> +7.5%	MODERATE	POOR	Medium benefit
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Medium benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	MODERATE	POOR	Medium benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	+1 to +4%	HIGH	MEDIUM	Medium benefit
MEDIUM	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
MEDIUM	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	Medium risk
MEDIUM	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	Risks & benefits
MEDIUM	> +7.5%	> +7.5%	HIGH	POOR	Medium risk
POOR	> +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk

GOOD	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	< +1%	LOW	POOR	Medium risk
GOOD	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
MEDIUM	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
MEDIUM	> +7.5%	< +1%	LOW	POOR	Limited impact
MEDIUM	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
MEDIUM	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
GOOD	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	+1 to +4%	HIGH	MEDIUM	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Medium risk
MEDIUM	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	< +1%	LOW	POOR	Medium risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	+1 to +4%	< +1%	LOW	POOR	Limited impact
GOOD	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
GOOD	> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
GOOD	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	+4 to +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	Medium risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
GOOD	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
MEDIUM	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	< +1%	MODERATE	POOR	High risk

POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
GOOD	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
MEDIUM	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
MEDIUM	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	> +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
MEDIUM	> +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	< +1%	< +1%	LOW	MEDIUM	Limited impact
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
GOOD	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	> +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	< +1%	LOW	POOR	Medium risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
GOOD	< +1%	< +1%	LOW	MEDIUM	Limited impact
MEDIUM	< +1%	< +1%	LOW	MEDIUM	Limited impact
MEDIUM	+1 to +4%	< +1%	LOW	POOR	Limited impact
GOOD	+1 to +4%	< +1%	LOW	POOR	High risk
MEDIUM	> +7.5%	+1 to +4%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	+1 to +4%	MODERATE	POOR	Medium risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
POOR	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	+4 to +7.5%	MODERATE	POOR	Medium benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit

POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	> +7.5%	MODERATE	POOR	Medium benefit
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
MEDIUM	+1 to +4%	+1 to +4%	MODERATE	MEDIUM	Medium benefit
MEDIUM	< +1%	+4 to +7.5%	LOW	MEDIUM	Limited impact
POOR	< +1%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
MEDIUM	+4 to +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	MEDIUM	Medium risk
MEDIUM	+4 to +7.5%	+4 to +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Medium risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	+1 to +4%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
MEDIUM	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	MODERATE	POOR	Medium benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	MODERATE	POOR	Medium risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	+4 to +7.5%	MODERATE	MEDIUM	Medium benefit
POOR	> +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	+4 to +7.5%	HIGH	POOR	Medium benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	MEDIUM	High risk
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	+4 to +7.5%	HIGH	POOR	Risks & benefits
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	MEDIUM	High benefit
POOR	< +1%	< +1%	LOW	MEDIUM	Limited impact
POOR	+1 to +4%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	MEDIUM	Medium risk

POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	Medium risk
MEDIUM	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	Medium risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	MEDIUM	High risk
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	+4 to +7.5%	HIGH	POOR	Medium risk
MEDIUM	+4 to +7.5%	+4 to +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	POOR	Medium benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	+1 to +4%	< +1%	LOW	MEDIUM	Medium risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
MEDIUM	> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	< +1%	< +1%	LOW	MEDIUM	Limited impact
POOR	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	HIGH	POOR	Risks & benefits
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	< +1%	MODERATE	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	MEDIUM	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	POOR	Medium benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	< +1%	LOW	MEDIUM	High risk

POOR	> +7.5%	> +7.5%	HIGH	POOR	Risks & benefits
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	+1 to +4%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
POOR	< +1%	< +1%	LOW	MEDIUM	Medium risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
MEDIUM	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	MEDIUM	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Medium risk
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
MEDIUM	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	MODERATE	POOR	Medium benefit
MEDIUM	> +7.5%	+1 to +4%	HIGH	POOR	Medium risk
POOR	+1 to +4%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	> +7.5%	< +1%	LOW	POOR	Medium risk
POOR	+1 to +4%	< +1%	LOW	MEDIUM	Medium risk
POOR	< +1%	< +1%	LOW	MEDIUM	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	MEDIUM	Medium risk
POOR	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	< +1%	> +7.5%	MODERATE	POOR	Medium benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	< +1%	> +7.5%	MODERATE	POOR	High risk
MEDIUM	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	HIGH	POOR	Risks & benefits
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
MEDIUM	+1 to +4%	< +1%	LOW	POOR	High risk

POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+1 to +4%	< +1%	LOW	MEDIUM	High risk
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	Medium risk
POOR	+1 to +4%	> +7.5%	HIGH	POOR	Medium benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	< +1%	MODERATE	POOR	High risk
POOR	> +7.5%	< +1%	LOW	POOR	Medium risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	Limited impact
POOR	> +7.5%	+4 to +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	< +1%	< +1%	LOW	MEDIUM	Medium risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	+4 to +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	+1 to +4%	HIGH	POOR	Risks & benefits
POOR	+4 to +7.5%	< +1%	LOW	POOR	Medium risk
MEDIUM	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	+4 to +7.5%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	+1 to +4%	MODERATE	POOR	Medium benefit
POOR	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk

POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
MEDIUM	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	+1 to +4%	> +7.5%	MODERATE	POOR	Medium benefit
POOR	< +1%	> +7.5%	MODERATE	POOR	Medium benefit
POOR	< +1%	> +7.5%	MODERATE	POOR	Medium benefit
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
MEDIUM	+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	< +1%	> +7.5%	MODERATE	POOR	Risks & benefits
MEDIUM	< +1%	< +1%	LOW	MEDIUM	High risk
MEDIUM	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	< +1%	> +7.5%	MODERATE	POOR	Medium benefit
POOR	> +7.5%	+1 to +4%	HIGH	POOR	Medium risk
MEDIUM	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	Risks & benefits
POOR	> +7.5%	+4 to +7.5%	VERY HIGH	POOR	Medium benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	+4 to +7.5%	HIGH	POOR	Medium risk
POOR	+4 to +7.5%	> +7.5%	HIGH	POOR	Medium risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	< +1%	< +1%	LOW	MEDIUM	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
MEDIUM	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	+4 to +7.5%	VERY HIGH	POOR	Risks & benefits
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	+1 to +4%	MODERATE	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	+4 to +7.5%	VERY HIGH	POOR	Risks & benefits
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
POOR	> +7.5%	+1 to +4%	HIGH	POOR	Risks & benefits
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Medium benefit
POOR	> +7.5%	+1 to +4%	HIGH	POOR	High benefit

POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	< +1%	LOW	POOR	High risk
POOR	+4 to +7.5%	< +1%	LOW	POOR	Limited impact
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	+1 to +4%	> +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	Medium benefit
POOR	+1 to +4%	< +1%	LOW	POOR	Limited impact
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	< +1%	LOW	POOR	High risk
POOR	> +7.5%	< +1%	LOW	MEDIUM	Limited impact
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	+1 to +4%	HIGH	POOR	Medium risk
POOR	> +7.5%	+4 to +7.5%	HIGH	POOR	Medium risk
POOR	< +1%	< +1%	LOW	MEDIUM	Medium risk
POOR	+1 to +4%	> +7.5%	HIGH	POOR	Medium benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	+4 to +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	HIGH	POOR	High benefit
POOR	+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
POOR	> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit

Group	Latin Name	English name	NERC species	Observed decline	Projected decline	Risk of decline	Associated confidence
Ants	<i>Formica cu</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Ants	<i>Formica fu</i>	Negro Ant	0	> -1%	> -1%	LOW	POOR
Ants	<i>Formica sa</i>	NA	0	-4 to -1%	< -7.5%	VERY HIGH	POOR
Ants	<i>Lasius ali</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Ants	<i>Lasius fla</i>	Yellow Mea	0	< -7.5%	> -1%	LOW	POOR
Ants	<i>Lasius mix</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Ants	<i>Lasius nig</i>	Small Blac	0	> -1%	> -1%	LOW	POOR
Ants	<i>Leptothora</i>	Slender An	0	> -1%	-4 to -1%	MODERATE	POOR
Ants	<i>Myrmica ru</i>	Red Ant	0	< -7.5%	> -1%	LOW	POOR
Ants	<i>Myrmica ru</i>	NA	0	> -1%	> -1%	LOW	POOR
Ants	<i>Myrmica sa</i>	NA	0	> -1%	> -1%	LOW	POOR
Ants	<i>Myrmica sc</i>	NA	0	-7.5 to -4%	> -1%	LOW	POOR
Ants	<i>Myrmica sc</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Bees	<i>Andrena ha</i>	Early Mini	0	-4 to -1%	> -1%	LOW	POOR
Bees	<i>Andrena ta</i>	Tormentil	1	< -7.5%	> -1%	LOW	POOR
Bees	<i>Andrena va</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Bees	<i>Bombus hum</i>	NA	1	> -1%	< -7.5%	HIGH	POOR
Bees	<i>Bombus mon</i>	Mountain B	0	> -1%	< -7.5%	HIGH	POOR
Bees	<i>Bombus mus</i>	Moss Carde	1	< -7.5%	> -1%	MODERATE	POOR
Bees	<i>Bombus rua</i>	NA	1	< -7.5%	> -1%	MODERATE	POOR
Bees	<i>Bombus syl</i>	NA	1	> -1%	> -1%	LOW	MEDIUM
Bees	<i>Colletes h</i>	Sea-aster	1	< -7.5%	> -1%	LOW	POOR
Bees	<i>Halictus c</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Bees	<i>Lasiogloss</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Bees	<i>Lasiogloss</i>	NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Bees	<i>Lasiogloss</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Bees	<i>Melecta al</i>	NA	0	-4 to -1%	> -1%	LOW	POOR
Bees	<i>Melitta ha</i>	NA	0	-7.5 to -4%	> -1%	LOW	POOR
Bees	<i>Nomada fla</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Bees	<i>Sphecodes</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Bees	<i>Sphecodes</i>	NA	0	-7.5 to -4%	> -1%	LOW	POOR
Bees	<i>Sphecodes</i>	NA	0	-7.5 to -4%	> -1%	LOW	POOR
Birds	<i>Accipiter</i>	Goshawk	0	> -1%	-7.5 to -4%	MODERATE	POOR
Birds	<i>Alauda arv</i>	Skylark	1	-4 to -1%	> -1%	LOW	MEDIUM
Birds	<i>Anas platy</i>	Mallard	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Anthus tri</i>	Tree Pipit	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Birds	<i>Asio flamm</i>	Short-eare	0	< -7.5%	-7.5 to -4%	VERY HIGH	MEDIUM
Birds	<i>Botaurus s</i>	Bittern	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Branta can</i>	Canada Goo	0	> -1%	< -7.5%	MODERATE	POOR
Birds	<i>Branta leu</i>	Barnacle G	0	< -7.5%	-7.5 to -4%	VERY HIGH	MEDIUM
Birds	<i>Burhinus o</i>	Stone-curl	1	< -7.5%	-4 to -1%	VERY HIGH	MEDIUM
Birds	<i>Buteo bute</i>	Buzzard	0	> -1%	-4 to -1%	LOW	POOR
Birds	<i>Caprimulgu</i>	Nightjar	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Carduelis</i>	Lesser Red	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Birds	<i>Carduelis</i>	Linnet	1	> -1%	> -1%	LOW	GOOD
Birds	<i>Carduelis</i>	Twite	1	< -7.5%	< -7.5%	VERY HIGH	MEDIUM

Birds	<i>Certhia fa</i> Treecreeper	0	< -7.5%	< -7.5%	MODERATE	POOR
Birds	<i>Cettia cet</i> Cetti's Warbler	0	> -1%	> -1%	LOW	GOOD
Birds	<i>Cinclus ci</i> Dipper	0	-7.5 to -4%	-7.5 to -4%	HIGH	POOR
Birds	<i>Circus cya</i> Hen Harrier	1	< -7.5%	-7.5 to -4%	HIGH	MEDIUM
Birds	<i>Coccothrau</i> Hawfinch	1	< -7.5%	> -1%	MODERATE	POOR
Birds	<i>Columba li</i> Rock Dove	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Columba oe</i> Stock Dove	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Corvus fru</i> Rook	0	> -1%	-4 to -1%	LOW	MEDIUM
Birds	<i>Crex crex</i> Corncrake	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Cuculus ca</i> Cuckoo	1	< -7.5%	-4 to -1%	VERY HIGH	POOR
Birds	<i>Delichon u</i> House Martin	0	> -1%	> -1%	LOW	GOOD
Birds	<i>Dendrocopo</i> Great Spotted Woodpecker	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Dendrocopo</i> Lesser Spotted Woodpecker	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Emberiza c</i> Corn Bunting	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Emberiza c</i> Cirl Bunting	1	-4 to -1%	> -1%	LOW	POOR
Birds	<i>Emberiza c</i> Yellowhammer	1	-7.5 to -4%	-4 to -1%	MODERATE	MEDIUM
Birds	<i>Emberiza s</i> Reed Bunting	1	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Falco colu</i> Merlin	0	> -1%	< -7.5%	MODERATE	POOR
Birds	<i>Falco pere</i> Peregrine	0	-7.5 to -4%	-7.5 to -4%	VERY HIGH	POOR
Birds	<i>Falco tinn</i> Kestrel	0	-4 to -1%	> -1%	LOW	POOR
Birds	<i>Fulica atr</i> Coot	0	-4 to -1%	> -1%	LOW	POOR
Birds	<i>Haematopus</i> Oystercatcher	0	-4 to -1%	> -1%	LOW	MEDIUM
Birds	<i>Lagopus la</i> Red Grouse	1	< -7.5%	< -7.5%	VERY HIGH	GOOD
Birds	<i>Lanius col</i> Red-backed Thrush	0	< -7.5%	> -1%	LOW	POOR
Birds	<i>Larus arge</i> Herring Gull	1	> -1%	> -1%	LOW	GOOD
Birds	<i>Larus fusc</i> Lesser Black-backed Gull	0	< -7.5%	> -1%	LOW	POOR
Birds	<i>Limosa lim</i> Black-tailed Godwit	1	> -1%	> -1%	LOW	GOOD
Birds	<i>Locustella</i> Savi's Warbler	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Locustella</i> Grasshopper Warbler	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Lullula ar</i> Woodlark	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Morus bass</i> Gannet	0	> -1%	-4 to -1%	MODERATE	POOR
Birds	<i>Motacilla</i> Yellow Wagtail	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Muscicapa</i> Spotted Flycatcher	1	-7.5 to -4%	> -1%	MODERATE	POOR
Birds	<i>Numenius a</i> Curlew	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Birds	<i>Passer dom</i> House Sparrow	1	> -1%	> -1%	LOW	GOOD
Birds	<i>Passer mon</i> Tree Sparrow	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Perdix per</i> Grey Partridge	1	< -7.5%	> -1%	HIGH	POOR
Birds	<i>Phalacroco</i> Cormorant	0	< -7.5%	> -1%	LOW	POOR
Birds	<i>Phoenicuru</i> Black Redstart	0	< -7.5%	> -1%	LOW	POOR
Birds	<i>Phoenicuru</i> Redstart	0	< -7.5%	> -1%	VERY HIGH	POOR
Birds	<i>Phylloscop</i> Wood Warbler	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Birds	<i>Picus viri</i> Green Woodpecker	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Podiceps c</i> Great Crested Newt	0	-4 to -1%	> -1%	LOW	POOR
Birds	<i>Poecile mo</i> Willow Tit	1	-4 to -1%	> -1%	VERY HIGH	POOR
Birds	<i>Poecile pa</i> Marsh Tit	1	-7.5 to -4%	-7.5 to -4%	HIGH	POOR
Birds	<i>Porzana po</i> Spotted Cranes	0	< -7.5%	> -1%	LOW	POOR
Birds	<i>Prunella m</i> Dunnock	1	> -1%	> -1%	LOW	GOOD
Birds	<i>Puffinus p</i> Manx Shearwater	0	< -7.5%	> -1%	LOW	POOR
Birds	<i>Pyrrhula p</i> Bullfinch	1	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Rallus aqu</i> Water Rail	0	< -7.5%	> -1%	LOW	POOR

Birds	<i>Saxicola r</i>	Whinchat	0	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Birds	<i>Saxicola t</i>	Stonechat	0	< -7.5%	-7.5 to -4%	LOW	MEDIUM
Birds	<i>Sitta eur</i>	Nuthatch	0	> -1%	> -1%	LOW	POOR
Birds	<i>Sterna dou</i>	Roseate Te	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Streptopel</i>	Collared D	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Streptopel</i>	Turtle Dov	1	< -7.5%	> -1%	LOW	POOR
Birds	<i>Sturnus vu</i>	Starling	1	> -1%	> -1%	LOW	GOOD
Birds	<i>Sylvia bor</i>	Garden War	0	-7.5 to -4%	-7.5 to -4%	VERY HIGH	POOR
Birds	<i>Sylvia com</i>	Whitethroa	0	> -1%	> -1%	LOW	GOOD
Birds	<i>Tachybaptu</i>	Little Gre	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Tetrao tet</i>	Black Grou	1	> -1%	< -7.5%	MODERATE	POOR
Birds	<i>Tringa tot</i>	Redshank	0	< -7.5%	> -1%	LOW	POOR
Birds	<i>Troglodyte</i>	Wren	0	> -1%	> -1%	LOW	GOOD
Birds	<i>Turdus mer</i>	Blackbird	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Turdus phi</i>	Song Thrus	1	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Turdus tor</i>	Ring Ouzel	1	< -7.5%	< -7.5%	VERY HIGH	GOOD
Birds	<i>Tyto alba</i>	Barn Owl	0	> -1%	> -1%	LOW	MEDIUM
Birds	<i>Vanellus v</i>	Lapwing	1	-7.5 to -4%	> -1%	HIGH	POOR
Bryophytes	<i>Calypogeia</i>	Bog Pouchw	0	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Bryophytes	<i>Dicranum s</i>	Rusty Fork	1	< -7.5%	-4 to -1%	HIGH	POOR
Bryophytes	<i>Didymodon</i>	Cylindric	0	> -1%	> -1%	LOW	MEDIUM
Bryophytes	<i>Didymodon</i>	Brown Bear	0	> -1%	-4 to -1%	MODERATE	POOR
Bryophytes	<i>Leucobryum</i>	Large Whit	0	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Bryophytes	<i>Plagiochil</i>	Western Fe	0	> -1%	< -7.5%	HIGH	POOR
Bryophytes	<i>Plagiomniu</i>	Many-fruit	0	< -7.5%	> -1%	LOW	POOR
Bryophytes	<i>Plagiothec</i>	Bright Sil	0	-4 to -1%	< -7.5%	HIGH	POOR
Bryophytes	<i>Porella pl</i>	Wall Scale	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Bryophytes	<i>Racomitriu</i>	Slender Fr	0	> -1%	< -7.5%	HIGH	POOR
Bryophytes	<i>Radula aqu</i>	Brown Scal	0	-7.5 to -4%	< -7.5%	VERY HIGH	POOR
Bryophytes	<i>Rhytidiade</i>	Little Sha	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Bryophytes	<i>Sphagnum p</i>	Golden Bog	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Bryophytes	<i>Zygodon vi</i>	NA	0	> -1%	< -7.5%	HIGH	POOR
Carbid bee	<i>Amara ovat</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Carbid bee	<i>Amara tibi</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Carbid bee	<i>Anatis oce</i>	Eyed Ladyb	0	< -7.5%	> -1%	LOW	POOR
Carbid bee	<i>Anthracus</i>	NA	0	-4 to -1%	> -1%	LOW	POOR
Carbid bee	<i>Bembidion</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Carbid bee	<i>Bembidion</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Carbid bee	<i>Bradycellu</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Carbid bee	<i>Calathus m</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Carbid bee	<i>Calosoma i</i>	NA	1	> -1%	< -7.5%	HIGH	POOR
Carbid bee	<i>Carabus mo</i>	Necklace G	1	< -7.5%	> -1%	MODERATE	POOR
Carbid bee	<i>Carabus pr</i>	NA	0	-4 to -1%	-4 to -1%	HIGH	POOR
Carbid bee	<i>Curtonotus</i>	NA	0	> -1%	> -1%	LOW	POOR
Carbid bee	<i>Laemostenu</i>	NA	0	> -1%	> -1%	LOW	POOR
Carbid bee	<i>Ocys harpa</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Carbid bee	<i>Philorhizu</i>	NA	1	< -7.5%	> -1%	LOW	POOR
Carbid bee	<i>Pterostich</i>	NA	0	> -1%	< -7.5%	HIGH	POOR
Carbid bee	<i>Syntomus f</i>	NA	0	> -1%	> -1%	LOW	POOR
Centipedes	<i>Cryptops h</i>	NA	0	> -1%	> -1%	LOW	MEDIUM

Centipedes <i>Geophilus</i> NA	0	< -7.5%	> -1%	LOW	POOR
Centipedes <i>Geophilus</i> NA	0	> -1%	> -1%	LOW	POOR
Centipedes <i>Geophilus</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Centipedes <i>Geophilus</i> NA	0	< -7.5%	> -1%	MODERATE	POOR
Centipedes <i>Henia vesu</i> NA	0	> -1%	> -1%	LOW	MEDIUM
Centipedes <i>Lithobius</i> NA	0	< -7.5%	> -1%	MODERATE	POOR
Centipedes <i>Lithobius</i> NA	0	> -1%	> -1%	LOW	MEDIUM
Centipedes <i>Lithobius</i> NA	0	> -1%	> -1%	LOW	POOR
Centipedes <i>Lithobius</i> NA	0	> -1%	> -1%	LOW	MEDIUM
Centipedes <i>Schendyla</i> NA	0	> -1%	> -1%	LOW	MEDIUM
Centipedes <i>Stigmatoga</i> NA	0	> -1%	> -1%	LOW	POOR
Centipedes <i>Strigamia</i> NA	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Adalia bip</i> Two-spot L	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Adalia dec</i> Ten-spot L	0	> -1%	> -1%	LOW	MEDIUM
Coccinelid <i>Anisostict</i> Water Lady	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Coccidula</i> NA	0	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Coccinelid <i>Coccinella</i> Seven-spot	0	-4 to -1%	> -1%	LOW	POOR
Coccinelid <i>Coccinella</i> Eleven-spo	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Halysia se</i> Orange Lad	0	< -7.5%	-4 to -1%	HIGH	POOR
Coccinelid <i>Hippodamia</i> Adonis' La	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Propylea q</i> Fourteen-s	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Psyllobora</i> Twentytwo-	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Scymnus su</i> NA	0	< -7.5%	> -1%	LOW	POOR
Coccinelid <i>Subcoccine</i> Twentyfour-	0	> -1%	> -1%	LOW	POOR
Craneflies <i>Nephrotoma</i> NA	0	-4 to -1%	> -1%	LOW	POOR
Craneflies <i>Ptychopter</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Craneflies <i>Ptychopter</i> NA	0	< -7.5%	> -1%	LOW	POOR
Craneflies <i>Ptychopter</i> NA	0	< -7.5%	> -1%	LOW	POOR
Craneflies <i>Tipula ful</i> NA	0	< -7.5%	> -1%	LOW	POOR
Craneflies <i>Tipula lat</i> NA	0	> -1%	-7.5 to -4%	HIGH	POOR
Craneflies <i>Tipula lun</i> NA	0	< -7.5%	> -1%	LOW	POOR
Craneflies <i>Tipula max</i> NA	0	> -1%	> -1%	LOW	MEDIUM
Craneflies <i>Tipula ole</i> NA	0	< -7.5%	> -1%	LOW	POOR
Craneflies <i>Tipula unc</i> NA	0	> -1%	< -7.5%	HIGH	POOR
Craneflies <i>Tipula var</i> NA	0	> -1%	-4 to -1%	MODERATE	POOR
Crickets a <i>Chorthippu</i> NA	0	-7.5 to -4%	> -1%	LOW	POOR
Crickets a <i>Conocephal</i> NA	0	> -1%	-7.5 to -4%	HIGH	POOR
Crickets a <i>Conocephal</i> NA	0	> -1%	> -1%	LOW	MEDIUM
Crickets a <i>Ectobius p</i> Tawny Cock	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Crickets a <i>Ectobius p</i> Lesser Coc	0	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Crickets a <i>Forficula</i> Common Ear	0	> -1%	> -1%	LOW	POOR
Crickets a <i>Forficula</i> Lesne's Ea	0	> -1%	> -1%	LOW	MEDIUM
Crickets a <i>Meconema t</i> NA	0	> -1%	-7.5 to -4%	HIGH	POOR
Crickets a <i>Omocestus</i> Woodland G	0	> -1%	< -7.5%	HIGH	POOR
Crickets a <i>Omocestus</i> Common Gre	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Crickets a <i>Platycleis</i> NA	0	< -7.5%	> -1%	LOW	POOR
Crickets a <i>Tetrix cep</i> Cepero's G	0	< -7.5%	> -1%	LOW	POOR
Crickets a <i>Tetrix sub</i> NA	0	> -1%	> -1%	LOW	MEDIUM
Hoverflies <i>Anasimyia</i> NA	0	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Hoverflies <i>Cheilosia</i> NA	0	< -7.5%	> -1%	LOW	POOR

Hoverflies	<i>Cheilosia</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Hoverflies	<i>Cheilosia</i>	NA	0	> -1%	> -1%	LOW	POOR
Hoverflies	<i>Cheilosia</i>	NA	0	-4 to -1%	< -7.5%	VERY HIGH	POOR
Hoverflies	<i>Eristalis</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Hoverflies	<i>Eupeodes</i>	bNA	0	< -7.5%	> -1%	MODERATE	POOR
Hoverflies	<i>Eupeodes</i>	nNA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Hoverflies	<i>Neoascia</i>	mNA	0	< -7.5%	> -1%	LOW	POOR
Hoverflies	<i>Platycheir</i>	NA	0	> -1%	< -7.5%	HIGH	POOR
Hoverflies	<i>Sphaeropho</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Hoverflies	<i>Sphagina</i>	eNA	0	< -7.5%	> -1%	LOW	POOR
Hoverflies	<i>Xylota</i>	segNA	0	-4 to -1%	> -1%	LOW	POOR
Millipedes	<i>Blaniulus</i>	Spotted Sn	0	> -1%	> -1%	LOW	MEDIUM
Millipedes	<i>Brachydesm</i>	NA	0	> -1%	> -1%	LOW	POOR
Millipedes	<i>Chordeuma</i>	NA	0	> -1%	< -7.5%	HIGH	POOR
Millipedes	<i>Cylindroi</i>	Blunt-tail	0	-4 to -1%	> -1%	LOW	POOR
Millipedes	<i>Glomeris</i>	mPill Milli	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Millipedes	<i>Melogona</i>	sNA	0	> -1%	< -7.5%	HIGH	POOR
Millipedes	<i>Nanogona</i>	pEyed Flat-	0	> -1%	-4 to -1%	MODERATE	POOR
Millipedes	<i>Nemasoma</i>	vNA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Millipedes	<i>Ommatoiulu</i>	Striped Mi	0	> -1%	-7.5 to -4%	HIGH	POOR
Millipedes	<i>Ophiulus</i>	NA	0	> -1%	< -7.5%	HIGH	POOR
Millipedes	<i>Polydesmus</i>	Common Fla	0	> -1%	> -1%	LOW	MEDIUM
Millipedes	<i>Polydesmus</i>	NA	0	> -1%	-7.5 to -4%	HIGH	POOR
Millipedes	<i>Tachypodoi</i>	White-legg	0	-4 to -1%	< -7.5%	VERY HIGH	POOR
Moths	<i>Acronicta</i>	Knot Grass	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Adscita</i>	stThe Forest	1	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Moths	<i>Agrochola</i>	Flounced C	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Agrochola</i>	Brown-spot	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Agrochola</i>	Beaded Che	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Alcis</i>	jubaDotted Car	0	> -1%	< -7.5%	HIGH	POOR
Moths	<i>Aleucis</i>	disSloe Carpe	1	< -7.5%	-4 to -1%	HIGH	POOR
Moths	<i>Allophyes</i>	Green-brin	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Moths	<i>Alsophila</i>	March Moth	0	< -7.5%	> -1%	MODERATE	POOR
Moths	<i>Apamea</i>	ancLarge Nutm	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Arctia</i>	cayGarden Tig	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Atethmia</i>	cCentre-bar	1	-4 to -1%	> -1%	LOW	POOR
Moths	<i>Blepharita</i>	Dark Broca	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Cabera</i>	exaCommon Wav	0	> -1%	-4 to -1%	MODERATE	POOR
Moths	<i>Caradrina</i>	Mottled Ru	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Celaena</i>	haHaworth's	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Moths	<i>Chesias</i>	ruBroom-tip	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Colotois</i>	pFeathered	0	< -7.5%	> -1%	LOW	POOR
Moths	<i>Cosmia</i>	affLesser-spo	0	< -7.5%	> -1%	LOW	POOR
Moths	<i>Cossus</i>	cosGoat Moth	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Cyclophora</i>	Dingy Moch	1	> -1%	> -1%	LOW	MEDIUM
Moths	<i>Cyclophora</i>	False Moch	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Cymatophor</i>	Oak Lutest	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Dasypolia</i>	Brindled O	1	< -7.5%	-4 to -1%	HIGH	POOR
Moths	<i>Diloba</i>	caeFigure of	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Eilema</i>	sorOrange Foo	0	> -1%	< -7.5%	HIGH	POOR

Moths	<i>Ennomos er</i>	September	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Ennomos qu</i>	August Tho	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Entephria</i>	Grey Mount	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Eugnorisma</i>	Autumnal R	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Eulithis m</i>	The Spinac	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Euxoa trit</i>	White-line	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Graphiphor</i>	Double Dar	1	< -7.5%	> -1%	MODERATE	POOR
Moths	<i>Hadena alb</i>	White Spot	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Heliophobu</i>	Bordered G	1	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Moths	<i>Hemistola</i>	Small Emer	1	-7.5 to -4%	> -1%	LOW	POOR
Moths	<i>Hoplodrina</i>	The Rustic	1	> -1%	> -1%	LOW	MEDIUM
Moths	<i>Idaea muri</i>	Purple-bor	0	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Moths	<i>Jodis lact</i>	Little Eme	0	< -7.5%	> -1%	LOW	POOR
Moths	<i>Lycia hirt</i>	Brindled B	1	< -7.5%	> -1%	MODERATE	POOR
Moths	<i>Macaria wa</i>	The V-Moth	1	< -7.5%	> -1%	MODERATE	POOR
Moths	<i>Malacosoma</i>	The Lackey	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Melanchra</i>	Dot Moth	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Melanthia</i>	Pretty Cha	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Mythimna c</i>	Shoulder-s	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Operophter</i>	Northern W	0	< -7.5%	> -1%	MODERATE	POOR
Moths	<i>Oria muscu</i>	Brighton W	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Orthosia g</i>	Powdered Q	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Paracolax</i>	Clay Fan-f	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Pelurga co</i>	Dark Spina	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Photedes c</i>	Least Mino	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Polia bomb</i>	Pale Shini	1	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Moths	<i>Rheumapter</i>	Argent & S	1	< -7.5%	-7.5 to -4%	VERY HIGH	MEDIUM
Moths	<i>Rhizedra l</i>	Large Wain	1	-7.5 to -4%	> -1%	LOW	POOR
Moths	<i>Scotoptery</i>	Chalk Carp	1	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Moths	<i>Selidosema</i>	Bordered G	0	< -7.5%	> -1%	LOW	POOR
Moths	<i>Shargacucu</i>	Striped Ly	1	< -7.5%	-4 to -1%	HIGH	MEDIUM
Moths	<i>Spilosoma</i>	Buff Ermin	1	> -1%	> -1%	LOW	MEDIUM
Moths	<i>Stilbia an</i>	The Anomal	1	< -7.5%	-4 to -1%	HIGH	POOR
Moths	<i>Tholera ce</i>	Hedge Rust	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Tholera de</i>	Feathered	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Trichiura</i>	Pale Eggar	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Trichopter</i>	Early Toot	0	> -1%	-4 to -1%	MODERATE	POOR
Moths	<i>Trichopter</i>	Barred Too	1	< -7.5%	> -1%	MODERATE	POOR
Moths	<i>Trisateles</i>	Olive Cres	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Tyta luctu</i>	The Four-s	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Watsonalla</i>	Barred Hoo	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Moths	<i>Xanthia gi</i>	Dusky-lemo	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Moths	<i>Xanthia ic</i>	The Sallow	1	< -7.5%	> -1%	LOW	POOR
Moths	<i>Xanthorhoe</i>	Red Carpet	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Moths	<i>Xestia aga</i>	Heath Rust	1	< -7.5%	> -1%	LOW	POOR
Odonata	<i>Aeshna cae</i>	Azure hawk	0	> -1%	< -7.5%	HIGH	POOR
Odonata	<i>Aeshna gra</i>	Brown hawk	0	< -7.5%	-4 to -1%	HIGH	POOR
Odonata	<i>Aeshna jun</i>	Common haw	0	< -7.5%	> -1%	LOW	POOR
Odonata	<i>Anax imper</i>	Emperor dr	0	> -1%	> -1%	LOW	MEDIUM
Odonata	<i>Ceriagrion</i>	Small red	0	< -7.5%	< -7.5%	VERY HIGH	POOR

Odonata	<i>Enallagma</i>	Common blu	0	-7.5 to -4%	-4 to -1%	HIGH	POOR
Odonata	<i>Erythromma</i>	Red-eyed d	0	>-1%	< -7.5%	HIGH	POOR
Odonata	<i>Ischnura e</i>	Blue-taile	0	-7.5 to -4%	> -1%	LOW	POOR
Odonata	<i>Orthetrum</i>	Black-tail	0	>-1%	> -1%	LOW	MEDIUM
Odonata	<i>Orthetrum</i>	Keeled ski	0	-7.5 to -4%	< -7.5%	VERY HIGH	POOR
Odonata	<i>Pyrrhosoma</i>	Large red	0	< -7.5%	> -1%	LOW	POOR
Odonata	<i>Sympetrum</i>	Ruddy dart	0	>-1%	> -1%	LOW	MEDIUM
Odonata	<i>Sympetrum</i>	Common dar	0	-7.5 to -4%	> -1%	LOW	POOR
Soldier be	<i>Cantharis</i>	NA	0	-4 to -1%	< -7.5%	HIGH	POOR
Soldier be	<i>Cantharis</i>	NA	0	< -7.5%	-4 to -1%	HIGH	POOR
Soldier be	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Soldier be	<i>Cantharis</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Soldier be	<i>Cantharis</i>	NA	0	-7.5 to -4%	-4 to -1%	HIGH	POOR
Soldier be	<i>Malthinus</i>	NA	0	>-1%	-4 to -1%	MODERATE	POOR
Soldier be	<i>Malthinus</i>	NA	0	>-1%	> -1%	LOW	POOR
Soldier be	<i>Podabrus a</i>	NA	0	>-1%	< -7.5%	HIGH	POOR
Soldier be	<i>Rhagonycha</i>	Common Red	0	-7.5 to -4%	> -1%	LOW	POOR
Soldier be	<i>Rhagonycha</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Soldier be	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Soldier be	<i>Rhagonycha</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Soldier be	<i>Rhagonycha</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Spiders	<i>Anelosimus</i>	NA	0	>-1%	> -1%	LOW	MEDIUM
Spiders	<i>Araneus ma</i>	NA	0	>-1%	< -7.5%	HIGH	POOR
Spiders	<i>Araneus qu</i>	NA	0	>-1%	> -1%	LOW	POOR
Spiders	<i>Bathyphant</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Spiders	<i>Ceratinella</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Spiders	<i>Clubiona n</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Spiders	<i>Dictyna pu</i>	Small Mesh	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Spiders	<i>Diploceph</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Spiders	<i>Haplodrass</i>	Heath Gras	1	< -7.5%	> -1%	LOW	POOR
Spiders	<i>Mecopisthe</i>	Peus' s Lon	1	< -7.5%	> -1%	LOW	POOR
Spiders	<i>Meioneta m</i>	Thin Weble	1	< -7.5%	> -1%	LOW	POOR
Spiders	<i>Monocephal</i>	Broad Groo	1	< -7.5%	-7.5 to -4%	VERY HIGH	MEDIUM
Spiders	<i>Porrhomma</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Spiders	<i>Porrhomma</i>	NA	0	< -7.5%	> -1%	MODERATE	POOR
Spiders	<i>Saaristoa</i>	Triangle H	1	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Spiders	<i>Salticus s</i>	NA	0	>-1%	> -1%	LOW	POOR
Spiders	<i>Sitticus c</i>	Sedge Jump	1	< -7.5%	> -1%	LOW	POOR
Spiders	<i>Tapinocyba</i>	NA	0	-4 to -1%	< -7.5%	VERY HIGH	POOR
Spiders	<i>Trichopter</i>	NA	0	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Spiders	<i>Walckenaer</i>	NA	0	>-1%	< -7.5%	HIGH	POOR
Vascular p	<i>Aceras ant</i>	Man Orchid	0	>-1%	< -7.5%	HIGH	POOR
Vascular p	<i>Ajuga pyra</i>	Pyramidal	1	>-1%	< -7.5%	HIGH	POOR
Vascular p	<i>Blysmus co</i>	Flat-sedge	1	< -7.5%	-7.5 to -4%	VERY HIGH	POOR
Vascular p	<i>Bupleurum</i>	Slender Ha	1	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Calamagros</i>	Purple Sma	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Calamagros</i>	Narrow Sma	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Calystegia</i>	Great Bind	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Carex eric</i>	Rare Sprin	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Carex viri</i>	Common Yel	0	>-1%	> -1%	LOW	POOR

Vascular p	<i>Centaurea</i>	Cornflower	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Cephalanth</i>	White Hell	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Chamaemelu</i>	Chamomile	1	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Chenopodiu</i>	Fig-leaved	0	> -1%	> -1%	LOW	MEDIUM
Vascular p	<i>Chenopodiu</i>	Stinking G	1	> -1%	> -1%	LOW	POOR
Vascular p	<i>Cicendia f</i>	Yellow Cen	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Dactylorhi</i>	Common Spo	0	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Dactylorhi</i>	Narrow-lea	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Dianthus a</i>	Deptford P	1	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Euphrasia</i>	Chalk Eyeb	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Euphrasia</i>	Cornish Ey	1	< -7.5%	< -7.5%	VERY HIGH	MEDIUM
Vascular p	<i>Fumaria pu</i>	Purple Ram	1	-7.5 to -4%	> -1%	LOW	POOR
Vascular p	<i>Galeopsis</i>	Red Hemp-n	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Herminium</i>	Musk Orchi	1	> -1%	< -7.5%	HIGH	POOR
Vascular p	<i>Illecebrum</i>	Coral-neck	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Luronium n</i>	Floating W	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Melittis m</i>	Bastard Ba	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Mentha pul</i>	Pennyroyal	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Minuartia</i>	Fine-leave	1	> -1%	< -7.5%	HIGH	POOR
Vascular p	<i>Muscari ne</i>	Grape-hyac	1	> -1%	> -1%	LOW	MEDIUM
Vascular p	<i>Najas flex</i>	Slender Na	1	> -1%	< -7.5%	HIGH	POOR
Vascular p	<i>Phyllitis</i>	Hart's-ton	0	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Pilularia</i>	Pillwort	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Potamogeto</i>	Grass-wrac	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Pulsatilla</i>	Pasqueflow	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Quercus ro</i>	Pedunculat	0	-4 to -1%	> -1%	LOW	POOR
Vascular p	<i>Ranunculus</i>	Corn Butte	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Ranunculus</i>	Three-lobe	1	> -1%	> -1%	LOW	MEDIUM
Vascular p	<i>Rubus saxa</i>	Stone Bram	0	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Rumex rupe</i>	Shore Dock	1	> -1%	> -1%	LOW	MEDIUM
Vascular p	<i>Salix lapp</i>	Downy Will	1	> -1%	< -7.5%	HIGH	POOR
Vascular p	<i>Scandix pe</i>	Shepherd's	1	> -1%	-4 to -1%	MODERATE	POOR
Vascular p	<i>Scrophular</i>	Water Figw	0	> -1%	> -1%	LOW	MEDIUM
Vascular p	<i>Silene gal</i>	Small-flow	1	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Sium latif</i>	Greater Wa	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Sparganium</i>	Branched B	0	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Spartina m</i>	Small Cord	1	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Stellaria</i>	Marsh Stit	1	< -7.5%	> -1%	LOW	POOR
Vascular p	<i>Torilis ar</i>	Spreading	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Valeriane1</i>	Broad-frui	1	< -7.5%	< -7.5%	VERY HIGH	POOR
Vascular p	<i>Veronica c</i>	Germander	0	-7.5 to -4%	> -1%	MODERATE	POOR
Vascular p	<i>Viola lact</i>	Pale Dog-v	1	> -1%	-4 to -1%	MODERATE	POOR
Wasps	<i>Agenioideu</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Ancistroce</i>	Wall Mason	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Cerceris r</i>	Ornate Tai	0	> -1%	> -1%	LOW	MEDIUM
Wasps	<i>Crabro pel</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Crossoceru</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Entomognat</i>	NA	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Hedychridi</i>	NA	0	> -1%	> -1%	LOW	MEDIUM
Wasps	<i>Nysson spi</i>	Large Spur	0	< -7.5%	< -7.5%	VERY HIGH	POOR

Wasps	<i>Odynerus m</i> NA	1	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Oxybelus a</i> Silver Spi	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Pemphredon</i> Mournful W	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Priocnemis</i> NA	0	< -7.5%	> -1%	LOW	POOR
Wasps	<i>Priocnemis</i> NA	0	>-1%	> -1%	LOW	MEDIUM
Wasps	<i>Smicromyrm</i> Small Velv	0	>-1%	> -1%	LOW	MEDIUM

Observed expansion	Projected expansion	Benefit from expansion	Associated confidence	Final outcome
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+4 to +7.5%	+1 to +4%	MODERATE	POOR	Medium benefit
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	+4 to +7.5%	MODERATE	POOR	Medium benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	+4 to +7.5%	HIGH	POOR	Risks & benefits
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	HIGH	POOR	Medium benefit
+4 to +7.5%	> +7.5%	HIGH	POOR	Medium benefit
< +1%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	> +7.5%	HIGH	MEDIUM	High benefit
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
> +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	+1 to +4%	HIGH	POOR	High benefit
+4 to +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+4 to +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	< +1%	HIGH	POOR	Medium benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	High risk
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	POOR	Medium risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	Risks & benefits
> +7.5%	> +7.5%	HIGH	MEDIUM	Medium risk
> +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	POOR	High risk

> +7.5%	< +1%	LOW	POOR	Medium risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	+1 to +4%	MODERATE	POOR	Medium benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	+1 to +4%	HIGH	MEDIUM	High benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	Medium risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	POOR	Medium risk
> +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
+4 to +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Medium risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	MODERATE	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit

+1 to +4%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
< +1%	< +1%	LOW	MEDIUM	Limited impact
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	POOR	Medium risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
< +1%	< +1%	LOW	MEDIUM	Limited impact
< +1%	< +1%	LOW	MEDIUM	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	High risk
> +7.5%	+4 to +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	Medium benefit
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+1 to +4%	+1 to +4%	MODERATE	MEDIUM	Risks & benefits
+4 to +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	MODERATE	POOR	Medium benefit
+1 to +4%	< +1%	LOW	POOR	High risk
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	HIGH	POOR	Medium risk
< +1%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	+4 to +7.5%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+1 to +4%	+4 to +7.5%	MODERATE	POOR	Medium benefit
+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	> +7.5%	HIGH	POOR	Medium benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	+1 to +4%	MODERATE	POOR	Medium benefit
< +1%	> +7.5%	MODERATE	POOR	Medium benefit
< +1%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit

+1 to +4%	< +1%	MODERATE	POOR	Medium benefit
+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	Medium risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+4 to +7.5%	< +1%	MODERATE	POOR	Risks & benefits
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	+1 to +4%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	+1 to +4%	MODERATE	MEDIUM	Medium benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	+4 to +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
+1 to +4%	+4 to +7.5%	MODERATE	POOR	Medium benefit
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	+4 to +7.5%	MODERATE	POOR	Medium benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	+4 to +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	High risk
+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	+1 to +4%	HIGH	POOR	Risks & benefits
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
< +1%	< +1%	LOW	MEDIUM	Limited impact
+1 to +4%	+1 to +4%	LOW	MEDIUM	Limited impact
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	POOR	Medium risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	Medium benefit
> +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	> +7.5%	HIGH	POOR	Medium risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	POOR	High risk
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	HIGH	POOR	High benefit

> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	+4 to +7.5%	MODERATE	MEDIUM	Medium benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
< +1%	< +1%	LOW	MEDIUM	Limited impact
< +1%	< +1%	LOW	MEDIUM	High risk
> +7.5%	+4 to +7.5%	HIGH	POOR	Risks & benefits
+1 to +4%	< +1%	LOW	MEDIUM	Medium risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
+4 to +7.5%	+1 to +4%	MODERATE	POOR	Medium risk
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	HIGH	POOR	Risks & benefits
+1 to +4%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	Medium risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
< +1%	< +1%	LOW	MEDIUM	Medium risk
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
> +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	< +1%	LOW	MEDIUM	Limited impact
+4 to +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
< +1%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	MEDIUM	High risk

+1 to +4%	+1 to +4%	MODERATE	POOR	Medium benefit
+4 to +7.5%	+4 to +7.5%	HIGH	MEDIUM	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	Medium risk
> +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+4 to +7.5%	< +1%	MODERATE	POOR	Medium benefit
> +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	+1 to +4%	MODERATE	MEDIUM	Medium benefit
> +7.5%	< +1%	LOW	POOR	Medium risk
+1 to +4%	< +1%	LOW	POOR	Medium risk
< +1%	< +1%	LOW	MEDIUM	Limited impact
+1 to +4%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	POOR	Medium risk
< +1%	< +1%	LOW	GOOD	High risk
+4 to +7.5%	< +1%	MODERATE	POOR	Medium benefit
< +1%	> +7.5%	MODERATE	POOR	Medium benefit
+4 to +7.5%	> +7.5%	HIGH	POOR	High benefit
< +1%	+1 to +4%	LOW	POOR	High risk
< +1%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	HIGH	MEDIUM	Medium risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	High risk
+4 to +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	Medium benefit
+1 to +4%	< +1%	LOW	POOR	Limited impact
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
+1 to +4%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	Medium risk
+1 to +4%	> +7.5%	MODERATE	POOR	Risks & benefits
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	Medium risk
> +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk

< +1%	< +1%	LOW	MEDIUM	High risk
> +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	< +1%	LOW	POOR	High risk
< +1%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	MODERATE	POOR	Medium risk
+4 to +7.5%	< +1%	LOW	MEDIUM	Medium risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	+1 to +4%	MODERATE	POOR	Medium benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	+1 to +4%	MODERATE	MEDIUM	Medium benefit
< +1%	< +1%	LOW	GOOD	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	High risk
> +7.5%	+4 to +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
+1 to +4%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	HIGH	POOR	High benefit
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
< +1%	> +7.5%	HIGH	POOR	High benefit
< +1%	> +7.5%	MODERATE	POOR	Medium benefit
+1 to +4%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
< +1%	> +7.5%	MODERATE	POOR	Risks & benefits
< +1%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
< +1%	> +7.5%	MODERATE	POOR	Medium benefit
> +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
+4 to +7.5%	> +7.5%	HIGH	MEDIUM	Risks & benefits
> +7.5%	+1 to +4%	HIGH	POOR	Risks & benefits
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
> +7.5%	< +1%	LOW	MEDIUM	High risk
< +1%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact

> +7.5%	> +7.5%	VERY HIGH	MEDIUM	Risks & benefits
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	Risks & benefits
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
> +7.5%	+1 to +4%	HIGH	POOR	Medium risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	+1 to +4%	MODERATE	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	+1 to +4%	HIGH	POOR	Medium risk
> +7.5%	< +1%	LOW	MEDIUM	High risk
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	Risks & benefits
> +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	+4 to +7.5%	HIGH	POOR	Risks & benefits
> +7.5%	+1 to +4%	HIGH	POOR	High benefit
> +7.5%	+4 to +7.5%	HIGH	POOR	Medium risk
> +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	< +1%	LOW	POOR	High risk
+4 to +7.5%	< +1%	LOW	POOR	Limited impact
+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	< +1%	LOW	MEDIUM	High risk
+1 to +4%	> +7.5%	HIGH	POOR	High benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
+1 to +4%	< +1%	LOW	MEDIUM	Limited impact
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	POOR	High risk
> +7.5%	< +1%	LOW	POOR	Limited impact
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	< +1%	LOW	MEDIUM	High risk
> +7.5%	< +1%	LOW	MEDIUM	High risk
< +1%	< +1%	LOW	MEDIUM	Medium risk
+1 to +4%	> +7.5%	HIGH	POOR	Medium benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
+4 to +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	HIGH	MEDIUM	High benefit
+4 to +7.5%	< +1%	LOW	POOR	High risk

> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	POOR	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit
> +7.5%	> +7.5%	VERY HIGH	MEDIUM	High benefit