

This is a repository copy of *Towards broad spectrum activity-based glycosidase probes: synthesis and evaluation of deoxygenated cyclophellitol aziridines*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/123810/>

Version: Accepted Version

Article:

Schroder, Sybrin P, van de Sande, Jasper W, Kallemeijn, Wouter W. et al. (13 more authors) (2017) Towards broad spectrum activity-based glycosidase probes: synthesis and evaluation of deoxygenated cyclophellitol aziridines. Chemical Communications. pp. 1-5. ISSN: 1364-548X

<https://doi.org/10.1039/C7CC07730K>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. P. Schröder, J. W. van de Sande, W. W. Kallemeijn, C. Kuo, M. Artola, E. J. van Rooden, J. Jiang, T. J. Beenakker, B. I. Florea, W. A. Offen, G. Davies, A. J. Minnaard, H. Aerts, J. D. Codee, G. van der Marel and H. S. Overkleef, *Chem. Commun.*, 2017, DOI: 10.1039/C7CC07730K.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Journal Name

COMMUNICATION

Towards broad spectrum activity-based glycosidase probes: synthesis and evaluation of deoxygenated cyclophellitol aziridines

Sybrin P. Schröder,^a Jasper W. van de Sande,^a Wouter W. Kallemeijn,^b Chi-Lin Kuo,^b Marta Artola,^a Eva J. van Rooden,^a Jianbing Jiang,^a Thomas J. M. Beenakker,^a Bogdan I. Florea,^a Wendy A. Offen,^c Gideon J. Davies,^c Adriaan J. Minnaard,^d Johannes M. F. G. Aerts,^b Jeroen D. C. Codée,^a Gijsbert A. van der Marel^a and Herman S. Overkleeft^{a*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Activity-based protein profiling has emerged as a powerful tool for visualizing glycosidases in complex biological samples. Several configurational cyclophellitol isomers have been shown to display high selectivity as probes for glycosidases processing substrates featuring the same configuration. Here, a set of deoxygenated cyclophellitols are presented which enable inter-class profiling of β -glucosidases and β -galactosidases.

Glycosidases are hydrolytic enzymes that catalyze the hydrolysis of oligosaccharides and glycoconjugates.¹ Glycosidases are involved in a broad array of human pathologies and are therefore promising therapeutic targets.² Activity-based protein profiling (ABPP) provides an effective tool for the analysis of enzyme composition, expression and activity, and can assist in the discovery of enzyme inhibitors.³ ABPP relies on the addition of a mechanism-based inhibitor to the sample of interest, which reacts with the catalytic machinery of the target enzyme in a covalent and irreversible manner. The inhibitor carries a reporter tag and is termed an activity-based probe (ABP). As the first example on the development of ABPP, Cravatt and co-workers have described irreversible fluorophosphonate based ABPs which display high reactivity towards a wide range of serine hydrolases (SH).⁴ These ABPs were used to reveal active SH composition and expression level in various tissues, and as well in the development of highly specific and potent inhibitors against, for instance, monoacylglycerol lipase in a competitive ABPP setting.⁵ Cyclophellitol (**1**, Fig. 1), first isolated from *Phellinus* sp. is a covalent irreversible retaining β -glucosidase inhibitor.⁶




Fig. 1. (a) Structures of cyclophellitols **1–3**, selective covalent and irreversible retaining glycosidase inhibitors. (b) Selective β -glucosidase ABPs **4–5**, 4-deoxy ABPs **6–7** and 2,4-deoxy ABPs **8–9**.

The configurational isomers, *galacto*-cyclophellitol (**2**)⁷ and *manno*-cyclophellitol (**3**)⁸ are inactivators for β -galactosidases and β -mannosidases, respectively, and their selectivity arises from their absolute configurations. We have previously reported several cyclophellitol based ABPs for selective glycosidase profiling.^{9–11} For example, treatment of mouse tissue lysates with acyl ABP **4** resulted in tissue-specific fluorescent labelling of the three retaining β -glucocerebrosidases (GBA1, 2 and 3) and lactase-phlorizin hydrolase (LPH) without non-specific cross-labelling. Later studies revealed that alkyl ABP **5** also effectively labels these retaining β -glucosidases.¹² While the selectivity of these probes is an attractive value, it also impedes the study of multiple enzyme classes with a single probe, as for example can be done by the broad-spectrum SH ABPs developed by Cravatt and co-workers.⁴ Therefore, the development of broad spectrum glycosidase ABPs would be of interest. Since the substrate specificity of glucosidases, mannosidases and galactosidases is dependent on the carbohydrate hydroxyl configuration at C2 and C4 (axial or equatorial), removal of these alcohols may abolish active site preference. We therefore synthesized four deoxygenated ABPs **6–9** and we set out to establish whether these compounds would be effective

^a Department of Bioorganic Synthesis, Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands.

^b Department of Medical Biochemistry, Leiden Institute of Chemistry.


^c Department of Chemistry, York Structural Biology Laboratory, The University of York, Heslington, York, YO10 5DD, U.K.

^d Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands.

* Corresponding author. E-mail: h.s.overkleeft@chem.leidenuniv.nl

Electronic Supplementary Information (ESI) available: [experimental procedures and compound characterization, SDS-page images, proteomics and kinetics data]. See DOI: 10.1039/x0xx00000x





Scheme 1 The synthesis of deoxygenated ABPs required two different synthetic strategies (**a** and **b**) to afford key intermediates **12** and **15**, which were further functionalized to ABPs **6-9** following similar synthetic transformations (**c**). Reagents and conditions i) PMBCl, 2-aminoethyl diphenylborinate (10 mol%), K₂CO₃, KI, MeCN, 60 °C, 95%; ii) TSCl, pyridine, 60 °C, 90%; iii) LiAlH₄, THF, reflux, 86%; iv) HCl (10 %), DCM/HMPf, 86%; v) (+)-Ipc₂B(allyl)borane, THF, -90 °C, 2. Grubbs II (5 %), DCM, reflux, 77%, *dr* 86:14; vi) BzCl, Et₃N, DMAP, DCM, 84%; vii) CSA, MeOH, DCM, 94%.

ABPs indiscriminate towards glucosidases and galactosidases (and for **8-9**, mannosidases).

The synthesis of compounds **6-9** was achieved from key-intermediates **12** and **15**, which were prepared as follows. The primary alcohol in diol **10** (available by chemistry developed by Madsen and co-workers, Scheme 1a)¹³ was protected as the *p*-methoxybenzyl ether (PMB) after which tosylation of the secondary alcohol afforded intermediate **11**. Treatment of **11** with lithium aluminum hydride followed by deprotection afforded **12**. For cyclohexene **15** (Scheme 1b), chiral aldehyde **13** (available by chemistry developed by Helmchen and co-workers)¹⁴ was asymmetrically allylated using Brown's reagent and subsequently subjected to ring-closing metathesis using the Grubbs 2nd generation catalyst to give **14**. Benzoylation and detritylation afforded cyclohexene **15**. Both cyclohexenes **12** and **15** could be reacted with trichloroacetonitrile to afford the corresponding primary imidates (Scheme 1c), which upon treatment with *N*-iodosuccinimide (NIS) cyclized to afford the cyclic imidates **16** and **19** stereospecifically. Acidolysis and base treatment resulted in formation of aziridines **17** and **20**, which were alkylated with an azidoalkyl spacer and deprotected to give alkyl aziridines **18** and **21**, and then ligated with a reporter tag using click chemistry to afford deoxygenated ABPs **6-9** (see ESI†).

The labelling efficiency and specificity towards recombinant β -glucosidases, β -galactosidases and β -mannosidases was investigated for fluorescent ABPs **6** and **8**, and compared to that of β -glucosidase selective probe **5**. After enzyme incubation with a range of ABP concentrations (30 minutes, 37 °C, optimal enzyme pH), recombinant *Homo sapiens* GBA1 is highly efficiently labelled by ABP **5**, with protein bands




Fig. 2. Fluorescent labelling of recombinant *H. sapiens* GBA1 and *C. japonicus* β -galactosidase with ABPs **5**, **6** and **8** (left panel), and competition experiments with selective cytophellitins **1** or **2** (right panel). (a) GBA1 is potentially labelled by **5** and **6**, but not **8**. All labelling was activity-based, indicated by competition experiments with GBA1 selective inhibitor **1**. (b) β -galactosidase is potentially labelled by **5**, in an activity-based fashion.

visualized at probe concentrations as low as 10 nM (Fig. 2a). Additionally, 4-deoxy ABP **6** also labelled GBA1 efficiently, although the potency was slightly reduced, likely due to lacking hydrogen-bonding interactions within the active site residue(s) as the result of the absence of OH-4. The labelling potency of 2,4-deoxy ABP **8** proved to be drastically reduced. All bands could be competed with selective β -glucosidase inactivator **1**. Also, denatured enzyme (1% SDS, 100 °C, 5 min) is not labelled by any of the probes indicating that all labelling is activity-based. As expected, binding kinetics studies revealed that **5** is a far more potent GBA1 inactivator ($k_{\text{inact}}/K_i = 27.51 \pm 0.85 \mu\text{M}^{-1}\text{min}^{-1}$) than **6** ($k_{\text{inact}}/K_i = 0.14 \pm 0.08 \mu\text{M}^{-1}\text{min}^{-1}$). Kinetic parameters for **8** could not be determined under our assay conditions reflecting its low potency. Labelling of *Cellvibrio japonicus* β -galactosidase from glycoside hydrolase family 35 (CjGH35A)¹⁵ could be detected at micromolar concentrations of **5** and **8**. Interestingly, 4-deoxy ABP **6** was significantly more potent and labelled the enzyme at concentrations down to 100 nM, similar to GBA1 labelling; which is surprising given the extensive interactions of O-4, from a *galacto*-cyclophellitol aziridine, with the enzyme (see ESI†). Indeed, kinetic parameters for **6** in CjGH35A ($k_{\text{inact}}/K_i = 0.16 \pm 0.07 \mu\text{M}^{-1}\text{min}^{-1}$) are comparable to that in GBA1 (kinetic parameters of **5** and **8** could not be determined under our assay conditions, due to low potency). Competition experiments indicated the labelling was activity-based, although full competition was not achieved which could indicate a fraction of non-specific binding. Lastly, we found that none of the ABPs showed significant labelling of GH2 β -mannosidase from *Helix pomatia* (see ESI†). Indeed, it is known that OH-2 is highly important for coordination to the glycosidase catalytic nucleophile and stabilizing the transition state (TS).¹⁶ Moreover, probes **5**, **6** and **8** mimic preferentially adopt a ⁴H₃ configuration, which is characteristic for β -glucosidase and β -galactosidase transition states, whereas β -mannosidases often favour a B_{2,5} TS.¹⁷ Next, the glycosidase profiling by probes **6** and **8** in lysates from mouse (C57bl/6j, Jackson's laboratories) kidney and liver tissue, and human wild-type fibroblast cells (Lonza) was investigated, and compared to that of β -glucosidase selective probe **5**. The samples were incubated with the ABPs under identical

conditions (1 μ M ABP, pH 5.0, 37 $^{\circ}$ C, 30 min). In mouse liver lysates, **5** labels a distinct band at \sim 65 kDa (Fig. 3a), corresponding to the molecular weight of GBA1 (58 - 66 kDa¹⁸), and this band is fully abolished when the sample was pre-incubated with GBA1 specific inhibitor **1**.¹⁹ Similarly, incubation with **6** resulted in a band at the same height. This band could only be partially competed with **1**, and pre-incubation by specific β -galactosidase inactivator **2** resulted in partial competition as well. Ultimately, full competition was achieved by pre-incubation with both cyclophellitols **1** and **2**. Thus, in contrast to labelling with **5**, it was concluded that incubation with **6** resulted in the concomitant labelling of GBA1 and a β -galactosidase, presumably acid β -galactosidase (GLB1, \sim 63 kDa).²⁰ To identify the bands labelled on gel, the proteins were enriched by incubation of the lysate with biotin probe **7** and subsequent pull-down with streptavidin beads. After tryptic digestion, the peptide fragments were identified by LC-MS/MS and the abundance of the protein hits was quantified as previously described,²¹ in unsupervised mode using the default settings of the PLGS (Waters) and IsoQuant software. Indeed, incubation with **7** resulted in enrichment of GBA1 and GLB1, corresponding to the mixed band at \sim 65 kDa on gel. In contrast, labelling with ABP **8** did not result in labelling of GBA1 or GLB1. Multiple bands were visible on gel, but no competition was observed except for one band at \sim 115 kDa of an unknown protein. We therefore conclude that the other bands were a result of non-specific binding. Regardless, the lysate was incubated with **9** and the labelled proteins were identified by proteomic analysis. Whereas no enrichment of GBA1 could be detected, the probe did show increased levels of GLB1 labelling. Additionally, LC-MS/MS analysis detected enrichment of β -galactosylceramidase (GALC), an 80 kDa protein which is processed in the lysosomes into two catalytically active subunits of \sim 30 and \sim 50 kDa,²² and is responsible for the hydrolysis of galactosylceramide. Interestingly, fluorescent labelling with **8** did not identify GALC on gel. However, it is known that expression levels of GALC in the liver are low.²³

In mouse kidney lysates (Fig. 3b), **5** only labels GBA1 minimally. In contrast, bands at \sim 65 kDa and \sim 45 kDa were observed with probe **6** which could be fully competed out by pre-incubation with **2**, but not **1**. This indicated the labelling of mainly β -galactosidases in this sample, presumably GLB1 and GALC, which was subsequently confirmed by proteomics experiments as described above. Enrichment of GBA1 was also detected but less pronounced, in line with the fluorescent assay. Labelling with probe **6** was also observed for GLB1-like protein 1 (GLB1L, \sim 68 kDa) and 2 (GLB1I2, processed mass unknown), which both possess the GLB1 active site motif.²⁴ While 2,4-deoxy ABP **8** did not significantly label GBA1 or GLB1, proteomics identified GLB1 and GLB1I2 by pull-down with **9**. Interestingly, a strong band at \sim 45 kDa was labelled, which could be competed with **2**. Indeed, proteomics revealed increased labelling of GALC by this probe and it was therefore concluded that **8** is a GALC specific ABP.

Lastly, the labelling of human fibroblast lysates was investigated (Fig. 3c). A pattern of bands between 55-70 kDa is




Fig. 3 Fluorescent labelling of mouse liver (a), kidney (b) and human fibroblast (c) lysates with probes **5**, **6** and **8**, complemented with competition experiments with β -glucosidase specific covalent inhibitor **1** and β -galactosidase specific **2**. The bar charts display the LC-MS/MS pull-down analyses with the biotinylated ABP analogues corresponding to the gels on the left, compared to the negative control (DMSO). (a) Probe **5** only labels GBA1, while **6** labels GBA1 and GLB1, and **8** only shows non-specific labelling of proteins in gel, although pull-down analysis shows enrichment of GLB1 and GALC with **9**. (b) Only low amounts of GBA1 could be detected by specific probe **5**. 4-Deoxy probe **6** labels GBA1, GLB1 (like) protein and GALC. ABP **8** shows specific fluorescent labelling of GALC on gel. (c) ABP **5** labels GBA1 only, while **6** labels both GBA1 and GLB1. ABP **8** does not significantly label proteins, however pull-down analysis shows enrichment of GLB1.

labelled by **5** which can subsequently be fully competed for with **1**, indicating the labelling of different isoforms¹⁸ of GBA1 in this lysate. Similarly, **6** labels bands between 55-70 kDa, however with increased signal intensity. Approximately 50% competition was achieved by pre-incubation with either **1** or **2**, whereas pre-incubation with both inhibitors completely abrogated labelling. Indeed, pull-down analysis identified both GBA1 and GLB1 enrichment by probe **7**. Finally, **8** did not show



significant labelling of proteins in the sample, whereas GLB1 was enriched by ABP **9** according to LC-MS/MS analysis. In summary, we have developed syntheses towards a set of deoxygenated probes which were modelled on cyclophellitol aziridine **5**. None of the probes display activity towards β -mannosidases. Deoxygenation at C2 and C4 (ABPs **8-9**) enables labelling of purified β -glucosidases and β -galactosidases, albeit with low potency. Nevertheless, it was found that **8** displayed unexpected specificity towards β -galactosylceramidase in mouse kidney lysate where the enzyme is abundant.²³ Deoxygenation at C4 (ABPs **6-7**) affects the binding kinetics for GBA1, however *in vitro* labelling proceeded with high efficiency and moreover these probes now identify retaining β -galactosidases as well. Based on the results presented herein we conclude that, indeed, probes designed for simultaneously profiling retaining glycosidases processing differently configured glycosides are within reach. We had earlier observed that probes designed for a specific class of retaining glycosidases may possess cross-reactivity towards other glycosidases,¹⁰ however these results could not be predicted. Here, we show that by substitution of the hydroxyl group that distinguishes glucose from galactose for hydrogen on an otherwise unaltered cyclophellitol aziridine yields probes labelling both glucosidases and galactosidases. The strategy is however not general, since stripping off OH-2 yields a largely inactive probe. Future research on differently configured, OH-deleted cyclophellitol aziridines is required to establish if and to what extent this strategy is viable to deliver a set of broad-spectrum glycosidase ABPs, for use in combination with the specific exo-glycosidase probes already in hand for dissecting large numbers of retaining glycosidases from various species. We thank The Netherlands Organization for Scientific Research (NWO-CW, ChemThem grant to JMFGA and HSO), the European Research Council (ERC-2011-AdG-290836 "Chembiosphing" to HSO, and ERC-2012-AdG-322942 "Glycopoise", to GJD and WAO), Sanofi Genzyme (research grant to JMFGA and HSO and postdoctoral contract to MA), ChemAxon for the Instant JChem software and Diamond Light Source for access to beamline I02 (proposal number mx-9948) that contributed to the results presented here. GJD is a Royal Society Ken Murray Research Professor.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- G. Davies and B. Henrissat, *Structure*, 1995, **3**, 853–859.
- K. Ohtsubo and J. D. Marth, *Cell*, 2006, **126**, 855–867.
- M. J. Evans and B. F. Cravatt, *Chem. Rev.*, 2006, **106**, 3279.
- Y. Liu, M. P. Patricelli and B. F. Cravatt, *Proc. Natl. Acad. Sci.*, 1999, **96**, 14694–14699.
- J. Z. Long, W. Li, L. Booker, J. J. Burston, S. G. Kinsey, J. E. Schlosburg, F. J. Pavón, A. M. Serrano, D. E. Selley, L. H. Parsons, A. H. Lichtman and B. F. Cravatt, *Nat. Chem. Biol.*,

- 2008, **5**, 37–44.
- S. Atsumi, K. Umezawa, H. Iinuma, H. Naganawa, H. Nakamura, Y. Iitaka and T. Takeuchi, *J. Antibiot. (Tokyo)*, 1989, **43**, 49–53.
- L. I. Willems, T. J. M. Beenakker, B. Murray, B. Gagestein, H. Van Den Elst, E. R. Van Rijssel, J. D. C. Codée, W. W. Kallemeyn, J. M. F. G. Aerts, G. A. Van Der Marel and H. S. Overkleeft, *European J. Org. Chem.*, 2014, 6044.
- V. W. F. T. Tony K. M. Shing, *J. Chem. Soc. Perkin Trans. 1*, 1994, 2017–2025.
- W. W. Kallemeyn, K. Y. Li, M. D. Witte, A. R. A. Marques, J. Aten, S. Scheij, J. Jiang, L. I. Willems, T. M. Voorn-Brouwer, C. P. A. A. Van Roomen, R. Ottenhoff, R. G. Boot, H. Van Den Elst, M. T. C. Walvoort, B. I. Florea, J. D. C. Codée, G. A. Van Der Marel, J. M. F. G. Aerts and H. S. Overkleeft, *Angew. Chemie - Int. Ed.*, 2012, **51**, 12529–12533.
- J. Jiang, C. L. Kuo, L. Wu, C. Franke, W. W. Kallemeyn, B. I. Florea, E. Van Meel, G. A. Van Der Marel, J. D. C. Codée, R. G. Boot, G. J. Davies, H. S. Overkleeft and J. M. F. G. Aerts, *ACS Cent. Sci.*, 2016, **2**, 351–358.
- L. I. Willems, T. J. M. Beenakker, B. Murray, S. Scheij, W. W. Kallemeyn, R. G. Boot, M. Verhoek, W. E. Donker-Koopman, M. J. Ferraz, E. R. Van Rijssel, B. I. Florea, J. D. C. Codée, G. A. Van Der Marel, J. M. F. G. Aerts and H. S. Overkleeft, *J. Am. Chem. Soc.*, 2014, **136**, 11622–11625.
- J. Jiang, T. J. M. Beenakker, W. W. Kallemeyn, G. A. Van Der Marel, H. Van Den Elst, J. D. C. Codée, J. M. F. G. Aerts and H. S. Overkleeft, *Chem. - A Eur. J.*, 2015, **21**, 10861.
- F. G. Hansen, E. Bundgaard and R. Madsen, *J. Org. Chem.*, 2005, **70**, 10139–10142.
- G. Franck, K. Brödner and G. Helmchen, *Org. Lett.*, 2010, **12**, 3886–9.
- J. Larsbrink, A. J. Thompson, M. Lundqvist, J. G. Gardner, G. J. Davies and H. Brumer, *Mol. Microbiol.*, 2014, **94**, 418.
- S. G. Withers, *Carbohydr. Polym.*, 2001, **44**, 325–337.
- G. J. Davies, A. Planas and C. Rovira, *Acc. Chem. Res.*, 2012, **45**, 308–316.
- S. van Weely, J. M. F. G. Aerts, M. B. van Leeuwen, J. C. Heikoop, W. E. Donker-Koopman, J. A. Barranger, J. M. Tager and A. W. Schram, *Eur. J. Biochem.*, 1990, **191**, 669–677.
- M. D. Witte, W. W. Kallemeyn, J. Aten, K.-Y. Li, A. Strijland, W. E. Donker-Koopman, A. M. C. H. van den Nieuwendijk, B. Bleijlevens, G. Kramer, B. I. Florea, B. Hooibrink, C. E. M. Hollak, R. Ottenhoff, R. G. Boot, G. A. van der Marel, H. S. Overkleeft and J. M. F. G. Aerts, *Nat. Chem. Biol.*, 2010, **6**, 907–913.
- S. Tomino and M. Meisler, *J. Biol. Chem.*, 1975, **250**, 7752.
- J. Kuharev, P. Navarro, U. Distler, O. Jahn and S. Tenzer, *Proteomics*, 2015, **15**, 3140–3151.
- S. Nagano, T. Yamada, N. Shinnoh, H. Furuya, T. Taniwaki and J. I. Kira, *Clin. Chim. Acta*, 1998, **276**, 53–61.
- P. Luzi, M. A. Rafi, M. Zaka, M. Curtis, M. T. Vanier and D. A. Wenger, *Mol. Genet. Metab.*, 2001, **73**, 211–23.
- J. Le Carré, D. F. Schorderet and S. Cottet, *Mol. Vis.*, 2011, **17**, 1287–97.

