



This is a repository copy of *Criteria for a ring to have a left Noetherian left quotient ring*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/124642/>

Version: Accepted Version

Article:

Bavula, V.V. (2018) Criteria for a ring to have a left Noetherian left quotient ring. *Journal of Pure and Applied Algebra*, 222 (7). pp. 1548-1564. ISSN 0022-4049

<https://doi.org/10.1016/j.jpaa.2017.07.011>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Criteria for a ring to have a left Noetherian left quotient ring

V. V. Bavula

Abstract

Two criteria are given for a ring to have a left Noetherian left quotient ring (to find a criterion was an open problem since 70's). It is proved that each such ring has only *finitely many* maximal left denominator sets.

Key Words: Goldie's Theorem, the left quotient ring of a ring, the largest left quotient ring of a ring, a maximal left denominator set, the prime radical.

Mathematics subject classification 2010: 16P50, 16P60, 16P20, 16U20.

Contents

1. Introduction.
2. Properties of rings with a left Noetherian left quotient ring.
3. Proof of Theorem 1.2. and the Second Criterion (via the associated graded ring).
4. Finiteness of $\max.\text{Den}_l(R)$ when $Q(R)$ is a left Noetherian ring.

1 Introduction

In this paper, module means a left module, and the following notation is fixed:

- R is a ring with 1;
- $\mathcal{C} = \mathcal{C}_R$ is the set of *regular* elements of the ring R (i.e., \mathcal{C} is the set of non-zero-divisors of the ring R);
- $Q = Q(R) := Q_{l,cl}(R) := \mathcal{C}^{-1}R$ is the *left quotient ring* (the *classical left ring of fractions*) of the ring R (if it exists) and Q^* is the group of units of Q ;
- $\mathfrak{n} = \mathfrak{n}_R$ is the prime radical of R , $\nu \in \mathbb{N} \cup \{\infty\}$ is its *nilpotency degree* ($\mathfrak{n}^\nu \neq 0$ but $\mathfrak{n}^{\nu+1} = 0$) and $\mathcal{N}_i := \mathfrak{n}^i / \mathfrak{n}^{i+1}$ for $i \in \mathbb{N}$;
- $\bar{R} := R/\mathfrak{n}$ and $\pi : R \rightarrow \bar{R}$, $r \mapsto \bar{r} = r + \mathfrak{n}$;
- $\bar{\mathcal{C}} := \mathcal{C}_{\bar{R}}$ is the set of regular elements of the ring \bar{R} and $\bar{Q} := \bar{\mathcal{C}}^{-1}\bar{R}$ is its left quotient ring;
- $\tilde{\mathcal{C}} := \pi(\mathcal{C})$, $\tilde{Q} := \tilde{\mathcal{C}}^{-1}\bar{R}$ and $\mathcal{C}^\dagger := \mathcal{C}_{\tilde{Q}}$ is the set of regular elements of the ring \tilde{Q} ;
- $S_l = S_l(R)$ is the *largest left Ore* of R that consists of regular elements and $Q_l = Q_l(R) := S_l(R)^{-1}R$ is the *largest left quotient ring* of R [5, Theorem 2.1];
- $\text{Ore}_l(R) := \{S \mid S \text{ is a left Ore set in } R\}$;
- $\text{Den}_l(R) := \{S \mid S \text{ is a left denominator set in } R\}$.

A question of describing rings such that their (left) quotient ring satisfies certain conditions tends to be a challenging one. A first step was done by Goldie (1960) [12] who gave a criterion for a ring R to have a semisimple Artinian quotient ring $Q(R)$. The case when $Q(R)$ is a simple Artinian ring is due to Goldie (1960) [10] and Lesieur and Croisot (1958) [15]. Criteria for a ring

to have a left Artinian left quotient ring were given by Small (1966) [20], [21]; Robson (1967) [19], Tachikawa (1971) [25], Hajarnavis (1972) [14], Warfield (1981) [26] and the author (2013) [2].

Remark. In the paper, a statement that ‘If $R \dots$ then $Q(R) \dots$ ’ means that ‘If R satisfies \dots then $Q(R)$ **exists** and satisfies \dots ’.

Criteria for a ring to have a left Noetherian left quotient ring. The aim of the paper is to give two criteria for a ring R to have a left Noetherian left quotient ring (Theorem 1.2 and Theorem 1.3). The case when R is a *semiprime* ring is a very easy special case.

Theorem 1.1 *Let R be a semiprime ring. Then the following statements are equivalent.*

1. $Q(R)$ is a left Noetherian ring.
2. $Q(R)$ is a semisimple ring.
3. R is a semiprime left Goldie ring.
4. $Q_l(R)$ is a left Noetherian ring.
5. $Q_l(R)$ is a semisimple ring.

If one of the equivalent conditions holds then $S_l(R) = \mathcal{C}_R$ and $Q(R) = Q_l(R)$. In particular, if the left quotient ring $Q(R)$ (respectively, $Q_l(R)$) is not a semisimple ring then the ring $Q(R)$ (respectively, $Q_l(R)$) is not left Noetherian.

The proof of Theorem 1.1 is given in Section 3.

Example, [1]. The ring $\mathbb{I}_1 := K\langle x, \frac{d}{dx}, f \rangle$ of polynomial integro-differential operators over a field K of characteristic zero is a semiprime ring but not left Goldie (as it contains infinite direct sums of non-zero left ideals). Therefore, the largest left quotient ring $Q_l(\mathbb{I}_1)$ is not a left Noetherian ring (moreover, the left quotient ring $Q(\mathbb{I}_1)$ does not exist). The ring $Q_l(\mathbb{I}_1)$ and the largest regular left Ore set $S_l(\mathbb{I}_1)$ of \mathbb{I}_1 were described explicitly in [1].

The first criterion for a ring to have a left Noetherian left quotient ring is below, its proof is given in Section 3.

Theorem 1.2 *Let R be a ring. The following statements are equivalent.*

1. The ring R has a left Noetherian left quotient ring $Q(R)$.
2. (a) $\tilde{\mathcal{C}} \subseteq \bar{\mathcal{C}}$.
 (b) $\tilde{\mathcal{C}} \in \text{Ore}_l(\bar{R})$.
 (c) $\tilde{Q} = \tilde{\mathcal{C}}^{-1}\bar{R}$ is a left Noetherian ring.
 (d) \mathfrak{n} is a nilpotent ideal of the ring R .
 (e) The \tilde{Q} -modules $\tilde{\mathcal{C}}^{-1}\mathcal{N}_i$, $i = 1, \dots, \nu$, are finitely generated (where ν is the nilpotency degree of \mathfrak{n} and $\mathcal{N}_i := \mathfrak{n}^i/\mathfrak{n}^{i+1}$).
 (f) For each element $\bar{c} \in \tilde{\mathcal{C}}$, the left \bar{R} -module $\mathcal{N}_i/\mathcal{N}_i\bar{c}$ is $\tilde{\mathcal{C}}$ -torsion for $i = 1, \dots, \nu$.

Remark. The conditions (a) and (b) above imply that the set $\tilde{\mathcal{C}}$ is a left denominator set of the ring R , and so the ring \tilde{Q} exists.

The powers of the prime radical \mathfrak{n} , $\{\mathfrak{n}^i\}_{i \geq 0}$, form a descending filtration on R . Let $\text{gr } R = \bar{R} \oplus \mathfrak{n}/\mathfrak{n}^2 \oplus \dots$ be the associated graded ring. The second criterion is given in terms of properties of the ring $\text{gr } R$, its proof is given in Section 3.

Theorem 1.3 *Let R be a ring. The following statements are equivalent.*

1. *The ring R has a left Noetherian left quotient ring $Q(R)$.*
2. *The set $\tilde{\mathcal{C}}$ is a left denominator set of the ring $\text{gr } R$, $\tilde{\mathcal{C}} \subseteq \bar{\mathcal{C}}$, $\tilde{\mathcal{C}}^{-1}\text{gr } R$ is a left Noetherian ring and \mathfrak{n} is a nilpotent ideal.*

If one of the equivalent conditions holds then $\text{gr } Q(R) \simeq \tilde{\mathcal{C}}^{-1}\text{gr } R$ where $\text{gr } Q(R) := \tilde{Q} \oplus \mathfrak{n}_Q / \mathfrak{n}_Q^2 \oplus \dots$ is the associated graded ring with respect to the prime radical filtration. In particular, the ring $\text{gr } Q(R)$ is a left Noetherian ring.

Finiteness of the set of maximal left denominators for a left Noetherian ring. For a ring R , the set $\text{max.Den}_l(R)$ of maximal left denominator sets (with respect to \subseteq) is a *non-empty* set, [5]. It was proved that the set $\text{max.Den}_l(R)$ is a finite set if the left quotient ring $Q(R)$ of R is a semisimple ring, [3], or a left Artinian ring, [6]. The next theorem extends this result for a larger class of rings that includes the class of left Noetherian rings for which the left quotient ring exists.

Theorem 1.4 *Let R be a ring with a left Noetherian left quotient ring $Q(R)$. Then $|\text{max.Den}_l(R)| < \infty$. Moreover, $|\text{max.Den}_l(R)| \leq s = |\text{max.Den}_l(\bar{R})|$ where $\bar{Q} \simeq \prod_{i=1}^s \bar{Q}_i$ and \bar{Q}_i are simple Artinian rings (see Theorem 2.4.(3)).*

The next corollary follows at once from Theorem 1.4.

Corollary 1.5 *If R is a left Noetherian ring such that its left quotient ring $Q(R)$ exists (i.e., $\mathcal{C} \in \text{Ore}_l(R)$) then $|\text{max.Den}_l(R)| < \infty$.*

In Corollary 1.5, the condition that ‘its left quotient ring $Q(R)$ exists’ is redundant.

Theorem 1.6 ([7].) *If R is a left Noetherian ring then $|\text{max.Den}_l(R)| < \infty$.*

The next corollary is an explicit description of the set $\text{max.Den}_l(R)$ for a ring R with a left Noetherian left quotient ring $Q(R)$.

Corollary 1.7 *Let R be a ring with a left Noetherian left quotient ring $Q(R)$. For each $i = 1, \dots, s$, let $p_i : R \rightarrow \bar{Q}_i$ be the natural projection (see (8) and Theorem 1.4), \bar{Q}_i^* be the group of units of the simple Artinian ring \bar{Q}_i , S'_i be the largest element (w.r.t. \subseteq) of the set $D_i = \{S' \in \text{Den}_l(R) \mid p_i(S') \subseteq \bar{Q}_i^*\}$. Then*

1. *$\text{max.Den}_l(R)$ is the set of maximal elements (w.r.t. \subseteq) of the set $\{S'_1, \dots, S'_s\}$.*
2. *For all $i = 1, \dots, s$, $\mathcal{C} \subseteq S'_i$.*
3. *The rings $S'_i{}^{-1}R$ are left Noetherian where $i = 1, \dots, s$.*

A criterion for a left Noetherian ring to have a left Noetherian left quotient ring. In [11], Goldie posed a problem of deciding when a Noetherian ring possesses a Noetherian quotient ring (i.e., when the set \mathcal{C}_R is a left and right Ore set). In [23], Stafford obtained a criterion to determine when a Noetherian ring is its own quotient ring. In [8], Chatters and Hajarnavis obtained necessary and sufficient conditions for a Noetherian ring which is a finite module over its centre to have a quotient ring.

The next corollary follows from Theorem 1.2, Theorem 1.3 and the fact that for a left Noetherian ring its the prime radical is a nilpotent ideal and $\tilde{\mathcal{C}} \subseteq \bar{\mathcal{C}}$ (see, [9, Lemma 11.8]).

Corollary 1.8 *Let R be a left Noetherian ring. The following statements are equivalent.*

1. *The set $\mathcal{C} = \mathcal{C}_R$ is a left Ore set (i.e., the left quotient ring $Q(R) = \mathcal{C}^{-1}R$ of R exists).*
2. *$\tilde{\mathcal{C}} \in \text{Ore}_l(\bar{R})$ and for each element $\bar{c} \in \tilde{\mathcal{C}}$ the left \bar{R} -module $N_i/N_i\bar{c}$ is $\tilde{\mathcal{C}}$ -torsion for $i = 1, \dots, \nu$.*

3. $\tilde{\mathcal{C}} \in \text{Den}_l(\text{gr } R)$.

Proof. (1 \Leftrightarrow 2) and (1 \Leftrightarrow 3) follow from Theorem 1.2 and Theorem 1.3, respectively. \square

Corollary 2.5 is a criterion for a ring to have a left Noetherian left quotient ring Q such that the factor ring Q/\mathfrak{n}_Q is a semisimple ring (or \tilde{Q} is a semisimple ring; or $\tilde{\mathcal{C}} = \bar{\mathcal{C}}$; or $\mathcal{C} = \pi^{-1}(\bar{\mathcal{C}})$). Theorem 4.2 is a criterion for a ring R to have a left Noetherian ring such that $|\text{max.Den}_l(R)| = |\text{max.Den}_l(\bar{R})|$ (recall that, in general, $|\text{max.Den}_l(R)| \leq |\text{max.Den}_l(\bar{R})|$, Theorem 1.4).

The set of maximal denominator sets for a commutative ring. In the commutative situation, the following concepts are identical: a multiplicatively closed set, an Ore set and a denominator set. For a commutative ring, the next proposition describes the set of maximal denominator sets (i.e., maximal Ore sets, i.e., maximal multiplicatively closed sets).

Proposition 1.9 *Let R be a commutative ring and $\text{Min}(R)$ be the set of minimal prime ideals of the ring R . Then $\text{max.Den}(R) = \{S_{\mathfrak{p}} := R \setminus \mathfrak{p} \mid \mathfrak{p} \in \text{Min}(R)\}$.*

The paper is organized as follows. In Section 2, many properties of a ring R with a left Noetherian left quotient ring are proven (Theorem 2.4). In particular, the implication (1 \Rightarrow 2) of Theorem 2.4 is proven.

In Section 3, proofs of Theorem 1.1, Theorem 1.2 and Theorem 1.3 are given. In Section 4, Theorem 1.4 is proven.

2 Properties of rings with a left Noetherian left quotient ring

In this section, we establish many properties of rings with left Noetherian left quotient ring (Theorem 2.4). In particular, the implication (1 \Rightarrow 2) of Theorem 1.2 is proven (Theorem 2.4.(2)). A criterion is given (Corollary 2.5) for the ring \tilde{Q} to be a semisimple ring or a left Artinian ring.

At the beginning of the section, we collect necessary results that are used in the proofs of this paper. More results on localizations of rings (and some of the missed standard definitions) the reader can find in [13], [24] and [16]. In this paper the following notation will remain fixed:

- $\text{Den}_l(R, \mathfrak{a}) := \{S \in \text{Den}_l(R) \mid \text{ass}(S) = \mathfrak{a}\}$ where \mathfrak{a} is an ideal of R ;
- $S_{\mathfrak{a}} = S_{\mathfrak{a}}(R) = S_{l,\mathfrak{a}}(R)$ is the *largest element* of the poset $(\text{Den}_l(R, \mathfrak{a}), \subseteq)$ and $Q_{\mathfrak{a}}(R) := Q_{l,\mathfrak{a}}(R) := S_{\mathfrak{a}}^{-1}R$ is the *largest left quotient ring associated with \mathfrak{a}* . $S_{\mathfrak{a}}$ exists (Theorem 2.1, [5]);
- In particular, $S_0 = S_0(R) = S_{l,0}(R)$ is the largest element of the poset $(\text{Den}_l(R, 0), \subseteq)$ and $Q_l(R) := S_0^{-1}R$ is the *largest left quotient ring of R* , [5].

The largest regular left Ore set and the largest left quotient ring of a ring. Let R be a ring. A *multiplicatively closed subset* S of R or a *multiplicative subset* of R (i.e., a multiplicative sub-semigroup of (R, \cdot) such that $1 \in S$ and $0 \notin S$) is said to be a *left Ore set* if it satisfies the *left Ore condition*: for each $r \in R$ and $s \in S$, $Sr \cap Rs \neq \emptyset$. Let $\text{Ore}_l(R)$ be the set of all left Ore sets of R . For $S \in \text{Ore}_l(R)$, $\text{ass}(S) := \{r \in R \mid sr = 0 \text{ for some } s \in S\}$ is an ideal of the ring R .

A left Ore set S is called a *left denominator set* of the ring R if $rs = 0$ for some elements $r \in R$ and $s \in S$ implies $tr = 0$ for some element $t \in S$, i.e., $r \in \text{ass}(S)$. Let $\text{Den}_l(R)$ be the set of all left denominator sets of R . For $S \in \text{Den}_l(R)$, the ring $S^{-1}R = \{s^{-1}r \mid s \in S, r \in R\}$ is called the *left localization* of the ring R at S (the *left quotient ring* of R at S). In Ore's method of localization one can localize *precisely* at left denominator sets.

In general, the set \mathcal{C} of regular elements of a ring R is neither left nor right Ore set of the ring R and as a result neither left nor right classical quotient ring ($Q_{l,cl}(R) := \mathcal{C}^{-1}R$ and $Q_{r,cl}(R) :=$

$R\mathcal{C}^{-1}$) exists. There exists the largest (w.r.t. \subseteq) regular left Ore set $S_0 = S_{l,0} = S_{l,0}(R)$, [5]. This means that the set $S_{l,0}(R)$ is an Ore set of the ring R that consists of regular elements (i.e., $S_{l,0}(R) \subseteq \mathcal{C}$) and contains all the left Ore sets in R that consist of regular elements. Also, there exists the largest regular right (respectively, left and right) Ore set $S_{r,0}(R)$ (respectively, $S_{l,r,0}(R)$) of the ring R . In general, the sets \mathcal{C} , $S_{l,0}(R)$, $S_{r,0}(R)$ and $S_{l,r,0}(R)$ are distinct, for example, when $R = \mathbb{I}_1 = K\langle x, \partial, f \rangle$ is the ring of polynomial integro-differential operators over a field K of characteristic zero, [1]. In [1], these four sets are explicitly described for $R = \mathbb{I}_1$.

Definition, [1], [5]. The ring

$$Q_l(R) := S_{l,0}(R)^{-1}R$$

(respectively, $Q_r(R) := RS_{r,0}(R)^{-1}$ and $Q(R) := S_{l,r,0}(R)^{-1}R \simeq RS_{l,r,0}(R)^{-1}$) is called the *largest left* (respectively, *right and two-sided*) *quotient ring* of the ring R .

In general, the rings $Q_l(R)$, $Q_r(R)$ and $Q(R)$ are not isomorphic, for example, when $R = \mathbb{I}_1$, [1].

Small and Stafford [22] have shown that any (left and right) Noetherian ring R possesses a uniquely determined set of prime ideals P_1, \dots, P_n such that $\mathcal{C}_R = \bigcap_{i=1}^n \mathcal{C}(P_i)$, an irreducible intersection, where $\mathcal{C}(P_i) := \{r \in R \mid r + P_i \in \mathcal{C}_{R/P_i}\}$. Michler and Müller [17] mentioned that the ring R contains a unique maximal (left and right) Ore set of regular elements $S_{l,r,0}(R)$ and called the ring $Q(R)$ the *total quotient ring* of R . For certain Noetherian rings, they described the set $S_{l,r,0}(R)$ and the ring $Q(R)$. For the class of affine Noetherian PI-rings, further generalizations were given by Müller in [18].

The next theorem gives various properties of the ring $Q_l(R)$. In particular, it describes its group of units.

Theorem 2.1 ([5].)

1. $S_0(Q_l(R)) = Q_l(R)^*$ and $S_0(Q_l(R)) \cap R = S_0(R)$.
2. $Q_l(R)^* = \langle S_0(R), S_0(R)^{-1} \rangle$, i.e., the group of units of the ring $Q_l(R)$ is generated by the sets $S_0(R)$ and $S_0(R)^{-1} := \{s^{-1} \mid s \in S_0(R)\}$.
3. $Q_l(R)^* = \{s^{-1}t \mid s, t \in S_0(R)\}$.
4. $Q_l(Q_l(R)) = Q_l(R)$.

The maximal left denominator sets and the maximal left localizations of a ring.

The set $(\text{Den}_l(R), \subseteq)$ is a poset (partially ordered set). In [5], it is proved that the set $\text{max.Den}_l(R)$ of its maximal elements is a *non-empty* set.

Definition, [5]. An element S of the set $\text{max.Den}_l(R)$ is called a *maximal left denominator set* of the ring R and the ring $S^{-1}R$ is called a *maximal left quotient ring* of the ring R or a *maximal left localization ring* of the ring R . The intersection

$$\mathfrak{l}_R := \text{l.rad}(R) := \bigcap_{S \in \text{max.Den}_l(R)} \text{ass}(S) \quad (1)$$

is called the *left localization radical* of the ring R , [5].

For a ring R , there is a canonical exact sequence

$$0 \rightarrow \mathfrak{l}_R \rightarrow R \xrightarrow{\sigma} \prod_{S \in \text{max.Den}_l(R)} S^{-1}R, \quad \sigma := \prod_{S \in \text{max.Den}_l(R)} \sigma_S, \quad (2)$$

where $\sigma_S : R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$.

Properties of the maximal left quotient rings of a ring. For a ring R , let $\text{Ass}_l(R) := \{\text{ass}(S) \mid S \in \text{Den}_l(R)\}$. The next theorem describes various properties of the maximal left quotient rings of a ring, in particular, their groups of units and their largest left quotient rings.

Theorem 2.2 ([5].) *Let $S \in \text{max.Den}_l(R)$, $A = S^{-1}R$, A^* be the group of units of the ring A ; $\mathfrak{a} := \text{ass}(S)$, $\pi_{\mathfrak{a}} : R \rightarrow R/\mathfrak{a}$, $a \mapsto a + \mathfrak{a}$, and $\sigma_{\mathfrak{a}} : R \rightarrow A$, $r \mapsto \frac{r}{1}$. Then*

1. $S = S_{\mathfrak{a}}(R)$, $S = \pi_{\mathfrak{a}}^{-1}(S_0(R/\mathfrak{a}))$, $\pi_{\mathfrak{a}}(S) = S_0(R/\mathfrak{a})$ and $A = S_0(R/\mathfrak{a})^{-1}R/\mathfrak{a} = Q_l(R/\mathfrak{a})$.
2. $S_0(A) = A^*$ and $S_0(A) \cap (R/\mathfrak{a}) = S_0(R/\mathfrak{a})$.
3. $S = \sigma_{\mathfrak{a}}^{-1}(A^*)$.
4. $A^* = \langle \pi_{\mathfrak{a}}(S), \pi_{\mathfrak{a}}(S)^{-1} \rangle$, i.e., the group of units of the ring A is generated by the sets $\pi_{\mathfrak{a}}(S)$ and $\pi_{\mathfrak{a}}^{-1}(S) := \{\pi_{\mathfrak{a}}(s)^{-1} \mid s \in S\}$.
5. $A^* = \{\pi_{\mathfrak{a}}(s)^{-1}\pi_{\mathfrak{a}}(t) \mid s, t \in S\}$.
6. $Q_l(A) = A$ and $\text{Ass}_l(A) = \{0\}$. In particular, if $T \in \text{Den}_l(A, 0)$ then $T \subseteq A^*$.

A bijection between $\text{max.Den}_l(R)$ and $\text{max.Den}_l(Q_l(R))$. The next theorem shows that there is a canonical bijection between the maximal left denominator sets of a ring R and of its largest left quotient ring $Q_l(R)$.

Proposition 2.3 ([3].) *Let R be a ring, S_l be the largest regular left Ore set of the ring R , $Q_l := S_l^{-1}R$ be the largest left quotient ring of the ring R , and \mathcal{C} be the set of regular elements of the ring R . Then*

1. $S_l \subseteq S$ for all $S \in \text{max.Den}_l(R)$. In particular, $\mathcal{C} \subseteq S$ for all $S \in \text{max.Den}_l(R)$ provided \mathcal{C} is a left Ore set.
2. Either $\text{max.Den}_l(R) = \{\mathcal{C}\}$ or, otherwise, $\mathcal{C} \notin \text{max.Den}_l(R)$.
3. The map

$$\text{max.Den}_l(R) \rightarrow \text{max.Den}_l(Q_l), \quad S \mapsto SQ_l^* = \{c^{-1}s \mid c \in S_l, s \in S\},$$

is a bijection with the inverse $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$ where $\sigma : R \rightarrow Q_l$, $r \mapsto \frac{r}{1}$, and SQ_l^* is the sub-semigroup of (Q_l, \cdot) generated by the set S and the group Q_l^* of units of the ring Q_l , and $S^{-1}R = (SQ_l^*)^{-1}Q_l$.

4. If \mathcal{C} is a left Ore set then the map

$$\text{max.Den}_l(R) \rightarrow \text{max.Den}_l(Q), \quad S \mapsto SQ^* = \{c^{-1}s \mid c \in \mathcal{C}, s \in S\},$$

is a bijection with the inverse $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$ where $\sigma : R \rightarrow Q$, $r \mapsto \frac{r}{1}$, and SQ^* is the sub-semigroup of (Q, \cdot) generated by the set S and the group Q^* of units of the ring Q , and $S^{-1}R = (SQ^*)^{-1}Q$.

A ring R is called a *left Goldie ring* if it satisfies ACC (the *ascending chain condition*) for left annihilators and contains no infinite direct sums of left ideals.

Theorem 2.4 *Let R be a ring such that its left quotient ring Q is a left Noetherian ring. Let $\sigma : R \rightarrow Q$, $r \mapsto \frac{r}{1}$, $(Q/\mathfrak{n}_Q)^*$ be the group of units of the ring Q/\mathfrak{n}_Q , and $\sigma' : Q/\mathfrak{n}_Q \rightarrow Q(Q/\mathfrak{n}_Q)$, $q + \mathfrak{n}_Q \mapsto \frac{q}{1} + \mathfrak{n}_Q$ (the existence of $Q(Q/\mathfrak{n}_Q)$ follows from Goldie's Theorem using part 2(c) of the theorem). Then*

1. $\mathfrak{n} = R \cap \mathfrak{n}_Q$, $\mathcal{C}^{-1}\mathfrak{n} = \mathfrak{n}_Q$, $(\mathcal{C}^{-1}\mathfrak{n})^i = \mathcal{C}^{-1}\mathfrak{n}^i$ for all $i \geq 1$, and $\nu = \nu_Q < \infty$ where ν and ν_Q are the nilpotency degrees of the prime radicals \mathfrak{n} and \mathfrak{n}_Q , respectively.

2. (a) $\mathcal{C} + \mathfrak{n} \subseteq \mathcal{C}$.
 (b) $\tilde{\mathcal{C}} \in \text{Den}_l(\overline{R}, 0)$. In particular, $\tilde{\mathcal{C}} \subseteq \overline{\mathcal{C}}$.
 (c) $\tilde{Q} := \tilde{\mathcal{C}}^{-1}\overline{R} \simeq Q/\mathfrak{n}_Q$ is a semiprime left Noetherian ring.
 (d) \mathfrak{n} is a nilpotent ideal of the ring R .
 (e) The \tilde{Q} -modules $\tilde{\mathcal{C}}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1})$, $i = 1, \dots, \nu$, are finitely generated (where ν is the nilpotency degree of \mathfrak{n}).
 (f) For each elements $\bar{c} \in \overline{\mathcal{C}}$, the left \overline{R} -module $\mathcal{N}_i/\mathcal{N}_i\bar{c}$ is $\overline{\mathcal{C}}$ -torsion where $\mathcal{N}_i := \mathfrak{n}^i/\mathfrak{n}^{i+1}$.
3. The ring \overline{R} is a semiprime left Goldie ring and its left quotient ring $\overline{Q} := Q(\overline{R}) \simeq Q(Q/\mathfrak{n}_Q) \simeq Q(\tilde{Q})$ is a semisimple ring.
4. $1 \rightarrow 1 + \mathfrak{n}_Q \rightarrow Q^* \xrightarrow{\pi_Q^*} (Q/\mathfrak{n}_Q)^* \rightarrow 1$ is a short exact sequence of group homomorphisms where $\pi_Q : Q \rightarrow Q/\mathfrak{n}_Q$, $q \mapsto q + \mathfrak{n}_Q$ and $\pi_Q^* := \pi_Q|_{Q^*}$.
5. $\mathcal{C} = \sigma^{-1}(Q^*) = (\pi_Q\sigma)^{-1}((Q/\mathfrak{n}_Q)^*) = \pi^{-1}(\tilde{\sigma}^{-1}((Q/\mathfrak{n}_Q)^*))$ where $\tilde{\sigma} : \overline{R} \rightarrow Q/\mathfrak{n}_Q$, $\bar{r} \mapsto \frac{\bar{r}}{1} + \mathfrak{n}_Q$, see (4).
6. Let $\mathcal{C}^\dagger := \mathcal{C}_{\tilde{Q}}$, i.e., $\mathcal{C}^\dagger = \mathcal{C}_{Q/\mathfrak{n}_Q}$ when we identify the rings \tilde{Q} and Q/\mathfrak{n}_Q via the isomorphism in statement 2(c). Then $\mathcal{C}^\dagger = \sigma'^{-1}(Q(Q/\mathfrak{n}_Q)^*)$ and $\overline{\mathcal{C}} = \tilde{\sigma}^{-1}(\mathcal{C}^\dagger) = (\sigma'\tilde{\sigma})^{-1}(Q(Q/\mathfrak{n}_Q)^*)$.

Proof. 1. The prime radical \mathfrak{n}_Q is a nilpotent ideal since the ring Q is a left Noetherian ring. Then the intersection $R \cap \mathfrak{n}_Q$ is a nilpotent ideal of the ring R , hence $R \cap \mathfrak{n}_Q \subseteq \mathfrak{n}$. To prove that the equality $R \cap \mathfrak{n}_Q = \mathfrak{n}$ holds it suffices to show that the factor ring $R/R \cap \mathfrak{n}_Q$ has no nonzero nilpotent ideals. Suppose that \bar{I} is a nilpotent ideal of the ring $R/R \cap \mathfrak{n}_Q$ then its preimage I under the epimorphism $R \rightarrow R/R \cap \mathfrak{n}_Q$, $r \mapsto r + R \cap \mathfrak{n}_Q$, is a nilpotent ideal of the ring R since the ideal $R \cap \mathfrak{n}_Q$ is a nilpotent ideal. We have to show that $\bar{I} = 0$. The left ideal $\mathcal{C}^{-1}I$ of Q is an ideal of the ring Q since the ring Q is a left Noetherian ring. Then $I\mathcal{C}^{-1} := \{ic^{-1} \mid i \in I, c \in \mathcal{C}\} \subseteq \mathcal{C}^{-1}I$, and so

$$(\mathcal{C}^{-1}I)^i \subseteq \mathcal{C}^{-1}I^i, \quad i \geq 1. \quad (3)$$

So, $\mathcal{C}^{-1}I$ is a nilpotent ideal of Q , and so $\mathcal{C}^{-1}I \subseteq \mathfrak{n}_Q$ and $I \subseteq R \cap \mathcal{C}^{-1}I \subseteq R \cap \mathfrak{n}_Q$. Therefore, $\bar{I} = 0$, as required. Clearly,

$$\mathcal{C}^{-1}\mathfrak{n} = \mathcal{C}^{-1}(R \cap \mathfrak{n}_Q) = \mathcal{C}^{-1}R \cap \mathcal{C}^{-1}\mathfrak{n}_Q = Q \cap \mathfrak{n}_Q = \mathfrak{n}_Q,$$

and $(\mathcal{C}^{-1}\mathfrak{n})^i = \mathcal{C}^{-1}\mathfrak{n}^i$ for $i \geq 1$. In particular, $\nu = \nu_Q < \infty$.

2(a) Let $c \in \mathcal{C}$ and $n \in \mathfrak{n}$. Then the element $c^{-1}n \in \mathcal{C}^{-1}\mathfrak{n} = \mathfrak{n}_Q$ (statement 1) is a nilpotent element of the ring Q and so the element $1 + c^{-1}n$ is a unit of the ring Q . Now,

$$c + n = c(1 + c^{-1}n) \in \mathcal{C}.$$

2(b), 2(c) Since $\mathfrak{n} = R \cap \mathfrak{n}_Q$ (statement 1), there is a commutative diagram of ring homomorphisms

$$\begin{array}{ccc} Q & \xrightarrow{\pi_Q} & Q/\mathfrak{n}_Q \\ \sigma \uparrow & & \uparrow \tilde{\sigma} \\ R & \xrightarrow{\pi} & \overline{R} \end{array} \quad (4)$$

where the horizontal maps are natural epimorphisms and the vertical maps are natural monomorphisms (where $\tilde{\sigma}(\bar{r}) := \frac{\bar{r}}{1} = \frac{\bar{r}}{1} + \mathfrak{n}_Q$). Since

$$Q/\mathfrak{n}_Q = \{\pi(c)^{-1}\bar{r} \mid c \in \mathcal{C}, \bar{r} \in \overline{R}\},$$

we see that $\tilde{\mathcal{C}} = \pi(\mathcal{C}) \in \text{Den}_l(\overline{R}, 0)$ and $Q/\mathfrak{n}_Q \simeq \tilde{\mathcal{C}}^{-1}\overline{R}$ is a semiprime left Noetherian ring.

2(d) This follows from statement 1.

2(e) By the statement (c), $\tilde{Q} \simeq Q/\mathfrak{n}_Q$ is a left Noetherian ring. Hence, the Q/\mathfrak{n}_Q -modules

$$\mathfrak{n}_Q^i/\mathfrak{n}_Q^{i+1} \stackrel{\text{st.1}}{\simeq} (\mathcal{C}^{-1}\mathfrak{n})^i/(\mathcal{C}^{-1}\mathfrak{n})^{i+1} \stackrel{\text{st.1}}{\simeq} \mathcal{C}^{-1}\mathfrak{n}^i/\mathcal{C}^{-1}\mathfrak{n}^{i+1} \simeq \tilde{\mathcal{C}}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1})$$

are finitely generated where $i = 1, \dots, \nu$.

2(f) For each $i = 1, \dots, \nu$, the left R -module $\mathcal{N}_i/\mathcal{N}_i\bar{c}$ is $\tilde{\mathcal{C}}$ -torsion since

$$\begin{aligned} \tilde{\mathcal{C}}^{-1}(\mathcal{N}_i/\mathcal{N}_i\bar{c}) &= \tilde{\mathcal{C}}^{-1}(\mathfrak{n}^i/(\mathfrak{n}^i\bar{c} + \mathfrak{n}^{i+1})) = \mathcal{C}^{-1}(\mathfrak{n}^i/(\mathfrak{n}^i\bar{c} + \mathfrak{n}^{i+1})) \\ &= \mathcal{C}^{-1}\mathfrak{n}^i/(\mathcal{C}^{-1}\mathfrak{n}^i\bar{c} + \mathcal{C}^{-1}\mathfrak{n}^{i+1}) = \mathcal{C}^{-1}\mathfrak{n}^i/\mathcal{C}^{-1}\mathfrak{n}^i = 0. \end{aligned}$$

4. Statement 4 follows from the fact that \mathfrak{n}_Q is a nilpotent ideal of the ring Q .

3. By statement 2(c), the ring $\tilde{Q} \simeq Q/\mathfrak{n}_Q$ is a semiprime left Noetherian ring. In particular, it is a semiprime left Goldie ring and, by Goldie's Theorem, its left quotient ring $Q(\tilde{Q}) \simeq Q(Q/\mathfrak{n}_Q)$ is a semisimple ring. Since $\bar{R} \subseteq Q/\mathfrak{n}_Q \simeq \tilde{\mathcal{C}}^{-1}\bar{R}$ (statement 2(c)) we have $Q(\bar{R}) = Q(\tilde{\mathcal{C}}^{-1}\bar{R})$ is a semisimple ring. By Goldie's Theorem, the ring \bar{R} is a semiprime left Goldie ring. So, we can extend the commutative diagram (4) to the commutative diagram (which is used in the proof of statement 6)

$$\begin{array}{ccccc} Q & \xrightarrow{\pi_Q} & Q/\mathfrak{n}_Q & \xrightarrow{\sigma'} & Q(Q/\mathfrak{n}_Q) \\ \sigma \uparrow & & \uparrow \tilde{\sigma} & & \downarrow \simeq \\ R & \xrightarrow{\pi} & \bar{R} & \xrightarrow{\bar{\sigma}} & \bar{Q} \end{array} \quad (5)$$

where the maps σ' and $\bar{\sigma}$ are monomorphisms, $\sigma'(q + \mathfrak{n}_Q) = \frac{q + \mathfrak{n}_Q}{1}$ and $\bar{\sigma}(\bar{r}) = \frac{\bar{r}}{1}$.

5. By Theorem 2.1.(1), $\mathcal{C} = \sigma^{-1}(Q^*)$. By statement 4, $Q^* = \pi_Q^{-1}((Q/\mathfrak{n}_Q)^*)$. Then, in view of the commutative diagram (4),

$$\begin{aligned} \mathcal{C} &= \sigma^{-1}\pi_Q^{-1}((Q/\mathfrak{n}_Q)^*) = (\pi_Q\sigma)^{-1}((Q/\mathfrak{n}_Q)^*) = (\tilde{\sigma}\pi)^{-1}((Q/\mathfrak{n}_Q)^*) \\ &= \pi^{-1}(\tilde{\sigma}^{-1}((Q/\mathfrak{n}_Q)^*)). \end{aligned}$$

6. By Theorem 2.1.(1), $\mathcal{C}^\dagger = \sigma'^{-1}(Q(Q/\mathfrak{n}_Q)^*)$ and $\bar{\mathcal{C}} = \bar{\sigma}^{-1}(\bar{Q}^*) = \bar{\sigma}^{-1}(Q(Q/\mathfrak{n}_Q)^*)$ (statement 3). Thus, the commutativity of the second square in the diagram (5) yields,

$$\bar{\mathcal{C}} = (\sigma'\tilde{\sigma})^{-1}(Q(Q/\mathfrak{n}_Q)^*) = \tilde{\sigma}^{-1}(\mathcal{C}^\dagger). \quad \square$$

The next corollary is a criterion for a ring to have a left Noetherian left quotient ring Q such that the factor ring Q/\mathfrak{n}_Q is a semisimple ring (or \tilde{Q} is a semisimple ring; or $\tilde{\mathcal{C}} = \bar{\mathcal{C}}$; or $\mathcal{C} = \pi^{-1}(\bar{\mathcal{C}})$).

Corollary 2.5 *Let R be a ring with a left Noetherian left quotient ring Q , we keep the notation of Theorem 2.4 and its proof. The following statements are equivalent (recall that $\tilde{Q} = Q/\mathfrak{n}_Q$, Theorem 2.4.(2c)).*

1. \tilde{Q} is a semisimple ring.
2. $\tilde{Q} = Q(\tilde{Q})$.
3. $\bar{\mathcal{C}} = \tilde{\sigma}^{-1}(\tilde{Q}^*)$.
4. $\mathcal{C} = \pi^{-1}(\bar{\mathcal{C}})$.
5. $\tilde{\mathcal{C}} = \bar{\mathcal{C}}$.
6. \tilde{Q} is a left Artinian ring.

Proof. (1 \Rightarrow 2) Trivial.

(2 \Rightarrow 3) If $\tilde{Q} = Q(\tilde{Q})$, i.e., the map σ' in (5) is an isomorphism (by Theorem 2.4.(3)), then the rings $\tilde{Q} \simeq Q/\mathfrak{n}_Q$ and \bar{Q} are isomorphic, see the commutative diagram (5). Now, $\bar{C} = \bar{\sigma}^{-1}(\bar{Q}^*) = \tilde{\sigma}^{-1}(\tilde{Q}^*)$ where the first equality holds by Theorem 2.1.(1).

(3 \Rightarrow 4) By Theorem 2.4.(5) and Theorem 2.4.(2c), $C = \pi^{-1}(\tilde{\sigma}^{-1}(\tilde{Q}^*)) = \pi^{-1}(\bar{C})$.

(4 \Rightarrow 5) $\tilde{C} = \pi(C) = \pi(\pi^{-1}(\bar{C})) = \bar{C}$ since the map π is an epimorphism.

(5 \Rightarrow 1) If $\tilde{C} = \bar{C}$ then $\tilde{Q} \simeq \bar{Q}$ is a semisimple ring, by Theorem 2.4.(3).

(1 \Rightarrow 6) Trivial.

(6 \Rightarrow 1) This implication follows from Theorem 2.4.(2c). \square

While the following three propositions are interesting results about localizations, they are not directly connected with the main results in this work. The reader may skip them without impairing their understanding of the remainder of the paper.

Proposition 2.6 *Let R be a ring and $S \in \text{Den}_l(R)$. Then the ring R is a left Noetherian ring iff the ring $S^{-1}R$ is a left Noetherian ring and for each left ideal I of R the left R -module $\text{tor}_S(R/I)$ is left Noetherian.*

Proof. (\Rightarrow) Trivial.

(\Leftarrow) Let I be a left ideal of the ring R . We have to show that I is a finitely generated left R -module. The ring $S^{-1}R$ is a left Noetherian ring. Then its left ideal $S^{-1}I$ is a finitely generated $S^{-1}R$ -module. We can find elements $u_1, \dots, u_n \in I$ such that $S^{-1}I = \sum_{i=1}^n S^{-1}Ru_i/1$. Let $I' := \sum_{i=1}^n Ru_i$. Then $I' \subseteq I$ and I/I' is a submodule of the left Noetherian R -module $\text{tor}_S(R/I')$ and as a result the left R -module I/I' is finitely generated. This fact implies that the left ideal I is finitely generated. \square

Let R be a ring, $S \in \text{Ore}_l(R)$ and M be a left R -module. Let $\ker(S, M) = \{\ker(s_M) \mid s \in S\}$ where $s_M : M \rightarrow M, m \mapsto sm$. The set $(\ker(S, M), \subseteq)$ is a poset. Let $\max \ker(S, M)$ be the set of its maximal elements and let $\max(S, M) := \{s \in S \mid \ker(s_M) \in \max \ker(S, M)\}$.

Proposition 2.7 *Let R be a ring, $S \in \text{Den}_l(R)$, M be a left R -module such that $\max \ker(S, M) \neq \emptyset$. Then $\max \ker(S, M) = \{\text{tor}_S(M)\}$, i.e., $\ker(s_M) = \text{tor}_S(M)$ for all elements $s \in \max(S, M)$. In particular, $\mathfrak{a} := \text{ann}_R(\text{tor}_S(M)) \neq 0$ and $\emptyset \neq \max(S, M) \subseteq S \cap \mathfrak{a}$.*

Proof. Suppose that $\ker(s_M) \neq \text{tor}_S(M)$ for some $s \in \max(S, M)$, we seek a contradiction. Fix $m \in \text{tor}_S(M) \setminus \ker(s_M)$ and $t \in S$ with $tm = 0$. Since $S \in \text{Ore}_l(R)$, we have $St \cap Rs \neq \emptyset$, i.e., $s' := s_1t \in St \cap Rs$ for some $s_1 \in S$. Clearly, $s' \in S$ and $\ker(s'_M) \supseteq \ker(t_M) + \ker(s_M) \not\supseteq \ker(s_M)$, a contradiction. \square

Proposition 2.8 *Let R be a ring, $\mathfrak{p}_1, \dots, \mathfrak{p}_n$ be prime ideals of the ring R such that the rings R/\mathfrak{p}_i ($i = 1, \dots, n$) are left Goldie rings. Let M be a left R -module such that $M = M_n \supset M_{n-1} \supset \dots \supset M_1 \supset M_0 = 0$ is a chain of its submodules such that \mathfrak{p}_i is maximal among left annihilators of non-zero submodules of M/M_{i-1} , $M_i = \text{ann}_{M/M_{i-1}}(\mathfrak{p}_i) := \{m \in M/M_{i-1} \mid \mathfrak{p}_i m = 0\}$, and $\max \ker(\mathcal{C}_{R/\mathfrak{p}_i}, R/M_{i-1}) \neq \emptyset$ for $i = 1, \dots, n$. Then $\bigcap_{i=1}^n \mathcal{C}(\mathfrak{p}_i) \subseteq \mathcal{C}_M(0) := \{r \in R \mid r_M \text{ is an injection}\}$ where $\mathcal{C}(\mathfrak{p}_i) := \{r \in R \mid r + \mathfrak{p}_i \in \mathcal{C}_{R/\mathfrak{p}_i}\}$.*

Proof. Let $c \in \bigcap_{i=1}^n \mathcal{C}(\mathfrak{p}_i)$. We have to show that $\ker(c_M) = 0$. Suppose that this is not true, i.e., $cm = 0$ for some $0 \neq m \in M$, we seek a contraction. Then $m \in M_i \setminus M_{i-1}$ for some i , and so $0 \neq \bar{m} := m + M_{i-1} \in \tau_i := \text{tor}_{\mathcal{C}_{R/\mathfrak{p}_i}}(M_i/M_{i-1})$ since $\bar{c} := c + \mathfrak{p}_i \in \mathcal{C}_{R/\mathfrak{p}_i}$. By Proposition 2.7, $\bar{s}\tau_i = 0$ for some $\bar{s} \in \mathcal{C}_{R/\mathfrak{p}_i}$ (since $\max \ker(\mathcal{C}_{R/\mathfrak{p}_i}, M_i/M_{i-1}) \neq \emptyset$), i.e., $s\tau_i = 0$ for some $s \in \mathcal{C}(\mathfrak{p}_i)$. This means that $\text{ann}_R(\tau_i) \not\supseteq \mathfrak{p}_i$, a contradiction. \square

3 Proof of Theorem 1.2 and a Second Criterion (via the associated graded ring)

The aim of this section is to give proofs of Theorem 1.2 and Theorem 1.3 which are criteria for a ring R to have a left Noetherian left quotient ring.

Proof of Theorem 1.1. The implications $(1 \Leftarrow 2)$ and $(4 \Leftarrow 5)$ are obvious and the equivalence $(2 \Leftrightarrow 3)$ is Goldie's Theorem.

$(1 \Rightarrow 2)$ Since R is a semiprime ring then so is the ring $Q = Q(R)$. In particular, the ring Q is a semiprime left Goldie ring, hence, by Goldie's Theorem, $Q(Q(R)) = Q(R)$ is a semisimple ring.

$(2 \Leftrightarrow 5)$ [5, Theorem 2.9].

$(4 \Rightarrow 2)$ Since R is a semiprime ring then so is the ring $Q_l(R)$ (since the ring $Q_l(R)$ is left Noetherian). The ring $Q_l(R)$ is a left Noetherian ring, hence $Q_l(R)$ is a semiprime left Goldie ring. By Goldie's Theorem, $Q(Q_l(R))$ is a semisimple ring. Then, by [5, Theorem 2.9], $Q(Q_l(R)) = Q_l(Q_l(R))$ is a semisimple ring. By [5, Theorem 2.8.(4)], $Q_l(Q_l(R)) = Q_l(R)$, and so $Q_l(R)$ is a semisimple ring. Then, by [5, Theorem 2.9], $Q(R) = Q_l(R)$, i.e., $Q(R)$ is a semisimple ring. \square

Proof of Theorem 1.2. $(1 \Rightarrow 2)$ Theorem 2.4.(1) and Theorem 2.4.(2).

$(1 \Leftarrow 2)$ (i) $\mathcal{C} \in \text{Ore}_l(R)$: We have to show that for given elements $c \in \mathcal{C}$ and $r \in R$ there are elements $c' \in \mathcal{C}$ and $r' \in R$ such that $c'r = r'c$. We can assume that $r \neq 0$ since otherwise take $c' = 1$ and $r' = 0$. To prove this fact we use a downward induction on the *degree* of the element $r \neq 0$:

$$\deg(r) := \max\{i \mid r \in \mathfrak{n}^i \text{ where } 0 \leq i \leq \nu\}.$$

Suppose that $\deg(r) = \nu$. By the condition (f), the \bar{R} -module $\mathfrak{n}^\nu/\mathfrak{n}^{\nu+1}$ is $\tilde{\mathcal{C}}$ -torsion, hence $c'r = r'\bar{c}$ for some elements $c' = c' + \mathfrak{n} \in \tilde{\mathcal{C}}$ (where $c' \in \mathcal{C}$) and $r' \in \mathfrak{n}^\nu$. Then $c'r = r'c$.

Suppose that $i = \deg(r) < \nu$, and the result is true for all elements $c \in \mathcal{C}$ and $r \in R$ with $\deg(r) > i$.

Suppose that $i = 0$, i.e., $r \in R \setminus \mathfrak{n}$. Since $\tilde{\mathcal{C}} \in \text{Den}_l(\bar{R}, 0)$ (by the conditions (a) and (b)), $c_1\bar{r} = \bar{r}_1\bar{c}$ for some elements $c_1 \in \mathcal{C}$ and $r_1 \in R$. The difference $a := c_1r - r_1c$ belongs to the ideal \mathfrak{n} , and so $\deg(a) > 0$. By induction, $c_2a = bc$ for some elements $c_2 \in \mathcal{C}$ and $b \in R$. Then

$$c_2c_1r = (c_2r_1 + b)c,$$

and it suffices to take $c' = c_2c_1$ and $r' = c_2r_1 + b$.

Suppose that $i > 0$. By the condition (f), the left \bar{R} -module $\mathcal{N}_i/\mathcal{N}_{i+1} = \mathfrak{n}^i/(\mathfrak{n}^i + \mathfrak{n}^{i+1})$ is $\tilde{\mathcal{C}}$ -torsion. Therefore, $sr = xc + y$ for some elements $s \in \mathcal{C}$, $x \in \mathfrak{n}^i$ and $y \in \mathfrak{n}^{i+1}$. Since $\deg(y) \geq i+1$, by induction, there are elements $t \in \mathcal{C}$ and $z \in R$ such that $ty = zc$. Therefore,

$$tsr = (tx + z)c$$

and it suffices to take $c' = ts$ and $r' = tx + z$.

(ii) $\mathcal{C}^{-1}\mathfrak{n}$ is an ideal of the ring Q such that $Q/\mathcal{C}^{-1}\mathfrak{n} \simeq \tilde{Q}$: By the statement (i), the left quotient ring $Q = \mathcal{C}^{-1}R$ exists. Let $\sigma : R \rightarrow Q, r \mapsto \frac{r}{1}$. By the universal property of left localization, there is a ring homomorphism $\pi_Q : Q \rightarrow \tilde{Q}, c^{-1}r \mapsto \bar{c}^{-1}\bar{r}$, where $\bar{c} = c + \mathfrak{n}$ and $\bar{r} = r + \mathfrak{n}$, and we have the commutative diagram of ring homomorphisms

$$\begin{array}{ccc} Q & \xrightarrow{\pi_Q} & \tilde{Q} \\ \sigma \uparrow & & \uparrow \tilde{\sigma} \\ R & \xrightarrow{\pi} & \bar{R} \end{array} \quad (6)$$

where $\tilde{\sigma} : \bar{r} \mapsto \frac{\bar{r}}{1}$ is a monomorphism and π_Q is an epimorphism (by the very definition of π_Q). Applying the exact functor $\mathcal{C}^{-1}(-)$ to the short exact sequence of R -modules $0 \rightarrow \mathfrak{n} \rightarrow R \xrightarrow{\pi} \bar{R} \rightarrow 0$ we obtain the short exact sequence of Q -modules

$$0 \rightarrow \mathcal{C}^{-1}\mathfrak{n} \rightarrow Q \xrightarrow{\pi_Q} \mathcal{C}^{-1}\bar{R} = \tilde{\mathcal{C}}^{-1}\bar{R} = \tilde{Q} \rightarrow 0.$$

Therefore, $\ker(\pi_Q) = \mathcal{C}^{-1}\mathfrak{n}$ is an ideal of Q (since π_Q is a ring homomorphism) such that $Q/\mathcal{C}^{-1}\mathfrak{n} \simeq \tilde{Q}$.

(iii) $(\mathcal{C}^{-1}\mathfrak{n})^i = \mathcal{C}^{-1}\mathfrak{n}^i$ for $i \geq 1$: This follows from the statement (ii).

(iv) *The ring Q is a left Noetherian ring*: By localizing at \mathcal{C} the descending chain of ideals of the ring R ,

$$R \supset \mathfrak{n} \supset \mathfrak{n}^2 \supset \cdots \supset \mathfrak{n}^i \supset \cdots \supset \mathfrak{n}^\nu \supset \mathfrak{n}^{\nu+1} = 0,$$

we obtain the descending chain of ideals of the ring Q (by the statement (iii)):

$$Q \supseteq \mathcal{C}^{-1}\mathfrak{n} \supseteq \mathcal{C}^{-1}\mathfrak{n}^2 \supseteq \cdots \supseteq \mathcal{C}^{-1}\mathfrak{n}^i \supseteq \cdots \supseteq \mathcal{C}^{-1}\mathfrak{n}^\nu \supseteq \mathcal{C}^{-1}\mathfrak{n}^{\nu+1} = 0. \quad (7)$$

By (ii), $\tilde{Q} \simeq Q/\mathcal{C}^{-1}\mathfrak{n}$ is a left Noetherian Q -module since the ring \tilde{Q} is a left Noetherian ring, by the condition (c). For each $i = 1, \dots, \nu$, the left Q -module

$$\mathcal{C}^{-1}\mathfrak{n}^i/\mathcal{C}^{-1}\mathfrak{n}^{i+1} \simeq (\mathcal{C}^{-1}\mathfrak{n})^i/(\mathcal{C}^{-1}\mathfrak{n})^{i+1}$$

is a $Q/\mathcal{C}^{-1}\mathfrak{n} = \tilde{Q}$ -module, by the statement (ii). The \tilde{Q} -modules

$$\mathcal{C}^{-1}\mathfrak{n}^i/\mathcal{C}^{-1}\mathfrak{n}^{i+1} \simeq \mathcal{C}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1}) \simeq \tilde{\mathcal{C}}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1}) = \tilde{\mathcal{C}}^{-1}\mathcal{N}_i$$

are finitely generated, by the condition (e), hence Noetherian since the ring \tilde{Q} is a left Noetherian. Since all the factors of the finite filtration (7) are Noetherian \tilde{Q} -modules/ Q -modules, the ring Q is a left Noetherian ring. \square

The minimal primes of the rings R , $Q(R)$, \bar{Q} and \tilde{Q} . For a ring R such that $Q(R)$ is a left Noetherian ring, the next corollary shows that the localizations of R at the maximal left denominator sets are left Noetherian rings and there are natural bijections between the sets of minimal primes of the rings R , $Q(R)$, \bar{Q} and \tilde{Q} .

Corollary 3.1 *Let R be a ring with a left Noetherian left quotient ring $Q(R)$. Then*

1. *For every $S \in \max.\text{Den}_l(R)$, the ring $S^{-1}R$ is a left Noetherian ring.*
2. (a) *The map $\text{Min}(R) \rightarrow \text{Min}(\bar{Q})$, $\mathfrak{p} \mapsto \bar{\mathcal{C}}^{-1}(\mathfrak{p}/\mathfrak{n})$, is a bijection with the inverse $\mathfrak{q} \mapsto (\bar{\sigma}\pi)^{-1}(\mathfrak{q})$ where the maps $\bar{\sigma}$ and π are defined in (8).*
 (b) *The map $\text{Min}(R) \rightarrow \text{Min}(\tilde{Q})$, $\mathfrak{p} \mapsto \tilde{\mathcal{C}}^{-1}(\mathfrak{p}/\mathfrak{n})$, is a bijection with the inverse $\mathfrak{q} \mapsto \tau^{-1}(\mathfrak{q})$ where $\tau : R \rightarrow \tilde{Q}$, $r \mapsto \frac{r+\mathfrak{p}}{1}$.*
 (c) *The map $\text{Min}(R) \rightarrow \text{Min}(Q)$, $\mathfrak{p} \mapsto \mathcal{C}^{-1}\mathfrak{p}$, is a bijection with the inverse $\mathfrak{q} \mapsto \mathfrak{q} \cap R$.*

Proof. 1. By Proposition 2.3.(1), $\mathcal{C} = \mathcal{C}_R \subseteq S$ for all $S \in \max.\text{Den}_l(R)$. The ring $Q := \mathcal{C}^{-1}R$ is a left Noetherian ring and the ring $S^{-1}R \simeq (SQ^*)^{-1}Q$ is a left localization of the ring Q (by Proposition 2.3.(3), the map $\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q)$, $S \mapsto SQ^*$, is a bijection with $S^{-1}R \simeq (SQ^*)^{-1}Q$). Therefore, the ring $S^{-1}R$ is a left Noetherian ring since the ring Q is so.

2(a) The map $\text{Min}(R) \rightarrow \text{Min}(\bar{R})$, $\mathfrak{p} \mapsto \mathfrak{p}/\mathfrak{n}$, is a bijection with the inverse $\mathfrak{p}' \mapsto \pi^{-1}(\mathfrak{p}')$ where $\pi : R \rightarrow \bar{R}$, $r \mapsto r + \mathfrak{p}$. The ring \bar{R} is a semiprime left Goldie ring such that $\bar{Q} \simeq Q(Q/\mathfrak{n}_Q)$ is a semisimple ring (Theorem 2.4.(3)). Hence, the map $\text{Min}(\bar{R}) \rightarrow \text{Min}(\bar{Q})$, $\mathfrak{p}' \mapsto \bar{\mathcal{C}}^{-1}\mathfrak{p}'$, is a bijection with the inverse $\mathfrak{q} \mapsto \bar{\sigma}^{-1}(\mathfrak{q})$ where $\bar{\sigma} : \bar{R} \rightarrow \bar{Q}$, $\bar{r} \mapsto \frac{\bar{r}}{1}$. Now, the statement (a) follows.

(b) The ring \tilde{Q} is a semiprime left Goldie ring (by Theorem 2.4.(2c)) and $Q(\tilde{Q}) \simeq \bar{Q}$ is a semisimple ring (Theorem 2.4.(3)). So, the map $\text{Min}(\tilde{Q}) \rightarrow \text{Min}(\bar{Q})$, $P \mapsto \bar{Q} \otimes_{\tilde{Q}} P$, is a bijection with the inverse $P' \mapsto P' \cap \tilde{Q}$. Now, the statement (b) follows from the statement (a).

(c) By Theorem 2.4.(2c), $\tilde{Q} \simeq Q/\mathfrak{n}_Q$ and the map $\text{Min}(Q) \rightarrow \text{Min}(Q/\mathfrak{n}_Q)$, $P \mapsto P/\mathfrak{n}_Q$, is a bijection. Now, the statement (c) follows from the statement (b). \square

Lemma 3.2 Let G be a monoid and $e \in G$ be its neutral element, $A = \bigoplus_{g \in G} A_g$ be a G -graded ring, $1 \in A_e$, $S \in \text{Den}_l(A)$ and $\mathfrak{a} = \text{ass}(S)$. If $S \subseteq A_e$ then the ring $S^{-1}A = \bigoplus_{g \in G} (S^{-1}A)_g$ is a G -graded ring where $(S^{-1}A)_g = S^{-1}A_g := \{s^{-1}a_g \mid s \in S, a_g \in A_g\}$, $S \in \text{Den}_l(A_e)$ and $\mathfrak{a} = \bigoplus_{g \in G} \mathfrak{a}_g$ is a G -graded ideal of the ring A , i.e., $\mathfrak{a}_g = \mathfrak{a} \cap A_g$ for all $g \in G$.

Proof. Straightforward. \square

Suppose that a ring R has a left Noetherian left quotient ring Q . By Theorem 2.4, the associated graded ring $\text{gr } Q := Q/\mathfrak{n}_Q \oplus \mathfrak{n}_Q/\mathfrak{n}_Q^2 \oplus \cdots$ is equal to

$$\text{gr } Q = \tilde{Q} \oplus \tilde{\mathcal{C}}^{-1}(\mathfrak{n}/\mathfrak{n}^2) \oplus \cdots \oplus \tilde{\mathcal{C}}^{-1}(\mathfrak{n}^\nu/\mathfrak{n}^{\nu+1}).$$

Proof of Theorem 1.3. (1 \Rightarrow 2) Suppose that the ring $Q = Q(R)$ is a left Noetherian ring, and so the conditions of Theorem 1.2 and Theorem 2.4 hold. In particular, $\tilde{\mathcal{C}} \subseteq \bar{\mathcal{C}}$ and \mathfrak{n} is a nilpotent ideal.

(i) $\tilde{\mathcal{C}} \in \text{Ore}_l(\text{gr } R)$: It suffices to show that for given elements $\bar{c} = c + \mathfrak{n} \in \tilde{\mathcal{C}}$ and $r + \mathfrak{n}^{i+1} \in \mathfrak{n}^i/\mathfrak{n}^{i+1}$ where $c \in \mathcal{C}$, $r \in \mathfrak{n}^i$ and $i = 0, 1, \dots, \nu$, there are elements $\bar{c}' = c' + \mathfrak{n} \in \tilde{\mathcal{C}}$ and $r' + \mathfrak{n}^{i+1} \in \mathfrak{n}^i/\mathfrak{n}^{i+1}$ where $c' \in \mathcal{C}$, $r' \in \mathfrak{n}^i$ such that $\bar{c}'(r + \mathfrak{n}^{i+1}) = (r' + \mathfrak{n}^{i+1})\bar{c}$.

The case $i = 0$ is obvious, by Theorem 1.2.(2b). So, we can assume that $i \geq 1$. By Theorem 2.4.(1),

$$\mathfrak{n}^i \mathcal{C}^{-1} := \{nc^{-1} \mid n \in \mathfrak{n}^i, c \in \mathcal{C}\} \subseteq \mathcal{C}^{-1} \mathfrak{n}^i.$$

So, $rc^{-1} = c'^{-1}r'$ for some elements $c' \in \mathcal{C}$ and $r' \in \mathfrak{n}^i$, hence $c'r = r'c$. This equality implies the required one.

(ii) $\tilde{\mathcal{C}} \in \text{Den}_l(\text{gr } R)$: We have to show that if $\bar{r}\bar{c} = 0$ for some elements $\bar{r} = r + \mathfrak{n}^{i+1} \in \mathfrak{n}^i/\mathfrak{n}^{i+1}$ and $\bar{c} = c + \mathfrak{n}$ where $r \in \mathfrak{n}^i$, $c \in \mathcal{C}$ and $i = 0, 1, \dots, \nu$, then $\bar{c}_1\bar{r} = 0$ for some element $\bar{c}_1 = c_1 + \mathfrak{n}$ where $c_1 \in \mathcal{C}$. The case $i = 0$ follows from Theorem 2.4.(2b). Suppose that $i \geq 1$. Then $n := rc \in \mathfrak{n}^{i+1}$, and so $r = nc^{-1} = c_1^{-1}n_1$ for some elements $c_1 \in \mathcal{C}$ and $n_1 \in \mathfrak{n}^{i+1}$, by Theorem 2.4.(1). Hence, $\bar{c}_1\bar{r} = 0$, as required.

(iii) $\text{gr } Q \simeq \tilde{\mathcal{C}}^{-1} \text{gr } R$: By Theorem 2.4.(1), $\mathfrak{n}_Q = \mathcal{C}^{-1} \mathfrak{n}$. By Theorem 2.4.(2c), $Q/\mathfrak{n}_Q \simeq \tilde{Q}$. Now, using Theorem 2.4.(1,2), we have

$$\begin{aligned} \text{gr } Q &= \tilde{Q} \oplus \cdots \oplus \mathfrak{n}_Q^i/\mathfrak{n}_Q^{i+1} \oplus \cdots = \tilde{Q} \oplus \cdots \oplus \mathcal{C}^{-1} \mathfrak{n}^i/\mathcal{C}^{-1} \mathfrak{n}^{i+1} \oplus \cdots \\ &= \tilde{Q} \oplus \cdots \oplus \mathcal{C}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1}) \oplus \cdots = \tilde{Q} \oplus \cdots \oplus \tilde{\mathcal{C}}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1}) \oplus \cdots \\ &\simeq \tilde{\mathcal{C}}^{-1} \text{gr } R. \end{aligned}$$

(iv) $\tilde{\mathcal{C}}^{-1} \text{gr } R$ is a left Noetherian ring: The ring \tilde{Q} is a left Noetherian ring (Theorem 2.4.(2c)) and the left \tilde{Q} -modules $\tilde{\mathcal{C}}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1}) \simeq (\mathcal{C}^{-1} \mathfrak{n})^i/(\mathcal{C}^{-1} \mathfrak{n})^{i+1}$ are finitely generated where $i = 1, \dots, \nu$ (since Q is a left Noetherian ring). Therefore, the left \tilde{Q} -module $\text{gr } Q = \tilde{Q} \oplus \cdots \oplus \tilde{\mathcal{C}}^{-1}(\mathfrak{n}^i/\mathfrak{n}^{i+1}) \oplus \cdots \oplus \tilde{\mathcal{C}}^{-1}(\mathfrak{n}^\nu/\mathfrak{n}^{\nu+1})$ is finitely generated, hence Noetherian. Since $\tilde{Q} \subseteq \text{gr } Q$, the ring $\text{gr } Q$ is a left Noetherian ring.

(1 \Leftarrow 2) It suffices to show that the conditions (a)-(f) of Theorem 1.2.(2) hold. The conditions (a) and (d) are given. The set $\tilde{\mathcal{C}}$ is a left denominator set of the \mathbb{N} -graded ring $\text{gr } R$ such that $\tilde{\mathcal{C}} \subseteq \bar{\mathcal{R}}$. By Lemma 3.2, the ring $\tilde{\mathcal{C}}^{-1} \text{gr } R = \tilde{Q} \oplus \cdots \oplus \tilde{\mathcal{C}}^{-1} \mathcal{N}_i \oplus \cdots$ is an \mathbb{N} -graded ring and $\tilde{\mathcal{C}} \in \text{Den}_l(\bar{\mathcal{R}}, 0)$, and so the condition (b) holds.

The ring \tilde{Q} is a factor ring of the left Noetherian ring $\tilde{\mathcal{C}}^{-1} \text{gr } R$, hence \tilde{Q} is a left Noetherian ring, i.e., the condition (c) holds.

The ring $\tilde{\mathcal{C}}^{-1} \text{gr } R$ is left Noetherian, hence the \tilde{Q} -modules $\tilde{\mathcal{C}}^{-1} \mathcal{N}_i$ are finitely generated, i.e., the condition (e) holds.

The ring $\tilde{\mathcal{C}}^{-1} \text{gr } R$ is an \mathbb{N} -graded ring. In particular, $\tilde{\mathcal{C}}^{-1} \mathcal{N}_i \tilde{Q} \subseteq \tilde{\mathcal{C}}^{-1} \mathcal{N}_i$ for all i . Therefore, $\mathcal{N}_i \tilde{\mathcal{C}}^{-1} = \{nc^{-1} \mid n \in \mathcal{N}_i, c \in \tilde{\mathcal{C}}\} \subseteq \tilde{\mathcal{C}}^{-1} \mathcal{N}_i$, i.e., the condition (f) holds. \square

Corollary 3.3 Let R be a ring with a left Noetherian left quotient ring Q and $\tilde{\mathfrak{a}} := \text{ass}_{\text{gr } R}(\tilde{\mathcal{C}})$. Then the largest left quotient ring $Q_l(\text{gr } R/\tilde{\mathfrak{a}})$ of the ring $\text{gr } R/\tilde{\mathfrak{a}}$ is a left Noetherian ring.

Proof. By Theorem 1.3.(2), the ring $\tilde{\mathcal{C}}^{-1}\text{gr } R$ is a left Noetherian ring. The ring $Q_l(\text{gr } R/\tilde{\mathfrak{a}})$ is a left localization of the ring $\tilde{\mathcal{C}}^{-1}\text{gr } R$, hence is a left Noetherian ring. \square

4 Finiteness of $\max.\text{Den}_l(R)$ when $Q(R)$ is a left Noetherian ring

The aim of this section is to give a proof of Theorem 1.4.

Let R be a ring. Let S and T be submonoids of the multiplicative monoid (R, \cdot) . We denote by ST the *submonoid* of (R, \cdot) generated by S and T . This notation should not be confused with the product of two sets which is *not* used in this paper. The next result is a criterion for the set ST to be a left Ore (denominator) set, it is used at the final stage in the proof of Theorem 1.4.

Lemma 4.1 ([4].)

1. Let $S, T \in \text{Ore}_l(R)$. If $0 \notin ST$ then $ST \in \text{Ore}_l(R)$.
2. Let $S, T \in \text{Den}_l(R)$. If $0 \notin ST$ then $ST \in \text{Den}_l(R)$.

Proof of Theorem 1.4. Let $S \in \max.\text{Den}_l(R)$ and $\pi : R \rightarrow \bar{R}$, $r \mapsto \bar{r} := r + \mathfrak{n}$.

(i) $\bar{S} := \pi(S) \in \text{Ore}_l(\bar{R})$: This obvious since $S \cap \mathfrak{n} = \emptyset$.

(ii) $\bar{\mathcal{C}} \subseteq \bar{S}$ (Proposition 2.3.(1)).

(iii) $S^{-1}R$ is a left Noetherian ring: By (ii), $S^{-1}R$ is a left localization of the left Noetherian ring Q , hence $S^{-1}R$ is a left Noetherian ring.

(iv) $S^{-1}\mathfrak{n}$ is an ideal of $S^{-1}R$ such that $(S^{-1}\mathfrak{n})^i = S^{-1}\mathfrak{n}^i$ for all $i \geq 1$: By (iii), $S^{-1}\mathfrak{n}$ is an ideal of the ring $S^{-1}R$. In particular, $\mathfrak{n}S^{-1} \subseteq S^{-1}\mathfrak{n}$, hence $(S^{-1}\mathfrak{n})^i = S^{-1}\mathfrak{n}^i$ for all $i \geq 1$.

(v) $\bar{S} \in \text{Den}_l(\bar{R})$: In view of (i), we have to show that if $\bar{r}\bar{s} = 0$ for some $\bar{r} \in \bar{R}$ and $\bar{s} \in \bar{S}$ then $\bar{t}\bar{r} = 0$ for some element $\bar{t} \in \bar{S}$. The equality $\bar{r}\bar{s} = 0$ means that $n := rs \in \mathfrak{n}$. Then $S^{-1}R \ni \frac{r}{1} = ns^{-1} = s_1^{-1}n_1$ for some $s_1 \in S$ and $n_1 \in \mathfrak{n}$, (by (iv)). Hence, $s_2s_1r = s_2n_1 \in \mathfrak{n}$ for some element $s_2 \in S$, and so $\bar{t}\bar{r} = 0$ where $\bar{t} = \bar{s}_2\bar{s}_1 \in \bar{S}$.

The ring \bar{R} is a semiprime left Goldie ring (Theorem 2.4.(3)), $\bar{Q} = \prod_{i=1}^s \bar{Q}_i$ and \bar{Q}_i are simple Artinian rings. By [3, Theorem 4.1], $|\max.\text{Den}_l(\bar{R})| = s$. Let $\max.\text{Den}_l(\bar{R}) = \{T_1, \dots, T_s\}$.

(vi) $|\max.\text{Den}_l(R)| \leq s$: By (v), $\bar{S} \subseteq T_i$ for some i . Then S is the largest element with respect to inclusion of the set $\{S' \in \text{Den}_l(R) \mid \pi(S') \subseteq T_i\}$ (as $0 \in SS_1$ for all *distinct* $S, S_1 \in \max.\text{Den}_l(R)$, by Lemma 4.1.(2)). Hence, $|\max.\text{Den}_l(R)| \leq s$. \square

Let R be a ring such that its left quotient ring $Q(R)$ is a left Noetherian ring. Consider the ring homomorphism

$$p_i : R \xrightarrow{\pi} \bar{R} \xrightarrow{\bar{\sigma}} \bar{Q} = \prod_{j=1}^s \bar{Q}_j \xrightarrow{\bar{p}_i} \bar{Q}_i. \quad (8)$$

Let $\bar{\sigma}_i := \bar{p}_i \bar{\sigma} : \bar{R} \rightarrow \bar{Q}_i$. By [3, Theorem 4.1], $\max.\text{Den}_l(\bar{R}) = \{T_1, \dots, T_s\}$ where $T_i := \bar{\sigma}_i^{-1}(\bar{Q}_i^*)$.

Proof of Corollary 1.7. 1. The set D_i is a non-empty set since $R^* \subseteq D_i$. By Lemma 4.1.(2), the set $S'_i := \bigcup_{S' \in D_i} S'$ is the largest element of the set D_i . In the proof of Theorem 1.4, we proved that $\max.\text{Den}_l(R) \subseteq \mathcal{M}$ where \mathcal{M} is the set of maximal elements of the set $\{S'_1, \dots, S'_s\}$ (see the statement (vi)). The reverse inclusion is obvious.

2. By Theorem 2.4.(2b), $\tilde{\mathcal{C}} \in \text{Den}_l(\bar{R}, 0)$ and $\tilde{\mathcal{C}} \subseteq \bar{\mathcal{C}}$. By Theorem 2.4.(3), the ring \bar{R} is a semiprime left Goldie ring and \bar{Q} is a semisimple ring. By Theorem 2.1.(1), $\bar{\mathcal{C}} = \bar{R} \cap \bar{Q}^*$. Hence, $\bar{\mathcal{C}} \subseteq S'_i$ for all $i = 1, \dots, s$.

3. Statement 3 follows from statement 2: The ring $S'_i{}^{-1}R$ is a localization of the left Noetherian ring $\mathcal{C}^{-1}R$ (see statement 2). Hence it is a left Noetherian ring. \square

Criterion for $|\max.\text{Den}_l(R)| = |\max.\text{Den}_l(\overline{R})|$. Let R be a ring with a left Noetherian left quotient ring $Q(R)$. In general, by Theorem 1.4, $|\max.\text{Den}_l(R)| \leq |\max.\text{Den}_l(\overline{R})|$. The next theorem is a criterion for $|\max.\text{Den}_l(R)| = |\max.\text{Den}_l(\overline{R})|$.

Theorem 4.2 *Let R be a ring with a left Noetherian left quotient ring $Q(R)$. Then $|\max.\text{Den}_l(R)| = |\max.\text{Den}_l(\overline{R})|$ iff for each pair of indices $i \neq j$ where $1 \leq i, j \leq s = |\max.\text{Den}_l(\overline{R})|$ there exist $S_i, S_j \in \text{Den}_l(R)$ such that $0 \in S_i S_j$ (where $S_i S_j$ is the multiplicative submonoid of R generated by S_i and S_j), $p_i(S_i) \subseteq \overline{Q}_i^*$ and $p_j(S_j) \subseteq \overline{Q}_j^*$. In this case, $\max.\text{Den}_l(R) = \{S'_1, \dots, S'_s\}$ where S'_i is the largest element in the set $\{S' \in \text{Den}_l(R) \mid p_i(S') \subseteq \overline{Q}_i^*\}$.*

Proof. By [3, Theorem 4.1], $\max.\text{Den}_l(\overline{R}) = \{T_1, \dots, T_s\}$ where $T_i = (\overline{p}_i \overline{\sigma})^{-1}(\overline{Q}_i^*)$ for $i = 1, \dots, s$.

(\Rightarrow) It suffices to take $S_i = \pi^{-1}(T_i)$, $i = 1, \dots, s$ since $0 \in S_i S_j$ for all $i \neq j$, by Lemma 4.1.(2).

(\Leftarrow) Suppose that S_1, \dots, S_s are as in the theorem. For each $i = 1, \dots, s$, let S'_i be the largest element with respect to inclusion of the set $\{S' \in \text{Den}_l(R) \mid \pi(S') \subseteq T_i\}$. Since $S_i \subseteq S'_i$ and $S_j \subseteq S'_j$ for each distinct pair $i \neq j$ and $0 \in S_i S_j$, the elements S'_1, \dots, S'_s are distinct (Lemma 4.1.(2)) and incomparable (i.e., $S'_i \not\subseteq S'_j$ for all $i \neq j$). In the proof of the statement (vi) of Theorem 1.4, we have seen that $\max.\text{Den}_l(\overline{R}) \subseteq \{S'_1, \dots, S'_s\}$. Hence, $\max.\text{Den}_l(R) = \{S'_1, \dots, S'_s\}$ since the sets S'_1, \dots, S'_s are incomparable. \square

The next theorem gives sufficient conditions for finiteness of the set of $\max.\text{Den}_l(R)$.

Theorem 4.3 *Let R be a ring. Suppose that I is an ideal of the ring $Q_l(R)$ such that $S + I \subseteq S$ and $S^{-1}I$ is an ideal of $S^{-1}Q_l(R)$ for all $S \in \max.\text{Den}_l(Q_l(R))$. Then $|\max.\text{Den}_l(R)| \leq |\max.\text{Den}_l(Q_l(R)/I)|$. In particular, if $|\max.\text{Den}_l(Q_l(R)/I)| < \infty$ then $|\max.\text{Den}_l(R)| < \infty$.*

Proof. By Proposition 2.3.(3), there is a bijection between the sets $\max.\text{Den}_l(R)$ and $\max.\text{Den}_l(Q_l(R))$. Let $Q_l := Q_l(R)$ and $f : Q_l \rightarrow \overline{Q}_l := Q_l/I$, $q \mapsto \overline{q} := q + I$.

(i) For all $S \in \max.\text{Den}_l(Q_l)$, $\overline{S} := f(S) \in \text{Den}_l(\overline{Q}_l)$: Since $S + I \subseteq S$, $S \cap I = \emptyset$ and $\overline{S} \in \text{Ore}_l(\overline{R})$. It remains to show that if $\overline{q}\overline{s} = 0$ for some elements $\overline{q} \in \overline{Q}_l$ and $\overline{s} \in \overline{S}$ then $\overline{t}\overline{q} = 0$ for some element $\overline{t} \in \overline{S}$. The equality $\overline{q}\overline{s} = 0$ means that $i := qs \in I$. Then $S^{-1}Q_l \ni \frac{q}{1} = is^{-1} = s_1^{-1}i_1$ for some elements $s_1 \in S$ and $i_1 \in I$ (since $S^{-1}I$ is an ideal of the ring $S^{-1}Q_l$). Hence, $s_2 s_1 q = s_2 i \in I$ for some $s_2 \in S$, and so $\overline{t}\overline{q} = 0$ where $\overline{t} = \overline{s_2 s_1} \in \overline{S}$.

(ii) $|\max.\text{Den}_l(R)| \leq |\max.\text{Den}_l(Q_l(R)/I)|$: Let $S \in \max.\text{Den}_l(Q_l)$. By (i), $\overline{S} \subseteq T$ for some $T = T(S) \in \max.\text{Den}_l(\overline{Q}_l)$. Then S is the largest element (w.r.t. \subseteq) of the set $\{S' \in \text{Den}_l(R) \mid f(S') \subseteq T\}$ (as $0 \in SS_1$ for all distinct $S, S_1 \in \max.\text{Den}_l(R)$, by Lemma 4.1.(2)), and the statement (ii) follows. \square

Corollary 4.4 *Let R be a ring. Suppose that I is an ideal of $Q_l(R)$ such that $S^{-1}I$ is an ideal of the ring $S^{-1}Q_l(R)$ with $S^{-1}I \subseteq \text{rad}(S^{-1}Q_l(R))$ for all $S \in \max.\text{Den}_l(Q_l(R))$. Then $|\max.\text{Den}_l(R)| \leq |\max.\text{Den}_l(Q_l(R)/I)|$.*

Proof. In view of Theorem 4.3 and its proof, it suffices to show that $S + I \subseteq S$ for all $S \in \max.\text{Den}_l(Q_l)$ where $Q_l = Q_l(R)$. By the assumption, $S^{-1}I \subseteq \text{rad}(S^{-1}Q_l)$. Hence, $1 + S^{-1}I \subseteq (S^{-1}Q_l)^*$. Now, for all $s \in S$ and $i \in I$, $s + i = s(1 + s^{-1}i) \in (S^{-1}Q_l)^*$. Let $\sigma_S : Q_l \rightarrow S^{-1}Q_l$, $q \mapsto \frac{q}{1}$. By Theorem 2.1.(1), $S = \sigma_S^{-1}((S^{-1}Q_l)^*)$, and so $s + i \in S$, as required. \square

Proof of Proposition 1.9 (description of the set $\max.\text{Den}_l(R)$ for a commutative ring R). For each minimal prime ideal \mathfrak{p} of R , $S_{\mathfrak{p}} \in \max.\text{Den}(R)$ since the ring $R_{\mathfrak{p}} := S_{\mathfrak{p}}^{-1}R$ is a local ring such that its maximal ideal $S^{-1}\mathfrak{p}$ (which is also the prime radical of $R_{\mathfrak{p}}$) is a nil ideal, so every element of $R_{\mathfrak{p}}$ is either a unit or a nilpotent element.

Conversely, let $S \in \max.\text{Den}(R)$. We have to show that $S = S_{\mathfrak{p}}$ for some $\mathfrak{p} \in \text{Min}(R)$. In the ring $S^{-1}R$, every element is either a unit or a nilpotent element. Hence, the ring $S^{-1}R$ is a local ring $(S^{-1}R, \mathfrak{m})$ where the maximal ideal \mathfrak{m} is a nil ideal. Let $\sigma : R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$. Then $S = \sigma^{-1}((S^{-1}R)^*) = S_{\mathfrak{p}}$, by Theorem 2.2.(3). \square

Proposition 4.5 *Let R be a commutative ring, I be a nil ideal of R and $\pi_I : R \rightarrow R/I$, $r \mapsto \bar{r} = r + I$. Then*

1. *The map $\text{Den}(R) \rightarrow \text{Den}(R/I)$, $S \mapsto \pi_I(S)$, is a surjection that respects inclusions. Moreover, the map $\text{Den}(R, I) := \{S \in \text{Den}(R) \mid S + I \subseteq S\} \rightarrow \text{Den}(R/I)$, $S \mapsto \pi_I(S)$, is a bijection with the inverse $T \mapsto \pi_I^{-1}(T)$.*
2. *The map $\text{max.Den}(R) \rightarrow \text{max.Den}(R/I)$, $S \mapsto \pi_I(S)$, is a bijection with the inverse $T \mapsto \pi_I^{-1}(T)$.*

Proof. 1. Straightforward.

2. Statement 2 follows from statement 1 since if $S \in \text{max.Den}(R)$ then $S + I \subseteq S$. \square

Let R be a commutative ring such that its quotient ring $Q = Q(R)$ is a Noetherian ring. We keep the notation of Theorem 1.2 and Theorem 4.2. For each $i = 1, \dots, s$, let $p_i : R \rightarrow \bar{Q}_i$ be as in (8). The next theorem shows that the second inequality of Theorem 1.4 is an equality, it also provides another characterization of the set $\text{max.Den}(R)$.

Theorem 4.6 *Let R be a commutative ring such that its quotient ring Q is a Noetherian ring. Then $\text{max.Den}(R) = \{S_i \mid i = 1, \dots, s\}$ where $S_i := p_i^{-1}(\bar{Q}_i^*)$. In particular, $|\text{max.Den}(R)| = s = \text{max.Den}(\bar{R})$.*

Proof. Notice that $\text{max.Den}(\bar{Q}) = \{\bar{p}_i^{-1}(\bar{Q}_i^*) \mid i = 1, \dots, s\}$. By Proposition 2.3.(4), the map $\text{max.Den}(\bar{Q}) \rightarrow \text{max.Den}(\bar{R})$, $S \mapsto \bar{\sigma}^{-1}(S)$, is a bijection. By Proposition 4.5.(2), the map $\text{max.Den}(\bar{R}) \rightarrow \text{max.Den}(R)$, $T \mapsto \pi^{-1}(T)$, is a bijection. Therefore, $\text{max.Den}(R) = \{S_i \mid i = 1, \dots, s\}$. \square

Proposition 4.7 *Let R be a ring, $S \in \text{Den}_l(R, \mathfrak{a})$, I be an ideal of R such that $I \cap S = \emptyset$ and $S^{-1}I$ is an ideal of $S^{-1}R$, $\pi : R \rightarrow R/I$, $r \mapsto \bar{r} := r + I$, and $\bar{S} := \pi(S)$. Then*

1. $\bar{S} \in \text{Den}_l(R/I, \bar{\mathfrak{b}})$ for some ideal $\bar{\mathfrak{b}}$ of R/I such that $(\mathfrak{a} + I)/I \subseteq \bar{\mathfrak{b}}$.
2. Let $\mathfrak{b} := \pi^{-1}(\bar{\mathfrak{b}})$. Then the map $\phi : S^{-1}R/S^{-1}I \rightarrow \bar{S}^{-1}(R/I)$, $s^{-1}r + S^{-1}I \mapsto \bar{s}^{-1}\bar{r}$, is an epimorphism with $\ker(\phi) = S^{-1}\mathfrak{b}/S^{-1}I$.

Proof. 1. Since $I \cap S = \emptyset$, $\bar{S} \in \text{Ore}_l(R/I)$. It remains to show that if $\bar{r}\bar{s} = 0$ for some elements $\bar{r} \in R/I$ and $\bar{s} \in \bar{S}$ then $\bar{t}\bar{r} = 0$ for some $\bar{t} \in \bar{S}$. The equality $\bar{r}\bar{s} = 0$ means that $i := rs \in I$. Hence, $S^{-1}R \ni \frac{r}{1} = is^{-1} = s_1^{-1}i_1$ for some elements $s_1 \in S$ and $i_1 \in I$ (since $S^{-1}I$ is an ideal of $S^{-1}R$). Therefore, $s_2s_1r = s_2i_1$ for some element $s_2 \in S$, and so $\bar{t}\bar{r} = 0$ where $\bar{t} = \bar{s}_2\bar{s}_1 \in \bar{S}$. Clearly, $(\mathfrak{a} + I)/I \subseteq \bar{\mathfrak{b}}$.

2. It is obvious that ϕ is an epimorphism. Since $\bar{S}^{-1}(R/I) \simeq S^{-1}(R/I)$ and $I \subseteq \mathfrak{b}$, we have $\ker(\phi) = S^{-1}\mathfrak{b}/S^{-1}I$. \square

Proposition 4.8 *Let R be a ring, $S \in \text{Den}_l(R, 0)$, $Q' := S^{-1}R$ and Q'^* be the group of units of Q' . Suppose that $S = R \cap Q'^*$ and one of the following statements holds:*

1. *Every left invertible element of Q' is invertible.*
2. *Every right invertible element of Q' is invertible.*
3. *The ring Q' is a left Noetherian ring.*
4. *The ring Q' is a right Noetherian ring.*
5. *The ring Q' satisfies ACC on left annihilators.*
6. *The ring Q' satisfies ACC on right annihilators.*
7. *The ring Q' does not contain infinite direct sums of nonzero left ideals.*

8. The ring Q' does not contain infinite direct sums of nonzero right ideals.

If $yz \in S$ for some elements $y, z \in R$ then $y, z \in S$.

Proof. If $q := yz \in S$ then $q \in Q'^*$. In particular, $y \cdot zq^{-1} = 1$ and $q^{-1}y \cdot z = 1$. If condition 1 (respectively, 2) holds then z (respectively, y) is a unit of Q'^* . So, if any of conditions 1 or 2 holds then $y, z \in Q'^*$, and so $y, z \in R \cap Q'^* = S$. Clearly, (3 \Rightarrow 5) and (4 \Rightarrow 6). So, it suffices to consider the case when one of the conditions 5-8 holds. Let $x := zq^{-1}$. Then $yx = 1$ in Q' . The idempotents $e_i := x^i y^i - x^{i+1} y^{i+1} = x^i e_0 y^i$, $i = 0, 1, \dots$, are orthogonal idempotents. Since one of conditions 5-8 holds we must have $e_i = 0$ for some i then $0 = y^i e_i x^i = y^i x^i e_0 y^i x^i = e_0 = 1 - xy$, i.e., $1 = xy = yx$. Hence, $x, y \in Q'^*$ and $x, y \in R \cap Q'^* = S$. \square

Corollary 4.9 1. Let R be a ring such that its largest left quotient ring $Q' := Q_l(R)$ satisfies one of conditions 1-8 of Proposition 4.8. If $yz \in S_l(R)$ for some elements $y, z \in R$ then $y, z \in S_l(R)$.

2. Let R be a ring such that its left quotient ring $Q' := Q(R)$ satisfies one of conditions 1-8 of Proposition 4.8. If $yz \in \mathcal{C}_R$ for some elements $y, z \in R$ then $y, z \in \mathcal{C}_R$.

Proof. 1. Statement 1 follows from Proposition 4.8 since $S_l(R) = R \cap Q_l(R)^*$ (Theorem 2.1.(1)).

2. Statement 2 is a particular case of statement 1. \square

Acknowledgements

I would like to thank the referee for comments.

References

- [1] V. V. Bavula, The algebra of integro-differential operators on an affine line and its modules, *J. Pure Appl. Algebra* **217** (2013) 495-529. (Arxiv:math.RA: 1011.2997).
- [2] V. V. Bavula, Characterizations of left orders in left Artinian rings, *J. Algebra Appl.* **13** (2014) no 6, 1-21. (Arxiv:math.RA:1212.3529).
- [3] V. V. Bavula, New criteria for a ring to have a semisimple left quotient ring, *J. Algebra Appl.* **6** (2015) no. 6, [28 pages] DOI: 10.1142/S0219498815500905. (Arxiv:math.RA:1303.0859).
- [4] V. V. Bavula, The largest strong left quotient ring of a ring, *J. Algebra* **439** (2015) 1-32. (Arxiv:math.RA:1310.1077).
- [5] V. V. Bavula, The largest left quotient ring of a ring, *Comm. Algebra* **44** (2016) no. 8, 3219-3261. (Arxiv:math.RA:1101.5107).
- [6] V. V. Bavula, Left localizations of left Artinian rings, *J. Algebra Appl.* **15** (2016) no. 9, 1650165, 38 pages. (Arxiv:math.RA:1405.0214).
- [7] V. V. Bavula, Criteria for a ring to have a left Noetherian largest left quotient ring, *Algebras Repr. Theory*, DOI:10.1007/s10468-017-9717-9, published online on 10 July 2017.
- [8] A. W. Chatters and C. R. Hajarnavis, Quotient rings of Noetherian module finite rings, *Proc. AMS* **121** (1994) no. 2, 335-341.
- [9] K. Goodearl and R. J. Warfield, *An Introduction to Noncommutative Noetherian rings*, Second Edition, London Mathematical Society Students Texts 61, Cambridge University Press, 2004.
- [10] A. W. Goldie, The structure of prime rings under ascending chain conditions. *Proc. London Math. Soc.* (3) **8** (1958) 589-608.
- [11] A. W. Goldie, Some aspects of ring theory, *Bull. London Math. Soc.* **1** (1969) 129-154.
- [12] A. W. Goldie, Semi-prime rings with maximum condition. *Proc. London Math. Soc.* (3) **10** (1960) 201-220.

- [13] A. V. Jategaonkar, Localization in Noetherian Rings, Londo Math. Soc. LMS 98, Cambridge Univ. Press, 1986.
- [14] C. R. Hajarnavis, On Small's Theorem, *J. London Math. Soc.* (2) **5** (1972) 596–600.
- [15] L. Lesieur and R. Croisot, Sur les anneaux premiers noethériens à gauche. *Ann. Sci. École Norm. Sup.* (3) **76** (1959) 161–183.
- [16] J. C. McConnell and J. C. Robson, Noncommutative Noetherian rings. With the cooperation of L. W. Small. Revised edition. Graduate Studies in Mathematics, 30. American Mathematical Society, Providence, RI, 2001. 636 pp.
- [17] G. Michler and B. Müller, The maximal regular Ore set of a Noetherian ring, *Arch. Math.* **43** (1984) 218–223.
- [18] B. Müller, Affine Noetherian PI-Rings Have Enough Clans, *J. Algebra* **97** (1985) 116–129.
- [19] J. C. Robson, Artinian quotient rings, *Proc. London Math. Soc.* (3) **17** (1967) 600–616.
- [20] L. W. Small, Orders in Artinian rings, *J. Algebra* **4** (1966) 13–41.
- [21] L. W. Small, Correction and addendum: Orders in Artinian rings, *J. Algebra* **4** (1966) 505–507.
- [22] L. W. Small and J. T. Stafford, Regularity of zero divisors, *Proc. London Math. Soc.* (3) **44** (1982) 405–419.
- [23] J. T. Stafford, Noetherian full quotient rings, *Proc. London Math. Soc.* (3) , **44** (1982) 385–404.
- [24] B. Stenström, Rings of Quotients, Springer-Verlag, Berlin, Heidelberg, New York, 1975.
- [25] H. Tachikawa, Localization and Artinian Quotient Rings, *Math. Z.* **119** (1971) 239–253.
- [26] R. B. Warfield, Quotient rings and localization for Noetherian rings. *J. Algebra* **72** (1981), no. 1, 166–182.

Department of Pure Mathematics
 University of Sheffield
 Hicks Building
 Sheffield S3 7RH
 UK
 email: v.bavula@sheffield.ac.uk