



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/126882/>

Version: Accepted Version

Article:

Good, P, Bamber, J, Halladay, K et al. (2018) Recent progress in understanding climate thresholds: ice sheets, the Atlantic meridional overturning circulation, tropical forests and responses to ocean acidification. *Progress in Physical Geography*, 42 (1). pp. 24-60. ISSN: 0309-1333

<https://doi.org/10.1177/0309133317751843>

(c) 2018, The Author(s). This is an author produced version of a paper published as, 'Good, P, Bamber, J, Halladay, K et al (2018). Recent progress in understanding climate thresholds: ice sheets, the Atlantic meridional overturning circulation, tropical forests and responses to ocean acidification. *Progress in Physical Geography*, 42 (1). pp. 24-60.' Reprinted by permission of SAGE Publications.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Recent progress in understanding climate thresholds: ice sheets, the Atlantic meridional overturning circulation, tropical forests and responses to ocean acidification

Peter Good¹, Jonathan Bamber², Kate Halladay¹, Anna Harper³, Laura Jackson¹, Gillian Kay¹, Bart Kruijt⁴, Jason Lowe¹, Oliver Phillips⁵, Jeff Ridley¹, Meric Srokosz⁶, Carol Turley⁷, Phillip Williamson⁸

¹Met Office Hadley Centre, Exeter, United Kingdom.

²School of Geographical Sciences, University of Bristol, UK

³College of Engineering, Mathematics, and Physical Sciences, University of Exeter, UK

⁴ALTERRA, Wageningen UR, PO box 47, 6700 AA Wageningen, Netherlands

⁵School of Geography, University of Leeds, Leeds, UK

⁶National Oceanography Centre, University of Southampton Waterfront Campus, Southampton, UK

⁷Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth PL1 3DH, UK

⁸School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK and Natural Environment Research Council, UK

Abstract

This article reviews recent scientific progress, relating to four major systems that could exhibit threshold behaviour: ice sheets, the Atlantic meridional overturning circulation (AMOC), tropical forests and ecosystem responses to ocean acidification. The focus is on advances since the Intergovernmental Panel on Climate Change Fifth Assessment Report (IPCC AR5). The most significant developments in each component are identified by synthesizing input from multiple experts from each field. For ice sheets, some degree of

irreversible loss (time-scales of millennia) of part of the West Antarctic Ice Sheet (WAIS) may have already begun, but the rate and eventual magnitude of this irreversible loss is uncertain. The observed AMOC overturning has decreased from 2004-2014, but it is unclear at this stage whether this is forced or is internal variability. New evidence from experimental and natural droughts has given greater confidence that tropical forests are adversely affected by drought. The ecological and socio-economic impacts of ocean acidification are expected to greatly increase over the range from today's annual value of around 400, up to 650 ppm CO₂ in the atmosphere (reached around 2070 under RCP8.5), with rapid development of aragonite undersaturation at high latitudes affecting calcifying organisms. Tropical coral reefs are vulnerable to the interaction of ocean acidification and temperature rise, and the rapidity of those changes, with severe losses and risks to survival at 2K warming above pre-industrial. Across the four systems studied, however, quantitative evidence for a difference in risk between 1.5K and 2K warming above pre-industrial levels is limited.

I Introduction

While some aspects of climate change can be viewed as becoming proportionately larger with increasing forcings, other aspects may feature more complex, nonlinear behaviour (e.g. Lenton et al., 2008). This can include abrupt and/or irreversible change, which may be associated with key thresholds. Such behaviour must be considered differently in assessments of the potential benefits of mitigation: it implies, for example, that certain impacts could be significantly different above certain levels of anthropogenic interference, although the impacts may not always be negative (Lenton, 2013). Further, thresholds involve systems moving out of the limits of currently observed behaviour, so they require deep

process understanding based on a broad range of observations (Kopp et al., 2016). In some cases, early warning of an approaching threshold may be possible (Lenton, 2011).

Clear evidence of threshold behaviour in the earth system is seen in the paleoclimate record (e.g. McNeall et al., 2011). For example, central Greenland temperatures inferred from ice cores show abrupt changes of 10°C within 100 years (Guillevic et al., 2013) during the last ice age, even though the forcing over this period has evolved smoothly. These changes are thought in part to be associated with changes in the ocean's thermohaline circulation (Broecker, 2003). Strong paleoclimate evidence also exists for threshold behaviour in major ice sheets and methane reservoirs (McNeall et al., 2011). Various different forms of abrupt shifts have been found in current climate models (Drijfhout et al., 2015), although primarily for different processes than explored in the present review.

This study focuses on four major systems that may feature threshold behaviour: ice sheets, the Atlantic Meridional Overturning Circulation (AMOC), tropical forests, and ecosystem responses to ocean acidification. The risk of significant change in these systems is not necessarily linked to large-scale warming alone. Patterns of precipitation can be important for the AMOC and tropical forests; tropical forests are also strongly affected by anthropogenic land-use and the direct effect of carbon dioxide, while ocean acidification arises directly from increased atmospheric CO₂ (although its impacts combine with those of ocean warming) and West Antarctic Ice Sheet (WAIS) stability is influenced by changes in ocean circulation.

Consequences of change in these systems range from amplified global warming through altered climate patterns, elevated sea-level and direct loss of biodiversity and ecosystem

services (see individual sections below for details). These systems can in principle interact with each other (Lenton et al., 2008), although this is explored in only a few studies.

Here we report primarily on new literature subsequent to that presented in the IPCC Fifth Assessment Report, AR5. As found by O’Neil et al. (2017) in an updated review of IPCC Reasons for Concern, the headline conclusions of AR5 still broadly hold, but there have been considerable advances in understanding. We also briefly consider (in the Conclusions) the difference in risk between 1.5K and 2K global mean warming above pre-industrial levels. This review was prepared using an iterative approach, by specialists both within and external to the Met Office Hadley Centre. Initial drafts of each section were prepared by the Met Office, then sent to external experts for review and editing (except that for Ocean Acidification; prepared by experts in Plymouth Marine Laboratory and University of East Anglia). The sections were revised accordingly by the Met Office, then sent to the external experts for a second review.

Each system is addressed in a separate section below, each with the following subsections: Introduction (the key issues for that system); Observations (relevant real-world observations); Potential for significant change (literature addressing the question of how likely substantial change is); Consequences (of significant change); Cautions (key scientific uncertainties); and Comparison with AR5. The key conclusions are summarised in Table 1.

104

105 **II Ice Sheets**

106

107 **1 Introduction**

108

109 Ice-sheet mass loss, from the Greenland and Antarctic ice sheets, is of concern due to its
110 potential impact on global and local sea level (Alley et al., 2005; Shepherd, 2012), and
111 potential amplification of global warming over long timescales as low-albedo land surface is
112 exposed (Hansen et al., 2008). The ice sheets are the largest potential source of future sea
113 level rise on decadal to millennial timescales. Greenland and Antarctica respectively contain
114 enough ice to raise mean sea level by 7.4 m and 58.3m (Vaughan et al., 2013). In addition,
115 rapid mass loss may have an influence on ocean circulation through a change in salinity and
116 hence density gradients (Yang et al., 2016).

117

118 In a state of equilibrium, an ice sheet loses mass (through surface melting, and the calving
119 and submarine melting of its outlet glaciers and ice shelves, Rignot et al., 2010; Depoorter et
120 al., 2013), at the same rate as it gains mass (through the accumulation of snowfall, Alley et
121 al., 2005). Increased ice sheet mass loss occurs through two main mechanisms. Increased
122 surface melt is largely driven by higher air temperatures and currently affects Greenland and
123 the Antarctic Peninsula. ‘Dynamic thinning’ (i.e. losses due to increased solid ice discharge
124 into the ocean) involves glacier acceleration and consequent increases in iceberg calving for
125 marine-terminating glaciers and ice shelves, occurring at the fringes of Greenland (Pritchard
126 et al., 2009), the Antarctic Peninsula (Wouters et al., 2015) and West Antarctica (Pritchard et
127 al., 2009; Bingham et al., 2012). This may be induced by increased temperature of the water
128 beneath floating ice or at the glacial terminus (Gille, 2014).

129

130 Ice shelves, the floating portions of outlet glaciers, play a key role in modulating the mass
131 balance of the Antarctic ice sheet. They buttress the inland glaciers, controlling the rate of
132 ice leaving the continent and entering the ocean (Dupont and Alley, 2005). Ice shelves are
133 exposed to the underlying ocean and may weaken (Furst et al., 2016) as ocean temperatures
134 rise (Depoorter et al., 2013). If they melt rapidly or break away, ice flow can accelerate,
135 causing net ice-sheet mass loss (De Angelis and Skvarca, 2003; Nick et al., 2009) which adds
136 to sea level rise. This impact of ice shelf break-up occurred on Larsen-B on the Antarctic
137 Peninsula in 2002, with the consequent acceleration of the glaciers as buttressing was
138 removed (De Rydt et al., 2015). Much of the West Antarctic Ice Sheet (WAIS) is grounded
139 on bedrock below sea level on retrograde slopes (deeper inland). This configuration is
140 inherently unstable and sensitive to small changes at the grounding line (where the ice begins
141 to float; Mercer, 1968; Schoof, 2007; Durand et al., 2011; Gudmundsson et al., 2012). A
142 small retreat of the grounding line resting on a retrograde slope thickens the ice at the
143 grounding line, in turn increasing the ice flux and inducing further retreat, and so on, until a
144 prograde slope is reached. Hence local thresholds exist (where the grounding line retreats to
145 a retrograde slope). This is known as the Marine Ice Sheet Instability (MISI), and
146 simulations have shown that this is a mechanism for rapid collapse of the WAIS (Gladstone
147 et al., 2012; Cornford et al., 2015; DeConto and Pollard, 2016; Arthern and Williams, 2017).

148

149 Iceberg calving has been implicated in the retreat and acceleration of glaciers where ice
150 shelves have disintegrated along the margins of the Greenland and Antarctic ice sheets,
151 indicating that they may be vulnerable to rapid ice loss through catastrophic disintegration
152 (Bassis and Jacobs, 2013). Processes such as fracture propagation in response to local stress
153 imbalances in the immediate vicinity of the glacier front; undercutting of the glacier terminus

by melting at or below the waterline; and bending at the junction between grounded and buoyant parts of an ice tongue combine to generate a feedback which accelerates mass loss through increased iceberg calving (Bassis and Walker, 2012). This is known as the Marine Ice Cliff Instability (MICI).

Surface meltwater stored in ponds and crevasses can weaken and fracture ice shelves, triggering their rapid disintegration (Scambos et al., 2004). This ice-shelf collapse results in an increased flux of ice from adjacent glaciers. This mechanism has been included in one model (Pollard et al., 2015), predicting a sea level rise from Antarctica of around 1m by 2100 (Deconto and Pollard, 2016). However, there is uncertainty in this process due to additional effects from surface transport of meltwater onto, across and away from ice shelves. The net result of this transport could either increase or decrease ice-shelf stability (Bell et al., 2017; Kingslake et al., 2017).

It is thought that no ice sheet would grow in Greenland if the current one were to be removed, even without human-induced warming, and hence it is a “relict” from the last Glacial Cycle that ended about 12 thousand years ago. The altitude of the ice sheet interior maintains the persistently cold temperatures required for the ice sheet to survive. There is a temperature threshold above which the Greenland ice sheet is no longer viable (Gregory and Huybrechts, 2006; Robinson et al., 2012). This is because, as temperatures increase, so does the area of summer melt, resulting in a lower surface elevation, causing further warming and increased melt (atmospheric temperature decreases with altitude). This positive feedback is known as the small ice cap instability, or melt-elevation feedback (e.g. Crowley and North, 1988; Levermann and Winkelmann, 2016).

Key issues addressed by recent studies include: what the observed ice-sheet loss implies for the rate of future global sea-level change, the potential long-term sea-level rise, and the possibility of abrupt or irreversible changes on timescales of a few hundred years.

2 Observed recent changes

Between 2002 and 2011 the West Antarctic Ice Sheet (WAIS) contributed 0.3 ± 0.1 mm yr⁻¹ to global sea-level rise (Peng et al., 2016). The majority of this loss has come from basal melt of ice shelves, and associated dynamical thinning, with half the basal melt arising from 10 small ice shelves in the Bellingshausen and Amundsen seas (Rignot et al., 2013). Of these, Pine Island glacier (Favier et al., 2014) and Thwaites glacier (Joughin et al., 2014) are the principal outlets of the WAIS that have rapidly thinned, retreated, and accelerated since the 1990's. Recent assessments indicate that Thwaites is contributing ~ 0.1 mm per year to sea-level rise, double that of the 1990s, and Pine Island glacier ~0.13 mm per year, however, there has been no acceleration in mass loss since 2008 (Medley et al., 2014). The spatial pattern of coincident changes in thickness across ice shelves of the Amundsen Sea suggests that the loss of grounded ice is the direct result of increased basal melting of the ice shelf, as a consequence of the inflow of warm water from the southern Pacific (Jacobs et al., 2011; Ha et al., 2014). Multi-decadal warming at the seabed in the Bellingshausen and Amundsen seas is linked to a shoaling of the mid-depth temperature maximum over the continental slope, allowing warmer, saltier water greater access to the continental shelf in recent years (Schmidtke et al., 2014). Before 2009, the glaciers of the Southern Antarctic Peninsula were in equilibrium, but have since been contributing significantly to sea level rise at a near-

constant rate of $0.16 \pm 0.02 \text{ mm yr}^{-1}$ (Wouters et al., 2015). The onset of this sudden and rapid mass loss appears to have a similar origin to that seen in the Amundsen Sea sector. The retrograde bedrock configuration is such that the mass loss is likely to be sustained for years to decades into the future, for this sector of Antarctica.

In addition there have been synchronous advances and retreats of the calving front of tide-water glaciers of the East Antarctic Ice Sheet (EAIS) (Miles et al., 2013). These appear to be associated with changes in the Southern Annular Mode (SAM), and consequently due to natural climate variability. However, there is evidence of ocean warming causing thinning of the Totten ice shelf, combined with a retreat of the grounding line (Silvano et al., 2016). A substantial area of ice sheet inland of Totten is below sea level, equivalent to 3.5m of sea level rise, and consequently the grounding line is potentially unstable due to the marine ice sheet instability. Evidence provided by Silvano et al. (2016) that warm circumpolar water does cross over the EAIS shelf break to cause rapid basal melt for several small ice shelves, suggests that EAIS could be more vulnerable to ocean heat fluxes than previously thought. Overall EAIS currently shows a gain in mass, through increased precipitation, with an implied sea level fall of 0.32 mm per year over 2009-2011 (Boening et al., 2012).

Estimates of overall sea level changes associated with net ice loss from Antarctica have been made by the gravity satellite, GRACE, between 2003 and 2014, at $0.25 \pm 0.2 \text{ mm}$ per year (Harig and Simons, 2015).

The Greenland ice sheet (GrIS) is losing mass as a result of both increased runoff due to surface melting and increased ice discharge from marine-terminating outlet glaciers (Rignot et al., 2008; Rignot et al., 2011; Sasgen et al., 2012; van den Broeke et al., 2009). The

Greenland mass loss over the period 2000-2005 contributed about 0.43 ± 0.09 mm yr⁻¹ of sea level rise. The rate has, however, been accelerating, with estimates for 2009-2012 of 1.05 ± 0.14 mm yr⁻¹ (Figure 2; Enderlin et al., 2014), and for 2011-2014 of 0.74 ± 0.14 mm yr⁻¹ (McMillan et al., 2016). The relative contribution of ice discharge (dynamic thinning) to total loss decreased from 58% before 2005 to 32% between 2009 and 2012. As such, 84% of the increase in mass loss after 2009 was due to increased surface runoff, as opposed to increased discharge (Enderlin et al, 2014). These observations support recent model projections that changes in surface mass balance driven, primarily, by increases in air temperature, rather than ice dynamics, will likely dominate the ice sheet's contribution to 21st century sea level rise (e.g. Goelzer et al., 2013; Vizcaino et al., 2015).

The glaciers in the southeast and northwest of Greenland sped up between 2000 and 2005 and have since stabilised or slowed (Enderlin et al, 2014). The slow down in the southeast has been compensated for by the northeast Greenland ice stream, which extends more than 600 km into the interior of the ice sheet, and is now undergoing sustained dynamic thinning, linked to regional warming, after more than a quarter of a century of stability (Khan et al., 2014). This sector of the Greenland ice sheet is of particular interest, because the drainage basin area covers 16% of the ice sheet, and numerical model predictions suggest no significant mass loss for this sector, leading to a possible under-estimation of future global sea-level rise (Khan et al, 2014). As for the Southern Antarctic Peninsula, the geometry of the bedrock and monotonic trend in glacier speed-up and mass loss suggests that dynamic loss of grounded ice in this region will continue in the near future (Khan et al., 2014).

3 Potential for significant change

The ice sheets can respond to climate change through accelerated discharge of freshwater to the ocean, and the associated sea level rise may be irreversible. Accelerated discharge, particularly from a marine ice sheet instability (MISI), has implications for the predictability of future sea level rise. The IPCC AR5 projections of sea level rise (Church et al., 2013) states that MISI may add tens of centimetres by 2100, but this mechanism was not quantified in the summary projections due to a lack of understanding. New studies here have focussed on the sea level contribution from the WAIS.

An ice flow model (Gagliardini et al., 2013) reveals that the Pine Island Glacier's grounding line is probably engaged in an unstable 40 km retreat (Figure 1; Favier et al., 2014). The associated mass loss increases substantially over the course of the simulations from an average value of 0.05 mm yr⁻¹ observed for the 1992-2011 period, up to and above 0.28 mm yr⁻¹, equivalent to 3.5-10 mm mean sea-level rise over the next 20 years (Favier et al., 2014). They find that mass loss remains elevated from then on, ranging from 0.16 to 0.33 mm yr⁻¹. Paleoclimate evidence (Johnson et al., 2014) for the early Holocene (a period of seasonal regional warming about 2K above pre-industrial) has revealed mass loss from the Pine Island Glacier at a rate comparable to present-day loss, but no collapse. Simulations for the adjacent Thwaites glacier, also in the Amundsen Sea embayment, indicate future mass losses are moderate (less than 0.25 mm yr⁻¹) over the 21st century but generally increase thereafter (Joughin et al., 2014). The likely time scale for collapse, based on various imposed ice shelf basal melt rates, is the time required for 100-200 km of grounding line retreat in the Thwaites Glacier system plus 200-1000 years for an actual collapse event (Joughin et al., 2014). Except possibly for the lowest-melt scenario used in the simulations, the results indicate that early-stage irreversible collapse has already begun (Joughin et al., 2014). One model includes the process of hydrofracture for Antarctic ice shelves (Pollard et al., 2015; DeConto and Pollard,

2016), associated with surface melt water forcing open crevasses, leading to ice shelf disintegration, and marine ice cliff instability. In this idealised simulation, hydrofracture caused a rapid deglaciation of WAIS on a timescale of only about 100 years (from the beginning of major retreat on the Antarctic Peninsula through to peak rate of sea-level rise around year 2140 – see their Figure 4c). The ice sheet collapse projected by Deconto and Pollard (2016) does not occur if the strong mitigation scenario of RCP2.6 is followed. Ice loss from the EAIS Wilkes basin may become substantial over timescales beyond a century (up to 3-4m sea-level rise after several millennia), with the loss irreversible above a threshold of regional ice loss (Mengel and Levermann, 2014).

Some new understanding of the paleo record has emerged. For Antarctica as a whole, there is evidence (Weber et al., 2014) for periods of relatively abrupt Antarctic mass loss following the Last Glacial Maximum (26-19 thousand years ago), possibly associated with a positive feedback involving ocean heat transport. It is likely that WAIS collapse occurred in the last interglacial (125 thousand years ago), when the Southern Ocean temperature anomaly exceeded 2-3°C (Sutter et al., 2016). However, the solar insolation is sufficiently different in this interglacial, that a similar spatial pattern of warming cannot be achieved through present-day increases in CO₂. One study (Levy et al., 2016) combined a range of regional and global paleoproxy information to further constrain the response of Antarctica during the early to mid-Miocene (23-14 million years ago), when CO₂ levels fluctuated between 280 and 500 ppm (equivalent to pre-industrial and a value that will be reached in the next few decades). They identify a peak warming period (16 million years ago) showing a consistent picture of global and regional warming, Antarctic ice sheet retreat, and a corresponding sea level rise of 10 to 20 m.

The Southern Ocean as a whole has not warmed significantly over the last decades (Armour et al., 2016). Instead, local warming has occurred, as deeper warm waters have been forced, by increased circumpolar winds, onto the Amundsen Sea continental shelf. Increasing winds are a consequence of global warming, the depletion of stratospheric ozone, or natural variability. Regardless of the cause of the increased windspeed, warm water has reached the continental shelf of the Bellingshausen and Amundsen seas. As a consequence it has been suggested that a critical threshold for grounding line retreat has already been passed for glaciers in the Amundsen Sea sector (Rignot et al., 2014). High ice shelf thinning rates for this and the Bellingshausen Sea sector of West Antarctica over the last two decades (Paolo et al., 2015) combined with the dramatic shift in mass imbalance of the Southern Antarctic Peninsula (Wouters et al., 2015) also point to a widespread shift in behaviour for this region.

It cannot be ruled out that the observed ice shelf thinning is a natural fluctuation rather than a consequence of anthropogenic forcing. Thus the likelihood of (partial) collapse of the WAIS has not yet been quantified, and requires improved modelling through ice sheet models fully coupled within global atmosphere-ocean climate models. Some progress has been made along these lines, with realistic ice shelf cavities now represented in ocean models (Beckmann et al., 1999; Dinniman et al., 2007; Losch, 2008; Mathiot et al., 2017), and the idealised simulations of MISOMIP (Asay-Davis et al., 2016).

4 Potential consequences

If a collapse of the WAIS were to occur, it would lead to a global sea level rise of up to 3.3 m (Bamber et al., 2009) on timescales (from the onset of collapse) of 100 years (Deconto and Pollard, 2016) to 400 years (Cornford et al., 2015; Golledge et al., 2015). This inference is supported by records of past sea level rise. Under the low emissions scenario, RCP2.6, sea level contributions remain small, and a collapse of WAIS does not occur in simulations (Golledge et al., 2015; DeConto and Pollard, 2016). For Antarctica as a whole, paleoproxy evidence from the Miocene (Levy et al., 2016) suggests potential sea-level rise of the order of 10-20m, for CO₂ levels near 500ppm.

Surface melt from the Greenland Ice Sheet may influence local ocean circulation, through stratification reducing convection in the Labrador Sea (Yang et al., 2016), and consequently local sea level change, perhaps by 5 cm, in the North-West Atlantic (Swingedouw et al., 2013; Howard et al., 2014).

On centennial to millennial time scales, Antarctic Ice Sheet melt can moderate warming in the Southern Hemisphere, by up to 10°C regionally, in a 4 x CO₂ scenario (Swingedouw et al., 2008). This behaviour stems from the formation of a cold halocline in the Southern Ocean, which limits sea-ice cover retreat under global warming and increases surface albedo, reducing local surface warming. In addition, Antarctic ice sheet melt, by decreasing Antarctic Bottom Water formation, restrains the weakening of the Atlantic meridional overturning circulation, which is an effect of the bi-polar oceanic seesaw (Pedro et al., 2011). Consequently, it appears that Antarctic ice sheet melting strongly interacts with climate and ocean circulation globally. It is therefore necessary to account for this coupling in future climate and sea-level rise scenarios.

5 Cautions (uncertainties).

While substantial progress in understanding has been made, it is still unclear what the recent observed changes imply for long-term future ice-sheet loss (Wouters et al., 2013), due to regional natural variability. Some observations suggest that there may be a natural cycle of increase and decrease in the rates of mass loss from coastal glaciers (Murray et al., 2010), so short-term trends should not necessarily be extrapolated into the future (Wouters et al., 2013). Indeed many Greenland glaciers, which accelerated in the early 2000s have since slowed (Moon et al., 2012; Enderlin et al., 2014). There is a possibility that solid earth movement, in response to ice loss, may influence the bedrock slopes, and so reduce further ice loss from the West Antarctic Ice Sheet (Konrad et al., 2015), delaying WAIS collapse by as much as 5000 years.

6 Comparison with AR5

Of the key findings summarised in Table 1, the main new points since AR5 are: observational evidence (Enderlin et al. 2014) that, from Greenland, the proportion of loss from surface melt has increased, becoming more consistent with long term model projections; evidence that some degree of irreversible loss from the WAIS may have begun (Favier et al. 2014, Joughin et al. 2014, Rignot et al. 2014, Wouters et al. 2015); and indications that the East Antarctic Ice Sheet (Miles et al. 2013) and northeast Greenland (Khan et al. 2014) may be more sensitive to climate change than previously expected.

378

379 **III AMOC**

380

381 **1 Introduction**

382

383 The Atlantic Meridional Overturning Circulation (AMOC) transports large amounts of heat
384 northwards in the Atlantic Ocean, resulting in a milder climate in northwest Europe and the
385 North Atlantic than would otherwise be experienced (for recent reviews of AMOC behaviour
386 and observations see Srokosz et al., 2012; Srokosz and Bryden, 2015; Buckley and Marshall,
387 2016). The IPCC AR5 report (Collins et al., 2013) concludes that it is very likely that the
388 AMOC will weaken over the 21st century, although there is a large spread in the predicted
389 weakening among climate models. A large or rapid (over a decadal time scale) reduction in
390 the AMOC would likely have substantial impacts on global climate, although a collapse
391 (rapid shutdown) of the AMOC by 2100, however, was judged as very unlikely (Collins et
392 al., 2013). These assessments have not changed since the previous IPCC assessment.

393

394 **2 Observed recent changes**

395

396 The RAPID-MOCHA array has been observing the AMOC at 26°N since 2004 and now has
397 acquired over a decade of data (Rayner et al., 2011; McCarthy et al., 2015b). This dataset has
398 revealed large variability on timescales from daily to interannual (see Figure 2). This
399 included a large (30%), temporary decrease in AMOC strength over 2009-2010 (McCarthy et
400 al., 2012; Bryden et al., 2014), which resulted in cooling in the upper North Atlantic Ocean in
401 2010 north of the latitude of the RAPID array and warming to the south (Cunningham et al.,
402 2013; Bryden et al., 2014). This decrease began with a strengthening of the upper mid-ocean

recirculation in early 2009 and was compounded by a slowdown in the northward Ekman transport and Gulf Stream flow in late 2009 and early 2010 (accounting for 61%, 27% and 12% of the slowdown, respectively; Bryden et al., 2014). This decrease was well outside the range predicted for interannual AMOC variability in coupled ocean-atmosphere models (McCarthy et al., 2012; Roberts et al., 2014); note that model resolution may be an issue here. Roberts et al. (2013b) reproduced this AMOC decrease using an initial condition ensemble of ocean simulations driven by observed surface forcing (albeit with too weak an AMOC), suggesting that the atmosphere may have had a dominant role in the temporary AMOC decrease. However, the origin of, and complete explanation for, the 2009-10 event remains uncertain. To-date no explanations have fully accounted for the changes in Lower North Atlantic Deep Water (LNADW at 3000 to 5000m depth) and the lack of change in the Upper North Atlantic Deep Water (UNADW between 1000 and 3000m depth) described by McCarthy et al. (2012).

The links between changes in the AMOC, upper ocean heat content and atmospheric response represent an active area of research. For example, the ocean has been implicated in the re-emergence of sea surface temperature anomalies from the winter of 2009-10 during the following early winter season of 2010-11, which contributed to the persistence of the negative winter North Atlantic Oscillation (NAO) and wintry conditions in northern Europe (Taws et al., 2011). Such behaviour may lead to improved predictions of the NAO and winter conditions (Maidens et al., 2013; Scaife et al., 2014).

The AMOC overturning has also decreased from 2004-2014 (Figure 2; Srokosz and Bryden, 2015; Frajka-Williams et al., 2016); the majority of this was due to a weakening of the geostrophic flow (Smeed et al., 2014: who analysed the first eight and a half years of data).

428 This trend has been associated with decreases in subsurface density in the subpolar gyre,
429 similar to those seen in climate models when there is a reduction in the AMOC (Robson et
430 al., 2014). It is unclear at this stage whether the decrease is forced (so part of a longer-term
431 downturn). Some recent work suggests that it may be part of a downturn after a previous
432 increase (Jackson et al., 2016; Frajka-Williams, 2015). Statistical tests on the observations
433 (Smeed et al. 2014) suggested that the AMOC decrease is statistically significant, even if the
434 low AMOC event of 2009-10 is excluded. Roberts et al (2014) found similar trends as part of
435 natural variability in 2 out of 14 global climate models, and in all models considered when
436 corrections are made to include more realistic high frequency variability. They concluded that
437 more than a decade of observations would be required to detect and attribute an
438 anthropogenic weakening of the same trend as observed over the period 2004-2012 (although
439 this rate does not appear to have been maintained since 2012, Figure 2). In an earlier model
440 study, Roberts et al. (2013a) estimated that a minimum of two decades of data would be
441 required to detect an anthropogenic trend in the AMOC, based on multimodel 1% per year
442 CO₂-forced experiments. This means that the existing AMOC observing system would need
443 to make measurements until at least 2024. Another study (Mercier et al., 2015) analysed
444 repeat hydrographic data along a Greenland to Portugal section from 1993 to 2010, finding an
445 overall decline in the AMOC over that period. Send et al. (2011) observed a decreasing trend
446 in the transport of the deep western boundary current at 16°N (one component of the AMOC)
447 over a similar period. In the South Atlantic, based on a combination of satellite altimeter and
448 hydrographic observations, Dong et al. (2015) note that “since 2010 the MOC has exhibited
449 low values when compared to the 1993–2011 mean values.” Linking the observations of the
450 AMOC obtained at different latitudes by different observational means remains a significant
451 challenge (Elipot et al., 2014; Elipot et al., 2017). Landerer et al. (2015) show that satellite
452 observations of ocean bottom pressure may provide a useful method for examining latitudinal

coherence of signals, however it is currently restricted to detrended data and regions of steep topography.

A recent paper (Rahmstorf et al., 2015) suggested that the trend detected at 26°N is part of an ‘exceptional slowdown’ of the AMOC. They find a relationship between sea surface temperatures and the AMOC in a climate model and then use reconstructions of surface temperatures from paleoclimate records to suggest that there has been a weakening that is unprecedented over the last 1000 years. There are, however, inherent uncertainties around both the relationship used and the temperature reconstructions, raising questions over whether the results are robust. In contrast, a recent reconstruction of the AMOC in the South Atlantic since 1870 (Lopez et al., 2017) suggests that it is presently in a stronger than normal phase. Ultimately, all proxies for the AMOC, such as temperature, coastal sea level (Ezer, 2015; McCarthy et al., 2015b; Frajka-Williams, 2015), or gravity measurements (Landerer et al., 2015) need to be tested and verified against direct observations of AMOC strength, over the time scales of interest, if they are to be used to infer robustly its behaviour over longer periods.

Future observations and research will improve our assessments of past and on-going AMOC changes. In this context note that the Overturning in the Subpolar North Atlantic Program (OSNAP; Lozier et al., 2017) deployed instruments in 2014 along a line from Canada to Greenland to Scotland, to observe the AMOC in the subpolar gyre, complementing the 26.5°N observations in the subtropical gyre. Meanwhile in the South Atlantic there are trans-basin observations of the AMOC beginning to be made at 34.5°S (SAMBA – South Atlantic MOC Basin-wide Array; Meinen et al., 2013; Ansorge et al., 2014). Recently, a new component of the AMOC, the so-called East Greenland spill jet, has been identified from a

year of mooring observations (von Appen et al., 2014), but its importance in the long-term for the overall AMOC remains to be confirmed.

3 Potential for significant change

Paleoclimate studies have suggested that some abrupt changes to climate may have been caused by the AMOC switching from an “on” state, where it transports heat northwards in the Atlantic, to an “off” state (Rahmstorf, 2002). Paleooceanographic studies of the AMOC and abrupt climate change over the last glacial cycle have been recently summarised by Lynch-Stieglitz (2017), who found that the evidence for changes in the AMOC associated with the Younger Dryas and many Heinrich events is strong, and there is some evidence for AMOC changes over many Dansgaard-Oeschger events. However, the ultimate causal links between the co-incident changes in the AMOC and climate are less clear. Further, these studies are hard to interpret in terms of future change, as the conditions in which past abrupt changes occurred were very different to the present. It is thought that abrupt changes may be related to the existence of bistability (where both “on” and “off” states of the AMOC can exist for a given forcing) as predicted by theoretical models of the Atlantic (e.g. Stommel, 1961), Earth system models of intermediate complexity (Rahmstorf et al., 2005) and studies with low resolution global circulation models (Hawkins et al., 2011; Manabe and Stouffer, 1988). Statistical properties of the timeseries of AMOC strength may give warning of approaching a threshold, however a new model study finds that centuries of data from a reliable proxy would be required (Boulton et al., 2014).

There have been many model studies suggesting that the stability of the AMOC might be affected, or even controlled, by whether the AMOC imports or exports fresh water from the

Atlantic, since this can indicate the presence of a positive or negative advective feedback. De Vries and Weber (2005) found that the fresh water transport by the AMOC into the Atlantic was an important indicator of stability in their experiments. The relative importance, for AMOC stability, of freshwater export/import by the AMOC itself, is unclear, however. Other factors have subsequently been found to be important in determining AMOC stability. For example, Jackson (2013) found that, while the overturning component of freshwater transport does partially indicate the sign of the advective feedback in a GCM, the transport of fresh water by the gyres can also play a crucial role. Swingedouw et al. (2013) also found that gyre transports can affect the magnitude of AMOC reduction. The presence of eddies is lacking in many models (due to low resolution) but studies with an eddy resolving model show that they can also affect the fresh water transport (den Toom et al., 2014). Mecking et al. (2016) found that the AMOC in an eddy-permitting model was very slow to recover from an input of fresh water. They found that the freshwater transport by the AMOC was important for maintaining the weak AMOC state, and hypothesised that this transport was changed by the eddy-permitting resolution. Understanding these controls on AMOC stability is crucial to constraining the likelihood of AMOC collapse. A recent paper (Liu et al., 2017) has noted that biases in the models may affect the estimated probability of an AMOC collapse.

4 Potential consequences

A collapse in the AMOC would cause a large relative cooling over the North Atlantic, which would have wide-ranging impacts, such as cooling in the northern hemisphere, warming in the southern hemisphere, and a southward shift in the Inter Tropical Convergence Zone, causing substantial changes in tropical precipitation, (Vellinga and Wood, 2008; Jackson et

al., 2015).

The Amazon is one region sensitive to change in the AMOC, but the impacts are uncertain. A recent study by Parsons et al. (2014) found that a reduction in the AMOC caused an increase in vegetation over the Amazon, due to a change in precipitation seasonality (despite a reduction in annual mean precipitation). This contrasts with an earlier study (Bozbiyik et al., 2011), which found that a reduction in the AMOC causes large reductions in Amazon vegetation due to precipitation reductions.

Other studies have concentrated on impacts over Europe. Jackson et al. (2015) confirmed an earlier study by Woollings et al. (2012), that a reduction in AMOC strength could drive an increase in the number of winter storms across Europe. Jackson et al. (2015) also showed that the increase in winter storms resulted in greater precipitation over western coasts in Northern Europe, despite a general reduction of precipitation over the northern hemisphere from a cooling-induced reduction in evaporation. They also found regional changes in summer precipitation across Europe, similar to those associated with Atlantic sea temperature found by Sutton and Dong (2012). Haarsma et al. (2015) examined the relationship between European atmospheric circulation and the AMOC across the CMIP5 ensemble. They also found an influence of AMOC strength on European summer precipitation and cloud cover.

One impact of the AMOC suggested recently is its possible role in the so-called global warming “hiatus” (Chen and Tung, 2014), though various other explanations for the hiatus have been proposed. Another recently observed impact is the reduction in uptake of CO₂ by the Atlantic Ocean due to the weakening of the AMOC over the period 1990 to 2006 (Perez et al., 2013).

553

554 Another recent focus of attention has been the role of the AMOC in sea level rise (SLR) on
555 the eastern seaboard of the USA (Ezer, 2015; Goddard et al., 2015; McCarthy et al., 2015a;
556 Yin et al., 2009). In particular, Goddard et al. (2015) demonstrate that the 2009-10 temporary
557 downturn in the AMOC led to an unprecedented 12.8 cm sea level rise along the coast north
558 of New York over the same period. They show that this rise was a 1-in-850 year event.
559 Furthermore, they note that, “Unlike storm surge, this event caused persistent and widespread
560 coastal flooding even without apparent weather processes. In terms of beach erosion, the
561 impact of the 2009–2010 SLR event is almost as significant as some hurricane events.” This
562 observed short-term change provides evidence for what has been previously suggested only
563 by modelling studies (Levermann et al., 2005), that a slowdown or collapse of the AMOC
564 would lead to significant sea level rise on the eastern seaboard of the USA.

565

566 **5 Cautions**

567

568 There are large inter-model differences in projections of future AMOC decline amongst
569 models used for the IPCC AR5 report. Reintges et al. (2016) found that uncertainties in
570 AMOC projections were dominated by uncertainties in fresh water changes amongst the
571 models, with contributions from uncertainties in both changes in surface fresh water fluxes
572 and ocean fresh water transports.

573

574 Several studies have shown that many GCMs have biases in the fresh water transport of the
575 AMOC (importing instead of exporting fresh water), and that this might affect the simulated
576 stability of the AMOC. The source of this bias is unclear. Jackson (2013) attributed the bias
577 to an over-evaporative Atlantic in the model and notes the difference from observations in

578 salinity profiles in the South Atlantic. Liu et al. (2014) suggested that the presence of a
579 double Atlantic ITCZ (a common GCM bias) results in a tropical salinity bias that stabilises
580 the AMOC. Another source of uncertainty is the transport of saline water from the Indian
581 Ocean to the Atlantic by eddies that are shed from the Agulhas current. Current GCMs do not
582 resolve the scales required to correctly represent these eddies, but a recent study by Biastoch
583 and Böning (2013) used a high resolution nested model to resolve this region. They found
584 that a southwards shift of the southern hemisphere westerlies (as is expected to occur under
585 anthropogenic climate change) results in a decrease in salinity transport into the Atlantic,
586 however this change in salinity is small and has little impact on the AMOC. The lack of
587 eddy-resolving resolutions in current GCMs might also have an impact on the transient
588 response of the AMOC to increased freshwater input (Weijer et al., 2012; Mecking et al.,
589 2016).

590
591 There is also substantial uncertainty about the future inputs of freshwater into the Atlantic,
592 particularly since the climate models lack dynamic ice sheet models which could
593 substantially speed up the input of freshwater from the Greenland ice sheet. Separate studies
594 including additional freshwater inputs from the Greenland ice sheet find that projected
595 changes do not have major impacts on the AMOC, although there is uncertainty about future
596 changes in freshwater fluxes from Greenland (Bamber et al., 2012). Böning et al. (2016)
597 concluded that meltwater from the Greenland ice sheet has resulted in a gradual freshening of
598 Labrador Sea, but that this has had no significant impact on the AMOC yet. A recent study
599 found that the MOC became less sensitive to fresh water inputs when CO₂ levels were high,
600 because of increases in stratification caused by warming and changes in the wind-driven
601 circulation (Swingedouw et al., 2015). Another study suggests that future increases in
602 precipitation over the Arctic, leading to increased freshwater flux into the North Atlantic

could also affect the AMOC (Bintanja and Selten, 2014: see Methods). The most recent GCM study that accounted for Greenland melting (Bakker et al., 2016) concluded that Greenland Ice Sheet "melting affects AMOC projections, even though it is of secondary importance."

6 Comparison with AR5

The main development since the publication of AR5 has been the updated observations of overturning from the RAPID-MOCHA array (Smeed et al., 2014; Srokosz and Bryden, 2015), which shows a decline over the period 2004-2014. Studies suggest that this was related to decadal variability. This does not preclude the presence of a longer-term decline, but the time series is too short to make definitive statements. Continuous observations from the existing AMOC observing system until at least around 2024, combined with further understanding of the past record from multiple proxy information, and more model studies, will be required to isolate a forced decline in the AMOC. Another key finding is the unprecedented rise in US east coast sea level associated with the 2009-10 downturn in the AMOC (both of which subsequently recovered). Although this is a change on a shorter time scale than the 100 year time scale associated with climate change, it shows that changes in the AMOC may have impacts on multiple time scales. Finally, the inclusion of Greenland melting in GCMs has been found to affect AMOC projections, but appears to be of secondary importance.

626

627 **IV Tropical forests (Amazon focus)**

628

629 **1 Introduction**

630

631 Tropical forests regulate and supply to society a range of services, which bring benefits at
632 global to local scales. As well as sustaining high biodiversity they influence climate through
633 biogeochemical (carbon cycle) and biophysical (water and energy) mechanisms. Over the
634 period 1990-2007, intact tropical forests took up carbon at the rate of $1.2 \pm 0.4 \text{ Pg C year}^{-1}$
635 (corresponding to about half the global land carbon sink), compared with $0.50 \pm 0.08 \text{ Pg C}$
636 year^{-1} by the boreal forests (Pan et al., 2011), and around $1.1 \pm 0.8 \text{ Pg C year}^{-1}$ losses of
637 forest carbon stocks to the atmosphere through land use change over 2000-2009 (Settele et
638 al., 2014). However, large droughts can cause elevated mortality rates, especially for larger
639 trees (Phillips et al., 2010; Nepstad et al., 2007; da Costa et al., 2010; McDowell and Allen,
640 2015) and temporary shifts from ecosystem carbon sink to carbon source (Phillips et al.,
641 2010; Lewis et al., 2011; Gatti et al., 2014). Estimates of the impact of the 2005 and 2010
642 Amazon droughts (mostly through increases in tree mortality during and lagging the
643 droughts) stand at 1.6 and 2.2 Pg C, respectively (Phillips et al., 2009; Lewis et al., 2011).

644

645 Tropical forests are subject to interacting effects from atmospheric CO_2 , climate and land-use
646 change (e.g. Coe et al., 2013). Land-use change effects include direct deforestation, and
647 accidental 'leakage' fires (where intentional fires spread accidentally over a wider forest area).
648 Forest fragmentation (an important by-product of deforestation) lengthens the forest edge.
649 Since most forest fires occur at the forest edge, because of greater human activity,
650 fragmentation accelerates the rate of forest erosion by fire. Deforestation increases albedo

and reduces evapotranspiration, altering climate both locally and downwind; aerosols from deforestation fires may also reduce rainfall (Marengo et al., 2011). Climate change could alter vegetation productivity and mortality, both directly, and indirectly by modifying fire behaviour. Increased atmospheric CO₂ may increase tree growth (where nutrients are not limiting), but also increase tree mortality from lianas (vines). The full vegetation response to CO₂ and climate changes may take decades to be completely realised (Jones et al., 2009), and the subsequent carbon release even longer.

The AR5 finds that large-scale dieback due to climate change alone is unlikely by the end of this century (*medium confidence*). However, it states with *medium confidence* that “severe drought episodes, land use, and fire interact synergistically to drive the transition of mature Amazon forests to low-biomass, low-statured fire-adapted woody vegetation” (Settele et al., 2014). New research has largely, but not exclusively, focused on the Amazon: due in part to early climate model projections of climate-driven Amazon dieback (Cox et al., 2000). Severe Amazonian droughts in the last decade have provided insights on forest responses to extreme dry conditions. In addition to forest and climate monitoring, throughfall exclusion and prescribed-burn experiments have allowed in-situ study of the effects of longer-term drought and fire. Numerical studies have also increased in number and progress has been made in putting the early results into context.

2 Observations

New studies have given greater confidence that the Amazon represents a long-term net carbon sink (Brienen et al., 2015; Espirito-Santo et al., 2015; Gatti et al., 2014), but also suggest (Brienen et al., 2015) that its strength has weakened progressively as tree mortality

rates increase (Figure 3). Potential drivers for the mortality increase include more frequent or more severe droughts, and feedbacks of faster growth on mortality, resulting in shortened tree longevity (Bugmann and Bigler, 2011).

The response of trees to elevated CO₂ remains uncertain. Some recent longer-term studies of tropical tree rings (van der Sleen et al., 2015; Battipaglia et al., 2015; Groenendijk et al., 2015) have found no evidence for sustained increases in tree growth or carbon uptake, but as Brien et al. (2012) point out, tree-ring studies are subject to biases which preclude robust statements about ecosystem-level changes. So far, multi-decadal plot data been used systematically to probe recent growth trends at continental scale only in Amazonia (Brien et al., 2015). Here they indicate a long-term increase in growth rates since the 1980's, as well as a lagging increase in mortality rates, consistent with a long-term growth stimulation, such as by CO₂.

There is greater confidence that Amazon forests are adversely affected by drought. There has been new work on the response to the 2010 drought, and also the 1997 and 2005 events (Tomasella et al., 2013). A new attribution study (Shiogama et al., 2013) of the 2010 drought showed that, while sea surface temperature anomalies in the tropical Pacific and Atlantic likely increased the probability of drought (in addition to biomass burning; Marengo et al., 2011), unforced atmospheric variability probably also played a large role. Atmospheric measurements (Gatti et al., 2014) confirmed earlier plot-based findings (Phillips et al., 2010; Lewis et al., 2011) that the Amazon switched from a temporarily from a net carbon sink to a source during the 2010 drought. Compared to these short-term natural droughts, the impact was seen to be much stronger in the long-term persistent experimental droughts induced by a forest throughfall exclusion experiment in eastern Amazonia (da Costa et al., 2014), and by

2014, 13 years of 50% throughfall exclusion at Caxiuana had caused a cumulative biomass loss of $45.0 \pm 2.7\%$ (Rowland et al., 2015). Consistent with previous suggestions that effects of a single drought persist for several years (Saatchi et al., 2013; Phillips et al., 2010), even during the anomalously wet year of 2011, the Amazon was still estimated to be carbon neutral overall (Gatti et al. 2014; possibly due to lagged effects of the 2010 drought).

Drought mortality, especially in larger trees, is a major pathway for carbon release (Phillips et al., 2010; Nepstad et al., 2007; da Costa et al., 2010), but underlying mechanisms are not well understood (Meir et al., 2015), and poorly represented in current vegetation models (Powell et al., 2013). However, hydraulic failure is suggested as the primary cause from the Caxiuana drought experiment (Rowland et al., 2015). A study of detailed plot-level responses to the 2010 drought in several sites, compared to other years (Doughty et al., 2015) suggested that trees may prioritise growth in response to reduced photosynthesis from short-term drought, leaving some trees more vulnerable to mortality. In contrast, the long-term (> 12 years) response to persistent experimental rainfall exclusion (Rowland et al., 2015), shows no decline in photosynthetic capacity (although photosynthesis may have declined if mean stomatal conductance declined), but an increase in leaf dark respiration in tree taxa vulnerable to drought mortality (possibly a sign of drought stress). It has been suggested that early warning of drought mortality events may be plausible based on observations of tree properties (Camarero et al., 2015). Overall, the AR5 viewpoint of persistent drought causing a shift towards lower statured, low-biomass forest is retained (Rowland et al., 2015).

Drought can also cause abrupt increases in fire-induced tree mortality over sub-seasonal timescales (Brando et al., 2014), driving a lagged increase in carbon emissions over subsequent years. More than 85,500 km² of the southern Amazon was burnt by understorey

fires during 1999-2010, with evidence for a strong climate control on fire (Morton et al., 2013).

Forest responses to warming remain uncertain, and more forest warming field experiments are needed (Cavaleri et al., 2015). In one such experiment (Slot et al., 2014), although respiration increased with warming, thermal acclimation did occur. A new meta study integrating experimental and observational results (Vanderwel et al., 2015) suggests that acclimation could potentially half increases in leaf dark respiration over the century, compared with null model expectations that ignore acclimation. On the other hand, a global-scale analysis of interannual variability has suggested (Anderegg et al., 2015) that nighttime respiration in tropical forests may be highly sensitive to warming.

Some new observational studies have found substantial reductions in evapotranspiration in some (Oliveira et al., 2014; da Silva et al., 2015; Panday et al., 2015), but not all (Rodriguez et al., 2010) deforested regions. The full effects of deforestation over the Xingu river basin (a southeast tributary of the Amazon) may have been masked by climate variability (Panday et al., 2015).

3 Potential for significant change

A recent review of wider sources of evidence (Coe et al., 2013) identified South/South-East Amazonia as particularly vulnerable: due to high deforestation rates locally and in the upwind savanna region; its susceptibility to small climate shifts (being in a transitional climate zone between forest and savanna); and greater climate model agreement on future rainfall reductions in this region (compared to the West Amazon). A review for African rainforests

(Malhi et al., 2013) highlighted similar points: while deforestation rates have historically been relatively low over Africa, there is potential for significant future increases; and African forests have climate close to the limit of rainforest sustainability. Models tend to predict rainfall reductions(increases) over western(central) equatorial Africa (James et al., 2014). Rainfall decreases over west equatorial Africa can be large in some models (James et al., 2014), although the forest response is hard to predict.

The observations and field experiments summarised above have given greater confidence that forests are significantly affected by drought, emphasising the importance of extreme climate events in causing extensive tree losses (through drought and heat mortality and increased fire). On the other hand, some acclimation of trees to warming has been demonstrated. These vegetation responses are not well understood or represented in current dynamic vegetation models (e.g. Powell et al., 2013).

A recent study using three terrestrial biosphere models (Zhang et al., 2015) found the direction and severity of precipitation change to be critical. A greater model consensus for a projected lengthening and deepening of the dry season in Amazonia was found in CMIP5 compared with CMIP3 (Joetzjer et al., 2013), although it is unclear whether this represents a statistically significant improvement in model performance. A new observationally constrained model study (Boisier et al., 2015) found a greater lengthening of dry seasons over the Amazon than projected by unconstrained models (as found, with a different method, by Shiogama et al., 2011). A key uncertainty in terms of impacts is the extent to which forest whole-ecosystem responses to climate change might be protected by the wide functional diversity in many tropical forests. The work of Fauset et al. (2012) from Ghana suggests that by not accounting for biological diversity, most vegetation models may underestimate forest

resilience.

In terms of land-use, it has become clearer that as well as direct deforestation, the indirect effects of deforestation on forest fragmentation, and on climate locally and downwind, must be considered in regulatory policies (e.g. Harper et al., 2014; Lawrence and Vandecar, 2015; Brinck et al., 2017; Wu et al., 2017). While a 70% decline has been reported in deforestation in the Brazilian Amazon between 2005 and 2013 (Nepstad et al., 2014), maintaining low levels of deforestation in a sustainable manner remains a challenge (Nepstad et al., 2014). Despite the reduction in deforestation since 2004, around half of the area burnt during 1999-2010 over the southern Amazon occurred during 2007 and 2010, when deforestation activity was relatively low, suggesting that fire-free land use needs to be encouraged as well as reducing direct deforestation (Morton et al., 2013). Achieving similar reductions in deforestation in other countries may be challenging due to issues with governance and monitoring capability (DeFries et al., 2013), but for Indonesia, accounting for spatial variation in costs and benefits of avoided deforestation does reveal low cost options (Graham et al., 2017).

Various positive feedbacks (fire-vegetation and climate-vegetation; eg. Hirota et al., 2011; Staver et al., 2011; Hoffmann et al., 2012) exist that could lead to abrupt reductions in forest cover, for relatively small change in external forcings, and inhibit reversibility, but the processes are poorly characterised. The timescale of any abrupt change depends on the processes and the spatial scale considered, and may be strongly dependent on stochastic climate variability. Very locally, loss of tree cover from fire or drought mortality can occur over seasonal timescales in the event of severe drought. However, one model study (Higgins and Scheiter, 2012) found that, while transitions between vegetation states may be abrupt

locally, over continental and larger scales the effect on the carbon cycle is much more gradual (because the timing of transitions varies with location). It has been suggested (Verbesselt et al., 2016) that temporal autocorrelation in satellite data provides evidence for threshold behaviour in forests – and potential for monitoring forest resilience. Evidence for alternative stable states has recently been reported in vegetation height (Xu et al., 2016) as well as in tree fractional cover (Staver et al., 2011; Pausas and Dantas, 2017; Hirota et al., 2011). However, the spatial scale over which abrupt or irreversible change might extend depends on the strength of these positive feedbacks compared to environmental control on vegetation cover, and demonstrating whether alternative stable states exist over large scales is challenging (Good et al., 2016; Staal et al., 2016). Spatial interaction between forest and savanna can reduce the area over which alternative stable states exist (a clear exploration is provided by Staal et al., 2016). Indeed, recent observational work has challenged the notion that savanna and forest represent ‘alternative stable states’ over large areas of the tropics (Veenendaal et al., 2015; Wuyts et al., 2017). Local fire-vegetation feedbacks are seen in prescribed burning experiments (Silverio et al., 2013), but over large scales, only 10% of the locations burnt in the 2005 drought showed repeated burning by 2010 (Morton et al., 2013). A model study (Moncrieff et al., 2014) found that the area over which alternative stable states are possible could be large in present-day conditions, but declined substantially with future CO₂ increases. Hoffmann et al. (2012) noted that forest-fire feedbacks themselves can be sensitive to tree growth-rates – and hence to climate change.

4 Potential consequences

The observations summarised above give greater confidence that the Amazon represents a net carbon sink, but this appears to have been declining at least for a decade, and the long-term

future of this sink is uncertain. Persistent drought would be likely to cause a transition to lower statured, lower biomass forest, from mortality of larger trees (Rowland et al., 2015), and severely threaten biodiversity (Esquivel-Muelbert et al., 2017). Extreme events over the Amazon could have a large impact on the global carbon cycle and offset or counteract potential regional increases in biomass (Reichstein et al., 2013).

While it is accepted that tropical deforestation tends to reduce evapotranspiration locally, consequent changes in rainfall are complex and depend on the scale and pattern of deforestation (Lawrence and Vandecar, 2015). Including deforestation feedback on climate (via precipitation) is key in assessing river runoff change (Stickler et al., 2013; Lima et al., 2014). Stickler et al. (2013) estimate that when feedbacks on climate are included, the sign of change in hydropower generation potential for the plants under construction on the Amazonian Xingu River is reversed, declining to 25% of maximum plant output by 2050 under business-as-usual land-use projections (with 40% deforestation by 2050). The net runoff response in the Amazon is basin-dependent (Lima et al., 2014) and is sensitive to the scale and pattern of deforestation (Lawrence and Vandecar, 2015). Deforestation may reduce the length of the wet season, such that large-scale expansion of agriculture in Amazonia may be unsustainable (Oliveira et al., 2013; Arvor et al., 2014). Land-use-driven stream warming of at least 3-4K (in mean daily maximum temperature) in southeastern Amazonian has also been observed (Macedo et al., 2013) - well above the ~1K threshold for changes in fish physiology, growth and behaviour. Overall, multiple ecosystem services need to be taken into account when considering optimal management (Donoso et al., 2014).

5 Cautions (uncertainties)

Accurate projections are partly limited by the availability of observations. Inaccessibility of tropical forests increases reliance on remote sensing data, but also makes verifying remote sensing data (notably, precipitation, biomass, and vegetation productivity data) challenging. New studies have shown that great caution is required in interpreting satellite retrievals of variability in greenness (Morton et al., 2014). The tropical forest biome constitutes one of the largest terrestrial carbon sinks, but it is also associated with relatively large uncertainties (Pan et al., 2011), because of its great ecological complexity, huge scale, and multiple anthropogenic processes affecting it (Lewis et al., 2015).

There is substantial uncertainty in the CMIP5 projections of future precipitation in tropical forest regions, (Collins et al., 2013), although there is greater degree of inter-model agreement in some seasonal changes, such as a lengthening and a deepening of the dry season in Amazonia (Joetzjer et al., 2013; Boisier et al., 2015). However, the representation of present-day Amazon precipitation still contains large biases. Large uncertainties are also associated with the modelled response of vegetation to temperature (Galbraith et al., 2010; Huntingford et al., 2013) and to CO₂ (Rammig et al., 2010). Processes of direct mortality from fire and drought (and effects of fire on aerosol) are often either unrealistic or absent from models (e.g. Powell et al., 2013), and the range of plant functional types is extremely limited in relation to the large biodiversity and hence range of potential tree-level responses in most tropical forests.

6 Comparison with AR5

The new literature has not altered the broad, general view given in AR5. Probably the

876 greatest advances lie in increased confidence that, at least over the Amazon, drought
877 adversely affects the forest carbon balance – and improved understanding of how this occurs.
878 Many uncertainties remain, and estimating the likelihood of basin-scale forest dieback
879 remains challenging.
880

881

882 **V Responses to Ocean Acidification**

883

884 **1 Introduction**

885 Increased concentrations of atmospheric CO₂ reduce seawater pH, increase the solubility of
886 calcium carbonate (reducing saturation state), and cause other chemical changes, together
887 known as ocean acidification. The biogeochemical, ecological and societal implications of
888 ocean acidification have received greatly increased research attention during the past decade
889 (Riebesell and Gattuso, 2015; Mathis et al., 2015). Ocean acidification risks and impacts
890 were included as a component of climate change in the IPCC's Fourth Assessment Report,
891 with more detailed analyses in the Fifth Assessment Report, particularly by Working Group
892 II (Portner et al., 2014).

893 Analyses of geological ocean acidification events and modelling studies show that physico-
894 chemical recovery from perturbations in ocean carbonate chemistry of similar magnitude to
895 projected changes takes many thousands of years (Zeebe and Ridgwell, 2011), due to slow
896 rates of deep ocean mixing and of chemical equilibration with seafloor sediments. The rate
897 of CO₂ increase today is estimated to be around 10 times faster than any natural ocean
898 acidification event during the past 66 million years (Honisch et al., 2012; Zeebe et al., 2016).
899 The longterm hysteresis effects are inherent in the response of global ocean chemistry to
900 atmospheric CO₂ forcing, and there is only very limited capacity to accelerate future recovery
901 by actively removing CO₂ from the atmosphere (Mathesius et al., 2015). Species' extinctions
902 are necessary irreversible.

903 Many different thresholds for ocean acidification impacts can be considered under conditions
904 of steadily increasing atmospheric CO₂ levels; the focus here is on increased solubility of

calcium carbonate (in particular, the saturation state for aragonite, the form of carbonate in the shells and structures of many marine organisms) and the risk of rapid loss of tropical corals.

2 Observed recent changes

The IPCC Fifth Assessment Report (Rhein et al., 2013) provided decadal measurements of ocean carbonate chemistry in near-surface waters at three oceanic monitoring sites; and other datasets are also now available (WMO, 2014; Bates, 2017). All these observations unequivocally show decreasing pH in the upper ocean at rates (-0.0011 to -0.0024 yr^{-1}) closely matching those expected from rising atmospheric CO_2 . Both physical and biological factors are responsible for the spatial and temporal variability in these datasets; whilst seasonality is usually smoothed-out for trend analyses (WMO, 2014), it is of high ecological importance, determining the conditions experienced by marine organisms (Sasse et al., 2015). There is much less temporal variability of pH in the ocean mid-waters and at greater depth; however, there are also fewer longterm measurements. Atlantic observations (Woosley et al., 2016) confirm an anthropogenically-driven decrease in surface pH of ~ 0.0021 yr^{-1} with greatest changes in the top $\sim 1000\text{m}$; however, some decrease also occurs at greater depths. Such changes are superimposed on a natural decrease of pH with depth, with North Atlantic seafloor values generally being in the range $7.70 - 7.75$ (Vazquez-Rodriguez et al., 2012) compared to a global mean surface value of ~ 8.1 , and typical seasonal ranges of $7.9 - 8.3$.

Correlations between observed ocean acidification and biological or ecosystem changes are not necessarily causal, since other environmental factors are also likely to be involved. The strongest observational evidence relates to ocean acidification effects on pteropods (planktonic snails) in the Southern Ocean and northeast Pacific (Bednarsek et al., 2014a;

929 Bednarsek et al., 2012; Bednarsek et al., 2017); on cultivated oysters (Barton et al., 2015); on
930 warm-water corals, and at natural CO₂ vents (discussed below).

931

932 Longterm reductions of up to ~30% in the natural calcification and growth rates of tropical
933 corals have been reported in several studies (e.g. Silverman et al., 2014). Linkage to ocean
934 acidification has been demonstrated by *in situ* treatments of a natural coral community in the
935 Great Barrier Reef (Albright et al., 2016). When water chemistry was restored to pre-
936 industrial conditions by short-term alkalinity enrichment, coral growth rates increased by
937 ~7%.

938 Observations at natural, shallow-water CO₂ vents consistently show marked decreases in
939 overall biodiversity as pH declines (Hall-Spencer et al., 2008; Fabricius et al., 2011; Gambi et
940 al., 2016). Microbes in sediment are also affected (Raulf et al., 2015). Non-calcifying
941 seaweeds and sea grasses out-compete calcifying organisms under such high CO₂, low pH,
942 conditions, although some genetic adaptation of the latter can occur (Garilli et al., 2015).

943 **3 Potential for significant change**

944 Experimental studies have shown that many marine species are likely to be negatively
945 affected from future ocean acidification if high CO₂ emissions continue, with risk of
946 ecosystem alterations at the global scale (CBD, 2014; Gattuso et al., 2015; Nagelkerken and
947 Connell, 2015). Taxonomic variability in biotic responses to ocean acidification is, however,
948 high. Furthermore, many interactions occur with temperature, food availability and other
949 stressors (Wittmann and Portner, 2013; Ramajo et al., 2016; Kroeker et al., 2013; Kroeker et
950 al., 2017); responses may be sex-specific (Ellis et al., 2017); and impacts on behaviour,
951 competition and predator-prey relationships are complex (Nagelkerken and Munday, 2016;
952 Nagelkerken et al., 2017). Whilst the potential for evolutionary adaptation is largely

unknown (Sunday et al., 2014) , the sensitivity of populations could be shaped by regional adaptation to local conditions causing differences between geographically separated populations of the same species (Calosi et al., 2017).

Marine ecosystems are susceptible to non-linear changes occurring over just a few years (regime shifts; Mollmann et al., 2015) that cannot be easily reversed once thresholds, that may be of different kinds, are exceeded (Mumby et al., 2011; Hughes et al., 2013; Plaganyi et al., 2014). Two such ocean acidification-related thresholds were projected (Steinacher et al., 2013) in the context of allowable carbon emissions: aragonite undersaturation in the Southern Ocean, and the carbonate chemistry conditions necessary for warm-water coral reef survival.

Hauri et al. (2016) used a multi-model ensemble to determine changes in aragonite saturation state (Ω) around Antarctica and southern South America in an unabated CO₂ emissions scenario (RCP 8.5). The monthly occurrence of aragonite undersaturation ($\Omega < 1.0$) at the surface and at 100m water depth increased rapidly in most of these areas (Figure 4), particularly between 2040 - 2070 when atmospheric CO₂ levels are projected to be 500 - 650 ppm.

Similar effects are projected for the Arctic Ocean, where all surface waters north of 66° are projected to be unsaturated for aragonite by 2100 under RCP 8.5 (Popova et al., 2014; Qi et al., 2017). Regional differences are, however, greater - with surface undersaturation expected to have already occurred in the Siberian shelves and Canadian Arctic Archipelago (i.e. with current atmospheric CO₂ values of ~400 ppm), but not until the 2080s in the Barents and Norwegian seas (at ~ 900 ppm). The ecological significance of aragonite unsaturation is that

such conditions are chemically corrosive to unprotected shells made of that form of carbonate, e.g. those of pteropods (Bednarsek et al., 2014b; Bednarsek et al., 2017).

Coral exoskeletons are also made of aragonite: the depth distribution of coldwater corals is closely correlated with the aragonite saturation horizon (Guinotte et al., 2006; Jackson et al., 2014), whilst the calcification rate of both coldwater and tropical corals is sensitive to saturation state, responding semi-linearly over a wide range of values (McCulloch et al., 2012; Comeau et al., 2013). The dead unprotected reef-like structures of coldwater corals are especially susceptible to dissolution (Hennige et al., 2015).

Most tropical coral reefs occur in waters where $\Omega > 3.0$ (Manzello et al., 2014; Mongin et al., 2016), and that value has been used as a threshold for modelling climate change impacts (Steinacher et al., 2013). Whilst tropical coral growth can continue where $\Omega < 3.0$ (Comeau et al., 2013; Shamberger et al., 2014), growth rates need to exceed bioerosion (Andersson and Gledhill, 2013) and to be sufficiently rapid to allow reef recovery between temperature-induced bleaching events (Frieler et al., 2013). In theory, tropical corals could avoid the risk of bleaching by colonizing new sites where water temperatures have previously been too cool (Couce et al., 2013). However, the rate of current change may be too rapid for that to occur – and there are many geological precedents for ‘coral reef crises’, involving mass extinctions during geological warming and/or ocean acidification events (Kiessling and Simpson, 2011). Based on these considerations, many coral researchers consider atmospheric levels of ~350 ppm CO₂ to be the ‘safe’ limit to ensure coral reef survival (ISRS, 2015).

4 Potential consequences

The potential consequences of future ocean acidification are extremely wide-ranging,

particularly for high emission scenarios. They include physico-chemical impacts (reduction in seawater capacity to absorb further CO₂); species-specific physiological and behavioural changes; perturbations in marine community processes, ecosystem functions and biogeochemical feedbacks; and changes in ocean ecosystem services, with societal effects on food security, coastal protection and climate regulation. The scale of the biological and socio-economic changes is, however, uncertain.

An overall reduction in marine diversity and abundances is expected to occur in a high CO₂ world (Nagelkerken and Connell, 2015); nevertheless, not all species will be negatively affected. Some marine species that may be favoured also provide societal benefits, e.g. seagrasses (Garrard and Beaumont, 2014), but not all. Thus ‘nuisance’ species, such as jellyfish, seem generally tolerant of ocean acidification (Hall-Spencer and Allen, 2015).

With regard to the carbonate undersaturation threshold identified above, the loss of pteropods from polar oceans would have wider consequences for food-webs, also affecting higher predators (fish, seabirds and sea mammals) of high commercial or conservation value, even if those groups are not directly affected by ocean acidification. Increasing acidification in the Southern Ocean represents a risk to another key pelagic species, Antarctic krill. The hatch-rate for krill eggs decreases markedly at $p\text{CO}_2$ values > 1000 μatm (Kawaguchi et al., 2013), and major reduction in their abundance could also jeopardise the entire ecosystem.

The potential loss of tropical coral reefs would have major consequences for coastal protection, tourism and fisheries, with the global economic value of those ecosystem services estimated to be up to ~ \$1000 billion per year (Brander et al., 2012). However uncertainties

1025 in economic costs are high, and many other factors, in addition to ocean acidification, are
1026 affecting the future health and survival of coral reefs.

1027 **5 Cautions (uncertainties).**

1028 Many uncertainties remain regarding ocean acidification impacts in the context of specific
1029 thresholds (Pandolfi, 2015) and interactions with other stressors (CBD, 2014; Gattuso et al.,
1030 2015). The scaling-up of impacts from organisms to communities, food webs, ecosystems
1031 and economic impacts is challenging (Andersson et al., 2015; Ekstrom et al., 2015; Turley,
1032 2017) – particularly since ocean acidification impacts do not act on their own, but co-occur
1033 with other stressors, both climate-related (warming, de-oxygenation and sea-level rise)
1034 (Gattuso et al., 2015; Howes et al., 2015; Kroeker et al., 2017) and non-climate-related
1035 (pollution, over-fishing and habitat loss) (Breitburg et al., 2015). Furthermore, coastal
1036 ecosystems seem likely to be at greatest risk from ocean acidification, but these are inherently
1037 complex and difficult to simulate in models because of interactions with sediment processes
1038 and riverine inputs (Artioli et al., 2014), and other factors causing local variability in
1039 carbonate chemistry(Chan et al., 2017).

1040 **6 Comparison with AR5**

1041 Since IPCC AR5, many ocean acidification studies have demonstrated variability in
1042 environmental conditions and biological responses, and the complexity of multi-stressor
1043 interactions. Such research therefore may seem to have increased, rather than reduced
1044 uncertainty. Nevertheless, understanding of ocean acidification and its impacts has
1045 significantly improved: observations have greater geographical coverage, integrating
1046 chemical and biological measurements, whilst new meta-analyses and assessments have
1047 confirmed previously-identified patterns and have also provided additional insights.
1048 Furthermore, greater attention has been given to important topics such as palaeo- ocean

1049 acidification events; socio-economic modelling; acclimatization and adaptation; and the
1050 vulnerability of cold-water corals.

1051 Many of those more recent studies relate to the thresholds outlined here. In particular, there
1052 is now greater confidence that extensive aragonite undersaturation, with major ecological
1053 consequences, would occur throughout the water column in high latitudes within a few years
1054 of atmospheric CO₂ exceeding 450-500 ppm, and that warming will need to be well below
1055 2K to avoid damaging interactions between ocean acidification and temperature for tropical
1056 coral reefs.

1057

1058

1059

1060 **VI Conclusions**

1061

1062 This report reviews the major new advances in understanding of four systems with potential
1063 for climate thresholds, focussing on progress since IPCC AR5. Advances are reported in the
1064 context of observed recent changes, the potential for significant change, and the associated
1065 consequences. The key findings are summarised in Table 1. Overall, compared to AR5, a
1066 large number of studies have added further detail to our understanding of these systems, but
1067 the broad headline summaries of AR5 have not greatly changed.

1068

1069 Declines have been observed (in the Greenland and West Antarctic ice sheets, the AMOC,
1070 and ocean corals) that could be partly driven by anthropogenic activity, although the role of
1071 natural variability is uncertain. For the West Antarctic Ice Sheet, some degree of irreversible
1072 collapse may already have begun. For tropical forests the picture is more mixed, with some
1073 long-term increases in carbon storage, but also evidence of a more recent weakening in the
1074 Amazon carbon sink.

1075

1076 For various reasons, long term maintenance of detailed observing systems is critical. Early
1077 warning of approaching thresholds may be possible, as well as attribution of change to
1078 anthropogenic or natural drivers. Further observations are also needed to improve the
1079 models. Current numerical models have improved, but still suffer from biases, and lack key
1080 processes or sufficient spatial resolution. Detailed process-based observations are needed, to
1081 separate different drivers, and mechanisms of response, and forced change from internal
1082 variability. In each of the four systems, there are a range of drivers of change (e.g. CO₂,
1083 atmospheric temperature, regional ocean temperatures - affected by ocean circulation as well

1084 as large scale warming, surface winds, precipitation, fire and atmospheric composition).
1085 Further, the systems can have different mechanisms of response (e.g. dynamical thinning of
1086 outlet glaciers versus surface mass balance for Greenland; or, for tropical forests, productivity
1087 versus mortality, and also changes in allocation of new carbon and inter-species competition).
1088 For tropical forests and ocean biological organisms, the potential for evolutionary adaptation
1089 is a key unknown. Field experiments have provided key information for tropical forests and
1090 ocean acidification, and more are required.

1091

1092 For these systems, there is only limited quantitative information about the difference, in
1093 likelihood of crossing a threshold, between futures reaching 1.5 and 2K global-mean
1094 warming above pre-industrial levels. For ice-sheets and the effects of ocean acidification
1095 (combined with warming) on marine ecosystems, it is reasonable to assume that the
1096 likelihood of crossing a critical threshold is higher for a 2K world than a 1.5K world. For
1097 Greenland, rates of mass loss and sea level rise are a non-linear function of the temperature
1098 increase because of the combined effect of dynamic thinning at the margins and the
1099 temperature-elevation feedback (Applegate et al., 2015). A simplified model study of this ice
1100 sheet suggested that the global-mean warming threshold for irreversible loss could be only
1101 0.8–3.2K (best estimate 1.6°C) above pre-industrial (Robinson et al., 2012); while one long-
1102 term coupled model simulation found the threshold of zero surface mass balance may be
1103 crossed somewhere between 2 and 3K above pre-industrial levels (Vizcaino et al., 2015). For
1104 ocean acidification, there is now greater confidence that extensive aragonite undersaturation
1105 (with major ecological consequences) will occur in high latitudes if atmospheric CO₂ exceeds
1106 450-500 ppm, and that warming will need to be well below 2K to avoid risk of damaging
1107 interactions between ocean acidification and temperature for tropical coral reefs.

1108

For the ice sheets, AMOC and tropical forests, the potential consequences of crossing a threshold (section 4 for each system - e.g. sea-level rise from decline in ice sheets) are in general better constrained than the likelihood (or timing) of crossing the threshold under particular forcing scenarios. Given this, we suggest that the risk of collapse in such systems could be managed by ongoing detailed monitoring, including of variables that might give early warning of collapse; and by assessment of the potential timescales and impacts of collapse using theory and models (however, for ocean acidification, while there is a real risk of crossing thresholds in ecosystems this century, the potential impacts are complex and poorly understood, due to possible interactions amongst different species). Ongoing model development and analysis will help target observations and will improve our understanding of the likelihood of collapse. These recommendations are similar to those of NRC (2013), with the additional focus on timescales and impacts of collapse.

Acknowledgements

This work was supported by the Joint UK DECC/Defra Met Office Hadley Centre Climate Programme (GA01101). MAS was funded by the NERC RAPID-AMOC programme. OLP was supported by the European Research Council Advanced Grant, ‘Tropical Forests in the Changing Earth System’. PW and CT were supported by the UK Ocean Acidification research programme funded by NERC, Defra and DECC. BK, KH and GK were funded by European Union’s Seventh Framework Programme AMAZALERT project (282664). AH was funded under NERC grant NE/K016016/1: Process-based Emergent Constraints on Global Physical and Biogeochemical Feedbacks and by the Joint Weather and Climate Research Programme (JWCRP)‘.

Figures




Figure 1. Evidence (from Favier et al. 2014) that the Pine Island Glacier's grounding line is probably engaged in an unstable 40 km retreat, due to the retrograde bedrock slope. Left: map of bedrock elevation, with the grounding lines for 2009 (white) and 2011 (purple) shown. Right: bedrock height (solid black line) and geometry of the glacier centreline produced by the Elmer/Ice ice-flow model at time (t) = 0 (dotted line) and after 50 years of a melting scenario (red line).





Figure 2. AMOC transport measured at 26.5°N (Smeed et al., 2016). The gray line represents the 10 day filtered measurements, while the red line was produced using a 180 day running mean. Clearly visible are the low AMOC event in 2009-10 and the overall decrease in strength over the measurement period.



1152

1153 Figure 3. Trends in net above-ground biomass change, productivity and mortality rates, for
 1154 321 plots, weighted by plot size (after Brien et al. 2015).

1155




Figure 4. Area-weighted ensemble mean duration (months per year) of aragonite undersaturation at the surface (solid lines) and at 100m depth (dashed lines) for three sectors of the Southern Ocean (Bellingshausen Sea, Weddell Sea and East Antarctica), the central Chilean coast and the Patagonian shelf over the period 1900-2100, with future projections based on RCP 8.5. From Hauri et al. (2016).

System	Key findings
Ice sheets	• From Greenland, the proportion of loss from surface melt has increased, becoming more consistent with long term model projections. The bedrock topography of the WAIS lends itself to an inherently unstable ice sheet. Some degree of irreversible loss may have begun, although the eventual magnitude and rate of this irreversible loss is uncertain. • There are indications that the East Antarctic Ice Sheet (EAIS) and the northeast Greenland ice stream may be more sensitive to climate change than previously expected. • New paleoclimate evidence for: 1) periods of relatively abrupt Antarctic mass loss following the last glacial maximum; 2) during the early Holocene (sustained warming ~2K above pre-industrial), WAIS mass loss rates comparable to present-day, but no WAIS collapse. • Modelling studies indicate that ice sheet mass loss can be largely avoided under the RCP2.6 scenario. • Significant loss from WAIS will occur on timescales of 100-1000 years.
AMOC	• The observed AMOC overturning has decreased from 2004-2014, linked with decreases in subsurface density in the subpolar gyre. It is unclear at this stage whether this AMOC decrease is forced or is internal variability.

	• There was an unprecedented rise in US east coast sea level associated with the 2009-10 downturn in the AMOC (both of which subsequently recovered).
Tropical forests	• Greater confidence that tropical forests are adversely affected by drought. • New climate models continue to suggest that basin-scale Amazon dieback from climate alone (as in an early study) is not typical. However, these studies lack some key processes. • There remains a high level of uncertainty regarding future changes
Ocean acidification	• Global trends in ocean acidification driven by increasing CO₂ concentrations are superimposed on a dynamic natural system • Many factors affect variability in biological response; these are now much better understood • Extensive aragonite undersaturation in high latitudes can be expected if atmospheric CO₂ exceeds 450-500 ppm, with effects on key zooplankton and marine food-webs • Tropical coral reefs seem highly vulnerable to the interaction of ocean acidification and warming, with major economic consequences relating to coastal erosion, storm protection, fisheries and tourism.

Table 1. Key new findings, for each system.

1169 **References**

- 1170 Albright R, Caldeira L, Hosfelt J, et al. (2016) Reversal of ocean acidification enhances net
1171 coral reef calcification. *Nature* 531: 362-365; doi: 310.1038/nature17155.
- 1172 Alley RB, Clark PU, Huybrechts P, et al. (2005) Ice-sheet and sea-level changes. *Science*
1173 310: 456-460.
- 1174 Anderegg WRL, Ballantyne AP, Smith WK, et al. (2015) Tropical nighttime warming as a
1175 dominant driver of variability in the terrestrial carbon sink. *Proceedings of the*
1176 *National Academy of Sciences of the United States of America* 112: 15591-15596.
- 1177 Andersson AJ and Gledhill D. (2013) Ocean Acidification and Coral Reefs: Effects on
1178 Breakdown, Dissolution, and Net Ecosystem Calcification. *Annual Review of Marine*
1179 *Science, Vol 5* 5: 321-348.
- 1180 Andersson AJ, Kline DI, Edmunds PJ, et al. (2015) Understanding Ocean Acidification
1181 Impacts on Organismal to Ecological Scales. *Oceanography* 28: 16-27.
- 1182 Ansorge IJ, Baringer M, Campos EJD, et al. (2014) Basin-Wide Oceanographic Array
1183 Bridges the South Atlantic. *Eos* 95: 53-54.
- 1184 Applegate PJ, Parizek BR, Nicholas RE, et al. (2015) Increasing temperature forcing reduces
1185 the Greenland Ice Sheet's response time scale. *Climate Dynamics* 45: 2001-2011.
- 1186 Armour KC, Marshall J, Scott JR, et al. (2016) Southern Ocean warming delayed by
1187 circumpolar upwelling and equatorward transport. *Nature Geoscience* 9: 549-+.
- 1188 Arthern RJ and Williams CR. (2017) The sensitivity of West Antarctica to the submarine
1189 melting feedback. *Geophysical Research Letters* 44: 2352-2359.
- 1190 Artioli Y, Blackford JC, Nondal G, et al. (2014) Heterogeneity of impacts of high CO₂ on the
1191 North Western European Shelf. *Biogeosciences* 11: 601-612.
- 1192 Arvor D, Dubreuil V, Ronchail J, et al. (2014) Spatial patterns of rainfall regimes related to
1193 levels of double cropping agriculture systems in Mato Grosso (Brazil). *International*
1194 *Journal of Climatology* 34: 2622-2633.
- 1195 Asay-Davis XS, Cornford SL, Durand G, et al. (2016) Experimental design for three
1196 interrelated marine ice sheet and ocean model intercomparison projects: MISMIP v. 3
1197 (MISMIP+), ISOMIP v. 2 (ISOMIP+) and MISOMIP v. 1 (MISOMIP1).
1198 *Geoscientific Model Development* 9: 2471-2497.
- 1199 Bakker P, Schmittner A, Lenaerts JTM, et al. (2016) Fate of the Atlantic Meridional
1200 Overturning Circulation: Strong decline under continued warming and Greenland
1201 melting. *Geophysical Research Letters* 43: 12252-12260.
- 1202 Bamber J, van den Broeke M, Ettema J, et al. (2012) Recent large increases in freshwater
1203 fluxes from Greenland into the North Atlantic. *Geophysical Research Letters* 39.
- 1204 Bamber JL, Riva REM, Vermeersen BLA, et al. (2009) Reassessment of the Potential Sea-
1205 Level Rise from a Collapse of the West Antarctic Ice Sheet. *Science* 324: 901-903.
- 1206 Barton A, Waldbusser GG, Feely RA, et al. (2015) Impacts of Coastal Acidification on the
1207 Pacific Northwest Shellfish Industry and Adaptation Strategies Implemented in
1208 Response. *Oceanography* 28: 146-159.
- 1209 Bassis JN and Jacobs S. (2013) Diverse calving patterns linked to glacier geometry. *Nature*
1210 *Geoscience* 6: 833-836.
- 1211 Bassis JN and Walker CC. (2012) Upper and lower limits on the stability of calving glaciers
1212 from the yield strength envelope of ice. *Proceedings of the Royal Society a-*
1213 *Mathematical Physical and Engineering Sciences* 468: 913-931.
- 1214 Bates NR. (2017) Twenty Years of Marine Carbon Cycle Observations at Devils Hole

1215 Bermuda Provide Insights into Seasonal Hypoxia, Coral Reef Calcification, and
 1216 Ocean Acidification. *Frontiers in Marine Science* 4: DOI:10.3389/fmars.2017.00036.
 1217 Battipaglia G, Zalloni E, Castaldi S, et al. (2015) Long Tree-Ring Chronologies Provide
 1218 Evidence of Recent Tree Growth Decrease in a Central African Tropical Forest. *Plos*
 1219 *One* 10.
 1220 Beckmann A, Hellmer HH and Timmermann R. (1999) A numerical model of the Weddell
 1221 Sea: Large-scale circulation and water mass distribution. *Journal of Geophysical*
 1222 *Research-Oceans* 104: 23375-23391.
 1223 Bednarsek N, Feely RA, Reum JCP, et al. (2014a) *Limacina helicina* shell dissolution as an
 1224 indicator of declining habitat suitability owing to ocean acidification in the California
 1225 Current Ecosystem. *Proceedings of the Royal Society B-Biological Sciences* 281.
 1226 Bednarsek N, Klinger T, Harvey CJ, et al. (2017) New ocean, new needs: Application of
 1227 pteropod shell dissolution as a biological indicator for marine resource management.
 1228 *Ecological Indicators* 76: 240-244,
 1229 <https://doi.org/210.1016/j.ecolind.2017.1001.1025>.
 1230 Bednarsek N, Tarling GA, Bakker DCE, et al. (2014b) Dissolution Dominating Calcification
 1231 Process in Polar Pteropods Close to the Point of Aragonite Undersaturation. *Plos One*
 1232 9.
 1233 Bednarsek N, Tarling GA, Bakker DCE, et al. (2012) Extensive dissolution of live pteropods
 1234 in the Southern Ocean. *Nature Geoscience* 5: 881-885.
 1235 Bell RE, Chu WN, Kingslake J, et al. (2017) Antarctic ice shelf potentially stabilized by
 1236 export of meltwater in surface river. *Nature* 544: 344-+.
 1237 Biastoch A and Boning CW. (2013) Anthropogenic impact on Agulhas leakage. *Geophysical*
 1238 *Research Letters* 40: 1138-1143.
 1239 Bingham RG, Ferraccioli F, King EC, et al. (2012) Inland thinning of West Antarctic Ice
 1240 Sheet steered along subglacial rifts. *Nature* 487: 468-471.
 1241 Bintanja R and Selten FM. (2014) Future increases in Arctic precipitation linked to local
 1242 evaporation and sea-ice retreat. *Nature* 509: 479-+.
 1243 Boening C, Lebsack M, Landerer F, et al. (2012) Snowfall-driven mass change on the East
 1244 Antarctic ice sheet. *Geophysical Research Letters* 39.
 1245 Boisier JP, Ciais P, Ducharne A, et al. (2015) Projected strengthening of Amazonian dry
 1246 season by constrained climate model simulations. *Nature Climate Change* 5: 656-+.
 1247 Boning CW, Behrens E, Biastoch A, et al. (2016) Emerging impact of Greenland meltwater
 1248 on deepwater formation in the North Atlantic Ocean. *Nature Geoscience* 9: 523-+.
 1249 Boulton CA, Allison LC and Lenton TM. (2014) Early warning signals of Atlantic
 1250 Meridional Overturning Circulation collapse in a fully coupled climate model. *Nature*
 1251 *Communications* 5.
 1252 Bozbiyik A, Steinacher M, Joos F, et al. (2011) Fingerprints of changes in the terrestrial
 1253 carbon cycle in response to large reorganizations in ocean circulation. *Climate of the*
 1254 *Past* 7: 319-338.
 1255 Brander LM, Rehdanz K, Tol RSJ, et al. (2012) The economic impacts of ocean acidification.
 1256 *Climate Change Economics* 3: 1250002 doi: 1250010.1251142/S2010007812500029.
 1257 Brando PM, Balch JK, Nepstad DC, et al. (2014) Abrupt increases in Amazonian tree
 1258 mortality due to drought-fire interactions. *Proceedings of the National Academy of*
 1259 *Sciences of the United States of America* 111: 6347-6352.
 1260 Breitburg DL, Salisbury J, Bernhard JM, et al. (2015) And on Top of All That... Coping with
 1261 Ocean Acidification in the Midst of Many Stressors. *Oceanography* 28: 48-61.
 1262 Brienens RJW, Gloor E and Zuidema PA. (2012) Detecting evidence for CO2 fertilization
 1263 from tree ring studies: The potential role of sampling biases. *Global Biogeochemical*
 1264 *Cycles* 26.

1265 Brienens RJW, Phillips OL, Feldpausch TR, et al. (2015) Long-term decline of the Amazon
1266 carbon sink. *Nature* 519: 344-+.

1267 Brinck K, Fischer R, Groeneveld J, et al. (2017) High resolution analysis of tropical forest
1268 fragmentation and its impact on the global carbon cycle. *Nature Communications* 8.

1269 Broecker WS. (2003) Does the trigger for abrupt climate change reside in the ocean or in the
1270 atmosphere? *Science* 300: 1519-1522.

1271 Bryden HL, King BA, McCarthy GD, et al. (2014) Impact of a 30% reduction in Atlantic
1272 meridional overturning during 2009-2010. *Ocean Science* 10: 683-691.

1273 Buckley MW and Marshall J. (2016) Observations, inferences, and mechanisms of the
1274 Atlantic Meridional Overturning Circulation: A review. *Reviews of Geophysics* 54: 5-
1275 63.

1276 Bugmann H and Bigler C. (2011) Will the CO₂ fertilization effect in forests be offset by
1277 reduced tree longevity? *Oecologia* 165: 533-544.

1278 Calosi P, Melatunan S, Turner LM, et al. (2017) Regional adaptation defines sensitivity to
1279 future ocean acidification. *Nature Communications* 8.

1280 Camarero JJ, Gazol A, Sanguesa-Barreda G, et al. (2015) To die or not to die: early warnings
1281 of tree dieback in response to a severe drought. *Journal of Ecology* 103: 44-57.

1282 Cavaleri MA, Reed SC, Smith WK, et al. (2015) Urgent need for warming experiments in
1283 tropical forests. *Global Change Biology* 21: 2111-2121.

1284 CBD. (2014) Convention on Biological Diversity: an Updated Synthesis of the Impacts of
1285 Ocean Acidification on Marine Biodiversity (S. Hennige, J.M. Roberts & P.
1286 Williamson; eds). CBD Technical Series No.75, Montreal, 99 pp. .

1287 Chan F, Barth JA, Blanchette CA, et al. (2017) Persistent spatial structuring of coastal ocean
1288 acidification in the California Current System. *Scientific Reports* 7.

1289 Chen XY and Tung KK. (2014) Varying planetary heat sink led to global-warming slowdown
1290 and acceleration. *Science* 345: 897-903.

1291 Church JA, Clark PU, Cazenave A, et al. (2013) Sea Level Change. In: Climate Change
1292 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth
1293 Assessment Report of the Intergovernmental Panel on Climate Change
1294 [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.
1295 Xia, V. Bex and P.M. Midgley (eds.)]. *Cambridge University Press, Cambridge,*
1296 *United Kingdom and New York, NY, USA.*

1297 Coe MT, Marthews TR, Costa MH, et al. (2013) Deforestation and climate feedbacks
1298 threaten the ecological integrity of south-southeastern Amazonia. *Philosophical*
1299 *Transactions of the Royal Society B-Biological Sciences* 368.

1300 Collins M, Knutti R, Arblaster J, et al. (2013) Long-term Climate Change: Projections,
1301 Commitments and Irreversibility. *Cambridge University Press, Cambridge, United*
1302 *Kingdom and New York, NY, USA.*

1303 Comeau S, Edmunds PJ, Spindel NB, et al. (2013) The responses of eight coral reef calcifiers
1304 to increasing partial pressure of CO₂ do not exhibit a tipping point. *Limnology and*
1305 *Oceanography* 58: 388-398.

1306 Cornford SL, Martin DF, Payne AJ, et al. (2015) Century-scale simulations of the response of
1307 the West Antarctic Ice Sheet to a warming climate. *Cryosphere* 9: 1579-1600.

1308 Couce E, Ridgwell A and Hendy EJ. (2013) Future habitat suitability for coral reef
1309 ecosystems under global warming and ocean acidification. *Global Change Biology*
1310 19: 3592-3606.

1311 Cox PM, Betts RA, Jones CD, et al. (2000) Acceleration of global warming due to carbon-
1312 cycle feedbacks in a coupled climate model (vol 408, pg 184, 2000). *Nature* 408: 750-
1313 750.

1314 Crowley TJ and North GR. (1988) Abrupt Climate Change and Extinction Events in Earth

History. *Science* 240: 996-1002.

Cunningham SA, Roberts CD, Frajka-Williams E, et al. (2013) Atlantic Meridional Overturning Circulation slowdown cooled the subtropical ocean. *Geophysical Research Letters* 40: 6202-6207.

da Costa ACL, Galbraith D, Almeida S, et al. (2010) Effect of 7 yr of experimental drought on vegetation dynamics and biomass storage of an eastern Amazonian rainforest. *New Phytologist* 187: 579-591.

da Costa ACL, Metcalfe DB, Doughty CE, et al. (2014) Ecosystem respiration and net primary productivity after 8-10 years of experimental through-fall reduction in an eastern Amazon forest. *Plant Ecology & Diversity* 7: 7-24.

da Silva BB, Wilcox BP, da Silva VDR, et al. (2015) Changes to the energy budget and evapotranspiration following conversion of tropical savannas to agricultural lands in SAo Paulo State, Brazil. *Ecohydrology* 8: 1272-1283.

De Angelis H and Skvarca P. (2003) Glacier surge after ice shelf collapse. *Science* 299: 1560-1562.

De Rydt J, Gudmundsson GH, Rott H, et al. (2015) Modeling the instantaneous response of glaciers after the collapse of the Larsen B Ice Shelf. *Geophysical Research Letters* 42: 5355-5363.

de Vries P and Weber SL. (2005) The Atlantic freshwater budget as a diagnostic for the existence of a stable shut down of the meridional overturning circulation. *Geophysical Research Letters* 32.

DeConto RM and Pollard D. (2016) Contribution of Antarctica to past and future sea-level rise. *Nature* 531: 591-597.

DeFries R, Herold M, Verchot L, et al. (2013) Export-oriented deforestation in Mato Grosso: harbinger or exception for other tropical forests? *Philosophical Transactions of the Royal Society B-Biological Sciences* 368.

den Toom M, Dijkstra HA, Weijer W, et al. (2014) Response of a Strongly Eddying Global Ocean to North Atlantic Freshwater Perturbations. *Journal of Physical Oceanography* 44: 464-481.

Depoorter MA, Bamber JL, Griggs JA, et al. (2013) Calving fluxes and basal melt rates of Antarctic ice shelves (vol 502, pg 89, 2013). *Nature* 502.

Dinniman MS, Klinck JM and Smith WO. (2007) Influence of sea ice cover and icebergs on circulation and water mass formation in a numerical circulation model of the Ross Sea, Antarctica. *Journal of Geophysical Research-Oceans* 112.

Dong SF, Goni G and Bringas F. (2015) Temporal variability of the South Atlantic Meridional Overturning Circulation between 20 degrees S and 35 degrees S. *Geophysical Research Letters* 42: 7655-7662.

Donoso PJ, Frene C, Flores M, et al. (2014) Balancing water supply and old-growth forest conservation in the lowlands of south-central Chile through adaptive co-management. *Landscape Ecology* 29: 245-260.

Doughty CE, Metcalfe DB, Girardin CAJ, et al. (2015) Drought impact on forest carbon dynamics and fluxes in Amazonia. *Nature* 519: 78-U140.

Drijfhout S, Bathiany S, Beaulieu C, et al. (2015) Catalogue of abrupt shifts in Intergovernmental Panel on Climate Change climate models. *Proceedings of the National Academy of Sciences of the United States of America* 112: E5777-E5786.

Dupont TK and Alley RB. (2005) Assessment of the importance of ice-shelf buttressing to ice-sheet flow. *Geophysical Research Letters* 32.

Durand G, Gagliardini O, Favier L, et al. (2011) Impact of bedrock description on modeling ice sheet dynamics. *Geophysical Research Letters* 38.

Ekstrom JA, Suatoni L, Cooley SR, et al. (2015) Vulnerability and adaptation of US

1365 shellfisheries to ocean acidification. *Nature Climate Change* 5: 207-214.
 1366 Elipot S, Frajka-Williams E, Hughes CW, et al. (2017) Observed Basin-Scale Response of
 1367 the North Atlantic Meridional Overturning Circulation to Wind Stress Forcing.
 1368 *Journal of Climate* 30: 2029-2054.
 1369 Elipot S, Frajka-Williams E, Hughes CW, et al. (2014) The Observed North Atlantic
 1370 Meridional Overturning Circulation: Its Meridional Coherence and Ocean Bottom
 1371 Pressure. *Journal of Physical Oceanography* 44: 517-537.
 1372 Ellis RP, Davison W, Queiros AM, et al. (2017) Does sex really matter? Explaining
 1373 intraspecies variation in ocean acidification responses. *Biology Letters* 13.
 1374 Enderlin EM, Howat IM, Jeong S, et al. (2014) An improved mass budget for the Greenland
 1375 ice sheet. *Geophysical Research Letters* 41: 866-872.
 1376 Espirito-Santo FDB, Gloor M, Keller M, et al. (2015) Size and frequency of natural forest
 1377 disturbances and the Amazon forest carbon balance (vol 5, 3434, 2014). *Nature*
 1378 *Communications* 6.
 1379 Esquivel-Muelbert A, Baker TR, Dexter KG, et al. (2017) Seasonal drought limits tree
 1380 species across the Neotropics. *Ecography* 40: 618-629.
 1381 Ezer T. (2015) Detecting changes in the transport of the Gulf Stream and the Atlantic
 1382 overturning circulation from coastal sea level data: The extreme decline in 2009-2010
 1383 and estimated variations for 1935-2012. *Global and Planetary Change* 129: 23-36.
 1384 Fabricius KE, Langdon C, Uthicke S, et al. (2011) Losers and winners in coral reefs
 1385 acclimatized to elevated carbon dioxide concentrations. *Nature Climate Change* 1:
 1386 165-169.
 1387 Fauset S, Baker TR, Lewis SL, et al. (2012) Drought-induced shifts in the floristic and
 1388 functional composition of tropical forests in Ghana. *Ecology Letters* 15: 1120-1129.
 1389 Favier L, Durand G, Cornford SL, et al. (2014) Retreat of Pine Island Glacier controlled by
 1390 marine ice-sheet instability. *Nature Climate Change* 4: 117-121.
 1391 Frajka-Williams E. (2015) Estimating the Atlantic overturning at 26 degrees N using satellite
 1392 altimetry and cable measurements. *Geophysical Research Letters* 42: 3458-3464.
 1393 Frajka-Williams E, Meinen CS, Johns WE, et al. (2016) Compensation between meridional
 1394 flow components of the Atlantic MOC at 26 degrees N. *Ocean Science* 12: 481-493.
 1395 Frieler K, Meinshausen M, Golly A, et al. (2013) Limiting global warming to 2 degrees C is
 1396 unlikely to save most coral reefs. *Nature Climate Change* 3: 165-170.
 1397 Furst JJ, Durand G, Gillet-Chaulet F, et al. (2016) The safety band of Antarctic ice shelves.
 1398 *Nature Climate Change* 6: 479-482.
 1399 Gagliardini O, Zwinger T, Gillet-Chaulet F, et al. (2013) Capabilities and performance of
 1400 Elmer/Ice, a new-generation ice sheet model. *Geoscientific Model Development* 6:
 1401 1299-1318.
 1402 Galbraith D, Levy PE, Sitch S, et al. (2010) Multiple mechanisms of Amazonian forest
 1403 biomass losses in three dynamic global vegetation models under climate change. *New*
 1404 *Phytologist* 187: 647-665.
 1405 Gambi MC, Musco L, Giangrande A, et al. (2016) Distribution and functional traits of
 1406 polychaetes in a CO2 vent system: winners and losers among closely related species.
 1407 *Marine Ecology Progress Series* 550: 121-134.
 1408 Garilli V, Rodolfo-Metalpa R, Scuderi D, et al. (2015) Physiological advantages of dwarfing
 1409 in surviving extinctions in high-CO2 oceans. *Nature Climate Change* 5: 678-+.
 1410 Garrard SL and Beaumont NJ. (2014) The effect of ocean acidification on carbon storage and
 1411 sequestration in seagrass beds; a global and UK context. *Marine Pollution Bulletin*
 1412 86: 138-146.
 1413 Gatti LV, Gloor M, Miller JB, et al. (2014) Drought sensitivity of Amazonian carbon balance
 1414 revealed by atmospheric measurements. *Nature* 506: 76-+.

1415 Gattuso JP, Magnan A, Bille R, et al. (2015) Contrasting futures for ocean and society from
 1416 different anthropogenic CO₂ emissions scenarios. *Science* 349.
 1417 Gille S. (2014) How ice shelves melt. *Science* 346: 1180-1181.
 1418 Gladstone RM, Lee V, Rougier J, et al. (2012) Calibrated prediction of Pine Island Glacier
 1419 retreat during the 21st and 22nd centuries with a coupled flowline model. *Earth and*
 1420 *Planetary Science Letters* 333: 191-199.
 1421 Goddard PB, Yin JJ, Griffies SM, et al. (2015) An extreme event of sea-level rise along the
 1422 Northeast coast of North America in 2009-2010. *Nature Communications* 6.
 1423 Goelzer H, Huybrechts P, Furst JJ, et al. (2013) Sensitivity of Greenland ice sheet projections
 1424 to model formulations. *Journal of Glaciology* 59: 733-749.
 1425 Golledge NR, Kowalewski DE, Naish TR, et al. (2015) The multi-millennial Antarctic
 1426 commitment to future sea-level rise. *Nature* 526: 421-+.
 1427 Good P, Harper A, Meesters A, et al. (2016) Are strong fire-vegetation feedbacks needed to
 1428 explain the spatial distribution of tropical tree cover? *Global Ecology and*
 1429 *Biogeography* 25: 16-25.
 1430 Graham V, Laurance SG, Grech A, et al. (2017) Spatially explicit estimates of forest carbon
 1431 emissions, mitigation costs and REDD plus opportunities in Indonesia. *Environmental*
 1432 *Research Letters* 12.
 1433 Gregory JM and Huybrechts P. (2006) Ice-sheet contributions to future sea-level change.
 1434 *Philosophical Transactions of the Royal Society a-Mathematical Physical and*
 1435 *Engineering Sciences* 364: 1709-1731.
 1436 Groenendijk P, van der Sleen P, Vlam M, et al. (2015) No evidence for consistent long-term
 1437 growth stimulation of 13 tropical tree species: results from tree-ring analysis. *Global*
 1438 *Change Biology* 21: 3762-3776.
 1439 Gudmundsson GH, Krug J, Durand G, et al. (2012) The stability of grounding lines on
 1440 retrograde slopes. *Cryosphere* 6: 1497-1505.
 1441 Guillevic M, Bazin L, Landaïs A, et al. (2013) Spatial gradients of temperature, accumulation
 1442 and delta O-18-ice in Greenland over a series of Dansgaard-Oeschger events. *Climate*
 1443 *of the Past* 9: 1029-1051.
 1444 Guinotte JM, Orr J, Cairns S, et al. (2006) Will human-induced changes in seawater
 1445 chemistry alter the distribution of deep-sea scleractinian corals? *Frontiers in Ecology*
 1446 *and the Environment* 4: 141-146.
 1447 Ha HK, Wahlin AK, Kim TW, et al. (2014) Circulation and Modification of Warm Deep
 1448 Water on the Central Amundsen Shelf. *Journal of Physical Oceanography* 44: 1493-
 1449 1501.
 1450 Haarsma RJ, Selten FM and Drijfhout SS. (2015) Decelerating Atlantic meridional
 1451 overturning circulation main cause of future west European summer atmospheric
 1452 circulation changes. *Environmental Research Letters* 10.
 1453 Hall-Spencer JM and Allen R. (2015) The impact of CO₂ emissions on 'nuisance' marine
 1454 species. *Research and Reports in Biodiversity Studies* 4: 33-46, doi:
 1455 10.2147/RRBS.S70357.
 1456 Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, et al. (2008) Volcanic carbon dioxide vents
 1457 show ecosystem effects of ocean acidification. *Nature* 454: 96-99.
 1458 Hansen J, Sato M, Kharecha P, et al. (2008) Target Atmospheric CO₂: Where Should
 1459 Humanity Aim? *The Open Atmospheric Science Journal* 2: 217-213, DOI:
 1460 210.2174/1874282300802010217.
 1461 Harig C and Simons FJ. (2015) Accelerated West Antarctic ice mass loss continues to
 1462 outpace East Antarctic gains. *Earth and Planetary Science Letters* 415: 134-141.
 1463 Harper A, Baker IT, Denning AS, et al. (2014) Impact of Evapotranspiration on Dry Season
 1464 Climate in the Amazon Forest. *Journal of Climate* 27: 574-591.

1465 Hauri C, Friedrich T and Timmermann A. (2016) Abrupt onset and prolongation of aragonite
 1466 undersaturation events in the Southern Ocean. *Nature Climate Change* 6: 172-+.
 1467 Hawkins E, Smith RS, Allison LC, et al. (2011) Bistability of the Atlantic overturning
 1468 circulation in a global climate model and links to ocean freshwater transport.
 1469 *Geophysical Research Letters* 38.
 1470 Hennige SJ, Wicks LC, Kamenos NA, et al. (2015) Hidden impacts of ocean acidification to
 1471 live and dead coral framework. *Proceedings of the Royal Society B-Biological*
 1472 *Sciences* 282.
 1473 Higgins SI and Scheiter S. (2012) Atmospheric CO₂ forces abrupt vegetation shifts locally,
 1474 but not globally. *Nature* 488: 209-+.
 1475 Hirota M, Holmgren M, Van Nes EH, et al. (2011) Global Resilience of Tropical Forest and
 1476 Savanna to Critical Transitions. *Science* 334: 232-235.
 1477 Hoffmann WA, Geiger EL, Gotsch SG, et al. (2012) Ecological thresholds at the savanna-
 1478 forest boundary: how plant traits, resources and fire govern the distribution of tropical
 1479 biomes. *Ecology Letters* 15: 759-768.
 1480 Honisch B, Ridgwell A, Schmidt DN, et al. (2012) The Geological Record of Ocean
 1481 Acidification. *Science* 335: 1058-1063.
 1482 Howard T, Ridley J, Pardaens AK, et al. (2014) The land-ice contribution to 21st-century
 1483 dynamic sea level rise. *Ocean Science* 10: 485-500.
 1484 Howes EL, Joos F, Eakin CM, et al. (2015) An updated synthesis of the observed and
 1485 projected impacts of climate change on the chemical, physical and biological
 1486 processes in the oceans. *Frontiers in Marine Science* 2: doi:
 1487 10.3389/fmars.2015.00036.
 1488 Hughes TP, Linares C, Dakos V, et al. (2013) Living dangerously on borrowed time during
 1489 slow, unrecognized regime shifts. *Trends in Ecology & Evolution* 28: 149-155.
 1490 Huntingford C, Zelazowski P, Galbraith D, et al. (2013) Simulated resilience of tropical
 1491 rainforests to CO₂-induced climate change. *Nature Geoscience* 6: 268-273.
 1492 ISRS. (2015) Climate Change Threatens the Survival of Coral Reefs. International Society
 1493 for Reef Studies Consensus Statement on Climate Change and Coral Bleaching.
 1494 [http://coralreefs.org/wp-content/uploads/2014/2003/ISRS-Consensus-Statement-on-](http://coralreefs.org/wp-content/uploads/2014/2003/ISRS-Consensus-Statement-on-Coral-Bleaching-Climate-Change-FINAL-2014Oct2015-HR.pdf)
 1495 [Coral-Bleaching-Climate-Change-FINAL-2014Oct2015-HR.pdf](http://coralreefs.org/wp-content/uploads/2014/2003/ISRS-Consensus-Statement-on-Coral-Bleaching-Climate-Change-FINAL-2014Oct2015-HR.pdf).
 1496 Jackson EL, Davies AJ, Howell KL, et al. (2014) Future-proofing marine protected area
 1497 networks for cold water coral reefs. *Ices Journal of Marine Science* 71: 2621-2629.
 1498 Jackson LC. (2013) Shutdown and recovery of the AMOC in a coupled global climate model:
 1499 The role of the advective feedback. *Geophysical Research Letters* 40: 1182-1188.
 1500 Jackson LC, Kahana R, Graham T, et al. (2015) Global and European climate impacts of a
 1501 slowdown of the AMOC in a high resolution GCM. *Climate Dynamics* 45: 3299-
 1502 3316.
 1503 Jackson LC, Peterson KA, Roberts CD, et al. (2016) Recent slowing of Atlantic overturning
 1504 circulation as a recovery from earlier strengthening. *Nature Geoscience* 9: 518-+.
 1505 Jacobs SS, Jenkins A, Giulivi CF, et al. (2011) Stronger ocean circulation and increased
 1506 melting under Pine Island Glacier ice shelf. *Nature Geoscience* 4: 519-523.
 1507 James R, Washington R and Rowell DP. (2014) African Climate Change Uncertainty in
 1508 Perturbed Physics Ensembles: Implications of Global Warming to 4 degrees C and
 1509 Beyond. *Journal of Climate* 27: 4677-4692.
 1510 Joetzjer E, Douville H, Delire C, et al. (2013) Present-day and future Amazonian
 1511 precipitation in global climate models: CMIP5 versus CMIP3. *Climate Dynamics* 41:
 1512 2921-2936.
 1513 Johnson JS, Bentley MJ, Smith JA, et al. (2014) Rapid Thinning of Pine Island Glacier in the
 1514 Early Holocene. *Science* 343: 999-1001.

1515 Jones C, Lowe J, Liddicoat S, et al. (2009) Committed terrestrial ecosystem changes due to
 1516 climate change. *Nature Geoscience* 2: 484-487.
 1517 Joughin I, Smith BE and Medley B. (2014) Marine Ice Sheet Collapse Potentially Under Way
 1518 for the Thwaites Glacier Basin, West Antarctica. *Science* 344: 735-738.
 1519 Kawaguchi S, Ishida A, King R, et al. (2013) Risk maps for Antarctic krill under projected
 1520 Southern Ocean acidification. *Nature Climate Change* 3.
 1521 Khan SA, Kjaer KH, Bevis M, et al. (2014) Sustained mass loss of the northeast Greenland
 1522 ice sheet triggered by regional warming. *Nature Climate Change* 4: 292-299.
 1523 Kiessling W and Simpson C. (2011) On the potential for ocean acidification to be a general
 1524 cause of ancient reef crises. *Global Change Biology* 17: 56-67.
 1525 Kingslake J, Ely JC, Das I, et al. (2017) Widespread movement of meltwater onto and across
 1526 Antarctic ice shelves. *Nature* 544: 349-+.
 1527 Konrad H, Sasgen I, Pollard D, et al. (2015) Potential of the solid-Earth response for limiting
 1528 long-term West Antarctic Ice Sheet retreat in a warming climate. *Earth and Planetary
 1529 Science Letters* 432: 254-264.
 1530 Kopp RE, Shwom RL, Wagner G, et al. (2016) Tipping elements and climate-economic
 1531 shocks: Pathways toward integrated assessment. *Earths Future* 4: 346-372.
 1532 Kroeker KJ, Kordas RL, Crim R, et al. (2013) Impacts of ocean acidification on marine
 1533 organisms: quantifying sensitivities and interaction with warming. *Global Change
 1534 Biology* 19: 1884-1896.
 1535 Kroeker KJ, Kordas RL and Harley CDG. (2017) Embracing interactions in ocean
 1536 acidification research: confronting multiple stressor scenarios and context
 1537 dependence. *Biology Letters* 13.
 1538 Landerer FW, Wiese DN, Bentel K, et al. (2015) North Atlantic meridional overturning
 1539 circulation variations from GRACE ocean bottom pressure anomalies. *Geophysical
 1540 Research Letters* 42: 8114-8121.
 1541 Lawrence D and Vandecar K. (2015) Effects of tropical deforestation on climate and
 1542 agriculture (vol 5, pg 27, 2015). *Nature Climate Change* 5: 174-174.
 1543 Lenton TM. (2011) Early warning of climate tipping points. *Nature Climate Change* 1: 201-
 1544 209.
 1545 Lenton TM. (2013) Environmental Tipping Points. *Annual Review of Environment and
 1546 Resources*, Vol 38 38: 1-29.
 1547 Lenton TM, Held H, Kriegler E, et al. (2008) Tipping elements in the Earth's climate system.
 1548 *Proceedings of the National Academy of Sciences of the United States of America*
 1549 105: 1786-1793.
 1550 Levermann A, Griesel A, Hofmann M, et al. (2005) Dynamic sea level changes following
 1551 changes in the thermohaline circulation. *Climate Dynamics* 24: 347-354.
 1552 Levermann A and Winkelmann R. (2016) A simple equation for the melt elevation feedback
 1553 of ice sheets. *Cryosphere* 10: 1799-1807.
 1554 Levy R, Harwood D, Florindo F, et al. (2016) Antarctic ice sheet sensitivity to atmospheric
 1555 CO₂ variations in the early to mid-Miocene. *Proceedings of the National Academy of
 1556 Sciences*.
 1557 Lewis SL, Brando P, Phillips OL, et al. (2011) The 2010 Amazon Drought. *Science* 331.
 1558 Lewis SL, Edwards DP and Galbraith D. (2015) Increasing human dominance of tropical
 1559 forests. *Science* 349: 827-832.
 1560 Lima LS, Coe MT, Soares BS, et al. (2014) Feedbacks between deforestation, climate, and
 1561 hydrology in the Southwestern Amazon: implications for the provision of ecosystem
 1562 services. *Landscape Ecology* 29: 261-274.
 1563 Liu W, Liu ZY and Brady EC. (2014) Why is the AMOC Monostable in Coupled General
 1564 Circulation Models? *Journal of Climate* 27: 2427-2443.

1565 Liu W, Xie SP, Liu Z, et al. (2017) Overlooked possibility of a collapsed Atlantic Meridional
 1566 Overturning Circulation in warming climate. *Science advances* 3: DOI:
 1567 10.1126/sciadv.1601666.
 1568 Lopez H, Goni G and Dong SF. (2017) A reconstructed South Atlantic Meridional
 1569 Overturning Circulation time series since 1870. *Geophysical Research Letters* 44:
 1570 3309-3318.
 1571 Losch M. (2008) Modeling ice shelf cavities in a z coordinate ocean general circulation
 1572 model. *Journal of Geophysical Research-Oceans* 113.
 1573 Lozier MS, Bacon S, Bower AS, et al. (2017) Overturning in the Subpolar North Atlantic
 1574 Program a New International Ocean Observing System. *Bulletin of the American*
 1575 *Meteorological Society* 98: 737-752.
 1576 Lynch-Stieglitz J. (2017) The Atlantic Meridional Overturning Circulation and Abrupt
 1577 Climate Change. *Annual Review of Marine Sciences*, Vol 9 9: 83-104.
 1578 Macedo MN, Coe MT, DeFries R, et al. (2013) Land-use-driven stream warming in
 1579 southeastern Amazonia. *Philosophical Transactions of the Royal Society B-Biological*
 1580 *Sciences* 368.
 1581 Maidens A, Arribas A, Scaife AA, et al. (2013) The Influence of Surface Forcings on
 1582 Prediction of the North Atlantic Oscillation Regime of Winter 2010/11. *Monthly*
 1583 *Weather Review* 141: 3801-3813.
 1584 Malhi Y, Adu-Bredu S, Asare RA, et al. (2013) African rainforests: past, present and future.
 1585 *Philosophical Transactions of the Royal Society B-Biological Sciences* 368.
 1586 Manabe S and Stouffer RJ. (1988) Two Stable Equilibria of a Coupled Ocean-Atmosphere
 1587 Model. *Journal of Climate* 1: 841-866.
 1588 Manzello DP, Enochs IC, Bruckner A, et al. (2014) Galapagos coral reef persistence after
 1589 ENSO warming across an acidification gradient. *Geophysical Research Letters* 41:
 1590 9001-9008.
 1591 Marengo JA, Tomasella J, Alves LM, et al. (2011) The drought of 2010 in the context of
 1592 historical droughts in the Amazon region. *Geophysical Research Letters* 38.
 1593 Mathesius S, Hofmann M, Caldeira K, et al. (2015) Long-term response of oceans to CO₂
 1594 removal from the atmosphere. *Nature Climate Change* 5: 1107-+.
 1595 Mathiot P, Jenkins A, Harris C, et al. (2017) Explicit representation and parametrised impacts
 1596 of under ice shelf seas in the z* coordinate ocean model NEMO 3.6. *Geoscientific*
 1597 *Model Development* 10.7: 2849-2874.
 1598 Mathis JT, Cooley SR, Yates KK, et al. (2015) Introduction to this Special Issue on Ocean
 1599 Acidification: THE PATHWAY FROM SCIENCE TO POLICY. *Oceanography* 28:
 1600 10-15.
 1601 McCarthy G, Frajka-Williams E, Johns WE, et al. (2012) Observed interannual variability of
 1602 the Atlantic meridional overturning circulation at 26.5 degrees N. *Geophysical*
 1603 *Research Letters* 39.
 1604 McCarthy GD, Haigh ID, Hirschi JJM, et al. (2015a) Ocean impact on decadal Atlantic
 1605 climate variability revealed by sea-level observations. *Nature* 521: 508-U172.
 1606 McCarthy GD, Smeed DA, Johns WE, et al. (2015b) Measuring the Atlantic Meridional
 1607 Overturning Circulation at 26 degrees N. *Progress in Oceanography* 130: 91-111.
 1608 McCulloch M, Falter J, Trotter J, et al. (2012) Coral resilience to ocean acidification and
 1609 global warming through pH up-regulation. *Nature Climate Change* 2: 623-633.
 1610 McDowell NG and Allen CD. (2015) Darcy's law predicts widespread forest mortality under
 1611 climate warming. *Nature Climate Change* 5: 669-672.
 1612 McMillan M, Leeson A, Shepherd A, et al. (2016) A high-resolution record of Greenland
 1613 mass balance. *Geophysical Research Letters* 43: 7002-7010.
 1614 McNeall D, Halloran PR, Good P, et al. (2011) Analyzing abrupt and nonlinear climate

1615 changes and their impacts. *Wiley Interdisciplinary Reviews-Climate Change* 2: 663-
 1616 686.
 1617 Mecking JV, Drijfhout SS, Jackson LC, et al. (2016) Stable AMOC off state in an eddy-
 1618 permitting coupled climate model. *Climate Dynamics* 47: 2455-2470.
 1619 Medley B, Joughin I, Smith BE, et al. (2014) Constraining the recent mass balance of Pine
 1620 Island and Thwaites glaciers, West Antarctica, with airborne observations of snow
 1621 accumulation. *Cryosphere* 8: 1375-1392.
 1622 Meinen CS, Speich S, Perez RC, et al. (2013) Temporal variability of the meridional
 1623 overturning circulation at 34.5 degrees S: Results from two pilot boundary arrays in
 1624 the South Atlantic. *Journal of Geophysical Research-Oceans* 118: 6461-6478.
 1625 Meir P, Mencuccini M and Dewar RC. (2015) Drought-related tree mortality: addressing the
 1626 gaps in understanding and prediction. *New Phytologist* 207: 28-33.
 1627 Mengel M and Levermann A. (2014) Ice plug prevents irreversible discharge from East
 1628 Antarctica. *Nature Climate Change* 4: 451-455.
 1629 Mercer JH. (1968) Antarctic ice and Sangamon sea level. *IAHS Publ.* 179: 217-225.
 1630 Mercier H, Lherminier P, Sarafanov A, et al. (2015) Variability of the meridional overturning
 1631 circulation at the Greenland-Portugal OVIDE section from 1993 to 2010. *Progress in*
 1632 *Oceanography* 132: 250-261.
 1633 Miles BWJ, Stokes CR, Vieli A, et al. (2013) Rapid, climate-driven changes in outlet glaciers
 1634 on the Pacific coast of East Antarctica. *Nature* 500: 563-+.
 1635 Mollmann C, Folke C, Edwards M, et al. (2015) Marine regime shifts around the globe:
 1636 theory, drivers and impacts. *Philosophical Transactions of the Royal Society B-*
 1637 *Biological Sciences* 370.
 1638 Moncrieff GR, Scheiter S, Bond WJ, et al. (2014) Increasing atmospheric CO2 overrides the
 1639 historical legacy of multiple stable biome states in Africa. *New Phytologist* 201: 908-
 1640 915.
 1641 Mongin M, Baird ME, Tilbrook B, et al. (2016) The exposure of the Great Barrier Reef to
 1642 ocean acidification. *Nature Communications* 7.
 1643 Moon T, Joughin I, Smith B, et al. (2012) 21st-Century Evolution of Greenland Outlet
 1644 Glacier Velocities. *Science* 336: 576-578.
 1645 Morton DC, Le Page Y, DeFries R, et al. (2013) Understorey fire frequency and the fate of
 1646 burned forests in southern Amazonia. *Philosophical Transactions of the Royal Society*
 1647 *B-Biological Sciences* 368.
 1648 Morton DC, Nagol J, Carabajal CC, et al. (2014) Amazon forests maintain consistent canopy
 1649 structure and greenness during the dry season. *Nature* 506: 221-+.
 1650 Mumby PJ, Iglesias-Prieto R, Hooten AJ, et al. (2011) Revisiting climate thresholds and
 1651 ecosystem collapse. *Frontiers in Ecology and the Environment* 9: 94-96.
 1652 Murray T, Scharrer K, James TD, et al. (2010) Ocean regulation hypothesis for glacier
 1653 dynamics in southeast Greenland and implications for ice sheet mass changes. *Journal*
 1654 *of Geophysical Research-Earth Surface* 115.
 1655 Nagelkerken I and Connell SD. (2015) Global alteration of ocean ecosystem functioning due
 1656 to increasing human CO2 emissions. *Proceedings of the National Academy of*
 1657 *Sciences of the United States of America* 112: 13272-13277.
 1658 Nagelkerken I, Goldenberg SU and Ferreira CM. (2017) Species Interactions Drive Fish
 1659 Biodiversity Loss in a High-CO₂ World. *Current Biology* 27: 2177-2184.e2174,
 1660 <https://doi.org/2110.1016/j.cub.2017.2106.2023>.
 1661 Nagelkerken I and Munday PL. (2016) Animal behaviour shapes the ecological effects of
 1662 ocean acidification and warming: moving from individual to community-level
 1663 responses. *Global Change Biology* 22: 974-989.
 1664 Nepstad D, McGrath D, Stickler C, et al. (2014) Slowing Amazon deforestation through

1665 public policy and interventions in beef and soy supply chains. *Science* 344: 1118-
 1666 1123.
 1667 Nepstad DC, Tohver IM, Ray D, et al. (2007) Mortality of large trees and lianas following
 1668 experimental drought in an amazon forest. *Ecology* 88: 2259-2269.
 1669 Nick FM, Vieli A, Howat IM, et al. (2009) Large-scale changes in Greenland outlet glacier
 1670 dynamics triggered at the terminus. *Nature Geoscience* 2: 110-114.
 1671 NRC. (2013) (National Research Council) Abrupt Impacts of Climate Change: Anticipating
 1672 Surprises. *Washington, DC: The National Academies Press:*
 1673 <https://doi.org/10.17226/18373>.
 1674 O'Neill BC, Oppenheimer M, Warren R, et al. (2017) IPCC reasons for concern regarding
 1675 climate change risks. *Nature Climate Change* 7: 28-37.
 1676 Oliveira LJC, Costa MH, Soares BS, et al. (2013) Large-scale expansion of agriculture in
 1677 Amazonia may be a no-win scenario. *Environmental Research Letters* 8.
 1678 Oliveira PTS, Nearing MA, Moran MS, et al. (2014) Trends in water balance components
 1679 across the Brazilian Cerrado. *Water Resources Research* 50: 7100-7114.
 1680 Pan YD, Birdsey RA, Fang JY, et al. (2011) A Large and Persistent Carbon Sink in the
 1681 World's Forests. *Science* 333: 988-993.
 1682 Panday PK, Coe MT, Macedo MN, et al. (2015) Deforestation offsets water balance changes
 1683 due to climate variability in the Xingu River in eastern Amazonia. *Journal of*
 1684 *Hydrology* 523: 822-829.
 1685 Pandolfi JM. (2015) Incorporating Uncertainty in Predicting the Future Response of Coral
 1686 Reefs to Climate Change. *Annual Review of Ecology, Evolution, and Systematics*, Vol
 1687 46 46: 281-303.
 1688 Paolo FS, Fricker HA and Padman L. (2015) Volume loss from Antarctic ice shelves is
 1689 accelerating. *Science* 348: 327-331.
 1690 Parsons LA, Yin JJ, Overpeck JT, et al. (2014) Influence of the Atlantic Meridional
 1691 Overturning Circulation on the monsoon rainfall and carbon balance of the American
 1692 tropics. *Geophysical Research Letters* 41: 146-151.
 1693 Pausas JG and Dantas VD. (2017) Scale matters: fire-vegetation feedbacks are needed to
 1694 explain tropical tree cover at the local scale. *Global Ecology and Biogeography* 26:
 1695 395-399.
 1696 Pedro JB, van Ommen TD, Rasmussen SO, et al. (2011) The last deglaciation: timing the
 1697 bipolar seesaw. *Climate of the Past* 7: 671-683.
 1698 Peng P, Zhu YZ, Zhong M, et al. (2016) Ice Mass Variation in Antarctica from GRACE Over
 1699 2002-2011. *Marine Geodesy* 39: 178-194.
 1700 Perez FF, Mercier H, Vazquez-Rodriguez M, et al. (2013) Atlantic Ocean CO2 uptake
 1701 reduced by weakening of the meridional overturning circulation. *Nature Geoscience*
 1702 6: 146-152.
 1703 Phillips OL, Aragao LEOC, Lewis SL, et al. (2009) Drought Sensitivity of the Amazon
 1704 Rainforest. *Science* 323: 1344-1347.
 1705 Phillips OL, van der Heijden G, Lewis SL, et al. (2010) Drought-mortality relationships for
 1706 tropical forests. *New Phytologist* 187: 631-646.
 1707 Plaganyi EE, Ellis N, Blamey LK, et al. (2014) Ecosystem modelling provides clues to
 1708 understanding ecological tipping points. *Marine Ecology Progress Series* 512: 99-
 1709 113.
 1710 Pollard D, DeConto RM and Alley RB. (2015) Potential Antarctic Ice Sheet retreat driven by
 1711 hydrofracturing and ice cliff failure. *Earth and Planetary Science Letters* 412: 112-
 1712 121.
 1713 Popova EE, Yool A, Aksenov Y, et al. (2014) Regional variability of acidification in the
 1714 Arctic: a sea of contrasts. *Biogeosciences* 11: 293-308.

1715 Portner HO, Karl DM, Boyd PW, et al. (2014) Ocean systems. In: Climate Change 2014:
 1716 Impacts, Adaptation, and Vulnerability. Part A: Global and
 1717 Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the
 1718 Intergovernmental
 1719 Panel on Climate Change [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D.
 1720 Mastrandrea, T.E. Bilir,
 1721 M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S.
 1722 MacCracken,
 1723 P.R. Mastrandrea, and L.L. White (eds.)]. *Cambridge University Press, Cambridge, United*
 1724 *Kingdom and New York, NY, USA.*
 1725 Powell TL, Galbraith DR, Christoffersen BO, et al. (2013) Confronting model predictions of
 1726 carbon fluxes with measurements of Amazon forests subjected to experimental
 1727 drought. *New Phytologist* 200: 350-364.
 1728 Pritchard HD, Arthern RJ, Vaughan DG, et al. (2009) Extensive dynamic thinning on the
 1729 margins of the Greenland and Antarctic ice sheets. *Nature* 461: 971-975.
 1730 Qi D, Chen LQ, Chen BS, et al. (2017) Increase in acidifying water in the western Arctic
 1731 Ocean. *Nature Climate Change* 7: 195-+.
 1732 Rahmstorf S. (2002) Ocean circulation and climate during the past 120,000 years. *Nature*
 1733 419: 207-214.
 1734 Rahmstorf S, Box JE, Feulner G, et al. (2015) Exceptional twentieth-century slowdown in
 1735 Atlantic Ocean overturning circulation. *Nature Climate Change* 5: 475-480.
 1736 Rahmstorf S, Crucifix M, Ganopolski A, et al. (2005) Thermohaline circulation hysteresis: A
 1737 model intercomparison. *Geophysical Research Letters* 32.
 1738 Ramajo L, Perez-Leon E, Hendriks IE, et al. (2016) Food supply confers calcifiers resistance
 1739 to ocean acidification. *Scientific Reports* 6.
 1740 Rammig A, Jupp T, Thonicke K, et al. (2010) Estimating the risk of Amazonian forest
 1741 dieback. *New Phytologist* 187: 694-706.
 1742 Raulf FF, Fabricius K, Uthicke S, et al. (2015) Changes in microbial communities in coastal
 1743 sediments along natural CO₂ gradients at a volcanic vent in Papua New Guinea.
 1744 *Environmental Microbiology* 17: 3678-3691.
 1745 Rayner D, Hirschi JJM, Kanzow T, et al. (2011) Monitoring the Atlantic meridional
 1746 overturning circulation. *Deep-Sea Research Part II-Topical Studies in Oceanography*
 1747 58: 1744-1753.
 1748 Reichstein M, Bahn M, Ciais P, et al. (2013) Climate extremes and the carbon cycle. *Nature*
 1749 500: 287-295.
 1750 Reintges A, Martin T, Latif M, et al. (2016) Uncertainty in twenty-first century projections of
 1751 the Atlantic Meridional Overturning Circulation in CMIP3 and CMIP5 models.
 1752 *Climate Dynamics* doi:10.1007/s00382-016-3180-x.
 1753 Rhein M, Rintoul SR, Aoki S, et al. (2013) Observations: Ocean. In: Climate Change 2013:
 1754 The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment
 1755 Report of the Intergovernmental Panel on Climate Change
 1756 [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.
 1757 Xia, V. Bex and P.M. Midgley (eds.)]. *Cambridge University Press, Cambridge,*
 1758 *United Kingdom and New York, NY, USA.*
 1759 Riebesell U and Gattuso JP. (2015) COMMENTARY: Lessons learned from ocean
 1760 acidification research. *Nature Climate Change* 5: 12-14.
 1761 Rignot E, Box JE, Burgess E, et al. (2008) Mass balance of the Greenland ice sheet from
 1762 1958 to 2007. *Geophysical Research Letters* 35.
 1763 Rignot E, Jacobs S, Mouginot J, et al. (2013) Ice-Shelf Melting Around Antarctica. *Science*
 1764 341: 266-270.

1765 Rignot E, Koppes M and Velicogna I. (2010) Rapid submarine melting of the calving faces of
1766 West Greenland glaciers. *Nature Geoscience* 3: 187-191.

1767 Rignot E, Mouginot J, Morlighem M, et al. (2014) Widespread, rapid grounding line retreat
1768 of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to
1769 2011. *Geophysical Research Letters* 41: 3502-3509.

1770 Rignot E, Velicogna I, van den Broeke MR, et al. (2011) Acceleration of the contribution of
1771 the Greenland and Antarctic ice sheets to sea level rise. *Geophysical Research Letters*
1772 38.

1773 Roberts CD, Garry FK and Jackson LC. (2013a) A Multimodel Study of Sea Surface
1774 Temperature and Subsurface Density Fingerprints of the Atlantic Meridional
1775 Overturning Circulation. *Journal of Climate* 26: 9155-9174.

1776 Roberts CD, Jackson L and McNeall D. (2014) Is the 2004-2012 reduction of the Atlantic
1777 meridional overturning circulation significant? *Geophysical Research Letters* 41:
1778 3204-3210.

1779 Roberts CD, Waters J, Peterson KA, et al. (2013b) Atmosphere drives recent interannual
1780 variability of the Atlantic meridional overturning circulation at 26.5 degrees N.
1781 *Geophysical Research Letters* 40: 5164-5170.

1782 Robinson A, Calov R and Ganopolski A. (2012) Multistability and critical thresholds of the
1783 Greenland ice sheet. *Nature Climate Change* 2: 429-432.

1784 Robson J, Hodson D, Hawkins E, et al. (2014) Atlantic overturning in decline? *Nature*
1785 *Geoscience* 7: 2-3.

1786 Rodriguez DA, Tomasella J and Linhares C. (2010) Is the forest conversion to pasture
1787 affecting the hydrological response of Amazonian catchments? Signals in the Ji-
1788 Parana Basin. *Hydrological Processes* 24: 1254-1269.

1789 Rowland L, da Costa ACL, Galbraith DR, et al. (2015) Death from drought in tropical forests
1790 is triggered by hydraulics not carbon starvation. *Nature* 528: 119-+.

1791 Saatchi S, Asefi-Najafabady S, Malhi Y, et al. (2013) Persistent effects of a severe drought
1792 on Amazonian forest canopy. *Proceedings of the National Academy of Sciences of the*
1793 *United States of America* 110: 565-570.

1794 Sasgen I, van den Broeke M, Bamber JL, et al. (2012) Timing and origin of recent regional
1795 ice-mass loss in Greenland. *Earth and Planetary Science Letters* 333: 293-303.

1796 Sasse TP, McNeil BI, Matear RJ, et al. (2015) Quantifying the influence of CO2 seasonality
1797 on future aragonite undersaturation onset. *Biogeosciences* 12: 6017-6031.

1798 Scaife AA, Arribas A, Blockley E, et al. (2014) Skillful long-range prediction of European
1799 and North American winters. *Geophysical Research Letters* 41: 2514-2519.

1800 Scambos TA, Bohlander JA, Shuman CA, et al. (2004) Glacier acceleration and thinning
1801 after ice shelf collapse in the Larsen B embayment, Antarctica. *Geophysical Research*
1802 *Letters* 31.

1803 Schmidtko S, Heywood KJ, Thompson AF, et al. (2014) Multidecadal warming of Antarctic
1804 waters. *Science* 346: 1227-1231.

1805 Schoof C. (2007) Ice sheet grounding line dynamics: Steady states, stability, and hysteresis.
1806 *Journal of Geophysical Research-Earth Surface* 112.

1807 Send U, Lankhorst M and Kanzow T. (2011) Observation of decadal change in the Atlantic
1808 meridional overturning circulation using 10 years of continuous transport data.
1809 *Geophysical Research Letters* 38.

1810 Settele J, Scholes RJ, Betts R, et al. (2014) Terrestrial and inland water systems. In: Climate
1811 Change 2014: Impacts, Adaptation, and Vulnerability.

1812 Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth
1813 Assessment Report of the
1814 Intergovernmental Panel on Climate Change [Field, C.B., V.R. Barros, D.J. Dokken, K.J.

1815 Mach,
 1816 M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma,
 1817 E.S. Kissel, A.N. Levy,
 1818 S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. *Cambridge University Press*,
 1819 *Cambridge, United*
 1820 *Kingdom and New York, NY, USA*: 271-359.
 1821 Shamberger KEF, Cohen AL, Golbuu Y, et al. (2014) Diverse coral communities in naturally
 1822 acidified waters of a Western Pacific reef. *Geophysical Research Letters* 41: 499-504.
 1823 Shepherd A. (2012) A reconciled estimate of ice-sheet mass balance (vol 57, pg 88, 2010).
 1824 *Science* 338: 1539-1539.
 1825 Shiogama H, Emori S, Hanasaki N, et al. (2011) Observational constraints indicate risk of
 1826 drying in the Amazon basin. *Nature Communications* 2.
 1827 Shiogama H, Watanabe M, Imada Y, et al. (2013) An event attribution of the 2010 drought in
 1828 the South Amazon region using the MIROC5 model. *Atmospheric Science Letters* 14:
 1829 170-175.
 1830 Silvano A, Rintoul SR and Herraiz-Borreguero L. (2016) Ocean-Ice Shelf Interaction in East
 1831 Antarctica. *Oceanography* 29: 130-143.
 1832 Silverio DV, Brando PM, Balch JK, et al. (2013) Testing the Amazon savannization
 1833 hypothesis: fire effects on invasion of a neotropical forest by native cerrado and
 1834 exotic pasture grasses. *Philosophical Transactions of the Royal Society B-Biological*
 1835 *Sciences* 368.
 1836 Silverman J, Schneider K, Kline DI, et al. (2014) Community calcification in Lizard Island,
 1837 Great Barrier Reef: A 33 year perspective. *Geochimica Et Cosmochimica Acta* 144:
 1838 72-81.
 1839 Slot M, Rey-Sanchez C, Gerber S, et al. (2014) Thermal acclimation of leaf respiration of
 1840 tropical trees and lianas: response to experimental canopy warming, and
 1841 consequences for tropical forest carbon balance. *Global Change Biology* 20: 2915-
 1842 2926.
 1843 Smeed D, McCarthy G, Rayner D, et al. (2016) Atlantic meridional overturning circulation
 1844 observed by the RAPID-MOCHA-WBTS (RAPID-Meridional Overturning
 1845 Circulation and Heatflux Array-Western Boundary Time Series) array at 26N from
 1846 2004 to 2015. British Oceanographic Data Centre - Natural Environment Research
 1847 Council, UK. doi:10.5285/35784047-9b82-2160-e053-6c86abc0c91b.
 1848 Smeed DA, McCarthy GD, Cunningham SA, et al. (2014) Observed decline of the Atlantic
 1849 meridional overturning circulation 2004-2012. *Ocean Science* 10: 29-38.
 1850 Srokosz M, Baringer M, Bryden H, et al. (2012) Past, Present, and Future Changes in the
 1851 Atlantic Meridional Overturning Circulation. *Bulletin of the American Meteorological*
 1852 *Society* 93: 1663-1676.
 1853 Srokosz MA and Bryden HL. (2015) Observing the Atlantic Meridional Overturning
 1854 Circulation yields a decade of inevitable surprises. *Science* 348.
 1855 Staal A, Dekker SC, Xu C, et al. (2016) Bistability, Spatial Interaction, and the Distribution
 1856 of Tropical Forests and Savannas. *Ecosystems* 19: 1080-1091.
 1857 Staver AC, Archibald S and Levin SA. (2011) The Global Extent and Determinants of
 1858 Savanna and Forest as Alternative Biome States. *Science* 334: 230-232.
 1859 Steinacher M, Joos F and Stocker TF. (2013) Allowable carbon emissions lowered by
 1860 multiple climate targets. *Nature* 499: 197-+.
 1861 Stickler CM, Coe MT, Costa MH, et al. (2013) Dependence of hydropower energy generation
 1862 on forests in the Amazon Basin at local and regional scales. *Proceedings of the*
 1863 *National Academy of Sciences of the United States of America* 110: 9601-9606.
 1864 Stommel H. (1961) Thermohaline convection with two stable regimes of flow. *Tellus* 13:

1865 224-230.

1866 Sunday JM, Calosi P, Dupont S, et al. (2014) Evolution in an acidifying ocean. *Trends in*

1867 *Ecology & Evolution* 29: 117-125.

1868 Sutter J, Gierz P, Grosfeld K, et al. (2016) Ocean temperature thresholds for Last Interglacial

1869 West Antarctic Ice Sheet collapse. *Geophysical Research Letters* 43: 2675-2682.

1870 Sutton RT and Dong BW. (2012) Atlantic Ocean influence on a shift in European climate in

1871 the 1990s. *Nature Geoscience* 5: 788-792.

1872 Swingedouw D, Fichefet T, Huybrechts P, et al. (2008) Antarctic ice-sheet melting provides

1873 negative feedbacks on future climate warming. *Geophysical Research Letters* 35.

1874 Swingedouw D, Rodehacke CB, Behrens E, et al. (2013) Decadal fingerprints of freshwater

1875 discharge around Greenland in a multi-model ensemble. *Climate Dynamics* 41: 695-

1876 720.

1877 Swingedouw D, Rodehacke CB, Olsen SM, et al. (2015) On the reduced sensitivity of the

1878 Atlantic overturning to Greenland ice sheet melting in projections: a multi-model

1879 assessment. *Climate Dynamics* 44: 3261-3279.

1880 Taws SL, Marsh R, Wells NC, et al. (2011) Re-emerging ocean temperature anomalies in

1881 late-2010 associated with a repeat negative NAO. *Geophysical Research Letters* 38.

1882 Tomasella J, Pinho PF, Borma LS, et al. (2013) The droughts of 1997 and 2005 in Amazonia:

1883 floodplain hydrology and its potential ecological and human impacts. *Climatic*

1884 *Change* 116: 723-746.

1885 Turley C. (2017) Chapter 5.6: The risk of ocean acidification to ocean ecosystems. In

1886 UNESCO IOC and UNEP (2016). *The Open Ocean: Status and Trends. United*

1887 *Nations Environment Programme, Nairobi*: 193-205.

1888 van den Broeke M, Bamber J, Ettema J, et al. (2009) Partitioning Recent Greenland Mass

1889 Loss. *Science* 326: 984-986.

1890 van der Sleen P, Groenendijk P, Vlam M, et al. (2015) No growth stimulation of tropical trees

1891 by 150 years of CO₂ fertilization but water-use efficiency increased. *Nature*

1892 *Geoscience* 8: 24-28.

1893 Vanderwel MC, Slot M, Lichstein JW, et al. (2015) Global convergence in leaf respiration

1894 from estimates of thermal acclimation across time and space. *New Phytologist* 207:

1895 1026-1037.

1896 Vaughan DG, Comiso JC, Allison I, et al. (2013) Observations: Cryosphere. In: Climate

1897 Change 2013: The Physical Science Basis. Contribution of Working Group I to the

1898 Fifth Assessment Report of the Intergovernmental Panel on Climate Change

1899 [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.

1900 Xia, V. Bex and P.M. Midgley (eds.)]. *Cambridge University Press, Cambridge,*

1901 *United Kingdom and New York, NY, USA*.

1902 Vazquez-Rodriguez M, Perez FF, Velo A, et al. (2012) Observed acidification trends in

1903 North Atlantic water masses. *Biogeosciences* 9: 5217-5230.

1904 Veenendaal EM, Torello-Raventos M, Feldpausch TR, et al. (2015) Structural, physiognomic

1905 and above-ground biomass variation in savanna-forest transition zones on three

1906 continents - how different are co-occurring savanna and forest formations?

1907 *Biogeosciences* 12: 2927-2951.

1908 Vellinga M and Wood RA. (2008) Impacts of thermohaline circulation shutdown in the

1909 twenty-first century. *Climatic Change* 91: 43-63.

1910 Verbesselt J, Umlauf N, Hirota M, et al. (2016) Remotely sensed resilience of tropical

1911 forests. *Nature Climate Change* 6: 1028-+.

1912 Vizcaino M, Mikolajewicz U, Ziemen F, et al. (2015) Coupled simulations of Greenland Ice

1913 Sheet and climate change up to AD 2300. *Geophysical Research Letters* 42: 3927-

1914 3935.

1915 von Appen WJ, Koszalka IM, Pickart RS, et al. (2014) The East Greenland Spill Jet as an
1916 important component of the Atlantic Meridional Overturning Circulation. *Deep-Sea*
1917 *Research Part I-Oceanographic Research Papers* 92: 75-84.

1918 Weber ME, Clark PU, Kuhn G, et al. (2014) Millennial-scale variability in Antarctic ice-
1919 sheet discharge during the last deglaciation. *Nature* 510: 134-+.

1920 Weijer W, Maltrud ME, Hecht MW, et al. (2012) Response of the Atlantic Ocean circulation
1921 to Greenland Ice Sheet melting in a strongly-eddy ocean model. *Geophysical*
1922 *Research Letters* 39.

1923 Wittmann AC and Portner HO. (2013) Sensitivities of extant animal taxa to ocean
1924 acidification. *Nature Climate Change* 3: 995-1001.

1925 WMO. (2014) The State of Greenhouse Gases in the Atmosphere Based on Global
1926 Observations through 2013. WMO Greenhouse Gas Bulletin 10, 8 pp.

1927 Woollings T, Gregory JM, Pinto JG, et al. (2012) Response of the North Atlantic storm track
1928 to climate change shaped by ocean-atmosphere coupling. *Nature Geoscience* 5: 313-
1929 317.

1930 Woosley RJ, Millero FJ and Wanninkhof R. (2016) Rapid anthropogenic changes in CO₂ and
1931 pH in the Atlantic Ocean: 2003-2014. *Global Biogeochemical Cycles* 30; doi:
1932 10.1002/2015GB005248.

1933 Wouters B, Bamber JL, van den Broeke MR, et al. (2013) Limits in detecting acceleration of
1934 ice sheet mass loss due to climate variability. *Nature Geoscience* 6: 613-616.

1935 Wouters B, Martin-Espanol A, Helm V, et al. (2015) Dynamic thinning of glaciers on the
1936 Southern Antarctic Peninsula. *Science* 348: 899-903.

1937 Wu M, Schurgers G, Ahlstrom A, et al. (2017) Impacts of land use on climate and ecosystem
1938 productivity over the Amazon and the South American continent. *Environmental*
1939 *Research Letters* 12.

1940 Wuyts B, Champneys AR and House JI. (2017) Amazonian forest-savanna bistability and
1941 human impact. *Nature Communications* 8.

1942 Xu C, Hantson S, Holmgren M, et al. (2016) Remotely sensed canopy height reveals three
1943 pantropical ecosystem states. *Ecology* 97: 2518-2521.

1944 Yang Q, Dixon TH, Myers PG, et al. (2016) Recent increases in Arctic freshwater flux
1945 affects Labrador Sea convection and Atlantic overturning circulation. *Nature*
1946 *Communications* 7.

1947 Yin JJ, Schlesinger ME and Stouffer RJ. (2009) Model projections of rapid sea-level rise on
1948 the northeast coast of the United States. *Nature Geoscience* 2: 262-266.

1949 Zeebe RE and Ridgwell A. (2011) Past changes in ocean carbonate chemistry. Chapter 2 in:
1950 Ocean Acidification. (Eds: J.P Gattuso & L. Hansson). Oxford University Press,
1951 Oxford. p 21-40.

1952 Zeebe RE, Ridgwell A and Zachos JC. (2016) Anthropogenic carbon release rate
1953 unprecedented during the past 66 million years. *Nature Geoscience* 9: 325-329.

1954 Zhang K, Castanho ADD, Galbraith DR, et al. (2015) The fate of Amazonian ecosystems
1955 over the coming century arising from changes in climate, atmospheric CO₂, and land
1956 use. *Global Change Biology* 21: 2569-2587.

1957

1958

1959

1960

1961 **Figure captions**

1962

1963 Figure 1. Evidence (from Favier et al. 2014) that the Pine Island Glacier's grounding line is
1964 probably engaged in an unstable 40 km retreat, due to the retrograde bedrock slope. Left:
1965 map of bedrock elevation, with the grounding lines for 2009 (white) and 2011 (purple)
1966 shown. Right: bedrock height (solid black line) and geometry of the glacier centreline
1967 produced by the Elmer/Ice ice-flow model at time (t) = 0 (dotted line) and after 50 years of a
1968 melting scenario (red line).

1969

1970 Figure 2. AMOC transport measured at 26.5°N (Smeed et al., 2016). The gray line represents
1971 the 10 day filtered measurements, while the red line was produced using a 180 day running
1972 mean. Clearly visible are the low AMOC event in 2009-10 and the overall decrease in
1973 strength over the measurement period.

1974

1975 Figure 3. Trends in net above-ground biomass change, productivity and mortality rates, for
1976 321 plots, weighted by plot size (after Brien et al. 2015).

1977

1978 Figure 4. Area-weighted ensemble mean duration (months per year) of aragonite
1979 undersaturation at the surface (solid lines) and at 100m depth (dashed lines) for three sectors
1980 of the Southern Ocean (Bellingshausen Sea, Weddell Sea and East Antarctica), the central
1981 Chilean coast and the Patagonian shelf over the period 1900-2100, with future projections
1982 based on RCP 8.5. From Hauri et al. (2016).

Figures





Figure 1. Evidence (from Favier et al. 2014) that the Pine Island Glacier's grounding line is probably engaged in an unstable 40 km retreat, due to the retrograde bedrock slope. Left: map of bedrock elevation, with the grounding lines for 2009 (white) and 2011 (purple) shown. Right: bedrock height (solid black line) and geometry of the glacier centreline produced by the Elmer/Ice ice-flow model at time (t) = 0 (dotted line) and after 50 years of a melting scenario (red line).

1996
1997




1998
1999
2000
2001
2002
2003
2004

Figure 2. AMOC transport measured at 26.5°N (Smeed et al., 2016). The gray line represents the 10 day filtered measurements, while the red line was produced using a 180 day running mean. Clearly visible are the low AMOC event in 2009-10 and the overall decrease in strength over the measurement period.

2005

2006



2007


2008 Figure 3. Trends in net above-ground biomass change, productivity and mortality rates, for

2009 321 plots, weighted by plot size (after Brien et al. 2015).

2010

2011

2012



2013 Figure 4. Area-weighted ensemble mean duration (months per year) of aragonite
2014 undersaturation at the surface (solid lines) and at 100m depth (dashed lines) for three sectors
2015 of the Southern Ocean (Bellingshausen Sea, Weddell Sea and East Antarctica), the central
2016 Chilean coast and the Patagonian shelf over the period 1900-2100, with future projections
2017 based on RCP 8.5. From Hauri et al. (2016).

2018

2019