

This is a repository copy of *Competing Pathways in the Photochemistry of Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub>*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/126914/>

Version: Accepted Version

---

**Article:**

Procacci, Barbara orcid.org/0000-0001-7044-0560, Duckett, Simon B. orcid.org/0000-0002-9788-6615, George, Michael W. et al. (6 more authors) (2018) *Competing Pathways in the Photochemistry of Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub>*. *Organometallics*. om-2017-00802b.R1. pp. 855-868. ISSN 0276-7333

<https://doi.org/10.1021/acs.organomet.7b00802>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

### Competing pathways in the photochemistry of $\text{Ru}(\text{H})_2(\text{CO})(\text{PPh}_3)_3$

Journal:	<i>Organometallics</i>
Manuscript ID	om-2017-00802b.R1
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Procacci, Barbara; University of York, Department of Chemistry Duckett, Simon; University of York, Chemistry George, Michael; Nottingham, Chemistry Hanson-Heine, Magnus; Nottingham, Chemistry Horvath, Raphael; University of Nottingham, School of Chemistry Perutz, Robin; University of York, Chemistry Sun, Xue Zhong; University of Nottingham, Chemistry Vuong, Khuong; UQ Dow Centre, School of Chemical Engineering Welch, Janet; University of York, Chemistry

SCHOLARONE™  
Manuscripts

revised Dec 2017

## Competing pathways in the photochemistry of $\text{Ru}(\text{H})_2(\text{CO})(\text{PPh}_3)_3$


Barbara Procacci,<sup>1,2</sup> Simon B. Duckett,<sup>\*1,2</sup> Michael W. George,<sup>\*3,4</sup> Magnus W. D. Hanson-Heine,<sup>3</sup> Raphael Horvath,<sup>3</sup> Robin N. Perutz,<sup>\*1</sup> Xue-Zhong Sun,<sup>3</sup> Khuong Q. Vuong,<sup>3</sup> and Janet A. Welch<sup>1</sup>

1. Department of Chemistry, University of York, York YO10 5DD, UK

2. Centre for Hyperpolarisation in Magnetic Resonance, Department of Chemistry, York Science Park, University of York, York, YO10 5NY, UK

3. School of Chemistry, University of Nottingham, Nottingham NG7 2RD, UK

4. Department of Chemical and Environmental Engineering, The University of Nottingham Ningbo China, 199 Taikang East Road Ningbo 315100, China



### Abstract

The photochemistry of  $\text{Ru}(\text{H})_2(\text{CO})(\text{PPh}_3)_3$  **1** has been re-investigated employing laser and conventional light sources in conjunction with NMR spectroscopy and IR spectroscopy. The sensitivity of NMR experiments was enhanced by use of para- $\text{H}_2$  induced polarization (PHIP) and a series of unexpected reactions observed. The photo-induced reductive elimination of  $\text{H}_2$  was demonstrated (a) *via* NMR spectroscopy by the observation of hyperpolarized **1** on pulsed laser photolysis in the presence of *p*- $\text{H}_2$  and (b) *via* nanosecond time-resolved infrared (TRIR) spectroscopy studies of the transient  $[\text{Ru}(\text{CO})(\text{PPh}_3)_3]$ . Elimination of  $\text{H}_2$  competes with photo-induced loss of  $\text{PPh}_3$  as demonstrated by formation of dihydrogen, triphenylarsine and pyridine substitution products which are detected by NMR spectroscopy. The corresponding coordinatively unsaturated 16-electron intermediate  $[\text{Ru}(\text{H})_2(\text{CO})(\text{PPh}_3)_2]$  exists in two isomeric forms according to TRIR spectroscopy that react with  $\text{H}_2$  and with pyridine on a nanosecond timescale. These two pathways, reductive elimination of  $\text{H}_2$  and

1  
2  
3 PPh<sub>3</sub> loss are shown to occur with approximately equal quantum yields upon 355 nm  
4 irradiation. Low-temperature photolysis in the presence of H<sub>2</sub> reveals the formation of the  
5 dihydrogen complex Ru(H)<sub>2</sub>(η<sup>2</sup>-H<sub>2</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub>, which is detected by NMR and IR  
6 spectroscopy. This complex reacts further within seconds at room temperature and its  
7 behavior provides a rationale to explain the PHIP results. Furthermore, photolysis in the  
8 presence of AsPh<sub>3</sub> and H<sub>2</sub> generates Ru(H)<sub>2</sub>(AsPh<sub>3</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub>. Two isomers of  
9 Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>(pyridine) are formed according to NMR spectroscopy on initial photolysis  
10 of **1** in the presence of pyridine under H<sub>2</sub>. Two further isomers are formed as minor products;  
11 the configuration of each isomer was identified by NMR spectroscopy. Laser pump-NMR  
12 probe spectroscopy was used to observe coherent oscillations in the magnetization of the  
13 one of the isomers of the pyridine complex; the oscillation frequency corresponds to the  
14 difference in chemical shift between the hydride resonances. Pyridine substitution products  
15 were also detected by TRIR spectroscopy.  
16  
17  
18  
19  
20  
21  
22  
23  
24

## 25 Introduction

26  
27 Photochemical methods offer low temperature routes to synthesis, excellent routes to study  
28 reaction mechanisms and great opportunities for catalysis. Indeed, there has been a great  
29 resurgence of interest in photocatalysis.<sup>1-3</sup> Metal hydrides play a major part in the  
30 photochemistry of organometallics and coordination complexes and we have recently  
31 reviewed their behavior systematically.<sup>4</sup> One of the earliest metal hydrides to be studied  
32 photochemically was Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub> **1** (Scheme 1), a compound of great interest for its  
33 catalytic properties both under illumination and thermally.<sup>5-13</sup> The original studies of its  
34 photochemistry demonstrated loss of H<sub>2</sub>, but not CO, on irradiation in benzene. Selectivity  
35 for loss of H<sub>2</sub> over loss of CO is now recognized for several metal carbonyl dihydrides.<sup>14</sup>  
36 When **1** was photolyzed in benzene solution under a CO atmosphere, Ru(CO)<sub>3</sub>(PPh<sub>3</sub>)<sub>2</sub> was  
37 formed leaving the sequence in which H<sub>2</sub> and PPh<sub>3</sub> had been dissociated and CO  
38 coordinated uncertain. In 1997, we identified **1** as an excellent target for investigating the  
39 photodissociation of H<sub>2</sub> by time-resolved UV/vis and time-resolved IR spectroscopy.<sup>15</sup> We  
40 observed a transient by both techniques that was assigned to [Ru(CO)(PPh<sub>3</sub>)<sub>3</sub>] and  
41 measured its rate of reaction with H<sub>2</sub> ((8.4 ± 0.4) × 10<sup>7</sup> dm<sup>3</sup> mol<sup>-1</sup> cm<sup>-1</sup> at room temperature).  
42 It was of particular importance that the transient exhibited a 95 cm<sup>-1</sup> shift to low frequency in  
43 its CO-stretching band, providing proof of H<sub>2</sub> loss. Moreover, ultrafast spectroscopy  
44 demonstrated that it was formed within the response time of the instrument of 6 ps. The only  
45 other transient observed by IR spectroscopy had a significant rise time suggesting that it was  
46 a secondary product. Thus we concluded that there was ultrafast and concerted  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 photochemical reductive elimination of H<sub>2</sub> from Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub>. As we now show, this  
4 study failed to find evidence for a further process because of limitations of the methodology  
5 and the techniques available to us at the time: neither the transient absorption nor the TRIR  
6 experiments probed the timescale of 0.5 - 500 ns and our steady state NMR experiments  
7 were conducted at room temperature and not at low temperature.  
8  
9

10 The advent of in-situ photochemistry at low temperature with NMR detection<sup>16-25</sup>  
11 offered a new method of probing reaction mechanism. In the case of a ruthenium complex  
12 related to **1** in which one phosphine has been replaced by an N-heterocyclic carbene,  
13 Ru(H)<sub>2</sub>(IEt<sub>2</sub>Me<sub>2</sub>)(PPh<sub>3</sub>)<sub>2</sub>(CO) (IEt<sub>2</sub>Me<sub>2</sub> = 1,3-bis(ethyl)-4,5-dimethylimidazol-2-ylidene), we  
14 showed that, the products are actually consistent with two competing pathways, reductive  
15 elimination of H<sub>2</sub> and dissociation of PPh<sub>3</sub>.<sup>26</sup> Loss of PPh<sub>3</sub> was revealed by the formation of a  
16 cyclometalation product in the absence of other substrates and  
17 Ru(H)<sub>2</sub>(IEt<sub>2</sub>Me<sub>2</sub>)(PPh<sub>3</sub>)(CO)(L) (L = pyridine or η<sup>2</sup>-H<sub>2</sub>) in the presence of H<sub>2</sub> or pyridine. The  
18 combination of this photochemical technique with *para*-hydrogen induced polarization (PHIP)  
19 showed weakly enhanced NMR resonances for the precursor and stronger enhancement for  
20 two of its isomers. This effect could be understood by photoinduced reductive elimination of  
21 H<sub>2</sub> followed by oxidative addition of *p*-H<sub>2</sub>. Photochemical loss of phosphine competing with  
22 loss of H<sub>2</sub> was also observed for Ru(H)<sub>2</sub>(PMe<sub>3</sub>)<sub>4</sub>.<sup>27</sup> The timescales available and the  
23 sensitivity of TRIR spectroscopy have improved greatly since our 1997 study and there are  
24 no longer any gaps in the nanosecond time-domain.<sup>28-31</sup>  
25  
26  
27  
28  
29  
30  
31  
32

33 Recently, we returned to the photochemistry of **1** and observed some unexpected  
34 features in its reactivity towards *p*-H<sub>2</sub>. We found that we could observe PHIP-enhanced  
35 hydride resonances if we used a pulsed laser for excitation (at 355 nm) synchronized to the  
36 NMR spectrometer under a pressure of *p*-H<sub>2</sub>.<sup>32</sup> Under these very dilute conditions with single  
37 laser shots, there was negligible photodecomposition. Notably, the observed signal  
38 enhancement decayed on a millisecond timescale and was only observed with  
39 synchronization. This behavior was very different from that of related ruthenium complexes.  
40 A paper published in 2017 has repeated these measurements with 308 nm irradiation and a  
41 hydrogen pressure of 1 atm. The results are similar but some photodecomposition is  
42 reported.<sup>33</sup>  
43  
44  
45  
46


47 In this paper, we report a series of studies of the photochemistry of **1** by NMR and IR  
48 methods; the NMR studies were performed using three types of experiment: (a) broadband  
49 (λ > 290 nm) *ex-situ* irradiation, (b) continuous wave (cw) laser (λ = 325 nm) *in situ*  
50 irradiation (NMR operating frequency 400 MHz); (c) pulsed laser (λ = 355 nm) *in situ*  
51 irradiation (NMR operating frequency 600 MHz). The studies were conducted at  
52 temperatures ranging from 220 -300 K. The infrared studies include time-resolved IR  
53  
54  
55  
56  
57  
58  
59  
60

spectroscopy at room temperature on timescales from nanoseconds to microseconds and FTIR methods at ca. 220 K. They reveal that photodissociation of PPh<sub>3</sub> occurs in addition to reductive elimination of H<sub>2</sub>, leading to substitution products. We show that it is the interplay of these two pathways that limits the use of PHIP methods when applied to **1**.


## Results

**1. Characterization of Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub> **1** and detection of hyperpolarized **1**.** Complex **1** (Scheme 1) has been characterized previously<sup>15,34</sup> but a summary of its main spectroscopic features are provided in Scheme 1 and a <sup>1</sup>H NMR spectrum of the key hydride region is illustrated in Figure 1a; a <sup>31</sup>P{<sup>1</sup>H} NMR spectrum is shown in the Supporting information.

**Scheme 1.** Structure of **1** and spectroscopic features.



When an optically dilute solution of **1** is irradiated with a single shot of the pulsed laser (355 nm) in the presence of *p*-H<sub>2</sub> (3 bar) and monitored through the response from a single, synchronized *r.f.* pulse that takes place up to 5 ms after laser irradiation, enhanced hydride signals at  $\delta$  -6.47 and  $\delta$  -8.30 are observed (Figure 1b). In this spectrum,  $J_{\text{HH}}$  corresponds to the peak separation of the antiphase features, while the other splittings arise from coupling to <sup>31</sup>P nuclei.<sup>32</sup> (The corresponding <sup>1</sup>H{<sup>31</sup>P} spectrum is shown in the SI). Similar hyperpolarization is observed on irradiation at 220 K. This result demonstrates that *p*-H<sub>2</sub> has been incorporated into **1**. Formation of hyperpolarized **1** can be understood by reductive elimination of H<sub>2</sub> upon 355 nm irradiation yielding [Ru(CO)(PPh<sub>3</sub>)<sub>3</sub>] which recombines with *p*-H<sub>2</sub> by oxidative addition. The re-addition of H<sub>2</sub> to [Ru(CO)(PPh<sub>3</sub>)<sub>3</sub>] is known to be fast under these conditions (the time constant for recombination under 3 atm H<sub>2</sub> is 1.4  $\mu$ s) precluding the detection of any transient species.



**Figure 1.** Hydride region of two  $^1\text{H}$  NMR spectra of **1** in  $\text{C}_6\text{D}_6$  after (a) 128 scans probing a concentrated solution of **1**; (b) one scan of an optically dilute solution after laser induced incorporation of  $p\text{-H}_2$  yielding hyperpolarized hydride peaks;  $\tau = 150 \mu\text{s}$ .


## 2. NMR studies of the photochemistry of **1**. (a) Formation of $\text{Ru}(\text{H})_2(\eta^2\text{-H}_2)(\text{CO})(\text{PPh}_3)_2$ ,

**2.** In contrast to the measurements described above, the steady-state photolysis of an optically dilute solution of **1** under a  $p\text{-H}_2$  atmosphere at 295 K in benzene- $d_6$  or toluene- $d_8$  undertaken either ex-situ ( $\lambda > 290 \text{ nm}$ ) or in situ with a laser ( $\lambda = 325 \text{ nm}$ , cw) resulted in no change in the observed NMR spectra of these solutions and PHIP was no longer observed in the hydride resonances of **1**. Notably, a very intense peak for *ortho*- $\text{H}_2$  was visible before irradiation was initiated indicating consumption of  $p\text{-H}_2$  via a thermal route.

When the in situ irradiation process with  $p\text{-H}_2$  was repeated at 223 K (toluene- $d_8$ ), there was still no PHIP enhancement of **1** upon photolysis (2 min,  $\lambda = 325 \text{ nm}$ ). However, a new broad resonance (fwhm ca. 63 Hz at 400 MHz) was observed at  $\delta -5.82$  in the  $^1\text{H}$  NMR spectrum along with new signals in the aromatic region indicating productive photochemistry and the formation of a new species **2** (Figure 2a).

The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum now displayed two new singlets at  $\delta -6.0$  and  $\delta 59.4$  in addition to the resonances of **1**. The new  $^{31}\text{P}\{^1\text{H}\}$  peak resonating at  $\delta -6.0$  was identified as free  $\text{PPh}_3$ , and an experiment with inverse-gated decoupling (quantitative phosphorus) showed that the integration ratio of the unknown peak to the free  $\text{PPh}_3$  is 2:1, suggesting that **2** contains two equivalent  $\text{PPh}_3$  ligands (Figure 2b). Further cooling to 193 K resulted in no change in these NMR features; however, warming to 295 K led to the disappearance of these signals and reformation of **1**. Complex **2** could also be generated at low temperatures with pulsed laser irradiation ( $\lambda = 355 \text{ nm}$ , 48 laser shots, 10 Hz repetition rate). On the basis of these results, we assign **2** as the dihydrogen complex  $\text{Ru}(\text{H})_2(\eta^2\text{-H}_2)(\text{CO})(\text{PPh}_3)_2$  formed by photochemically induced  $\text{PPh}_3$  loss from **1** and coordination of  $\text{H}_2$  (Scheme 2). Upon cooling to 193 K the associated NMR spectra of **2** show minimal change. Moreover, when a  $\{^{31}\text{P}\}^1\text{H}$  NMR spectrum was recorded at 220 K using a broadband decoupling sequence, the peak for **2** remained unaltered whereas the resonances for **1** simplified to doublets retaining


only  $J_{\text{HH}}$  of 6 Hz between the two hydrides. These observations are fully consistent with rapid exchange between hydride and dihydrogen ligands in conjunction with quick relaxation and the averaging of hydride-phosphorus couplings to a small and hence invisible value. The exchange is likely to occur by the  $\sigma$ -CAM mechanism,<sup>35</sup> as has been shown in detail by Crabtree.<sup>36</sup> Furthermore, the disappearance of **2** upon warming is consistent with the well-known lability of dihydrogen as a two electron donor ligand.<sup>37</sup> Complex **2** is an analogue of  $\text{Ru}(\text{H})_2(\eta^2\text{-H}_2)(\text{PPh}_3)_3$  formed from  $\text{Ru}(\text{H})_2(\text{PPh}_3)_4$  thermally and is also closely related to  $[\text{Ru}(\text{IPr})_2(\text{CO})(\eta^2\text{-H}_2)\text{H}]\text{BARF}_4$  (IPr = 1,3-bis(2,6-diisopropylphenyl)imidazol-2-ylidene).<sup>38,39,40</sup> Comparisons may also be made to Ru and Ir complexes studied with partial deuteration.<sup>41,42</sup>



**Figure 2.** NMR spectra after the photolysis of **1** under  $p\text{-H}_2$  in toluene- $d_8$  at 220 K. a)  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum showing the formation of **2** and free  $\text{PPh}_3$ . b) Hydride region of the  $^1\text{H}$  spectrum displaying signals for **1** and a broad resonance for **2**.

**Scheme 2.** Competing photochemistry of **1** under  $\text{H}_2$  atmosphere.





In order to confirm the presence of a dihydrogen ligand in **2**, data for  $T_1$  were recorded at 400 MHz between 212 and 242 K for the resonance at  $\delta -5.82$ . The value of  $T_{1(\text{min})}$  for **2** proved to be  $35 \pm 2$  ms at ca. 225 K, whereas the corresponding value of  $T_1$  for the hydride resonances of **1** was much longer (390 ms at 298 K). This is consistent with a dihydrogen arrangement as such ligands are known to relax much faster than their hydride analogues.<sup>37,43</sup> Further analysis of the  $T_1$  values is given in the Supporting Information.

We have already shown that PHIP was observed for the hydride resonances of **1** when a solution of **1** ( $\text{C}_6\text{D}_6$ ) under  $p\text{-H}_2$  (~4 bar) was irradiated at 298 K with a pulsed laser (355 nm) synchronized to the spectrometer.<sup>32</sup> However, if this sample was irradiated with multiple laser shots (4 sets of 32 laser shots, 10 Hz repetition rate – 1 NMR scan), **2** was detected, although its signal disappeared after the first NMR scan due to its low stability. As complex **2** disappeared, the signals for **1** and a new hydride resonance start to grow in a thermal reaction. This new product signal appears at  $\delta -6.34$ , as a triplet with  $J = 23$  Hz, that connects to a single  $^{31}\text{P}\{^1\text{H}\}$  resonance at  $\delta 57.1$  and simplifies into a singlet upon  $^{31}\text{P}$  decoupling; it is assigned to known  $\text{Ru}(\text{H})_2(\text{PPh}_3)_2(\text{CO})_2$ .<sup>44,45</sup>

Additional evidence for formation of **2** was obtained by photolysis of **1** under a  $\text{D}_2$  atmosphere at 298 K using either a pulsed or continuous laser. Depletion of the hydride signal for **1** during photolysis along with the appearance of peaks for  $\text{H}_2$  ( $\delta 4.46$  in  $\text{C}_6\text{D}_6$ , s) and HD ( $\delta 4.41$  in  $\text{C}_6\text{D}_6$ , t,  $J_{\text{HD}} = 43$  Hz). There was also evidence of partial deuteration of **1** through broadening of the hydride resonances. Direct reductive elimination accounts for formation of  $\text{H}_2$ . HD is logically formed through loss of  $\text{PPh}_3$  and initial  $\eta^2$ -coordination of  $\text{D}_2$  to form the deuterium analogue of **2** followed by intramolecular exchange with the hydride ligands to yield  $\text{Ru}(\text{H})(\text{D})(\text{HD})(\text{CO})(\text{PPh}_3)_2$  and ultimately HD. Complex **1-d<sub>2</sub>** was generated by a sequence of three cycles of photolysis of **1** under  $\text{D}_2$  (3 atm), followed by replacement of the atmosphere by fresh  $\text{D}_2$  and further photolysis. At this stage, no HD or hydride resonances for **1** could be detected. Photo-reaction (325 nm) of the preformed **1-d<sub>2</sub>** with  $\text{H}_2$  at 223 K was then followed by  $^1\text{H}$  NMR and, as expected, the hydride signals of both **1** and **2**


1  
2  
3 were seen. Although this sample should contain partially deuterated **2**, no  $J_{\text{HD}}$  splitting was  
4 observed, even at 193 K. The lack of a visible HD splitting is consistent with rapid exchange  
5 between the hydride and dihydrogen ligands.<sup>37</sup>  
6

7 Although the two hydride ligand signals of **1** are distinct at room temperature,  
8 heteronuclear NOESY experiments have previously demonstrated that **1** undergoes  
9 intramolecular ligand exchange at 335 K where the two hydrides exchange with each other  
10 as well as with the inequivalent phosphines.<sup>46</sup> The exchange was postulated to occur *via* a  
11 trigonal twist mechanism, but alternative mechanisms involving dihydrogen isomers are also  
12 possible.<sup>47</sup> We undertook a series of 1D EXSY experiments to probe for the thermal  
13 exchange of **1** with free H<sub>2</sub> or free PPh<sub>3</sub>. The two hydride peaks and free hydrogen peak  
14 were monitored at mixing times between 50 and 800 ms in C<sub>6</sub>D<sub>6</sub> solution. As expected, slow  
15 intramolecular exchange between the two hydride resonances was observed at room  
16 temperature and faster exchange at 333 K as previously reported,<sup>46</sup> but no exchange with  
17 free H<sub>2</sub> was detected up to 333 K demonstrating that no thermal H<sub>2</sub> elimination occurs on the  
18 EXSY timescale of ca. 1 s (see Supporting Information). In similar experiments, exchange  
19 between free PPh<sub>3</sub> and the phosphine ligands was monitored at 333 K (see Supporting  
20 Information). Once more, no intermolecular exchange was observed in agreement with the  
21 literature.<sup>46</sup>  
22  
23  
24  
25  
26  
27  
28

29 The photoreactions of **1** under H<sub>2</sub> provide evidence for H<sub>2</sub> reductive elimination and re-  
30 addition through PHIP of **1**; evidence for competing PPh<sub>3</sub> dissociation is seen *via* the  
31 observation of **2** and the formation of HD on photolysis under D<sub>2</sub>.  
32  
33


34 **(b) Formation of Ru(H)<sub>2</sub>(AsPh<sub>3</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub>, **3**.** In order to further probe the  
35 phosphine loss pathway, a sample of **1** (C<sub>6</sub>D<sub>6</sub>) was prepared under *p*-H<sub>2</sub> (~4 bar) in the  
36 presence of a 10 fold excess AsPh<sub>3</sub>. A <sup>1</sup>H{<sup>31</sup>P} NMR spectrum measured prior to photolysis  
37 revealed slight conversion of **1** into a new species **3** with two very weak hydride resonances  
38 at  $\delta$  -6.28 and  $\delta$  -9.46 ( $J_{\text{HH}} = 6$  Hz) and a new singlet in the <sup>31</sup>P{<sup>1</sup>H} spectrum at  $\delta$  58. When  
39 the sample was exposed to a single laser shot at 295 K, the hydride resonances of both **1**  
40 and **3** became PHIP-enhanced (Figure 3). The intensities of the new product peaks were  
41 also found to increase on irradiation with more laser shots. These peaks are split into triplets  
42 of antiphase doublets, in an overall 1:1 intensity ratio, in the <sup>1</sup>H NMR spectrum and simplify  
43 into antiphase doublets upon <sup>31</sup>P decoupling. The triplet splitting is due to coupling to two  
44 equivalent <sup>31</sup>P nuclei that lie *cis* to the hydrides ( $\delta$  -6.28,  $J_{\text{PH}} = 30$  Hz ;  $\delta$  -9.46,  $J_{\text{PH}} = 26$  Hz).  
45 We therefore assign **3** to Ru(H)<sub>2</sub>(AsPh<sub>3</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub>, formed by PPh<sub>3</sub> loss and AsPh<sub>3</sub>  
46 coordination, with the geometry shown in Scheme 3. Free PPh<sub>3</sub> was detected in the <sup>31</sup>P{<sup>1</sup>H}  
47 spectrum as expected. Some thermal loss of PPh<sub>3</sub> occurs as is evident from the presence of  
48 a little **3** before irradiation. Additionally, these observations also confirm that photolysis  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

causes PPh<sub>3</sub> loss along with H<sub>2</sub> reductive elimination. Complex **3** is relatively stable and can still be detected at room temperature after the *p*-H<sub>2</sub> response has decayed.



**Figure 3.** Hydride region of a series of <sup>1</sup>H NMR spectra of a solution of **1** in C<sub>6</sub>D<sub>6</sub> in the presence of 10 fold excess AsPh<sub>3</sub> under *p*-H<sub>2</sub> at room temperature showing hyperpolarized signals for **1** and **3** after: a) 1 laser shot, b) 2 laser shots and c) 4 laser shots.

**Scheme 3.** Photochemistry of **1** in the presence of excess AsPh<sub>3</sub> and *p*-H<sub>2</sub>.




**(c) Formation of Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>(pyridine).** An analogous reaction was then undertaken where AsPh<sub>3</sub> was replaced with pyridine. The thermal reaction of **1** with pyridine under *p*-H<sub>2</sub> (4 bar) at 298 K was first monitored in toluene-*d*<sub>8</sub> with added pyridine (10 μL). No reaction was observed by conventional NMR spectroscopy at 295 K; notably the presence of pyridine suppressed the conversion of *p*-H<sub>2</sub> to *o*-H<sub>2</sub> that had been observed in the absence of pyridine. However, when this process was repeated using the high sensitivity Only Para-hydrogen Spectroscopy (OPSY)<sup>47</sup> approach, two weak hydride resonances were observed that we assign to complex Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>(pyridine) **4-CN**. (We label the isomers of **4** according to the ligands that lie *trans* to the mutually *cis* hydride ligands. Thus **4-CN** indicates that the *cis* hydride ligands are *trans* to C and N, see below). Conversion to this product improved on increasing the temperature to 345 K, confirming a thermal route to its formation. The same product could be generated more efficiently by photochemical reaction in neat pyridine; after 3 h irradiation (λ > 290 nm, 295 K), 80% of the starting material **1** was converted to **4-CN** along with some minor by-products (see later).

1  
2  
3 The hydride region of the  $^1\text{H}$  NMR spectrum, measured without hyperpolarization,  
4 shows two major resonances for **4-CN** as triplet of doublets at  $\delta$ -4.59 ( $J_{\text{PH}} = 27$  Hz,  $J_{\text{HH}} =$   
5 6.5 Hz) and  $\delta$ -14.32 ( $J_{\text{PH}} = 23$  Hz,  $J_{\text{HH}} = 6.5$  Hz) in a 1:1 ratio. These signals simplify into  
6 doublets when  $^{31}\text{P}$  is decoupled. The corresponding  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum displayed a  
7 singlet at  $\delta$  65.4 for the main product. On the basis of these results we assign **4-CN** to  
8  $\text{Ru}(\text{H})_2(\text{CO})(\text{PPh}_3)_2(\text{NC}_5\text{H}_5)$ , an analogue of **3**, with mutually *trans*  $\text{PPh}_3$  groups formed by  
9 photochemical  $\text{PPh}_3$  loss and reaction with pyridine. The hydride resonance at  $\delta$  -14.32  
10 displayed a chemical shift characteristic of a hydride *trans* to nitrogen<sup>48</sup> while the remaining  
11 hydride is assigned as *trans* to CO; free  $\text{PPh}_3$  was also detected in the  $^{31}\text{P}\{^1\text{H}\}$  spectrum.  
12 Attempts to isolate **4-CN** were unsuccessful due to the lability of the pyridine ligand; when  
13 the sample was pumped to dryness and redissolved in benzene, **1** was regenerated almost  
14 quantitatively (see Supporting Information). When this reaction was repeated with a 10-fold  
15 excess  $\text{C}_5\text{D}_5\text{N}$  in benzene, the same product distribution was observed.

16  
17  
18 Irradiation of **1** in  $\text{C}_6\text{D}_6$  under  $p\text{-H}_2$  (4 bar) in the presence of 10 fold excess of pyridine-  
19  $d_5$  with the pulsed laser at 295 K led to the detection of strongly hyperpolarized **1** (Figure 4a)  
20 after a single laser shot with a single NMR scan.<sup>32</sup> When the laser was allowed to fire two  
21 shots consecutively (10 Hz repetition rate), PHIP-enhanced resonances of **4-CN** also  
22 became visible after one NMR scan, observed as triplets of antiphase doublets in the  $^1\text{H}$   
23 NMR spectrum. A weakly hyperpolarized resonance at  $\delta$ -4.82 with a shape that indicates a  
24 product with an AA'XX' spin system was also observed (Figure 4b). We assign it to **4-PP**, an  
25 isomer of **4-CN** with a square planar  $\text{Ru}(\text{H})_2(\text{PPh}_3)_2$  skeleton where the two hydrides and two  
26 phosphines are mutually *cis* (Scheme 4a). In this situation, the two hydrides are chemically  
27 equivalent but magnetically inequivalent because of *cis* and *trans* phosphorus couplings,  
28 which satisfies the symmetry-breaking conditions needed to detect PHIP. The  $^{31}\text{P}$  resonance  
29 of **4-PP** was found at  $\delta$  47.8.


30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




**Figure 4.** Hydride region of a series of  $^1\text{H}$  spectra of **1** under  $p\text{-H}_2$  in the presence of excess  $\text{C}_5\text{D}_5\text{N}$  at room temperature. a) After a single laser shot in  $\text{C}_6\text{D}_6$  solution showing hyperpolarized **1**; b) after two consecutive laser shots in  $\text{C}_6\text{D}_6$  solution showing the formation of hyperpolarized **4-PP** and **4-CN**; Inset: expansion of the hydride peak for hyperpolarized **4-PP** (the inset spectrum was acquired with 4 laser shots; the doublet of antiphase doublets marked by an asterisk is due to another isomer **4-NP** see below) and c) spectrum acquired at 237 K in toluene- $d_8$  without hyperpolarization, showing signals for the primary photochemical product **4-PP** with a typical AA'XX' appearance (the change in chemical shift is due to use of a different solvent).

**Scheme 4.** Photoreactions of **1** with pyridine; (a) pump-probe photochemistry of **1** under *p*-H<sub>2</sub> atmosphere; (b) low-temperature photochemistry

(a) Pump-probe with *p*-H<sub>2</sub>




(b) Low temperature photochemistry



The identities of **4-CN** and **4-PP** were confirmed by *in situ* irradiation of a C<sub>6</sub>D<sub>6</sub> solution of **1** in the presence of 10 fold excess <sup>15</sup>N-pyridine under *p*-H<sub>2</sub> pressure with the pulsed laser. <sup>1</sup>H-<sup>15</sup>N HMQC spectroscopy showed a cross peak between the hydride peak at  $\delta$  – 14.32 for complex **4-CN** and a resonance at  $\delta$  270.4 in the <sup>15</sup>N spectrum which now exhibits a *trans* <sup>15</sup>N coupling of  $J_{\text{NH}} = 12$  Hz (see Supporting Information). A cross peak was also observed between the hydride resonance for complex **4-PP** and a signal in the <sup>15</sup>N NMR spectrum at  $\delta$  256.3 indicating a small *cis*  $J_{\text{NH}}$  coupling (< 2 Hz).

The irradiation of a solution of **1** in C<sub>6</sub>D<sub>6</sub> in the presence of 10 fold excess C<sub>5</sub>D<sub>5</sub>N under *p*-H<sub>2</sub> pressure at room temperature (Figure 5a) with 32 pulsed laser shots (355 nm, 10 Hz repetition rate) resulted in almost quantitative conversion to **4-CN** (Figure 5b). Given that we are only irradiating a small sample cross section but monitoring across the whole sample, we can confirm that diffusional mixing is rapid on this timescale in order to allow for the quantitative conversion. When this sample was shaken to refresh the dissolved *p*-H<sub>2</sub> to test thermal exchange between free H<sub>2</sub> and the hydrides of **4-CN**, PHIP was detected on the hydride resonances (Figure 5c) leading to the conclusion that the hydride ligands of the pyridine complex **4-CN** exchange thermally with free H<sub>2</sub> at room temperature. These observations contrast with the behavior of **1** for which no direct thermal exchange with free H<sub>2</sub> was observed at room temperature. When the same solution of **4-CN** is left overnight and checked again by NMR, quantitative regeneration of the starting complex **1** is observed highlighting the lability of the pyridine ligand which is slowly displaced by the stoichiometric


amount of  $\text{PPh}_3$  present in solution after photosubstitution.




**Figure 5.** Hydride region of a series of  $^1\text{H}$  NMR spectra of a solution of **1** in  $\text{C}_6\text{D}_6$  in the presence of a 10 fold excess of  $\text{C}_5\text{D}_5\text{N}$  under  $\text{H}_2$  pressure at room temperature. a) Hydride resonances of **1** before laser irradiation; b) after 32 laser shots at 355 nm showing mainly **4-CN**; c) same sample after being shaken with  $p\text{-H}_2$  that displays PHIP of **4-CN**.

The photoreaction of **1** with pyridine (10  $\mu\text{L}$ ) under  $p\text{-H}_2$  was also studied with cw laser irradiation (325 nm) at 223 K in toluene- $d_8$ . Following photolysis (2 min), hyperpolarized signals were observed for **4-CN** and **4-PP** but not for **1**. (Note that no hyperpolarization of **1** is observed unless the pulsed laser is used and synchronized with the *r.f.* pulse of the spectrometer.) There was no evidence for the formation of **2** in these spectra, but the persistence of hyperpolarization after the photolysis source has been switched off confirms that **4-CN** and **4-PP** continue to undergo rapid thermal hydride ligand exchange with  $p\text{-H}_2$  as demonstrated above.

In order to probe the isomerization of **4-CN** and **4-PP**, *in-situ* photolysis of **1** in neat  $\text{C}_5\text{D}_5\text{N}$  was conducted at 237 K without a  $\text{H}_2$  atmosphere. After 24 laser shots, 2.5% conversion was observed with the initial ratio of **4-PP** to **4-CN** being 2:1. Further photolysis increased the overall conversion level without change in the ratio of **4-PP** to **4-CN** (Figure 6). The proportion of **4-CN** increased, however, with increase in temperature suggesting thermal equilibration between the two isomers (Figure 7). When the solution was allowed to warm up to room temperature **4-PP** converted to **4-CN** quantitatively. Totally analogous behavior was found when the same experiment was repeated under a  $\text{H}_2$  atmosphere (4 bar). Similarly, no changes in behavior were observed when the reaction was run in toluene- $d_8$  with 10-fold excess pyridine. These experiments lead to the surprising conclusion that **4-PP** is the first product of this reaction to be formed at low temperature that is stable on an NMR timescale, i.e. the primary photoproduct (Scheme 4b).



**Figure 6.** Hydride region of a series of  $^1\text{H}\{^{31}\text{P}\}$  spectra of a solution of **1** in neat  $\text{C}_5\text{D}_5\text{N}$  at 237 K. a) Starting solution at photolysis time = 0; b) after 2.5 s photolysis at 355 nm; c) after 35 s photolysis;




**Figure 7.** Hydride region of a series of  $^1\text{H}\{^{31}\text{P}\}$  spectra of a solution of **1** in neat  $\text{C}_5\text{D}_5\text{N}$  after 140 s of photolysis showing equilibration between **4-PP** and **4-CN** with increasing temperature. a)  $T = 237$ ; b)  $T = 258$  K; c)  $T = 273$  K; d)  $T = 295$  K.

We then used the PHIP effect to search for the formation of minor products in this reaction. When a sample of **1** under  $p\text{-H}_2$  in  $\text{C}_6\text{D}_6$  with a 10-fold excess of pyridine was exposed to multiple shots (up to 8) of the pulsed laser (355 nm) two new hyperpolarized minor species appeared in addition to those of **4-CN** and **4-PP** (Figure 8). Furthermore, under these higher signal-to-noise conditions hyperpolarization was clearly visible in the *ortho* aromatic protons of **4-PP** and the  $^{31}\text{P}$  resonance of this complex at  $\delta$  47.8 (see Supporting Information). Analogous behavior has been observed before for a complex with chemically equivalent but magnetically inequivalent hydrides.<sup>32</sup> The hydride resonances for the first of these species, complex **4-NP**, appear at  $\delta$  -6.91 and -15.28 (Scheme 5). The hydride at  $\delta$  -6.91 was assigned to lie *trans* to a  $\text{PPh}_3$  group because of the large  $J_{\text{PH}}$  coupling (79 Hz, *cis*  $J_{\text{PH}}$  was measured as 40 Hz). The other hydride ligand of **4-NP** was detected by photochemical hyperpolarized COSY and assigned to lie *trans* to pyridine ( $J_{\text{PH}} = 16, 30$  Hz, see Supporting Information for COSY). The second product, **4-CP**, exhibits a much weaker pair of hydride ligand signals whose chemical shifts and coupling constants



( $\delta$  -3.47;  $J_{\text{PH}} = 23, 30$  Hz and  $-5.35$ ;  $J_{\text{PH}} = 32, 98$  Hz) indicated that one hydride is *trans* to CO and the other *trans* to PPh<sub>3</sub>. With <sup>15</sup>N labeled pyridine, a very weak cross peak between the hydride peak at  $\delta$  -15.28 of complex **4-NP** and a signal in the <sup>15</sup>N NMR spectrum at  $\delta$  266.2 was also detected ( $J_{\text{NH}} = 14$  Hz) confirming that the pyridine lies *trans* to the hydride, as indicated by the characteristic chemical shift. No <sup>15</sup>N-<sup>1</sup>H cross peaks were observed for **4-CP**. A change of the solvent to neat pyridine did not provide additional information.

**Scheme 5.** Structures of the minor photolysis products with added pyridine.




**Figure 8.** Hydride region of the hyperpolarized <sup>1</sup>H NMR spectrum of a C<sub>6</sub>D<sub>6</sub> solution of **1** under *p*-H<sub>2</sub> in the presence of excess C<sub>5</sub>D<sub>5</sub>N. The solution was monitored after 8 laser shots.

In order to quantify the ratio of H<sub>2</sub> reductive elimination to PPh<sub>3</sub> loss, a sample of **1** was irradiated (355 nm) with the same concentration of D<sub>2</sub> and pyridine-*d*<sub>5</sub> (both 0.012 M, 10 fold excess) at 300 K. Under these conditions, H<sub>2</sub> loss is followed by D<sub>2</sub> oxidative addition to form **1-d**<sub>2</sub>; loss of PPh<sub>3</sub> leads to conversion to pyridine complex **4-CN**. Pentafluoroanisole was used as an internal standard after ensuring that the compound was inert to it under these reaction conditions. <sup>1</sup>H NMR spectra were acquired at different photolysis time intervals; the reaction was taken to 50% conversion. The overall quantity of **1** and **1-d**<sub>2</sub> was measured by integration of the *ortho* protons (12 H) ( $\delta$  7.46) for the axial phenyl rings of **1** that are well resolved. The integration of the hydride peak of **1** reported on the amount of **1** that remained undeuterated. The *ortho* protons (12 H) of **4-CN** ( $\delta$  7.94) quantified the production of **4-CN**.

1  
2  
3 Because of thermal exchange with  $D_2$  (see above), the hydride resonances for complex **4-CN**  
4 were not detected. This experiment established that at 50% conversion of **1**,  $H_2$   
5 reductive elimination accounts for  $\leq 40\%$  of the photochemical reactivity while  $PPh_3$  loss  
6 takes place with 60% probability. We were concerned that under these conditions formation  
7 of **2-d<sub>2</sub>** could compete with formation of **4-CN** following loss of  $PPh_3$ . We therefore repeated  
8 the experiment under exactly the same conditions but with a large excess of pyridine with  
9 respect to  $D_2$  in order to turn formation of **2-d<sub>2</sub>** into an uncompetitive route. Under these  
10 conditions, the ratio of  $PPh_3$  loss to  $H_2$  reductive elimination products were 52:48. These  
11 results indicate approximately the same quantum yields for the two photochemical  
12 processes.  
13  
14  
15  
16  
17

18 We have previously shown that it is possible to probe the hydride signal intensities  
19 under hyperpolarization conditions in a series of NMR measurements recorded as a function  
20 of the delay,  $\tau$ , between the laser irradiation step and the *r.f.* pulse. When an optically dilute  
21 solution of **1** in  $C_6D_6$  is used and the appropriate laser energy applied there is no  
22 decomposition of the sample due to photolysis, this allows the same solution to be used  
23 multiple times. The response is expected to oscillate at the frequency difference between the  
24 two coupled hydride resonances ( $\Delta\nu$  in  $C_6D_6$  9.75 ppm = 5851 Hz).<sup>32,49</sup> The evolution of the  
25 PHIP-enhanced NMR signals for **4-CN** was investigated in this way. A  $C_6D_6$  solution of **1**  
26 in the presence of excess pyridine under *p*- $H_2$  (4 bar) was therefore irradiated with the pulsed  
27 laser (355 nm). The resonances of **1** oscillate in intensity as reported before. Analogous  
28 coherent oscillations in intensity were found for the hydride resonances of **4-CN** at a  
29 frequency of  $5850 \pm 5$  Hz. Figure 9 illustrates the oscillation in the form of a 2D spectrum  
30 fourier-transformed with respect to the pump-probe delay,  $\tau$ .  
31  
32  
33  
34  
35  
36  
37  
38  
39




40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53 **Figure 9.** 2D  $^1H$  pump-probe NMR spectrum of a  $C_6D_6$  solution of **4-CN** where the vertical  
54 dimension corresponds to evolution frequency encoded by  $\tau$ . Measurements are made with  
55 four laser pulses followed at delay  $\tau$  by a single NMR pulse. Hydrides  $\Delta\nu$  in  $C_6D_6$  9.75 ppm =  
56  
57

5851 Hz, oscillation frequency  $5850 \pm 5$  Hz.

The NMR results demonstrate clearly that two competing pathways occur during the photochemistry of complex **1**. One photochemical process is H<sub>2</sub> reductive elimination, as demonstrated by the observation of PHIP in the associated hydride peaks after irradiation under a *p*-H<sub>2</sub> atmosphere and incorporation of deuterium if photolyzed under a deuterium pressure. *In-situ* photolysis at low temperature led to the discovery of a competing photochemical pathway involving PPh<sub>3</sub> loss, first identified through the formation of the dihydrogen complex **2**. Subsequently, the triphenylarsine complex **3** and the pyridine complexes, **4-CN** and **4-PP** were formed without the need for low temperatures. We determined that quantum yields for H<sub>2</sub> reductive elimination and PPh<sub>3</sub> loss at 300 K to be approximately equal. The use of *p*-H<sub>2</sub> is critical to revealing some of these species. However, the formation of the dihydrogen complex **2** causes *p*-H<sub>2</sub> relaxation, but this conversion can be suppressed by addition of other ligands such as pyridine. We observed no Ru(0) products either at room temperature or low temperature. We therefore wished to know if evidence from TRIR spectroscopy is compatible with these conclusions.

**2. IR studies.** In the earlier experiments, the transient [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] was detected by both time resolved UV-vis and IR spectroscopy of **1**, [ $\lambda_{\text{max}} = 380$  nm,  $\nu(\text{CO}) = 1845$  cm<sup>-1</sup>] and found to react with H<sub>2</sub> to regenerate **1** on a microsecond timescale [ $k_2 = (8.4 \pm 0.4) \times 10^7$  dm<sup>3</sup> mol<sup>-1</sup>s<sup>-1</sup>]. The unsaturated species [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] was found to be formed within 6 ps by ultrafast TRIR indicating that reductive elimination of H<sub>2</sub> also takes place on this timescale.<sup>15</sup> Our new results suggested that it should also be possible to detect PPh<sub>3</sub> loss by TRIR spectroscopy. We have therefore undertaken new TRIR experiments with modern equipment with higher sensitivity and nanosecond time-resolution. We first report investigations of the photochemistry of **1** by low temperature FTIR spectroscopy.

**(a) Photolysis of 1 monitored by FTIR spectroscopy.** Figure 10a shows the IR difference spectra obtained following irradiation of a solution of **1** in toluene-*d*<sub>8</sub> at 220 K under Ar. The parent  $\nu(\text{CO})$  band at 1937 cm<sup>-1</sup> is bleached and a new band at 1980 cm<sup>-1</sup> is formed. The band at 1980 cm<sup>-1</sup> can be assigned to a dimeric product, as was proposed in the previous TRIR studies.<sup>15</sup>




**Figure 10.** Difference spectra of **1** after photolysis at 220 K in toluene- $d_8$  (a) under argon; (b) under hydrogen (ca. 1 atm) (c) with added pyridine in toluene- $d_8$  (0.1 mL in 5.0 mL) under argon. The  $\Delta$ Absorbance values have been normalized.

A similar experiment, performed under  $H_2$  (1 atm), led to bleach of the parent band and the production of a broad band at ca. 1975  $cm^{-1}$  (Figure 10b). These spectra were fitted using a multi-Lorentzian function indicating the presence of two overlapping bands at ca. 1980 and 1971  $cm^{-1}$ . The solution was warmed up to 240 K and FTIR spectra were acquired every 3 min. The band at 1971  $cm^{-1}$  decayed much faster than the 1980  $cm^{-1}$  band providing further support for the presence of two bands. The band at 1971  $cm^{-1}$  can be assigned to the hydride dihydrogen species **2** which arises from phosphine loss and reaction of the  $[Ru(PPh_3)_2(CO)(H)_2]$  fragment with  $H_2$  in agreement with the result from NMR spectroscopy.


The photolysis of **1** at 220 K was carried out in the presence of added pyridine in toluene under Ar and again the dimeric band was observed together with a new band at 1911  $cm^{-1}$  (Figure 10c). We also carried out a similar experiment at room temperature in benzene with added pyridine- $d_5$  and observed the product at 1921  $cm^{-1}$  (see SI). We assign the band at 1921  $cm^{-1}$  to complex **4-CN** and the band at 1911  $cm^{-1}$  observed at 220 K to **4-PP**, based on the isomer distribution observed by NMR spectroscopy.

**TRIR spectroscopy.** Earlier TRIR studies were conducted in  $C_6D_6$  at room temperature and showed that  $H_2$  elimination from **1** ( $\nu(CO)$  1940  $cm^{-1}$ ) formed the 16e<sup>-</sup> Ru(0) species  $[Ru(CO)(PPh_3)_3]$  with a  $\nu(CO)$  band at 1840  $cm^{-1}$ .<sup>15</sup> These results were obtained covering a time window of 1 to 1000  $\mu s$  employing a lower resolution instrument based on a CO laser.<sup>15</sup> We have repeated these measurements using more sensitive instrumentation with faster time resolution. All the IR experiments were carried out in optically dilute solutions of **1** in  $C_6D_6$  for consistency with the NMR experiments.

1  
2  
3 **TRIR of 1 under an Ar atmosphere in C<sub>6</sub>D<sub>6</sub>.** Figure 11 shows the TRIR difference  
4 spectra at 298 K of **1** in C<sub>6</sub>D<sub>6</sub> obtained at 1 ns (a) and 23 μs (b) following irradiation at 355  
5 nm. The TRIR spectrum obtained after 1 ns shows bleach of the parent band and a product  
6 band at 1840 cm<sup>-1</sup> (not shown in Figure 11) which was observed in the previous TRIR  
7 experiments<sup>15</sup> and can be assigned to the Ru(0) species [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] arising from H<sub>2</sub>  
8 reductive elimination. Two new bands are also produced at ca. 1951 and 1911 cm<sup>-1</sup> which  
9 were not observed in the previous TRIR measurements. The TRIR spectrum obtained 28 μs  
10 after the flash shows that these two bands have decayed and a new absorption is formed  
11 due to the dimeric product at 1971 cm<sup>-1</sup>. The bands at 1951 and 1911 cm<sup>-1</sup> are tentatively  
12 assigned to two different isomers of [Ru(PPh<sub>3</sub>)<sub>2</sub>(CO)H<sub>2</sub>] produced by PPh<sub>3</sub> loss. These  
13 bands decay with two components, a fast component (1.7 ns) that approximately matches  
14 rapid reformation of the parent (ca. 4 ns, a,b) and a slower decay (ca. 10 μs, c,d) which is  
15 similar to the formation of the dimeric band and the further bleaching of the parent. The  
16 precise nature of the fast component is not known but could be due to in-cage recombination  
17 of photoejected PPh<sub>3</sub>. The slower kinetics show further bleaching of **1** as the dimer band  
18 forms at 1971 cm<sup>-1</sup>. This is consistent with formation of the dimer by reaction of the fragment  
19 [Ru(PPh<sub>3</sub>)<sub>2</sub>(CO)H<sub>2</sub>] with the parent **1** (Figure 12c,d).  
20  
21  
22  
23  
24  
25  
26  
27  
28




**Figure 11.** TRIR difference spectra obtained in the range (a) 1-20 ns and 1900-2000  $\text{cm}^{-1}$ ; and (b) 0.4-28  $\mu\text{s}$  and 1800-2000  $\text{cm}^{-1}$ , after 355 nm laser flash of a solution of **1** in  $\text{C}_6\text{D}_6$  under an atmosphere of argon.




**Figure 12.** Kinetic plots observed after photolysis of **1** under Ar. Plots a and b on nanosecond timescale, plots c and d on microsecond timescale. (a) 1939  $\text{cm}^{-1}$  assigned to **1** which shows a rapid increase, followed by a bleach recovery,  $\tau = 4.8 \pm 0.4$  ns; (b) 1951  $\text{cm}^{-1}$ ,  $\tau = 1.7 \pm 0.4$  ns, (c) 1938  $\text{cm}^{-1}$  assigned to **1**,  $\tau_1 = 1.1 \pm 0.4$   $\mu\text{s}$ ,  $\tau_2 = 9 \pm 2$   $\mu\text{s}$  and (d) 1970  $\text{cm}^{-1}$  assigned to the dimeric species,  $\tau = 12 \pm 2$   $\mu\text{s}$ .  $\Delta$ Absorbances have been normalized.

**TRIR of **1** under  $\text{H}_2$  in  $\text{C}_6\text{D}_6$ .** Irradiation of a solution of **1** in the presence of  $\text{H}_2$  (1 atm) displays, at early time (Figure 13a), a negative band for **1** and new product bands to both higher and lower frequency of the parent. In agreement with the previous results, the formation of  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  at 1840  $\text{cm}^{-1}$  is also observed within the time resolution of 0.5 ns; the transient  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  was stable up to 28 ns. A product band at 1911  $\text{cm}^{-1}$  was formed within 1 ns which decayed with time constant  $4.5 \pm 0.5$  ns and the bleach recovered partially with a time constant  $3.0 \pm 0.5$  ns (Figure 14). The spectrum under  $\text{H}_2$  at later times,  $\Delta t = 20$   $\mu\text{s}$ , (Figure 13b) superficially resembles the spectrum obtained under argon; however, close inspection shows that the broad band at  $\sim 1971$   $\text{cm}^{-1}$  is much more intense relative to the parent bleach in the spectrum under hydrogen. Furthermore, the band at 1971  $\text{cm}^{-1}$  grew in significantly faster (ca. 700 ns) under hydrogen compared to the results obtained under argon (ca. 11  $\mu\text{s}$ ). These results are consistent with the formation of **2** and its

isomers as a result of the reaction of  $[\text{Ru}(\text{PPh}_3)_2(\text{CO})(\text{H})_2]$  with dihydrogen after  $\text{PPh}_3$  loss. The formation of some of the dimeric ruthenium species under these conditions cannot be excluded since the  $\nu(\text{CO})$  bands of **2** and the dimeric species have been shown to overlap in low temperature FTIR experiments above.





**Figure 13.** TRIR difference spectra obtained in the range (a) 1-30 ns and  $1900\text{-}2000\text{ cm}^{-1}$ ; and (b)  $0.1\text{-}15\text{ }\mu\text{s}$  and  $1800\text{-}2000\text{ cm}^{-1}$  after 355 nm laser flash of a solution of **1** in  $\text{C}_6\text{D}_6$  under  $\text{H}_2$  (ca. 1.5 atm).



**Figure 14.** Kinetic plots at early times for (a) bleach recovery of **1** at  $1938\text{ cm}^{-1}$ ,  $\tau = 3.0 \pm 0.5$  ns; (b) loss of  $1911\text{ cm}^{-1}$  band assigned to  $[\text{Ru}(\text{CO})(\text{PPh}_3)_2(\text{H})_2]$ ,  $\tau = 4.5 \pm 0.5$  ns.

1  
2  
3  
4 **TRIR of 1 in the presence of added pyridine under Ar.** The TRIR spectra obtained  
5 after 355 nm irradiation solution of **1** in C<sub>6</sub>D<sub>6</sub> with added pyridine (ca. 10<sup>-2</sup> M) do not show the  
6 presence of the band at 1971 cm<sup>-1</sup> indicating that neither the dimeric product nor **2** are  
7 formed under these conditions. However, a clear band at 1915-1920 cm<sup>-1</sup> with shoulders  
8 both to lower and higher wavenumbers grows in over 100 ns and is assigned to complex **4-CN**  
9 or its isomers arising from the reaction of [Ru(PPh<sub>3</sub>)<sub>2</sub>(CO)(H)<sub>2</sub>] with pyridine. This band  
10 remains unchanged out to 28 μs (see Supporting Information).  
11  
12

13  
14  
15 **TRIR of 1 in the presence of added pyridine under H<sub>2</sub>.** The TRIR spectra obtained  
16 after irradiation at 355 nm of a solution of **1** in C<sub>6</sub>D<sub>6</sub> with added pyridine (ca. 10<sup>-2</sup> M) under a  
17 H<sub>2</sub> pressure (ca. 1 atm) showed a negative band for **1** at the earliest time (0.5 ns) and a band  
18 of [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] as observed in all the earlier experiments. Additionally, the band at 1951  
19 cm<sup>-1</sup> is detected at early times. Two new bands grew in over 70 ns at 1912 cm<sup>-1</sup> and 1975  
20 cm<sup>-1</sup> (Figure 15a). The band at 1912 cm<sup>-1</sup> is assigned to the formation of complex **4-PP** or  
21 isomers while the remaining band coincides with those of complex **2** and the dimer. The  
22 band at 1840 cm<sup>-1</sup> decreases and gives way to a band at 1851 cm<sup>-1</sup> which may be assigned  
23 to Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)(py). On a longer timescale, these bands are stable (Figure 15b).  
24  
25  
26  
27  
28





1  
2  
3  
4 **Figure 15.** TRIR difference spectra obtained in the range (a) 1-70 ns and 1800-2000  $\text{cm}^{-1}$  (b)  
5 0.4 - 28  $\mu\text{s}$  after 355 nm laser flash of a solution of **1** in  $\text{C}_6\text{D}_6$  with added pyridine (ca.  $10^{-2}$  M)  
6 under 1 atm  $\text{H}_2$ .  
7

## 8 **Discussion**

9  
10 The NMR results discussed above differ from our earlier studies in that we see  
11 evidence for photochemical loss of both  $\text{H}_2$  and of  $\text{PPh}_3$ . In the presence of  $\text{H}_2$  we see  
12 formation of dihydrogen complex **2** while in the presence of triphenylarsine we observe  
13 formation of **3**. Use of pyridine as the incoming ligand yields **4-CN** and its isomers. The low  
14 temperature FTIR and room temperature TRIR experiments show both intermediates and  
15 products consistent with this evidence.  
16  
17

18 **NMR results.** The behavior of the pyridine isomers is unusual; it might be expected  
19 that the initial product would be **4-CN** with *trans*  $\text{PPh}_3$  ligands, but the major product at low  
20 temperature is **4-PP**. Formation of **4-PP** requires dissociation of  $\text{PPh}_3$  and rearrangement of  
21 the 5-coordinate intermediate  $[\text{Ru}(\text{H})_2(\text{CO})(\text{PPh}_3)_2]$  to bring the  $\text{PPh}_3$  ligands into a *cis*  
22 orientation. The rearrangement is likely to take place either in an electronic excited state or  
23 in a vibrational excited state, as has been documented in detail for  $\text{Fe}(\text{CO})_4$  and  
24  $\text{Cr}(\text{CO})_5$ .<sup>14,50-55</sup> Initial loss of  $\text{PPh}_3$  *trans* to hydride followed by two successive Berry  
25 pseudorotations and coordination of pyridine would generate **4-PP**. The surprising feature is  
26 that this process is selective for **4-PP**. The isomer **4-PP** rearranges thermally to **4-CN**. Two  
27 further isomers of the pyridine complexes **4-NP** and **4-CP** are formed as minor  
28 photoproducts. All the isomers are accessible by a single trigonal twist starting from **4-PP**  
29 (see Supporting Information).<sup>56,57</sup> Alternatively, they may occur by isomerization to a  
30 dihydrogen complex, followed by a Berry pseudo-rotation.  
31  
32  
33  
34  
35  
36  
37


38 It may be asked why parallel isomerization processes are not observed for **2** and **3**. In  
39 the case of **2**, we have evidence for rapid exchange of dihydride and dihydrogen ligands  
40 which causes rapid conversion to the most stable isomer. For **3**, there is evidence for two  
41 further isomers on irradiation with multiple laser shots that are analogues of **4-NP** and **4-CP**  
42 (see Supporting Information), but the experiments are limited by the solubility of  $\text{AsPh}_3$  at  
43 low temperature.  
44  
45  
46

47 The observation of hyperpolarized hydride ligand signals for **3** and **4** on photolysis  
48 requires both the substitution of  $\text{PPh}_3$  by incoming ligand and reductive elimination of  $\text{H}_2$   
49 followed by oxidative addition of *p*- $\text{H}_2$ . Considering the short lifetime of hyperpolarized **1**, it is  
50 likely that  $\text{PPh}_3$  substitution occurs first. We have shown that **3** and **4** undergo thermal  
51 exchange with *p*- $\text{H}_2$  and they may also undergo photochemical exchange. The mechanism  
52 of this thermal exchange may involve loss of  $\text{AsPh}_3$  or pyridine from **3** or **4**, respectively,  
53 followed by formation of **2**, intramolecular exchange of hydrogens and recoordination of  
54  
55  
56  
57  
58  
59  
60

ligand, as proposed for related iridium complexes.<sup>58</sup> Alternatively and more simply, **3** and **4** may be converted to dihydrogen isomers which undergo exchange with *p*-H<sub>2</sub> and regeneration of the initial structure.

**IR results.** The IR spectra and reactivity of the Ru(0) intermediate, [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)], are consistent with a single isomer of this species. On the other hand, the IR spectra point to the formation of at least two isomers of the Ru(II) intermediate [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>] (Scheme 6).

**Scheme 6.** Summary of TRIR observations with experimental IR wavenumbers for  $\nu(\text{CO})$



We have investigated the assignments further by use of Timney ligand effect constants<sup>59</sup> and by estimating the CO stretching frequencies by DFT methods (Table 1). The ligand effect constant for PPh<sub>3</sub> *cis* to CO is negative while that for *trans*-PPh<sub>3</sub> is positive; thus values of  $\nu(\text{CO})$  will be very sensitive to the OC–Ru–P angle. The values for hydride are positive for both *cis* and *trans* geometries and greatly exceed the values for the other ligands. The Timney calculations are consistent with a butterfly structure for [Ru(PPh<sub>3</sub>)<sub>3</sub>CO] and two isomers of square pyramidal [Ru(CO)(PPh<sub>3</sub>)<sub>2</sub>(H)<sub>2</sub>]. Loss of PPh<sub>3</sub> *trans* to hydride without rearrangement of the skeleton of **1** should cause a high frequency shift in  $\nu(\text{CO})$  with

respect to **1**, whereas rearrangement to an isomer with mutually *cis* hydrides and CO *trans* to a vacancy should cause a shift lower frequency than that of **1**. The corresponding DFT calculations are summarized in Table 1. In light of these calculations, we deduce that PPh<sub>3</sub> photo-ejection does indeed generate two isomers of [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>].

The  $\nu(\text{CO})$  band of dihydrogen complex **2** is reproduced very accurately by both calculation methods; the high frequency shift is consistent with the expected  $\pi$ -acceptor characteristics. The values for pyridine complexes **4-CN** and **4-PP** pose greater problems (see SI).

The intermediate [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>] is much more reactive than [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] and we anticipate that it will react with incoming substrates to generate isomeric products. The nanosecond TRIR spectra showed strong evidence for partial in-cage recombination of PPh<sub>3</sub> with [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>] with time constant of ca. 5 ns. The reaction of this species with H<sub>2</sub> under 1.5 atm (the solubility of H<sub>2</sub> in benzene is  $2.9 \times 10^{-3} \text{ mol dm}^{-3} \text{ atm}^{-1}$ )<sup>60</sup> took place with  $\tau \sim 700 \text{ ns}$ , while the reaction with pyridine occurred with  $\tau \sim 70 \text{ ns}$  at  $10^{-2} \text{ mol dm}^{-3}$  pyridine, indicating considerably higher reactivity toward pyridine than toward H<sub>2</sub>. The reaction of [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] with H<sub>2</sub> is considerably slower ( $\tau$  ca. 3  $\mu\text{s}$  at the same pressure).

**Table 1. Wavenumbers of  $\nu(\text{CO})$  bands/cm<sup>-1</sup>**

Complex	$\nu(\text{CO})$ exptl in C <sub>6</sub> D <sub>6</sub>	$\nu(\text{CO})$ calcd by Timney method <sup>a</sup>	$\nu(\text{CO})$ calcd by DFT <sup>b</sup>
<b>1</b>	1939	1947	1927
<b>2</b>	1971	1971	1974
spy Ru(H) <sub>2</sub> (CO)(PPh <sub>3</sub> ) <sub>2</sub> vacancy <i>trans</i> to H <sup>c</sup>	1951	1960	1973
spy Ru(H) <sub>2</sub> (CO)(PPh <sub>3</sub> ) <sub>2</sub> vacancy <i>trans</i> to CO <sup>c</sup>	1911	1925	
Ru(CO)(PPh <sub>3</sub> ) <sub>3</sub> sq planar <sup>c</sup>	1840	1923	
Ru(CO)(PPh <sub>3</sub> ) <sub>3</sub> butterfly	1840	1842	1903

<sup>a</sup> Reference <sup>59</sup>

<sup>b</sup> M06/LACVP(d) with 0.96 scaling

<sup>c</sup> spy = square pyramidal; sq = square

## Conclusions

The evidence from *in-situ* photochemistry of **1** with both NMR and IR detection that PPh<sub>3</sub> loss competes with H<sub>2</sub> reductive elimination is very strong indeed. When we first obtained

1  
2  
3 NMR evidence for PPh<sub>3</sub> substitution by H<sub>2</sub> to form Ru(H)<sub>2</sub>(η<sup>2</sup>-H<sub>2</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub> **2**, the  
4 experiments seemed to conflict with our earlier TRIR experiments.<sup>15</sup> However with the  
5 improved performance of modern TRIR experiments, it is now evident that we missed this  
6 reaction because of the need for NMR spectroscopy at low temperature and TRIR  
7 spectroscopy with greater sensitivity and time resolution. Our NMR experiments indicate that  
8 the quantum yields for PPh<sub>3</sub> loss and H<sub>2</sub> reductive elimination are approximately equal on  
9 355 nm photolysis.

10  
11  
12  
13 The TRIR experiments allow characterization of coordinatively unsaturated  
14 [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] and two isomers of [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>] via their ν(CO) bands. They also  
15 demonstrate that [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>] is considerably more reactive toward H<sub>2</sub> than  
16 [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)]. The reaction of [Ru(PPh<sub>3</sub>)<sub>3</sub>(CO)] with *p*-H<sub>2</sub> generates **1** in hyperpolarized  
17 form, but this species relaxes fast and does not exchange with *p*-H<sub>2</sub> and consequently can  
18 only be detected if the laser pulse is synchronized with the *r.f.* pulse from the NMR  
19 spectrometer. The reaction of [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>] with H<sub>2</sub> generates the dihydrogen  
20 complex Ru(H)<sub>2</sub>(η<sup>2</sup>-H<sub>2</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub> **2**. Its short T<sub>1</sub> provides a route to relaxation of *p*-H<sub>2</sub> to  
21 thermal equilibrium of *o*-H<sub>2</sub> and *p*-H<sub>2</sub>; however, its formation and the fast relaxation process  
22 can be suppressed by addition of other ligands. Reaction of AsPh<sub>3</sub> in the presence of *p*-H<sub>2</sub>  
23 generates hyperpolarized Ru(H)<sub>2</sub>(AsPh<sub>3</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub> **3**. The reaction of [Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>]  
24 with pyridine is more complex. Use of low temperature methods shows that the initial product  
25 is Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>2</sub>(pyridine) **4-PP** with a planar Ru(H)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> core and pyridine *trans* to  
26 CO. This species rearranges on warming to **4-CN**, the pyridine analogue of complexes **2** and  
27 **3**. Two more isomers, **4-NP** and **4-CP**, are formed in low yield. Under an atmosphere of *p*-  
28 H<sub>2</sub>, all four isomers are detected in hyperpolarized form. Unlike **1**, complexes **3** and **4-CN**  
29 undergo thermal exchange at room temperature with *p*-H<sub>2</sub>.

### 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

## Experimental Section

Complex **1** was prepared using the old procedure reported in the literature.<sup>61</sup> Chemical  
manipulations and sample preparations were carried out under inert atmosphere conditions  
using standard Schlenk (vacuum 10<sup>-2</sup> mbar) or high vacuum (10<sup>-4</sup> mbar). *p*-H<sub>2</sub> was generated  
by cooling hydrogen gas over charcoal in a copper block at ca. 30 K. The proportion of *p*-H<sub>2</sub>  
at 30 K was calculated as > 99%. Pressures of *p*-H<sub>2</sub> were measured with an MKS Baratron  
capacitance manometer. Deuterated solvents for use in NMR were degassed and dried  
before use; benzene-*d*<sub>6</sub> and toluene-*d*<sub>8</sub> were obtained from Aldrich and dried on potassium  
mirrors in ampoules fitted with a greaseless tap. <sup>15</sup>N-pyridine (Aldrich) was used as received.

**NMR measurements. Ex-situ UV photolysis setup.** Photochemical reactions were  
performed with an Oriel 350 Watt high pressure xenon arc. Unwanted heating was  
minimized by use of a water filter of 10 cm length between the sample and the lamp.

1  
2  
3       **Continuous wave laser experiments.** The *in-situ* photolysis equipment has been  
4 described before.<sup>62</sup> In summary, *in-situ* photolysis experiments were performed using a  
5 wide-bore Bruker DRX-400 NMR spectrometer equipped with a 63 mW Kimmon IK series  
6 He-Cd laser at 325 nm. The laser beam is directed into the sample in the spectrometer  
7 through the air-gap between the magnet bore and a narrow-bore probe. The sample is  
8 cooled using a liquid N<sub>2</sub> boil-off evaporator. The laser was mounted on a specially designed  
9 platform situated in front of the NMR spectrometer that could be adjusted both horizontally  
10 and vertically. The laser alignment was optimized by using an NMR sample of  
11 Ru(CO)<sub>3</sub>(dppe) (dppe = Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>) in deuterated toluene, pressurized with *p*-H<sub>2</sub>.  
12 When exposed to UV light the sample reacts with *p*-H<sub>2</sub> to generate enhanced resonances  
13 arising from Ru(H)<sub>2</sub>(CO)<sub>3</sub>(dppe). Continuous recording of the spectra allowed the optimal  
14 position of the laser beam to be identified.  
15  
16

17       **Pulsed laser experiments.** All NMR spectra were recorded on a Bruker Avance II 600  
18 MHz spectrometer with a 14 T wide-bore magnet fitted with a 5 mm BBO probe. In situ laser  
19 photolysis was carried out with a pulsed Nd:YAG laser (Continuum Surelite II) fitted with a  
20 frequency tripling crystal (output 355 nm). Operating conditions were typically: 10 Hz  
21 repetition rate, flash lamp voltage 1.49 kV, and Q-switch delay increased from the standard  
22 to 320 μs yielding a laser power of 75 mW in internal mode. The energy of a single laser  
23 pulse was measured using an energy meter calibrated for 355 nm to be ~ 29.8 mJ at our  
24 operating conditions (external triggering with Q-switch delay set to 150 μs). The unfocused  
25 laser beam is directed at the base of the spectrometer and reflected up into the probe *via*  
26 a mirror as previously reported.<sup>32</sup> Adjustment screws control the vertical and horizontal  
27 position of the mirror which is on a kinematic mount. The system is fully shielded from the  
28 operator and the screws of the kinematic mount can be adjusted from outside the shield. The  
29 laser radiation is incident on a fixed mirror that is level with the sample and passes through a  
30 hole in the probe onto the NMR tube. Standard NMR tubes fitted with Young's taps were  
31 used. The samples contained 1–2 mg of compound (Abs<sub>355</sub> ~ 0.7) and approximately 0.4 mL  
32 of solvent. A sample of Ru(dppe)<sub>2</sub>H<sub>2</sub> in C<sub>6</sub>D<sub>6</sub> was used for laser alignment with *p*-H<sub>2</sub>  
33 amplification in real time. Standard NMR pulse sequences were modified for use with *p*-H<sub>2</sub>  
34 by including a synchronized laser initiation sequence prior to NMR excitation. A purpose-  
35 written program was used to control the laser firing from the NMR console with the laser set  
36 on external triggering. The program sets the laser to fire one warm-up shot before the fire  
37 signal. The NMR pulse is initiated at a set delay time (τ) following the fire signal. The intrinsic  
38 time delay between sending the fire signal from the spectrometer and the actual firing of the  
39 laser pulse was measured with a photodiode and an oscilloscope to be 150 μs (equal to the  
40 Q-switch delay for the generation of the pulse). This signal delay was incorporated into the  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 pulse sequence such that synchronized measurements with a time delay,  $\tau$ , were achieved  
4 by setting the spectrometer delay to:  $\tau + 150 \mu\text{s}$ . The precision of this delay between the  
5 laser and radio frequency (*r.f.*) pulses is controlled by the 200 ns clock of the spectrometer.  
6

7 **TRIR measurements.** Our time-resolved infrared (TRIR) measurements were based  
8 on pump-probe method that has been described in detail previously.<sup>30,63</sup> In brief, the probe  
9 source is a broadband mid-IR pulse with 100 fs duration (fwhm) and around 1  $\mu\text{J}$  energy at 1  
10 kHz. The pump source is the third harmonic output (355 nm) of a Q-switched Nd:YVO laser  
11 with 0.5 ps duration (fwhm) and around 3  $\mu\text{J}$  energy and is synchronized to the probe pulse.  
12 The delay between pump and probe pulses can be controlled with a pulse generator  
13 (DG535, Stanford Research System) from 0.5 ns to 100  $\mu\text{s}$ . The broadband transmitted  
14 probe pulse is detected with a 128-element HgCdTe array detector. The sample solutions  
15 were flowed through a  $\text{CaF}_2$  windowed IR cell, path length (0.25 mm), which was rastered in  
16 two dimensions at high speed to reduce overheating and degradation of the sample solution.  
17 All spectral and kinetic fitting procedures were carried out using Microcal Origin 7 software.  
18

19 **Low Temperature Photolysis.** The low temperature photolysis IR experiments were  
20 performed in a similar manner to those described previously.<sup>64</sup> The high pressure-low  
21 temperature (HP-LT) cell used in the low temperature experiments has been described in  
22 detail elsewhere.<sup>65</sup> Briefly, the HP-LT cell was filled with a solution of **1** dissolved in toluene-  
23  $d_8$ . Toluene- $d_8$  was used to minimize IR solvent background issues and to permit low  
24 temperature experiments in solution because of its lower freezing point than that of benzene-  
25  $d_6$ . The cell was attached to a cold finger of a cryogenic cooling system and cooled to the  
26 required temperature (220 K). The photolysis source used in these experiments was a  
27 Philips HPK 125W medium-pressure mercury arc lamp, which provided broadband UV and  
28 visible light. IR spectra were recorded using a Nicolet 730 interferometer linked to a  
29 computer running Omnic software.  
30

31 **IR spectroscopy.** Other IR spectra were recorded on a Mattson Unicam RS FTIR  
32 instrument, connected to a PC running WINFIRST software, or a Nicolet Nexus, connected  
33 to a PC running Omnic E.S.P. 5.2 software.  
34

35 **UV-Vis spectroscopy.** UV-Vis spectra were recorded on a Perkin-Lambda 7  
36 spectrometer, using a PC running PECSS software.  
37

38 **DFT calculations.** Density functional theory calculations were used to optimize  
39 geometries and simulate harmonic vibrational frequencies for several of the complexes (see  
40 Supporting information). Preliminary calculations of the parent species performed with the  
41 Stuttgart relativistic small core pseudopotential and basis set mixed with the 6-311G(d) basis  
42 set for non-metal atoms,<sup>66</sup> and using the B3LYP<sup>67,68</sup> PBE0,<sup>69</sup> and M06<sup>70</sup> functionals,  
43 representing 20, 25, and 27 % HF exchange, respectively, found that M06 gave the closest  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 match with experiment. Consequently, the M06 level of theory within the Q-Chem quantum-  
4 chemical software package was adopted for fuller calculations.<sup>71</sup> Calculations were  
5 converged then using the double- $\zeta$  6-31G(d) basis set for C, N, O, P, and H atoms, and the  
6 quasi-relativistic LANL2DZ pseudo-potential and basis set for Ru.<sup>72</sup> This combination has  
7 been labelled as LACVP(d). Optimized geometries were confirmed by the absence of  
8 imaginary frequencies, and these harmonic frequencies have been scaled by a uniform  
9 factor of 0.96 to account for anharmonic and experimental effects.<sup>73</sup>  
10  
11  
12

## 13 **ASSOCIATED CONTENT**

### 14 **Supporting Information**

15 The Supporting Information is available free of charge on the ACS Publications website at  
16 DOI: 10.1021/acs/organomet.xxxxx  
17

18 NMR spectra, pulse sequences used in laser pump-NMR probe experiments,  $T_{1(\text{min})}$   
19 relaxation, TRIR difference spectra, FTIR spectra, DFT calculated structures and CO-  
20 stretching frequencies, calculated and observed  $\nu(\text{CO})$  values for pyridine complexes.  
21  
22  
23  
24

## 25 **AUTHOR INFORMATION**

### 26 **Corresponding Authors**

27 \*E-mail for S.B.D.: [simon.duckett@york.ac.uk](mailto:simon.duckett@york.ac.uk)

28 \*E-mail for M.W.G.: [mike.george@nottingham.ac.uk](mailto:mike.george@nottingham.ac.uk)

29 \*E-mail for R.N.P.: [robin.perutz@york.ac.uk](mailto:robin.perutz@york.ac.uk)  
30

### 31 **ORCID**

32 Simon B. Duckett: 0000-0002-9788-6615

33 Michael W. George: 0000-0002-7844-1696

34 Robin N. Perutz: 0000-0001-6286-0282

35 Barbara Procacci: 0000-0001-7044-0560

36 Magnus Hanson-Heine 0000-0002-6709-297X

37 Raphael Horvath 0000-0002-2695-4895

38 Khuong Vuong 0000-0001-9558-9273  
39

### 40 **Notes**

41 The authors declare no competing financial interest.  
42  
43

## 44 **ACKNOWLEDGEMENTS**

45 We are grateful for financial support from EPSRC (grants EP/K022792/1 and EP/D058031).  
46 We acknowledge assistance from Dr. Pedro Aguiar. MWG gratefully acknowledges receipt  
47 of a Wolfson Merit Award.  
48  
49  
50  
51  
52  
53

## 54 **REFERENCES**

55  
56  
57  
58  
59  
60


- 1
- 2
- 3 (1) Crossley, S. W. M.; Obradors, C.; Martinez, R. M.; Shenvi, R. A. *Chem. Rev.* **2016**,
- 4 *116*, 8912-9000.
- 5
- 6 (2) Maity, A.; Teets, T. S. *Chem. Rev.* **2016**, *116*, 8873-911.
- 7
- 8 (3) Rossin, A.; Peruzzini, M. *Chem. Rev.* **2016**, *116*, 8848-72.
- 9
- 10 (4) Perutz, R. N.; Procacci, B. *Chem. Rev.* **2016**, *116*, 8506-44.
- 11
- 12 (5) Geoffroy, G. L.; Bradley, M. G. *Inorg. Chem.* **1977**, *16*, 744-8.
- 13
- 14 (6) Wakatsuki, Y.; Yamazaki, H.; Kumegawa, N.; Satoh, T.; Satoh, J. Y. *J. Am. Chem.*
- 15 *Soc.* **1991**, *113*, 9604-10.
- 16
- 17 (7) Wakatsuki, Y.; Yamazaki, H.; Maruyama, Y.; Shimizu, I. *J. Chem. Soc., Chem.*
- 18 *Commun.* **1991**, 261-3.
- 19
- 20 (8) Du, H. G.; Liu, Q.; Shi, S. J.; Zhang, S. W. *J. Organomet. Chem.* **2001**, *627*, 127-31.
- 21
- 22 (9) Samouei, H.; Grushin, V. V. *Organometallics* **2013**, *32*, 4440-3.
- 23
- 24 (10) Arockiam, P. B.; Bruneau, C.; Dixneuf, P. H. *Chem. Rev.* **2012**, *112*, 5879-918.
- 25
- 26 (11) Kakiuchi, F.; Murai, S. *Acc. Chem. Res.* **2002**, *35*, 826-34.
- 27
- 28 (12) Nixon, T. D.; Whittlesey, M. K.; Williams, J. M. J. *Dalton Trans.* **2009**, 753-62.
- 29
- 30 (13) Naota, T.; Takaya, H.; Murahashi, S. I. *Chem. Rev.* **1998**, *98*, 2599-660.
- 31
- 32 (14) Perutz, R. N.; Torres, O.; Vlček, A. J. *Photochemistry of Metal Carbonyls,*
- 33 *Comprehensive Inorganic Chemistry II Reedijk, J., Poeppelmeier, K., Eds.; Elsevier: Oxford,*
- 34 *U.K., 2013; Vol. 8, Ch. 8.06* **2013**, 229-53.
- 35
- 36 (15) Colombo, M.; George, M. W.; Moore, J. N.; Pattison, D. I.; Perutz, R. N.; Virrels, I. G.;
- 37 Ye, T. Q. *J. Chem. Soc., Dalton Trans.* **1997**, 2857-9.
- 38
- 39 (16) Torres, O.; Calladine, J. A.; Duckett, S. B.; George, M. W.; Perutz, R. N. *Chem. Sci.*
- 40 **2015**, *6*, 418-24.
- 41
- 42 (17) Calladine, J. A.; Torres, O.; Anstey, M.; Ball, G. E.; Bergman, R. G.; Curley, J.;
- 43 Duckett, S. B.; George, M. W.; Gilson, A. I.; Lawes, D. J.; Perutz, R. N.; Sun, X. Z.; Vollhardt,
- 44 K. P. C. *Chem. Sci.* **2010**, *1*, 622-30.
- 45
- 46 (18) Calladine, J. A.; Duckett, S. B.; George, M. W.; Matthews, S. L.; Perutz, R. N.;
- 47 Torres, O.; Khuong, Q. V. *J. Am. Chem. Soc.* **2011**, *133*, 2303-10.
- 48
- 49 (19) Godard, C.; Callaghan, P.; Cunningham, J. L.; Duckett, S. B.; Lohman, J. A. B.;
- 50 Perutz, R. N. *Chem. Commun.* **2002**, 2836-7.
- 51
- 52 (20) Blazina, D.; Dunne, J. P.; Aiken, S.; Duckett, S. B.; Elkington, C.; McGrady, J. E.;
- 53 Poli, R.; Walton, S. J.; Anwar, M. S.; Jones, J. A.; Carteret, H. A. *Dalton Trans.* **2006**, 2072-
- 54 80.
- 55
- 56 (21) Geftakis, S.; Ball, G. E. *J. Am. Chem. Soc.* **1998**, *120*, 9953-4.
- 57
- 58 (22) Lawes, D. J.; Geftakis, S.; Ball, G. E. *J. Am. Chem. Soc.* **2005**, *127*, 4134-5.
- 59
- 60 (23) Ball, G. E.; Brookes, C. M.; Cowan, A. J.; Darwish, T. A.; George, M. W.; Kawanami,
- H. K.; Portius, P.; Rourke, J. P. *Proc. Natl. Acad. Sci.* **2007**, *104*, 6927-32.



- 1  
2  
3 (24) Lawes, D. J.; Darwish, T. A.; Clark, T.; Harper, J. B.; Ball, G. E. *Angew. Chem., Int.*  
4 *Ed.* **2006**, *45*, 4486-90.  
5  
6 (25) Young, R. D.; Hill, A. F.; Hillier, W.; Ball, G. E. *J. Am. Chem. Soc.* **2011**, *133*, 13806-  
7 9.  
8  
9 (26) Ampt, K. A. M.; Burling, S.; Donald, S. M. A.; Douglas, S.; Duckett, S. B.; Macgregor,  
10 S. A.; Perutz, R. N.; Whittlesey, M. K. *J. Am. Chem. Soc.* **2006**, *128*, 7452-3.  
11  
12 (27) Montiel-Palma, V.; Perutz, R. N.; George, M. W.; Jina, O. S.; Sabo-Etienne, S.  
13 *Chem. Commun.* **2000**, 1175-6.  
14  
15 (28) Calladine, J. A.; Horvath, R.; Davies, A. J.; Wriglesworth, A.; Sun, X. Z.; George, M.  
16 *W. Appl. Spectrosc.* **2015**, *69*, 519-24.  
17  
18 (29) Greetham, G. M.; Burgos, P.; Cao, Q. A.; Clark, I. P.; Codd, P. S.; Farrow, R. C.;  
19 George, M. W.; Kogimtzis, M.; Matousek, P.; Parker, A. W.; Pollard, M. R.; Robinson, D. A.;  
20 Xin, Z. J.; Towrie, M. *Appl. Spectrosc.* **2010**, *64*, 1311-9.  
21  
22 (30) Towrie, M.; Grills, D. C.; Dyer, J.; Weinstein, J. A.; Matousek, P.; Barton, R.; Bailey,  
23 P. D.; Subramaniam, N.; Kwok, W. M.; Ma, C. S.; Phillips, D.; Parker, A. W.; George, M. W.  
24 *Appl. Spectrosc.* **2003**, *57*, 367-80.  
25  
26 (31) Banister, J. A.; George, M. W.; Grubert, S.; Howdle, S. M.; Jobling, M.; Johnson, F.  
27 P. A.; Morrison, S. L.; Poliakoff, M.; Schubert, U.; Westwell, J. R. *J. Organomet. Chem.*  
28 **1994**, *484*, 129-35.  
29  
30 (32) Torres, O.; Procacci, B.; Halse, M. E.; Adams, R. W.; Blazina, D.; Duckett, S. B.;  
31 Eguillor, B.; Green, R. A.; Perutz, R. N.; Williamson, D. C. *J. Am. Chem. Soc.* **2014**, *136*,  
32 10124-31.  
33  
34 (33) Pravdivtsev, A. N.; Yurkovskayaa, A. V.; Petrov, P. A.; Vieth, H. M. *Phys. Chem.*  
35 *Chem. Phys.* **2017**, *19*, 25961-69.  
36  
37 (34) Levison, J. J. *J. Chem. Soc. A* **1970**, 639-43.  
38  
39 (35) Perutz, R. N.; Sabo-Etienne, S. *Angew. Chem., Int. Ed.* **2007**, *46*, 2578-92.  
40  
41 (36) Luo, X. L.; Crabtree, R. H. *J. Am. Chem. Soc.* **1990**, *112*, 6912-8.  
42  
43 (37) Crabtree, R. H. *Chem. Rev.* **2016**, *116*, 8750-69.  
44  
45 (38) Gusev, D. G.; Vymenits, A. B.; Bakhmutov, V. I. *Inorg. Chim. Acta* **1991**, *179*, 195-  
46 201.  
47  
48 (39) Samouei, H.; Miloserdov, F. M.; Escudero-Adan, E. C.; Grushin, V. V.  
49 *Organometallics* **2014**, *33*, 7279-83.  
50  
51 (40) Riddlestone, I. M.; McKay, D.; Gutmann, M. J.; Macgregor, S. A.; Mahon, M. F.;  
52 Sparkes, H. A.; Whittlesey, M. K. *Organometallics* **2016**, *35*, 1301-12.  
53  
54 (41) Grellier, M.; Mason, S. A.; Albinati, A.; Capelli, S. C.; Rizzato, S.; Bijani, C.; Coppel,  
55 Y.; Sabo-Etienne, S. *Inorg. Chem.* **2013**, *52*, 7329-37.  
56  
57  
58  
59  
60

- 1  
2  
3 (42) Hebden, T. J.; Goldberg, K. I.; Heinekey, D. M.; Zhang, X. W.; Emge, T. J.; Goldman,  
4 A. S.; Krogh-Jespersen, K. *Inorg. Chem.* **2010**, *49*, 1733-42.  
5  
6 (43) Morris, R. H. *Coord. Chem. Rev.* **2008**, *252*, 2381-94.  
7  
8 (44) Sleigh, C. J.; Duckett, S. B.; Mawby, R. J.; Lowe, J. P. *Chem. Commun.* **1999**, 1223-  
9 4.  
10  
11 (45) The dimer observed by TRIR spectroscopy offers a potential route to formation of  
12  $\text{Ru}(\text{H})_2(\text{PPh}_3)_2(\text{CO})_2$  see section on IR studies.  
13  
14 (46) Ball, G. E.; Mann, B. E. *J. Chem. Soc., Chem. Commun.* **1992**, 561-3.  
15  
16 (47) Aguilar, J. A.; Elliott, P. I. P.; Lopez-Serrano, J.; Adams, R. W.; Duckett, S. B. *Chem.*  
17 *Commun.* **2007**, 1183-5.  
18  
19 (48) Abdur-Rashid, K.; Abbel, R.; Hadzovic, A.; Lough, A. J.; Morris, R. H. *Inorg. Chem.*  
20 **2005**, *44*, 2483-92.  
21  
22 (49) Halse, M. E.; Procacci, B.; Henshaw, S. L.; Perutz, R. N.; Duckett, S. B. *J. Magn.*  
23 *Reson.* **2017**, *278*, 25-38.  
24  
25 (50) Besora, M.; Carreon-Macedo, J. L.; Cimas, A.; Harvey, J. N. *Adv. Inorg. Chem., Vol*  
26 *61: Metal Ion Controlled Reactivity* **2009**, *61*, 573-623.  
27  
28 (51) Kosma, K.; Trushin, S. A.; Fuss, W.; Schmid, W. E.; Schneider, B. M. R. *Phys.*  
29 *Chem. Chem. Phys.* **2010**, *12*, 13197-214.  
30  
31 (52) Trushin, S. A.; Fuss, W.; Kompa, K. L.; Schmid, W. E. *J. Phys. Chem. A* **2000**, *104*,  
32 1997-2006.  
33  
34 (53) Paterson, M. J.; Hunt, P. A.; Robb, M. A.; Takahashi, O. *J. Phys. Chem. A* **2002**, *106*,  
35 10494-504.  
36  
37 (54) Portius, P.; Yang, J. X.; Sun, X. Z.; Grills, D. C.; Matousek, P.; Parker, A. W.; Towrie,  
38 M.; George, M. W. *J. Am. Chem. Soc.* **2004**, *126*, 10713-20.  
39  
40 (55) Burdett, J. K.; Grzybowski, J. M.; Perutz, R. N.; Poliakoff, M.; Turner, J. J.; Turner, R.  
41 *F. Inorg. Chem.* **1978**, *17*, 147-54.  
42  
43 (56) Soubra, C.; Oishi, Y.; Albright, T. A.; Fujimoto, H. *Inorg. Chem.* **2001**, *40*, 620-7.  
44  
45 (57) Rodger, A.; Johnson, B. F. G. *Inorg. Chem.* **1988**, *27*, 3061-2.  
46  
47 (58) Ruddlesden, A. J.; Mewis, R. E.; Green, G. G. R.; Whitwood, A. C.; Duckett, S. B.  
48 *Organometallics* **2015**, *34*, 2997-3006.  
49  
50 (59) Timney, J. A. *Inorg. Chem.* **1979**, *18*, 2502-6.  
51  
52 (60) Wilhelm, E.; Battino, R. *Chem. Rev.* **1973**, *73*, 1-9.  
53  
54 (61) Ahmad, N.; Uttley, M. F.; Robinson, S. D. *J. Chem. Soc., Dalton Trans.* **1972**, 843-7.  
55  
56 (62) Clark, J. L.; Duckett, S. B. *Dalton Trans.* **2014**, *43*, 1162-71.  
57  
58 (63) Brennan, P.; George, M. W.; Jina, O. S.; Long, C.; McKenna, J.; Pryce, M. T.; Sun,  
59 X. Z.; Vuong, K. Q. *Organometallics* **2008**, *27*, 3671-80.  
60

- 1  
2  
3 (64) Childs, G. I.; Colley, C. S.; Dyer, J.; Grills, D. C.; Sun, X. Z.; Yang, J. X.; George, M.  
4 *W. J. Chem. Soc., Dalton Trans.* **2000**, 1901-6.  
5  
6 (65) Cooper, A. I.; Poliakoff, M. *Chem. Phys. Lett.* **1993**, 212, 611-6.  
7  
8 (66) Dolg, M.; Wedig, U.; Stoll, H.; Preuss, H. *J. Chem. Phys.* **1987**, 86, 866-72.  
9  
10 (67) Becke, A. D. *J. Chem. Phys.* **1993**, 98, 5648-52.  
11  
12 (68) Stephens, P. J.; Devlin, F. J.; Chabalowski, C. F.; Frisch, M. J. *J. Phys. Chem.* **1994**,  
13 98, 11623-7.  
14  
15 (69) Adamo, C.; Barone, V. *J. Chem. Phys.* **1999**, 110, 6158-70.  
16  
17 (70) Zhao, Y.; Truhlar, D. G. *Theor. Chem. Acc.* **2008**, 120, 215-41.  
18  
19 (71) Shao, Y. H.; Gan, Z. T.; Epifanovsky, E.; Gilbert, A. T. B.; Wormit, M.; Kussmann, J.;  
20 Lange, A. W.; Behn, A.; Deng, J.; Feng, X. T.; Ghosh, D.; Goldey, M.; Horn, P. R.; Jacobson,  
21 L. D.; Kaliman, I.; Khaliullin, R. Z.; Kus, T.; Landau, A.; Liu, J.; Proynov, E. I.; Rhee, Y. M.;  
22 Richard, R. M.; Rohrdanz, M. A.; Steele, R. P.; Sundstrom, E. J.; Woodcock, H. L.;  
23 Zimmerman, P. M.; Zuev, D.; Albrecht, B.; Alguire, E.; Austin, B.; Beran, G. J. O.; Bernard,  
24 Y. A.; Berquist, E.; Brandhorst, K.; Bravaya, K. B.; Brown, S. T.; Casanova, D.; Chang, C.  
25 M.; Chen, Y. Q.; Chien, S. H.; Closser, K. D.; Crittenden, D. L.; Diedenhofen, M.; DiStasio,  
26 R. A.; Do, H.; Dutoi, A. D.; Edgar, R. G.; Fatehi, S.; Fusti-Molnar, L.; Ghysels, A.; Golubeva-  
27 Zadorozhnaya, A.; Gomes, J.; Hanson-Heine, M. W. D.; Harbach, P. H. P.; Hauser, A. W.;  
28 Hohenstein, E. G.; Holden, Z. C.; Jagau, T. C.; Ji, H. J.; Kaduk, B.; Khistyayev, K.; Kim, J.;  
29 Kim, J.; King, R. A.; Klunzinger, P.; Kosenkov, D.; Kowalczyk, T.; Krauter, C. M.; Lao, K. U.;  
30 Laurent, A. D.; Lawler, K. V.; Levchenko, S. V.; Lin, C. Y.; Liu, F.; Livshits, E.; Lochan, R. C.;  
31 Luenser, A.; Manohar, P.; Manzer, S. F.; Mao, S. P.; Mardirossian, N.; Marenich, A. V.;  
32 Maurer, S. A.; Mayhall, N. J.; Neuscamman, E.; Oana, C. M.; Olivares-Amaya, R.; O'Neill, D.  
33 P.; Parkhill, J. A.; Perrine, T. M.; Peverati, R.; Prociuk, A.; Rehn, D. R.; Rosta, E.; Russ, N.  
34 J.; Sharada, S. M.; Sharma, S.; Small, D. W.; Sodt, A. *Mol. Phys.* **2015**, 113, 184-215.  
35  
36 (72) Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, 82, 299-310.  
37  
38 (73) Merrick, J. P.; Moran, D.; Radom, L. *J. Phys. Chem. A* **2007**, 111, 11683-700.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



graphical abstract

38x17mm (300 x 300 DPI)