



UNIVERSITY OF LEEDS

This is a repository copy of *Reduced inter-hemispheric interference in ageing: Evidence from a divided field Stroop-like paradigm*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/126929/>

Version: Accepted Version

---

**Proceedings Paper:**

Castronovo, J and Delvenne, J-F [orcid.org/0000-0002-1990-9738](https://orcid.org/0000-0002-1990-9738) (2017) Reduced inter-hemispheric interference in ageing: Evidence from a divided field Stroop-like paradigm. In: Journal of Vision. Vision Sciences Society Seventeenth Annual Meeting, 19-24 May 2017, St. Pete Beach, Fla., USA. Association for Research in Vision and Ophthalmology .

<https://doi.org/10.1167/17.10.449>

---

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License [<https://creativecommons.org/licenses/by-nc-nd/4.0/>].

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Reduced inter-hemispheric interference in ageing: Evidence from a divided field Stroop-like paradigm

Jean-François Delvenne

University of Leeds

Julie Castronovo

University of Hull

## Abstract

One of the most important structural changes that occur in the brain during the course of life relates to the corpus callosum, the largest neural pathway that connects the two cerebral hemispheres. Whether the corpus callosum has an inhibitory or excitatory function in interhemispheric connection is still debated in the literature. It has been shown that the corpus callosum, and in particular its anterior sections, endures a process of degeneration in ageing. Hence, a primary question is whether such structural changes in the brain of older adults have functional consequences on inter-hemispheric communication. To investigate this question, we asked healthy older adults to perform a modified version of the classic Stroop paradigm in which the target and distracter were spatially separated. Across two experiments, we found that the Stroop effect was significantly reduced when the two stimuli were distributed in two different hemifields as opposed to the same single hemifield. This new finding suggests that age-related callosal thinning reduces inter-hemispheric interference by facilitating the two hemispheres to process information in parallel. The reduced inter-hemispheric interference found in older adults provides strong direct support for the excitatory theory of the corpus callosum.

Meeting abstract presented at VSS 2017