



This is a repository copy of *How primary care can contribute to good mental health in adults*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/129027/>

Version: Published Version

---

**Article:**

Gupta, S., Jenkins, R., Spicer, J. et al. (16 more authors) (2018) How primary care can contribute to good mental health in adults. *London Journal of Primary Care*, 10 (1). pp. 3-7. ISSN 1757-1472

<https://doi.org/10.1080/17571472.2017.1410043>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



## How primary care can contribute to good mental health in adults

Sunjai Gupta, Rachel Jenkins, John Spicer, Marina Marks, Nigel Mathers, Lise Hertel, Laura Calamos Nasir, Fiona Wright, Baljeet Ruprah-Shah, Brian Fisher, David Morris, Kurt C. Stange, Robert White, Gina Giotaki, Tony Burch, Catherine Millington-Sanders, Steve Thomas, Ricky Banarsee & Paul Thomas

To cite this article: Sunjai Gupta, Rachel Jenkins, John Spicer, Marina Marks, Nigel Mathers, Lise Hertel, Laura Calamos Nasir, Fiona Wright, Baljeet Ruprah-Shah, Brian Fisher, David Morris, Kurt C. Stange, Robert White, Gina Giotaki, Tony Burch, Catherine Millington-Sanders, Steve Thomas, Ricky Banarsee & Paul Thomas (2018) How primary care can contribute to good mental health in adults, *London Journal of Primary Care*, 10:1, 3-7, DOI: [10.1080/17571472.2017.1410043](https://doi.org/10.1080/17571472.2017.1410043)

To link to this article: <https://doi.org/10.1080/17571472.2017.1410043>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 13 Dec 2017.



Submit your article to this journal [↗](#)



Article views: 231



View related articles [↗](#)



View Crossmark data [↗](#)

## How primary care can contribute to good mental health in adults\*

Sunjai Gupta<sup>a,b</sup>, Rachel Jenkins<sup>c</sup>, John Spicer<sup>d</sup>, Marina Marks<sup>e</sup>, Nigel Mathers<sup>f</sup>, Lise Hertel<sup>g</sup>, Laura Calamos Nasir<sup>h</sup>, Fiona Wright<sup>i</sup>, Baljeet Ruprah-Shah<sup>j</sup>, Brian Fisher<sup>k</sup>, David Morris<sup>l</sup>, Kurt C. Stange<sup>m</sup>, Robert White<sup>n</sup>, Gina Giotaki<sup>o</sup>, Tony Burch<sup>p</sup>, Catherine Millington-Sanders<sup>q</sup>, Steve Thomas<sup>r</sup>, Ricky Banarsee<sup>s</sup> and Paul Thomas<sup>t</sup>

<sup>a</sup>Department of Health Services and Population Research, Institute of Psychology, Psychiatry and Neuroscience, Kings College, London, UK; <sup>b</sup>Honorary Consultant Psychiatrist, South London and Maudsley NHS Trust, London, UK; <sup>c</sup>Epidemiology and Mental Health Policy, Kings College London, ETHICS, London, UK; <sup>d</sup>Primary Care Education, Health Education England, London, UK; <sup>e</sup>Educational Trust for Health Improvement through Cognitive Strategies (ETHICS), London, UK; <sup>f</sup>Primary Medical Care, University of Sheffield, Sheffield, UK; <sup>g</sup>Integrated Care University College London Hospitals NHS Foundation Trust, London, UK; <sup>h</sup>School of Nursing, University of North Carolina at Chapel Hill, Chapel Hill, USA; <sup>i</sup>Public Health Barking and Dagenham Council and Greater London Authority, London, UK; <sup>j</sup>Accomplish Consultancy, London, UK; <sup>k</sup>Health Empowerment Leverage Project, New NHS Alliance, London, UK; <sup>l</sup>Mental Health, Inclusion and Community, University of Central Lancashire, Preston, UK; <sup>m</sup>Family Medicine and Community Health, Epidemiology and Biostatistics, Oncology and Sociology, Center for Community Health Integration, Case Western Reserve University, Cleveland, OH, USA; <sup>n</sup>Primary Care Mental Health Services, West London Mental Health Trust, London, UK; <sup>o</sup>Dance Practices, Liverpool John Moores University, Liverpool, UK; <sup>p</sup>Health Education England North London, London, UK; <sup>q</sup>Marie Curie National Clinical End of Life Care Champion, London, UK; <sup>r</sup>Mental Health & Learning Disabilities NHS Sheffield Clinical Commissioning Group, Sheffield, UK; <sup>s</sup>Public Health England, London, UK; <sup>t</sup>London Journal of Primary Care, London, UK

### ABSTRACT

The need for support for good mental health is enormous. General support for good mental health is needed for 100% of the population, and at all stages of life, from early childhood to end of life. Focused support is needed for the 17.6% of adults who have a mental disorder at any time, including those who also have a mental health problem amongst the 30% who report having a long-term condition of some kind. All sectors of society and all parts of the NHS need to play their part. Primary care cannot do this on its own. This paper describes how primary care practitioners can help stimulate such a grand alliance for health, by operating at four different levels – as individual practitioners, as organisations, as geographic clusters of organisations and as policy-makers.

### KEYWORDS

Mental health; primary care; collaboration; Accountable care organisations

### Why this matters to us

London Journal of Primary Care (LJPC) publishes articles on the multi-dimensional aspects of primary care that make it so human and vibrant. It is a network of people who want to develop holistic, community-oriented integrated care and health promotion as a force for whole-society health. We hope that this paper will help general practice and more broadly primary care, to take a strong role in developing this.

### Key message

Primary care can have an enhanced effect on the good mental health of the population by collaborating with others within local communities for health.

In 2015, LJPC contributed to a Think Tank, in partnership with the mental health charity ETHICS and the Royal College of General Practitioners (RCGP) to identify things that primary care can do to promote good mental health throughout the population [1]. This translated into national RCGP policy; an LJPC paper describes its 12 recommendations for action (Figure 1) [2].

In this paper, we describe things that General Practices in the UK National Health (NHS) can do to translate these

recommendations into practice. Different practices will want to do different things because of different health needs in different places, different skills of practice staff and different local services and resources. So the action areas are described as general aims. We invite you to apply them in ways that are relevant to your specific context.

The need is enormous. General support for good mental health is needed for 100% of the population, and at all stages of life, from early childhood to end of life. Focused

**CONTACT** Sunjai Gupta  [Sunjai.Gupta@kcl.ac.uk](mailto:Sunjai.Gupta@kcl.ac.uk)

\*Promoting Mental Health and Preventing Mental Illness in General Practice. Steve Thomas, Rachel Jenkins, Tony Burch, Laura Calamos Nasir, Brian Fisher, Gina Giotaki, Shamini Gnani, Lise Hertel, Marina Marks, Nigel Mathers, Catherine Millington-Sanders, David Morris, Baljeet Ruprah-Shah, Kurt Stange, Paul Thomas, Robert White & Fiona Wright. London Journal of Primary Care Vol. 8, Issue 1, 2016

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

- (1) Mental health promotion and prevention are too important to wait.
- (2) Work with your community to map risk factors, resources and assets.
- (3) Good health care, medicine and best practice are biopsychosocial rather than purely physical.
- (4) Integrate mental health promotion and prevention into your daily work.
- (5) Boost resilience in your community.
- (6) Identify people at increased risk of mental disorder for support and screening.
- (7) Support early intervention for people of all ages with signs or symptoms of illness—early recognition and intervention is crucial.
- (8) Maintain your biopsychosocial skills.
- (9) Ensure good communication, interdisciplinary team working and inter-sectoral working with other staff, teams and agencies.
- (10) Lead by example, taking action to promote the resilience of the general practice workforce.
- (11) Ensure mental health is appropriately included in the strategic agenda for your cluster, at the Clinical Commissioning Groups, and the Health and Wellbeing Board.
- (12) Be aware of national mental health strategies and localise them, including action to destigmatise mental illness.

**Figure 1.** RCGP Recommendations as set out in Thomas et al. [2].

support is needed for the 17.6% of adults who have a mental disorder at any time [1], including those who also have a mental health problem amongst the 30% who report having a long-term condition of some kind [3]. All sectors of society and all parts of the NHS need to play their part. Primary care cannot do this on its own.

Whatever the diagnosis and context, good mental health of adults is helped by good bio-psycho-social habits – good habits for physical fitness, for emotional centredness and for participation in initiatives that develop communities.

Making a significant improvement to the mental health of all citizens requires inter-sectoral collaboration at all levels, national, regional, district and local, in order to promote mental health. If we want to improve the mental health of the whole population, we need to develop a grand alliance for health and care. General practice, and more broadly primary care, is only one contributor, but well placed to have a coordinating role. Others are public health, mental health and social care services, voluntary care, schools, businesses and pretty much everyone else. To have effects which are more than the sum of the parts, these contributions need to be integrated, or at least aligned, in ways that help create and sustain environments that promote good mental health.

Aligning multiple efforts for good mental health is helped by a shared geographical locality. Working from a specific organisation (e.g. a general practice) helps those who belong to that organisation but it does not easily enable complex collaborations that have mutually enhancing effects. Recognition of the need for local collaboration for health dates from the 1978 World Health Organisation (WHO) international conference at Alma Ata [4], and reaffirmed in 2008 when the WHO Director-General explained why such localism is ‘more important than ever’ [5]. In the 2008 document, WHO specifically advocated community-based *hubs of coordination* to achieve comprehensive collaboration for health and care. More recently, the WHO’s Mental Health Action Plan 2013–2020 [6] aims to ‘implement strategies for promotion and prevention in mental

1. As a member of their community
2. **Primary care as an individual practitioner**
3. **Practice level action**
4. **Primary care clusters or localities**
5. **Clinical Commissioning Groups; Sustainability & Transformation Partnerships; Accountable Care Organisations**

**Figure 2.** Levels at which primary care can take action [1,9,10].

health...’ and the ‘development of comprehensive community-based mental health and social care services’.

In England, the WHO vision for community-based coordinating hubs is being translated into policy through ‘New Care Models’ [7] (which bring together professionals to work collaboratively including GP, hospital, community and mental health services), and the ‘Primary Care Home’ [8] which aims to improve personalised and preventative care for the local population. The Primary Care Home provides care to a geographic locality of between 30,000 and 50,000. This is small enough to feel part of and large enough to have a political impact. In the horizontal direction, those involved develop a local community for health. In the vertical direction, they coordinate large numbers of care pathways. In future, there will be a shift in some activities from the Clinical Commissioning Group (CCG) to the Accountable Care Organisation (ACO), though CCGs will continue to be responsible and accountable for the delivery of their functions [9].

It is to leaders of community-based coordinating hubs that we target this paper. Figure 2 describes five different levels at which primary care practitioners can make a difference – as a citizen, as an individual practitioner, through the practice, through localities and through CCGs, Sustainability and Transformation Partnerships and ACOs. The first we leave to you to think about – how you personally contribute to your local community is your concern. We target this paper at the other four levels. We hope that this paper stimulates local discussion about how to make a collective impact that is more than any individual practitioner or discipline can achieve on its own.

## The case for action

- Each year, 70 million sick days are lost to mental disorders which are the leading cause of sickness absence in the UK [1]
- Only 43% of all those with mental health problems are in employment, in contrast to 74% of the general population and up to 65% of those with other health problems [11]
- Forty-four per cent of employment and support allowance benefit claimants report having a mental disorder as their primary diagnosis [1]
- It has been estimated that mental health-related job loss costs employers £82 million per year [12]

- Most people who commit suicide have seen their GP in the recent past [13]
- The life expectancy of adults with severe mental illness lags behind by an average of 10–20 years [14] and those with major depression and schizophrenia have a 40–60% elevated risk of dying prematurely [6]
- Medically unexplained symptoms are involved in 15–30% of all primary care consultations [14]
- About 1% of adults have autistic spectrum disorder but most are undetected [12]

These present a compelling case for primary care to contribute to a broad local collaboration to maintain good mental health of adults in the local population.

### General practice can

- (1) Use initiatives like New Care Models and Primary Care Home to work with local general practices, public health, social care, mental health services, voluntary groups and others to *align plans for positive mental health* in the locality. This should encourage self-care, including use of digital technologies [15,16], and promote the 'Five ways to mental wellbeing' [17] (Connect, Be Active, Take Notice, Keep Learning, Give).
- (2) Work in clusters of practices and their extended primary care teams to integrate mental health care into routine care, and plan with mental health and public health practitioners, social workers and voluntary groups. This makes it easier to persuade a range of other organisations and disciplines to contribute to an integrated strategy for good mental health. Shared planning in this way can improve:
  - Training for clinicians to be skilled at discerning 'teachable moments' when patients are ready to learn and change
  - Strategically planned access to psychological therapies
  - Employers taking responsibility for employees who have mental health problems
  - Inter-organisational collaboration that builds local communities for health [18]
- (3) Write *live manuals*, tailored to local needs, to inform practitioner decisions, self-help and care pathways [19]. It is easy to forget simple things that can promote good mental health and improve mental illness. Web-based 'live manuals' that are regularly updated by multidisciplinary leadership teams and used every day

by practitioners help avoid this problem and empower good practice when they:

- Describe things that practitioners can do during consultations, including opportunistically assessing mental health of patients and using an empowering consulting style to discern 'teachable moments' when patients are open to learning and change
- Signpost self-help options that help people know how common it is to feel anxious and depressed and how to improve their skills at reducing anxiety and avoiding projecting negative feelings onto others
- Show how to refer into care pathways, supported by in-the-moment specialist advice and a facility to fast-track to specialist care those who need it

- (4) Develop *multidisciplinary teams*, for the care of mentally unwell people, whose members are skilled at integrating mental health care into their everyday care [20,21].

Care plans for people with mental illness are a well-established and effective way to share care. Care planning teams should include more than mental health practitioners and include primary care clinicians, specialists, carers and many others. Each of these could contribute with links to resources that help develop a sense of positive connection with self and others, and enhance physical fitness, emotional centred-ness and participation in initiatives that develop social networks. Such resources could include participatory forms of art that build communities and enhance health and well-being of individuals and groups. Together, care planning teams could:

- Identify relapse at an early stage
- Share records and agree 'Special Patient Notes' to guide the actions of others who have occasional contact with patients (e.g. Out of Hours Services)
- Promote incremental improvements in lifestyle, well-being and relationships as the opportunities arise

- (5) Highlight *wider determinants of mental illness*

Anxiety and depression commonly have a range of contributing factors including relationships difficulties, unresolved old conflict, grief, debt, alcohol misuse, lack of physical fitness and loss of purpose. These same factors worsen all mental illnesses. Highlighting them can help patients address them. Primary care practitioners can:

- Encourage patients to focus on controlling things they have the power to control and be less anxious about things they cannot control

- Refer into services for debt management, housing improvement, lifestyle improvement, domestic violence, anger management, alcohol misuse; and argue the need for continued funding of such services
- Support initiatives that build resilience, including helping people develop networks of trusted relationships which enable the community become more engaged and which build social capital

(6) Consider mental health needs of those who have *long-term conditions*

In England, 15.4 million people suffer from a long-term condition [3]. Many of those who have a long-term condition of one kind or another have Care Plans that include patient goals for self-care. The care planning process could highlight mental health issues by:

- Routinely asking about anxiety and depression
- Providing literature and courses to help people realise that it is common to have negative emotions to long-term illnesses and these can often be addressed through simple actions
- Reminding that medically unexplained symptoms and everyday complaints like backache and headache often have associated psychosocial problems that can be improved with purposeful actions

(7) *Signpost ways for people to self-care and make useful contributions to society*

Good mental health includes a positive sense of personal identity, including a coherent and positive sense of their life story in which the person is able to rise above the difficulties they encounter. Mental illness disturbs this coherence through internal things like anxiety and fear, and external things like weak relationships and contracted life options. A partnership of organisations in a locality can promote ways to reduce inner turbulence and increase external resilience. They can:

- Encourage everyone to make useful contributions to society, including paid and voluntary work that helps strengthen the local community, appreciate those around them and increase their webs of trusted relationships
- Encourage everyone to be skilled at being personally centred and able to listen positively to others, including things like mindfulness, yoga, meditation, music and gardening
- Encourage *healthy lifestyles* for everyone, including physical fitness, sport, dancing and weight control

Some of these actions that primary care can undertake are summarised in Figure 3.

- *Align plans for positive mental health* and promote the 'five ways to mental wellbeing'
- Write *live manuals* to inform practitioner decisions, and care pathways
- Develop *multidisciplinary teams* for the care of mentally unwell people
- Highlight *wider determinants of mental illness*
- Consider mental health needs of those who have *long-term conditions*
- *Signpost ways for people to self-care and make useful contributions to society*

**Figure 3.** Examples of actions that primary care can undertake to improve mental health.

### Please contribute to LJPC discussions about mental health promotion in primary care

The 2014 Commonwealth Fund Report reveals that the UK enjoys some of the best health care in the world. When compared with 10 other developed countries, it ranked first overall, scored highest for quality, access and efficiency; and second equal for equity (along with Switzerland and behind Sweden). This is despite having the second lowest per capita spend (New Zealand was lower). However, in the category of '*healthy lives*', it scored 10th – 2nd only to the USA for poor outcomes [22]. The category of '*healthy lives*' includes: (1) mortality amenable to medical care, (2) infant mortality and (3) healthy life expectancy at age 60. These are all things that require many different organisations to align their efforts to promote health – public health, local authorities, schools, community groups, families and many others. We need to develop and evaluate models of complex collaborations for whole society health, including mental health.

Primary and community care practitioners don't need the Commonwealth Fund to see the need for integrated working between different agencies for good mental health. Every time we see a patient who is anxious, or depressed, or lonely, or has difficult life circumstances, we recognise what's needed – families, friends, neighbours and other social support; getting meals on tables, keeping active, getting out of doors. It is obvious, surely, that sometimes people need a variety of help to do these things, beyond specific therapies, and to take part in activities that put a smile on a face – music, dance, photos, chats, gardening....

LJPC intends to continue this debate about locality-based collaboration for positive mental health. We want to do this through a network of collaborating sites that support the evaluation of case studies of community-oriented integrated care. Their published papers (in LJPC and elsewhere) can fuel debate, including social media to gain deeper understanding of how modern-day primary care can support mentally healthy societies.

We want to publish evaluations of things like social prescribing and primary care navigators, Primary Care Homes and Vanguard Sites and other models that build local communities for health. This paper describes things that primary care can do to improve the mental health of adults. In time we hope to publish similar papers focused

on children, parents and families, retirement, end of life care and refugees.

Please get involved.

## Governance

LJPC Board overviewed this work.

## Acknowledgements

We wish to thank the ETHICS Foundation for funding the Think Tank that led to this work, and Dr Victoria Tzortziou Brown, Dr John Sanfey, Dr Tony Coggins and Dr Shekhar Saxena for their assistance and advice.

## References

- [1] ETHICS Initiative for Integration of Mental Health Promotion into Primary Care. In collaboration with the London Journal of Primary Care and the Ionian University of Corfu The ETHICS/LJPC/RCGP Think Tank. Available from: <https://ethicscharity.files.wordpress.com/2015/09/ethics-ljpc-thinktank-9-9-15-v5.pdf>
- [2] Thomas S, Jenkins R, Burch T, et al. Promoting mental health and preventing mental illness in general practice. *London J Prim Care*. 2016;8:3–9.
- [3] Investing in emotional and psychological wellbeing for patients with long-term conditions. Mental Health Network. NHS CONFEDERATION. Available from: <http://www.nhsconfed.org/~media/Confederation/Files/Publications/Documents/Investing%20in%20emotional%20and%20psychological%20wellbeing%20for%20patients%20with%20long-Term%20conditions%2016%20April%20final%20for%20website.PDF>
- [4] McDonald JJ. Primary health care: medicine in its place. Abingdon: Earthscan; 1992.
- [5] World Health Organisation. The World Health Report 2008- primary health care (now more than ever). Available from: [www.who.int/whr/2008/](http://www.who.int/whr/2008/)
- [6] World Health Organization. Mental health action plan, 2013–2020. Geneva; 2013. Available from: [http://apps.who.int/iris/bitstream/10665/89966/1/9789241506021\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/89966/1/9789241506021_eng.pdf?ua=1)
- [7] New Care Models: Vanguard – developing a blueprint for the future of NHS and care services. September 2016; Available from: [https://www.england.nhs.uk/wp-content/uploads/2015/11/new\\_care\\_models.pdf](https://www.england.nhs.uk/wp-content/uploads/2015/11/new_care_models.pdf)
- [8] NHS England. Primary care home model. Available from: <https://www.england.nhs.uk/ourwork/new-care-models/pchamm/>
- [9] NHS England. Local partnerships to improve health and care. Sustainability and Transformation Partnerships (STPs) and accountable care systems (ACSS). Available from: <https://www.england.nhs.uk/stps/>
- [10] ACOs and the NHS commissioning system. Accountable Care Organisation (ACO) Contract package - supporting document; 2017 Aug. Available from: [https://www.england.nhs.uk/wp-content/uploads/2016/12/1693\\_DraftMCP-6\\_A.pdf](https://www.england.nhs.uk/wp-content/uploads/2016/12/1693_DraftMCP-6_A.pdf)
- [11] The five year forward view for mental health. A report from the independent Mental Health Taskforce to the NHS in England; 2016 Feb. Available from: <https://www.england.nhs.uk/wp-content/uploads/2016/02/Mental-Health-Taskforce-FYFV-final.pdf>
- [12] Background briefing prepared for the ETHICS /RCGP/LJPC Think Tank on integration of mental health promotion into primary care and multi-sectoral community services; 2015 Apr 20–24. Available from: [www.ethicsfoundation.org](http://www.ethicsfoundation.org)
- [13] Cultivating Mental Health Promotion and Prevention in General Practice. A collaborative report from Educational Trust for Health Improvement through Cognitive Strategies London Journal of Primary Care Royal College of General Practitioners. Available from: [https://ethicscharity.files.wordpress.com/2015/09/rcgp\\_keymsg\\_150925\\_v5.pdf](https://ethicscharity.files.wordpress.com/2015/09/rcgp_keymsg_150925_v5.pdf)
- [14] Naylor C, Das P, Ross S, et al. Bringing together physical and mental health A new frontier for integrated care. Kings Fund; 2016 Mar. Available from: [https://www.kingsfund.org.uk/sites/default/files/field/field\\_publication\\_file/Bringing-together-Kings-Fund-March-2016\\_1.pdf](https://www.kingsfund.org.uk/sites/default/files/field/field_publication_file/Bringing-together-Kings-Fund-March-2016_1.pdf)
- [15] The Sheffield Mental Health Guide. Available from: <https://www.sheffieldmentalhealth.org.uk/>
- [16] Sheffield Flourish. Available from: <https://sheffieldflourish.co.uk>
- [17] Five ways to wellbeing. The New Economics Foundation; 2008. Available from: [http://neweconomics.org/search/?\\_sft\\_project=five-ways-to-wellbeing](http://neweconomics.org/search/?_sft_project=five-ways-to-wellbeing)
- [18] Butler F, Wheeler J. New Models of Primary Care Mental Health – learning across London; 2017 Jun. Available from: [https://www.kingsfund.org.uk/sites/default/files/media/Fiona\\_Butler\\_Jane\\_Wheeler.pdf](https://www.kingsfund.org.uk/sites/default/files/media/Fiona_Butler_Jane_Wheeler.pdf)
- [19] Thomas P. Collaborating for Health. Abingdon: Routledge; 2017.
- [20] Cohen DJ, Balasubramanian BA, Davis M, et al. Understanding care integration from the ground up: five organizing constructs that shape integrated practices. *J Am Board Fam Med*. 2015;28(Suppl 1):S7–S20.
- [21] Davis M, Balasubramanian BA, Waller E, et al. Integrating behavioral and physical health care in the real world: early lessons from advancing care together. *J Am Board Fam Med*. 2013;26:588–602.
- [22] Davis K, Stremikis K, Squires D, et al. Mirror, mirror on the wall. How the U.S. Health Care System Compares Internationally. The Commonwealth Fund; [updated 2014]. Available from: [http://www.commonwealthfund.org/~media/files/publications/fund-report/2014/jun/1755\\_davis\\_mirror\\_mirror\\_2014\\_exec\\_summ.pdf](http://www.commonwealthfund.org/~media/files/publications/fund-report/2014/jun/1755_davis_mirror_mirror_2014_exec_summ.pdf)