

This is a repository copy of *Arbuscular Mycorrhizal Fungi and Plant Chemical Defence: Effects of Colonisation on Aboveground and Belowground Metabolomes*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/129480/>

Version: Accepted Version

Article:

Hill, Elizabeth M, Robinson, Lynne A., Abdul-Sada, Ali et al. (3 more authors) (2018) *Arbuscular Mycorrhizal Fungi and Plant Chemical Defence: Effects of Colonisation on Aboveground and Belowground Metabolomes*. *Journal of chemical ecology*. pp. 199-208. ISSN 0098-0331

<https://doi.org/10.1007/s10886-017-0921-1>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Journal of Chemical Ecology

ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND BELOWGROUND METABOLOMES

--Manuscript Draft--

Manuscript Number:	JOCE-D-17-00277R
Full Title:	ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND BELOWGROUND METABOLOMES
Article Type:	Original Research
Keywords:	Arbuscular mycorrhizal symbiosis, Above-belowground interactions, Metabolomics, Pyrrolizidine alkaloids, Blumenols, Senecio jacobaea, Rhizophagus irregularis.
Corresponding Author:	Sue Hartley University of York UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of York
Corresponding Author's Secondary Institution:	
First Author:	Elizabeth Hill
First Author Secondary Information:	
Order of Authors:	Elizabeth Hill Lynne Robinson Ali Abdul-Sada Adam Vanbergen Angela Hodge Sue Hartley
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	<p>Arbuscular mycorrhizal fungal (AMF) colonisation of plant roots is one of the most ancient and widespread interactions in ecology, yet the systemic consequences for plant secondary chemistry remain unclear. We performed the first metabolomic investigation into the impact of AMF colonisation, by <i>Rhizophagus irregularis</i> on the chemical defences, spanning above- and below-ground tissues, in its host-plant ragwort (<i>Senecio jacobaea</i>). We used a non-targeted metabolomics approach to profile, and where possible identify, compounds induced by AMF colonisation in both roots and shoots. Metabolomics analyses revealed that 33 compounds were significantly increased in the root tissue of AMF-colonised plants, including seven blumenols, plant-derived compounds known to be associated with AMF colonisation. One of these was a novel structure conjugated with a malonyl-sugar and uronic acid moiety, hitherto an unreported combination. Such structural modifications of blumenols could be significant for their previously reported functional roles associated with the establishment and maintenance of AM colonisation. Pyrrolizidine alkaloids (PAs), key anti-herbivore defence compounds in ragwort, dominated the metabolomic profiles of root and shoot extracts. Analyses of the metabolomic profiles revealed an increase in four PAs in roots (but not shoots) of AMF colonised plants, with the potential to protect colonised plants from below-ground organisms.</p>

Dear Prof Hartley:

Your manuscript, JOCE-D-17-00277, "ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND BELOWGROUND METABOLOMES", that you submitted to the JOURNAL OF CHEMICAL ECOLOGY has been reviewed by the handling editor. Some minor edits are needed before final acceptance.

Please notify all co-authors of the decision.

To submit your revised manuscript, please log onto the following:

Your username is: SHartley-928

If you forgot your password, you can click the 'Send Login Details' link on the EM Login page at <http://joce.edmgr.com/>.

Please click "Author Login" to get into the site. Be sure to delete the old version of your paper before uploading the revised version.

Please also be sure to upload a letter in which you explain in detail how you have responded to the reviewers' concerns.

Please find specific formatting instructions in the Author Instructions at:

http://www.springer.com/west/home/life+sci?SGWID=4-10027-70-356744610&detailsPage=contentItemPage&contentItemId=143542&CIPageCounter=CI_FOR_AUTHORS_AND_EDITORS_PAGE0&SHORTCUT=www.springer.com/sgw/cda/frontpage/0,11855,4-10027-70-35674461-0,00.html

(they can also be found below the reviewer comments at the end of this email). Please read these over before you prepare the final manuscript and follow them precisely. Papers are not accepted by the Journal until they have been edited after revision. Papers requiring major revisions are subject to additional review. If you anticipate that it will take longer than 2 months to complete the revision, please let us know.

Please make sure to submit your editable source files (i. e. Word, TeX).

Thank you for submitting your work to the JOURNAL OF CHEMICAL ECOLOGY for appraisal. If you have questions or concerns, please do not hesitate to email me.

Sincerely,

Nicole M. van Dam
Associate Editor
Journal of Chemical Ecology

Comments for the author:

The changes made this a great contribution to J Chem Ecology. It was a pleasure to read the revised MS. Thanks for taking the effort!

There is one minor detail that you need to revise: could you please align the floating points of all the values in the tables?

All the points have been aligned in the tables.

Furthermore I think the word: Plants in line 309 should be removed,

Removed

and in the legend of figure 1, it should be :Open and closed circles (move the word circles).

Corrected.

The title "Table 3" (was 4) still contains a correction marking. Can you please make it final?

Corrected.

Thanks again for submitting your work to JCE

[Click here to view linked References](#)

1 1 ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL
2
3
4 2 DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND
5
6
7 3 BELOWGROUND METABOLOMES
8
9

10 4
11
12
13 5 ELIZABETH M. HILL¹, LYNNE A. ROBINSON^{1,2}, ALI ABDUL-SADA¹,
14
15
16 6 ADAM J. VANBERGEN², ANGELA HODGE³ AND SUE E. HARTLEY^{3,4*}
17
18

19 7
20 8 ¹School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK.
21

22 9 ²Centre for Ecology and Hydrology, Bush Estate, Penicuik, Midlothian EH26 0QB, UK.
23
24

25 10 ³Department of Biology, University of York, Wentworth Way, York YO10 5DD, UK.
26

27 11 ⁴York Environment and Sustainability Institute, Department of Biology, University of York,
28
29
30 12 Wentworth Way, York YO10 5DD, UK.
31

32 13 * Corresponding author: sue.hartley@york.ac.uk
33
34
35 14
36
37 15
38
39
40 16
41
42
43 17
44
45 18
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

19 **Abstract** - Arbuscular mycorrhizal fungal (AMF) colonisation of plant roots is one of the
20 most ancient and widespread interactions in ecology, yet the systemic consequences for plant
21 secondary chemistry remain unclear. We performed the first metabolomic investigation into
22 the impact of AMF colonisation, by *Rhizophagus irregularis* on the chemical defences,
23 spanning above- and below-ground tissues, in its host-plant ragwort (*Senecio jacobaea*). We
24 used a non-targeted metabolomics approach to profile, and where possible identify,
25 compounds induced by AMF colonisation in both roots and shoots. Metabolomics analyses
26 revealed that 33 compounds were significantly increased in the root tissue of AMF-colonised
27 plants, including seven blumenols, plant-derived compounds known to be associated with
28 AMF colonisation. One of these was a novel structure conjugated with a malonyl-sugar and
29 uronic acid moiety, hitherto an unreported combination. Such structural modifications of
30 blumenols could be significant for their previously reported functional roles associated with
31 the establishment and maintenance of AM colonisation. Pyrrolizidine alkaloids (PAs), key
32 anti-herbivore defence compounds in ragwort, dominated the metabolomic profiles of root
33 and shoot extracts. Analyses of the metabolomic profiles revealed an increase in four PAs in
34 roots (but not shoots) of AMF colonised plants, with the potential to protect colonised plants
35 from below-ground organisms.

36
37 **Key Words** – Arbuscular mycorrhizal symbiosis, Above-belowground interactions,
38 Metabolomics, Pyrrolizidine alkaloids, Blumenols, *Senecio jacobaea*, *Rhizophagus*
39 *irregularis*.

41 INTRODUCTION

42 Plant-mediated interactions between organisms above- and below-ground are
43 increasingly recognised as a structuring force in ecology, though in many systems there is

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

still a need for a more mechanistic understanding of the chemical basis of these effects (Johnson et al. 2012, 2016; Wardle et al. 2004). Microbial symbionts have significant impacts on the chemical composition of their host plant, so are potentially key drivers of these indirect interactions (Hartley and Gange 2009). One of the most ancient and widely occurring plant-symbiont relationships involves arbuscular mycorrhizal fungi (AMF) (Smith and Read 2008). AMF are obligate symbionts that, via specialised structures (arbuscules) within the roots of their hosts, enhance plant uptake of macronutrients, particularly phosphorus and nitrogen (Hodge 2016; Hodge and Storer 2015; Javot et al. 2007; Leigh et al. 2009; Thirkell et al. 2016). In return, AMF are wholly dependent on the host plant for their carbon provision, usually acquiring around 10% of host photosynthate (Bago et al. 2000). These nutrient and carbon exchanges modify the nutrient status of the host plant, but also affect the allocation to secondary metabolites and other resistance mechanisms (Cameron et al. 2013; Minton et al. 2016). For example, colonisation by AMF has been shown to increase plant resistance to below-ground antagonists such as parasitic plants (Li et al. 2012) and plant parasitic nematodes (Rodriguez-Echeverria et al. 2009), as well as to generalist below-ground herbivores (Vannette and Rasmann 2012). Plant interactions with above-ground herbivores can also be affected by AMF colonisation (van der Putten et al. 2001), although usually to a lesser degree than below-ground (Van Wees et al. 2008), and outcomes vary with AMF species identity, herbivore diet breadth and/or feeding mode (Koricheva et al. 2009).

Early studies examining the impact of AMF colonisation on the chemical protection of plant tissues usually targeted specific chemical groups, limiting their power to detect the full range of induced chemical responses to AMF presence (de Deyn et al. 2009; Eftekhari et al. 2012). More recently, non-targeted metabolomic approaches have been used to investigate the effects of AMF on a wide range of metabolites (the metabolome) in foliar

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
69 tissues (Fester et al. 2011; Schweiger et al. 2014; Schweiger and Muller 2015). Again, several
70 of these did not measure effects on secondary metabolites, so much remains to be discovered
71 about the potential mechanisms by which AMF colonisation of plant roots can affect above-
72 ground plant-herbivore interactions (Gange et al. 2012; Schweiger and Muller 2015).
73 Similarly, the impact of AMF colonisation on the root metabolome has been little studied,
74 although Schliemann et al (2008a) used a metabolomic approach to show that both primary
75 and secondary metabolites differed between the roots of *Medicago truncatula* L. individuals
76 with and without AMF colonisation, and Saia et al (2015) studied impacts of colonisation on
77 the amino acid content of wheat. However, neither of these studies examined above-ground
78 foliar impacts, whilst none of the foliar studies (Fester et al. 2011; Schweiger et al. 2014;
79 Schweiger and Muller 2015) examined root tissue. Therefore, hitherto there has been no
80 metabolomic investigation into the concomitant above- and below-ground changes in plant
81 secondary chemistry following AMF colonisation, an important omission given the growing
82 interest in above-belowground links and systemic defence responses (Johnson et al. 2016).

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
83 This study addresses this knowledge gap by using non-targeted metabolomics
84 profiling techniques to examine the simultaneous effects of arbuscular mycorrhizal (AM)
85 colonisation on plant secondary chemistry in foliar and root tissues of ragwort (*Senecio*
86 *jacobaea* L.), a species which forms symbiotic associations with AMF (Gange et al. 2002;
87 Reidinger et al. 2012). We used experimental and untargeted metabolomic approaches to test
88 the effect of AMF (*Rhizophagus irregularis*) colonisation on the secondary chemistry of
89 ragwort roots and shoots. We aimed to identify the chemicals induced or repressed in host
90 root and shoot tissues as a result of AM colonisation. Effects of colonisation on host
91 metabolites might be expected to differ between the two tissue types given the established
92 differences in local vs systemic responses to AMF (Van Wees et al. 2008), and, whilst recent

1
2 93 research suggests high specificity in leaf metabolomic responses to AMF (Schweiger et al.
3 94 2014), this is as yet untested in roots. Hence, we address two questions:

4
5 95 1) How does colonisation by AMF affect the profiles of secondary metabolites in ragwort and
6
7 96 are the impacts more pronounced in root vs. foliar tissue?

8
9
10 97 2) Are the metabolites induced by AM colonisation similar to those identified in other plant-
11
12 98 fungal systems, or is there evidence of novel compounds, potentially indicating a degree of
13
14 99 host specificity?
15
16

17 100

18 19 20 101 MATERIALS AND METHODS

21 22 23 102 Microcosm experiment

24
25
26 103 Ragwort plants were grown from seeds collected from a single plant on the University of
27
28 104 Sussex campus (Lat: 50°52'18.6366"N, Long: 000°04'59.6860"W), sterilised in a 5% bleach
29
30 105 solution and germinated on double autoclaved vermiculite. Thirty-one pots (1.7 L) received a
31
32 106 double autoclaved (121°C; 60 min) growth medium comprising a 1:1 mixture of silver
33
34 107 sand:Terragreen® (an attapulgite clay; Turf-Pro Ltd, UK) and bone meal (0.25g/L⁻¹), a
35
36 108 complex N and P source to encourage mycorrhizal development (Hodge and Fitter 2010).
37
38 109 After mixing the growth media, all pots were twice flushed to saturation with water, aiming
39
40 110 to remove the pulse of nutrients released following sterilisation (Troelstra et al. 2001).
41
42

43
44
45 111 Seedlings were assigned to replicate pots and randomly assigned to AMF (n = 16) and control
46
47 112 (n = 15) treatments. Each replicate in the AMF treatment was inoculated with 50 g (dry
48
49 113 weight) of granular *Rhizophagus irregularis* (see Redecker et al. 2013 for recent changes in
50
51 114 nomenclature; obtained from PlantWorks, Kent, UK). Those replicates assigned to the
52
53 115 control treatment also had 50 g of this granular inoculate added, but double autoclaved
54
55 116 (121°C; 60 min) to kill the *R. irregularis*, to control for the input of organic material with the
56
57 117 inoculation (see Hodge 2001). In addition, non-AM pots received 100 ml of filtered
58
59
60
61
62
63
64
65

118 washings of the *R. irregularis* inoculum, but with AMF propagules removed to limit initial
119 differences among pots in starter microbial communities (as Atkin et al. 2009). To account
120 for any differences caused by the addition of water, 100 ml of deionised water was applied to
121 the AMF-treated microcosms. Each microcosm received a weekly dose of 50 ml of half-
122 strength Rorison's solution (Hewitt 1966), but with the plant-available phosphate removed to
123 encourage AMF colonisation. The experiment ran for 10 weeks in a greenhouse maintained
124 at 15-25°C with supplementary lighting (400W, high pressure sodium lamps) on a 16:8 L:D
125 photoperiod. Microcosms were watered with tap water ad libitum.

126 Plants were destructively harvested after 10 weeks in treatment. From each replicate
127 three new leaves (fully mature closest to the centre of the rosette) and two old leaves (closest
128 to the edge of the rosette with no visible signs of senescence) were excised, weighed,
129 immediately snap frozen in liquid nitrogen to halt enzymatic processes and stored in a -80°C
130 freezer for metabolomic analysis. Two representative root samples (each ~10% of the total
131 root mass) per plant were weighed and used for: i) plant metabolomic analysis (snap frozen
132 then stored at -80°C) and ii) to measure AMF colonisation (stored at 4°C until processed).
133 Root sections were cleared in a 10% potassium hydroxide solution, ink-stained (Vierheilig et
134 al. 1998), slide mounted and the presence or absence of AMF structures was recorded for a
135 minimum of 100 root intersections using a compound microscope (x 200 magnification).
136 Numbers of arbuscules, vesicles and root length colonized (RLC; the percentage of total
137 intercepts where hyphae or other AM fungal structures were present) were recorded for each
138 intersection (as Hodge 2003). The remaining shoot and root material was weighed and then
139 oven-dried (60°C, 72 hours) to obtain dry biomass.

140

141 Plant extraction and non-targeted metabolomic profiling

142 A homogenised sub-sample ($0.1 \pm 0.01\text{g}$), of the root and shoot tissue taken from each
143 replicate was mixed with 2mL of a 3:3:2 solvent mix of isopropanol, acetonitrile and water
144 using a ball mill (Pulverisette 23. Fritsch, Germany). This solvent mixture extracts both polar
145 metabolites (carbohydrates, amino acids) and non-polar metabolites (lipids) (Sana et al.
146 2010). Samples were spiked with two deuterated internal standards; 17β -estradiol 2,4,16,16-
147 d_4 sodium 3-sulfate (E2- d_4 -S, >99% D atom) and progesterone-2,2,4,6,6,17 α ,21,21,21- d_9 (P-
148 d_9 , 98% D atom) (Cambridge Isotope Laboratories Inc. MA and CDN isotopes, Quebec,
149 Canada, respectively). The sample was vortexed (1 min) and after overnight extraction at -
150 20°C , samples were centrifuged, the supernatant removed and the pellet extracted with 2 mL
151 of 80% methanol for 12 hours. The supernatants were combined and a 3 mL aliquot was
152 evaporated to dryness under vacuum and redissolved in 160 μl of methanol:water (3:1, v/v).
153 The extract was filtered (0.2 μm) prior to MS analysis. All extraction chemicals used were
154 purchased from Rathburn Chemicals Ltd, Walkerburn, UK.

155 Metabolites were profiled using ultraperformance liquid chromatography quadrupole
156 time-of-flight mass spectrometry (UPLC-QTOFMS) (Waters, Manchester, UK). Aliquots of
157 plant extracts were injected on to an Acquity UPLC BEH C18 column (1.7 μm particle size,
158 2.1 x 100 mm, Waters, UK). Metabolites were separated using a water-formic acid and
159 acetonitrile gradient as follows: 0-9.0 min, from 0 to 30.0% acetonitrile; 9.0-15.0 min, from
160 30.0 to 100% acetonitrile; 15.0 to 23.0 min, 100% acetonitrile. The flow rate was 0.2 mL
161 min^{-1} and the column temperature was 30°C . The injection volume of plant extracts was 20
162 μl .

163 Metabolites were detected in both positive and negative ESI modes using a
164 Micromass TOF-MS system (Waters, Manchester, UK). The mass spectrometer was tuned to
165 9000 mass resolution and data collected in full scan mode from 100 to 1200 m/z. The

166 collision gas used was argon, a constant collision energy of 10eV was used for all
167 experiments, and the TOF penning pressures ranged from 4.63×10^{-7} to 4.83×10^{-7} mbar.
168 Capillary voltage was 2.6 in positive mode and -2.9 in negative mode. In positive mode, the
169 cone voltage was set at 36V, and the multiplier voltage was set at 654V. In negative mode
170 the cone and multiplier voltage were set at 35V and 550V respectively. Desolvation N₂ gas
171 flow was set at 401L h⁻¹ for both ionisation modes.

172 MarkerLynx software (V 4.1, Waters, Manchester, UK) was used to align, normalise
173 and remove isotopic peaks from the metabolomic profiles. Each metabolite signal was the
174 description of an analyte using its specific retention time (r.t.) and mass-to-charge ratio (m/z).
175 Using SIMCA-P multivariate analysis software (Umetrics UK Ltd, Windsor, UK) the data
176 were pareto scaled, log-transformed and modelled using partial least square-discriminate
177 analysis (PLS-DA) for > 2 classes, or orthogonal partial least-square discriminate analysis
178 (OPLS-DA) for comparison of just 2 classes followed by examination of the loading plots to
179 detect MS signals associated with AMF colonisation (Liland 2011; Trygg and Wold 2002).
180 For all multivariate analyses, the explained variation (R²Y) and the predictive power of the
181 model (Q²) were examined to assess the performance of the models.

182 Biochemical markers associated with AM colonisation were extracted from 'S'-plots
183 derived from the OPLS-DA models (Wiklund et al. 2008). The identity of biochemical
184 markers was determined from their accurate mass composition and isotopic fit using the
185 elemental composition tool from the MassLynx software (V 4.1, Waters, Manchester, UK).
186 Fragmentation data, obtained from collision induced dissociation (CID) using QTOFMS
187 (collision energy between 20 and 50eV) was used to confirm the putative identity of the
188 markers.

189
190 Analyses of pyrrolizidine alkaloids in metabolomic profiles of tissue extracts.

191 Metabolomic profiles of root extracts acquired in positive electrospray ionization (ESI) mode
192 were dominated by highly abundant pyrrolizidine alkaloids (PAs) saturating the detector
193 response. PAs are the principal secondary metabolites in *Senecio* species, synthesised in the
194 roots (Hartmann et al. 1989) before being transported to the shoots (Hartmann and Dierich
195 1998) where they are effective defences against generalist herbivores (Macel et al. 2005;
196 Narberhaus et al. 2005; Thoden et al. 2009). In order to determine whether PA profiles
197 changed as a result of AM colonisation, an additional analysis of a 0.5 µl injection of root
198 extracts was undertaken to reduce the highly abundant PA signals. The chromatograms (of
199 both root and shoot extracts) were manually searched, and the relative abundance of observed
200 signals corresponding to PA metabolites was determined relative to the internal standard
201 using MassLynx peak integration software. PAs were assigned putative identities as
202 described above. Where possible, authenticated standards were purchased to confirm PA
203 identity: a mixed standard of PAs seneciphylline and senecionine (Carl Roth, GmbH & Co,
204 Karlsruhe, Germany) and a retrorsine N-oxide standard (PhytoLab GmbH & Co. KG,
205 Nürnberg, Germany). Details of the MS fragmentation data used for the identification of
206 each metabolite are outlined in the supporting information.

207

208 Data analysis

209 Adjusted fresh weights $((\text{total fresh weight}/100) \times \text{water content})$ of plant tissues were
210 calculated to account for a significant between-treatment difference in root and shoot water
211 content ($\text{water content} = (\text{fresh weight} - \text{dry weight}) / (\text{fresh weight}) \times 100$). Statistical analyses
212 of differences in plant biomass, biochemical markers and AM colonisation between
213 treatments were performed in SPSS v11. Where the residuals of the data met the
214 assumptions, either before or following transformations, of parametric models (normality,
215 homogeneity of variance), then one-way ANOVA (for plant biomass) or t-tests (for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

216 percentage RLC, metabolite and PA concentrations) were used. For non-parametric data,
217 Mann-Whitney U tests were employed. Bonferroni corrections were used to account for the
218 increased false discovery rate (FDR) associated with the multiple testing of metabolic
219 datasets containing many thousands of metabolite signals (Broadhurst and Kell 2006). A less
220 conservative Benjamini and Hochberg FDR correction was applied to the univariate tests of
221 PA concentrations (Benjamini et al. 2001), as this was more appropriate for these much
222 smaller datasets. The association between blumenol metabolites and percentage RLC were
223 investigated using Spearman rank-order correlations, as the data did not meet the assumptions
224 of parametric tests.

225 226 RESULTS

227 AMF colonisation and plant biomass

228 Mean percent RLC was $47 \pm 4\%$ in the *R. irregularis* colonised plants. Frequency of
229 arbuscules was $16 \pm 1\%$ and vesicles $13 \pm 2\%$ respectively. There was no AMF colonisation
230 of the non-AM controls. Shoot and root water content were higher in control plants than in *R.*
231 *irregularis* colonised plants, though plant biomass was unaffected by *R. irregularis*
232 colonisation (Table S1).

233 234 Root metabolome

235 The metabolite profiles of root extracts from AM colonised and non-AM control plants were
236 clearly separated, indicating a shift in chemical composition of plants in the different
237 treatment groups; (see Figure 1a for the positive ESI mode dataset, and a similar scores plot
238 was obtained for the negative ESI mode in Figure S1a). OPLS-DA of both datasets (positive
239 and negative ESI modes) produced models that explained a large amount of the variation in
240 metabolite profiles between treatments (negative: $R^2Y = 0.99$, positive: $R^2Y = 0.98$) with good

241 predictability (negative: $Q^2= 0.53$, positive: $Q^2= 0.48$). This separation between treatment
242 classes was driven by increased concentrations of 33 metabolites associated with AM
243 colonisation (threshold values after Bonferroni adjustment: positive mode: $P < 5.20 \times 10^{-7}$;
244 negative mode: $P < 5.01 \times 10^{-7}$). The fold increase in concentration of the 33 metabolites
245 associated with AM roots was between two to many thousands (up to 37,995-fold in one
246 instance). Eight of these metabolites were assigned identities (Table 1); the remaining
247 metabolites ($n = 25$) are listed in Table S2. Blumenol standards are not commercially
248 available, so identifications are derived from comparisons with published mass spectra (cited
249 in Strack and Fester 2006; Peipp et al. 1997; Schliemann et al. 2008b); see supporting
250 information (Note 1, Figure S2 and Table S3) for further details.

251 Seven of the identified metabolite signals were conjugates of C_{13} cyclohexenones, i.e.
252 blumenol apocarotenoids (metabolites 1-7, Table 1, Figure 2). Metabolites 1 and 4 were
253 identified as blumenols with a 13-hydroxyblumenol C moiety ($C_{13}H_{23}O_3$). Metabolites 2, 3,
254 5, 6 and 7 were identified as blumenols with a blumenol C moiety ($C_{13}H_{23}O_2$). Whilst
255 identified metabolites 1-7 share common blumenol structures, they differed in the nature of
256 conjugation (Figure 2). Metabolites 1, 2 and 5 were conjugated with glycoside sugars,
257 whereas metabolites 4 and 7 were conjugated with malonylglycoside sugars. Metabolites 3
258 and 6 were conjugated with an uronide group consistent with a glucuronide (Strack and
259 Fester 2006), and so were identified as a glycosyl-glucuronide and a malonylglycoside-
260 glucuronide conjugate respectively. Metabolite 6, blumenol C malonylglycosyl-glucuronide,
261 is a hitherto unreported structure, whereas metabolites 1-5 and 7 have been found to be
262 associated with AM colonisation in other species (Peipp et al. 1997; Schliemann et al.
263 2008b). All blumenols were present as their molecular ions, which were either $[M+H]^+$ or
264 $[M-H]^-$ in positive or negative ESI, respectively. In positive ESI, some metabolites were also
265 present as the sodium adduct of the molecular ion (Table S3). Of those blumenols identified,

266 two were observed in both positive and negative ESI modes (metabolites 1 & 6, Table 1).

267 The concentration changes of the identified blumenols corresponded to large differences,

268 with fold increases ranging from 250 to over 8000 in AM plants. The final identified

269 metabolite (number 8) was determined to be hexadecenoyl-glycero-phosphocholine, a

270 lysophospholipid.

271 All blumenol apocarotenoids were positively associated with increasing percentage

272 RLC in the AM colonised plants (Spearman correlation coefficient 0.523-0.711, $P \leq 0.03$ -

273 0.002, Table 2). Levels of hexadecenoyl-glycero-phosphocholine were not significantly

274 correlated with RLC ($P \geq 0.11$). A number of unidentified metabolites were detected that

275 were associated with AM colonisation (Table S2) and these were likely to be conjugated

276 phenolic compounds, two additional conjugated blumenols and (in one case) a phospholipid.

277 Based on their empirical formula, none of the metabolites (except for the phospholipid) were

278 present in the plant metabolite databases and so potentially represent new structures.

279 However, their concentrations were too low to allow further structural characterisation.

280 Changes in the concentrations of primary metabolites were not observed as a result of AM

281 colonisation, but this was most likely due to the fact that some structures such as lipids and

282 carotenoids were not have been detected by ESI and small organic molecules, including

283 alkaloid precursors, would not have been retained on the HPLC column.

284

285 Shoot metabolome

286 PLS-DA models of the shoot dataset from positive ESI analysis revealed good predictability

287 ($Q^2 = 0.66$) and a high degree of explained variation (total $R^2Y = 0.99$ for six latent variables)

288 (Figure 1b). PLS-DA recorded in negative ESI mode revealed similar models that explained

289 a high level of variation in shoot chemistry ($R^2Y = 0.62$, 4 latent variables) with lower

290 predictability ($Q^2 = 0.36$) than in positive mode (Figure S1b). In both positive and negative

291 modes, the first latent variable separated the metabolomes of old and new leaves. The
292 second, whilst separating AM and non-AM control groups, only explained a relatively small
293 amount of the observed variation in metabolite profiles between these treatments. OPLS-DA
294 models with the highest predictability ($R^2Y= 0.989$, $Q^2=0.431$) were associated with the
295 positive ESI profiles of old leaves from the different treatment groups. However, in all
296 OPLS-DA models of shoot tissues, no metabolites were found to vary significantly in
297 association with AMF colonisation of ragwort.

299 Concentrations of PA metabolites in the root and shoot metabolome.

300 Analysis of the UPLC-QTOFMS chromatograms from the metabolomics study
301 revealed that signals of PA metabolites were saturated in samples of the root extracts.
302 Analysis of lower volumes of samples revealed 19 PA signals that could be resolved in the
303 metabolomic profiles of roots, however more PA structures (potentially up to 33 metabolites,
304 Kostkeno et al. 2013) were likely to be present, which were not detected due to masking of
305 their signals by the most abundant PA signals. The concentration of individual PAs within a
306 species or genotype varies greatly in both roots and shoots (Cheng et al. 2011a). Of the
307 detected PAs, four were significantly increased in AM plants (Table 3). The PAs that
308 responded to AMF colonisation were identified as senecionine, jacoline N-oxide, jaconine N-
309 oxide, and usaramine N-oxide. Of these four PAs, only levels of senecionine were
310 significantly correlated with RLC (Spearman correlation coefficient -0.560 , $P \leq 0.02$).

311 Details of the identification of the different PA structures are given in supporting information.

312 The relative abundance of PAs was also determined in the metabolomics datasets of
313 the shoot extracts. A total of 17 abundant PA structures were resolved in the metabolomics
314 profiles of the shoot extracts out of a potential of 29 PA metabolites (Kostkeno et al. 2013).
315 However, there were no significant differences in concentrations (after FDR corrections) of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

316 PA signals between control and AMF-colonised treatment groups which was consistent with
317 the lack of discriminating metabolites observed in the above study on the shoot metabolome
318 (Table S4).

319

320 DISCUSSION

321 This study, the first to use a non-targeted metabolomic approach to assess the effect of AMF
322 colonisation on host plant secondary chemistry in both roots and shoots, showed that
323 colonisation by AMF could increase the concentrations of 33 metabolites in roots by between
324 two and many thousand fold. The extent of AMF colonisation was positively correlated with
325 levels of compounds thought to be associated with establishing and maintaining AMF
326 colonisation, specifically seven C₁₃ cyclohexenone blumenol structures, including one with a
327 hitherto unidentified structure. In addition, AMF colonisation affected the levels of the main
328 anti-herbivore defences in root tissues, with four PAs significantly increased in concentration.
329 Despite these below-ground changes, no significant changes in the metabolome of above-
330 ground tissues were detected in response to AMF colonisation of roots, although there were
331 clear metabolomic differences between leaf tissues of different ages. This is noteworthy as
332 had changes been found in aboveground tissues then it would indicate the AM fungus had a
333 direct impact upon the plant. However, as differences were only found in the ‘mycorrhizal’
334 (literally, ‘fungus-root’) root tissue, an unknown fraction of the metabolomics response
335 observed here, and in other studies (e.g. Rivero et al. 2015; Saia et al. 2015), may be due
336 directly to the fungal partner. Experimentally distinguishing between responses driven by the
337 AMF, the root and the mycorrhizal root is hampered by the fact that many of these fungal
338 symbionts cannot be grown in the absence of a host plant (Hodge et al. 2010). Nevertheless,
339 our results demonstrate that colonisation of roots by AMF results in clear metabolomics
340 differences compared to the un-colonised condition, a significant finding given that c. two-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

341 thirds of land plants form this type of close symbiotic association (Hodge and Fitter 2013;
342 Smith and Read 2008).

343 The process of initiating and maintaining the plant-AMF relationship involves a range
344 of metabolic changes within the host (Fester et al. 2011; Schliemann et al. 2008a), and the
345 blumenol metabolites we detected in our non-targeted analysis are of particular significance,
346 as they are thought to be involved in the chemical signalling underpinning the plant-AMF
347 interaction (Walter et al. 2010; Maier et al. 1995). It has been suggested that blumenols and
348 C14 polyene apocarotenoids have a role in maintaining the plant-AMF interaction once
349 formed (Fester et al. 2002). For instance, concentrations of root C14 polyenes and blumenols
350 increase (Peipp et al. 1997; Schliemann et al. 2006; 2008b) as the plant-AMF interaction
351 progresses, particularly in root cells hosting AMF structures (Fester et al. 2002). Mutant
352 plants, where blumenol production is reduced, are associated with an increased amount of
353 dead and degenerating mycorrhizal structures (Floss et al. 2008), providing further evidence
354 that these blumenol C13 cyclohexenones may have a role in the persistence of AMF
355 colonisation (Walter et al. 2010).

356 In support of this hypothesis, and in accordance with other studies (Maier et al. 2000;
357 Schliemann et al. 2008a), the concentrations of the seven blumenols we identified were all
358 positively related to levels of AMF colonisation. One of these induced blumenols (metabolite
359 6) was conjugated with a malonyl-sugar and glucuronic acid moiety, a combination that has
360 never been reported before. Species-specific blumenol metabolites have been observed in a
361 range of plant taxa (Fester et al. 2002, 2005; Maier et al. 2000; Peipp et al. 1997; Schliemann
362 et al. 2006, 2008a), and have been suggested as a mechanism of species recognition between
363 the AMF and its host (Strack and Fester 2006). This is a hypothesis we raise for further
364 testing, but current data is sparse. So far, only a few other plant species, such as the legume
365 *Medicago truncatula* colonised by *Glomus intraradices* (Schliemann et al. 2008a), tomato

1 366 colonised by *Rhizophagus irregularis* and *Funneliformis mossae* (Rivero et al. 2015) and
2 367 wheat colonised by multiple AM species (Saia et al. 2015) have had the root tissue response
3
4 368 to AMF colonisation been examined in an untargeted metabolomic approach. Until a wider
5
6
7 369 variety of plant-AMF species combinations are tested, it is too early to conclude whether any
8
9
10 370 species-specific signalling exists in AMF-plant interactions. It is notable, however, that one
11
12 371 recent study (Schweiger et al. 2014) which compared the impacts of the AMF *Rhizophagus*
13
14 372 *irregularis* on the leaf metabolome of five different plant species found that species-specific
15
16 373 metabolic changes following colonisation far out-numbered more generic cross-species
17
18 374 responses. Similarly, Rivero et al. (2015) detected fungus-specific aspects of the AM-
19
20
21 375 associated changes in the tomato metabolome.

22
23
24 376 Despite substantial changes to the root metabolome induced by AMF colonisation,
25
26 377 there were no concomitant shifts in the aboveground metabolome. Conceivably, changes in
27
28 378 the shoot metabolome may have been too small or inconsistent to have been detected, but this
29
30 379 seems unlikely given that our UPLC-QTOFMS methodology clearly discriminated between
31
32 380 the metabolomes of old and new leaves (Figure 1b), and that we have used the same
33
34 381 technique to successfully detect differences in the metabolomes of above-ground tissues in
35
36 382 ragwort previously (Hartley et al. 2012). Previous studies have demonstrated that AMF-
37
38 383 mediated changes occur in above-ground plant tissues, with effects on gene expression
39
40 384 (Taylor and Harrier 2003) and on herbivore performance (Koricheva et al. 2009), but changes
41
42 385 in foliar secondary metabolites in response to colonisation seem more inconsistent. It has
43
44 386 been suggested that induction of chemical defences by beneficial microbes are localised
45
46 387 rather than systemic (Van Wees et al. 2008), which may explain why the increases in PA
47
48 388 concentration in AMF colonised ragwort we observed were confined to the roots. Increases in
49
50 389 PA concentrations in the roots may even reduce PA levels in the shoots. Although our study
51
52 390 did not detect such a reduction in foliar allocation to PAs, Reidinger et al. (2012) showed that
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

391 leaf concentrations of both total PAs and jacoline were negatively correlated with AMF
392 colonisation of ragwort in the field.

393 The increased PA levels that we found in AMF colonised roots could impact on
394 trophic interactions. AMF colonisation has been shown to have a negative influence on
395 generalist below-ground herbivores such as parasitic nematodes (Li et al. 2006; Rodriguez-
396 Echeverria et al. 2009) and beetle larvae (Gange et al. 1994), but positive effects on the root-
397 feeding larvae of specialist weevils (Currie et al. 2011). Overall, generalist herbivores,
398 whether above or below ground, seem more affected by AMF colonisation of their host plants
399 than specialists (Gange et al. 2012; Koricheva et al. 2009). In ragwort, the highly toxic PAs
400 deter generalist herbivores (Joosten and van Veen 2011; Thoden et al. 2009), whilst specialist
401 herbivores have the capacity to use PAs to locate and select hosts (Cheng et al. 2013), to
402 detoxify them, and even sequester these compounds for their own defence against predators
403 (Beuerle et al. 2007; Narberhaus et al. 2003). Most previous studies have demonstrated these
404 effects on above-ground herbivores (e.g. Cheng et al. 2011b; Wei et al. 2015); far less is
405 known about the impacts of PAs on root-feeding herbivores (Cheng et al. 2011a) and we have
406 found no published studies on how AMF colonisation modifies those effects. It has been
407 shown that root damage increases PA levels (Hol et al. 2004) and that PAs affect root fungi
408 and the composition of the rhizosphere (Hol and van Veen 2002; Kowalchuk et al. 2006), so
409 there is potential for significant ecological consequences deriving from the changes in PAs
410 we observed in AM roots.

411 In summary, this study has shown that colonisation by the AMF species *R.*
412 *irregularis* causes numerous and significant changes in the root metabolome of ragwort, but
413 not in the shoot metabolome. In root tissues, the key changes were in groups of chemicals
414 associated with AMF-plant signalling (blumenols) and anti-herbivore defence (PAs). We
415 showed strong correlations between the levels of blumenols and AMF colonisation,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

416 supporting the idea that these metabolites play a key role in the interactions between AMF
417 and their hosts. The discovery that one of these was a novel structure not previously reported
418 raises questions about the significance of structural modifications in species recognition and
419 signalling that would repay further study.

420
421 Acknowledgements - For advice and practical assistance with the metabolomic analysis
422 we would like to thank Dr Raghad Al-Salhi and Ms Julia Horwood. Dr Stefan Reidinger and
423 Dr Kate Storer provided mycorrhizal cultures and advice on experimental procedures. We
424 also thank the University of Sussex and the Centre for Ecology and Hydrology (project
425 NEC03605) for co-funding this project.

426
427

428 REFERENCES

- 429 Atkin OK, Sherlock D, Fitter AH, Jarvis S, Hughes JK, Campbell C, Hurry V, Hodge A,
430 (2009) Temperature dependence of respiration in roots colonized by arbuscular
431 mycorrhizal fungi. *New Phytol* 182:188-199. doi:10.1111/j.1469-8137.2008.02727.x
- 432 Bago B, Pfeffer PE, Shachar-Hill Y (2000) Carbon metabolism and transport in arbuscular
433 mycorrhizas. *Plant Physiol* 124:949-957. doi:10.1104/pp.124.3.949
- 434 Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I (2001) Controlling the false discovery
435 rate in behavior genetics research. *Behav Brain Res* 125:279-284. doi:10.1016/s0166-
436 4328(01)00297-2
- 437 Beuerle T, Theuring C, Klewer N, Schulz S, Hartmann T (2007) Absolute configuration of
438 the creatonotines and callimorphines, two classes of arctiid-specific pyrrolizidine
439 alkaloids. *Insect Biochem Mol Biol* 37:80-89. doi:10.1016/j.ibmb.2006.10.005

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 440 Broadhurst DI, Kell DB (2006) Statistical strategies for avoiding false discoveries in
441 metabolomics and related experiments. *Metabolomics* 2:171-196.
442 doi:10.1007/s11306-006-0037-z
- 443 Cameron DD, Neal AL, Van Wees SCM, Ton J (2013) Mycorrhiza-induced resistance: more
444 than the sum of its parts? *Trends Plant Sci* 18:539-545.
445 doi:10.1016/j.tplants.2013.06.004
- 446 Cheng D, Kirk H Mulder PPJ Vrieling K Klinkhamer PGL (2011a) Pyrrolizidine alkaloid
447 variation in shoots and roots of segregating hybrids between *Jacobaea vulgaris* and
448 *Jacobaea aquatica*. *New Phytol* 192: 1010-1023 doi: 10.1111/j.1469-
449 8137.2011.03841.x
- 450 Cheng D, Kirk H, Vrieling K, Mulder PPJ, Klinkhamer PGL (2011b) The relationship
451 between structurally different pyrrolizidine alkaloids and western flower thrips
452 resistance in F-2 hybrids of *Jacobaea vulgaris* and *Jacobaea aquatica*. *Journal of*
453 *Chemical Ecology* 37:1071-1080. doi:10.1007/s10886-011-0021-6
- 454 Cheng D, van der Meijden E, Mulder PPJ, Vrieling K, Klinkhamer PGL (2013) Pyrrolizidine
455 alkaloid composition influences cinnabar moth oviposition preferences in *Jacobaea*
456 hybrids. *Journal of Chemical Ecology* 39:430-437. doi:10.1007/s10886-013-0257-4
- 457 Currie AF, Murray PJ, Gange AC (2011) Is a specialist root-feeding insect affected by
458 arbuscular mycorrhizal fungi? *Appl Soil Ecol* 47:77-83.
459 doi:10.1016/j.apsoil.2010.12.002
- 460 de Deyn GB, Biere A, van der Putten WH, Wagenaar R, Klironomos JN (2009) Chemical
461 defense, mycorrhizal colonization and growth responses in *Plantago lanceolata* L.
462 *Oecologia* 160:433-442. doi:10.1007/s00442-009-1312-2
- 463 Eftekhari M, Alizadeh M, Ebrahimi P (2012) Evaluation of the total phenolics and quercetin
464 content of foliage in mycorrhizal grape (*Vitis vinifera* L.) varieties and effect of

1 465 postharvest drying on quercetin yield. *Ind Crop Prod* 38:160-165.
2 466 doi:10.1016/j.indcrop.2012.01.022
3
4 467 Fester T, Hause B, Schmidt D, Halfmann K, Schmidt J, Wray V, Hanse G, Strack D (2002)
5
6 468 Occurrence and localization of apocarotenoids in arbuscular mycorrhizal plant roots.
7
8 469 *Plant Cell Physiol* 43:256-265. doi:10.1093/pcp/pcf029
9
10 470 Fester T, Fetzer I, Buchert S, Lucas R, Rillig MC, Hartig C (2011) Towards a systemic
11
12 471 metabolic signature of the arbuscular mycorrhizal interaction. *Oecologia* 167:913-
13
14 472 924. doi:10.1007/s00442-011-2037-6
15
16 473 Fester T, Wray V, Nimtz M, Strack D (2005) Is stimulation of carotenoid biosynthesis in
17
18 474 arbuscular mycorrhizal roots a general phenomenon? *Phytochem* 66:1781-1786.
19
20 475 doi:10.1016/j.phytochem.2005.05.009
21
22 476 Floss DS, Hause B, Lange PR, Kuster H, Strack D, Walter MH (2008) Knock-down of the
23
24 477 MEP pathway isogene 1-deoxy-D-xylulose 5-phosphate synthase 2 inhibits formation
25
26 478 of arbuscular mycorrhiza-induced apocarotenoids, and abolishes normal expression of
27
28 479 mycorrhiza-specific plant marker genes. *Plant J* 56:86-100. doi:10.1111/j.1365-
29
30 480 313X.2008.03575.x
31
32 481 Gange AC, Bower E, Brown VK (2002) Differential effects of insect herbivory on arbuscular
33
34 482 mycorrhizal colonization. *Oecologia* 131:103-112. doi:10.1007/s00442-001-0863-7
35
36 483 Gange AC, Brown VK, Sinclair GS (1994) Reduction of black vine weevil larval growth by
37
38 484 vesicular-arbuscular mycorrhizal infection. *Entomol Exp Appl* 70:115-119.
39
40 485 Gange AC, Eschen R, Schroeder V (2012) The soil microbial community and plant foliar
41
42 486 defences against insects. In: Iason GR, Dicke M and Hartley SE (eds) *The ecology of*
43
44 487 *plant secondary metabolites: from genes to global processes*. Cambridge University
45
46 488 Press, New York, pp 170-189
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 489 Hartley SE, Gange AC (2009) Impacts of plant symbiotic fungi on insect herbivores:
1
2 490 mutualism in a multitrophic context. In: Annual Review of Entomology, vol 54, Palo
3
4 491 Alto, pp 323-342. doi:10.1146/annurev.ento.54.110807.090614
5
6
7 492 Hartley SE, Eschen R, Horwood JM, Robinson LA, Hill EM (2012) Plant secondary
8
9 493 metabolites and the interactions between plants and other organisms: the potential of a
10
11 494 metabolomic approach. In: Iason GR, Dicke M and Hartley SE (eds) The ecology of
12
13 495 plant secondary metabolites: from genes to global processes, Cambridge University
14
15 496 Press, New York, pp. 204-225
16
17
18
19 497 Hartmann T, Dierich B (1998) Chemical diversity and variation of pyrrolizidine alkaloids of
20
21 498 the senecionine type: biological need or coincidence? *Planta* 206:443-451.
22
23 499 doi:10.1007/s004250050420
24
25
26 500 Hartmann T, Ehmke A, Eilert U, Vonborstel K, Theuring C (1989) Sites of synthesis,
27
28 501 translocation and accumulation of pyrrolizidine alkaloid N-oxides in *Senecio vulgaris*
29
30 502 L. *Planta* 177:98-107. doi:10.1007/bf00392159
31
32
33
34 503 Hewitt EJ (1966) Sand and water culture methods used in the study of plant nutrition. CAB,
35
36 504 Farnham Royal, UK.
37
38
39 505 Hodge A (2001) Arbuscular mycorrhizal fungi influence decomposition of, but not plant
40
41 506 nutrient capture from, glycine patches in soil. *New Phytol* 151:725-734.
42
43 507 doi:10.1046/j.0028-646x.2001.00200.x
44
45
46 508 Hodge A (2003) N capture by *Plantago lanceolata* and *Brassica napus* from organic
47
48 509 material: the influence of spatial dispersion, plant competition and an arbuscular
49
50 mycorrhizal fungus. *J Exp Bot* 54:2331-2342. doi:10.1093/jxb/erg249
51
52
53 511 Hodge A (2016) Accessibility of inorganic and organic nutrients for mycorrhizas. In: Johnson
54
55 512 NC, Gehring C and Jansa J (eds) *Mycorrhizal mediation of soil: fertility, structure,*
56
57 513 *and carbon storage.* Elsevier, Amsterdam, pp 129-148
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 514 Hodge A, Fitter AH (2010) Substantial nitrogen acquisition by arbuscular mycorrhizal fungi
515 from organic material has implications for N cycling. *P Natl Acad Sci USA*
516 107:13754-13759. doi:10.1073/pnas.1005874107
- 517 Hodge A, Fitter AH (2013) Microbial mediation of plant competition and community
518 structure. *Funct Ecol* 27:865-875. doi:10.1111/1365-2435.12002
- 519 Hodge A, Storer K (2015) Arbuscular mycorrhiza and nitrogen: implications for individual
520 plants through to ecosystems. *Plant Soil* 386:1-19. doi:10.1007/s11104-014-2162-1
- 521 Hodge A, Helgason T, Fitter AH (2010) Nutritional ecology of arbuscular mycorrhizal fungi.
522 *Fungal Ecol* 3:267-273. doi: 10.1016/j.funeco.2010.02.002
- 523 Hol WHG, Macela M, van Veen JA, van der Meijden E (2004) Root damage and
524 aboveground herbivory change concentration and composition of pyrrolizidine
525 alkaloids of *Senecio jacobaea* *Basic Appl Ecol* 5: 253–260. doi:
526 10.1016/j.baae.2003.12.002
- 527 Hol WHG, Van Veen JA (2002) Pyrrolizidine alkaloids from *Senecio jacobaea* affect fungal
528 growth. *J Chem Ecol* 28: 1763-1772. doi: 10.1023/A:1020557000707
- 529 Javot H, Pumplin N, Harrison MJ (2007) Phosphate in the arbuscular mycorrhizal symbiosis:
530 transport properties and regulatory roles. *Plant Cell Environ* 30:310-322.
531 doi:10.1111/j.1365-3040.2006.01617.x
- 532 Johnson SN, Clark KE, Hartley SE, Jones TH, McKenzie SW, Koricheva J (2012)
533 Aboveground-belowground herbivore interactions: a meta-analysis. *Ecology* 93:2208-
534 2215. doi: 10.1890/11-2272.1
- 535 Johnson SN, Erb M, Hartley SE (2016) Roots under attack: contrasting plant responses to
536 below- and aboveground insect herbivory. *New Phytol* 210:413-418.
537 doi:10.1111/nph.13807

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 538 Joosten L, van Veen JA (2011) Defensive properties of pyrrolizidine alkaloids against
539 microorganisms. *Phytochem Rev* 10:127-136. doi:10.1007/s11101-010-9204-y
540 Koricheva J, Gange AC, Jones T (2009) Effects of mycorrhizal fungi on insect herbivores: a
541 meta-analysis. *Ecology* 90:2088-2097. doi:10.1890/08-1555.1
542 Kostenko O, Mulder PJ, Martijn Bezemer T (2013) Effects of Root Herbivory on
543 Pyrrolizidine Alkaloid Content and Aboveground Plant-Herbivore-Parasitoid
544 Interactions in *Jacobaea Vulgaris*. *J Chem Ecol* 39:109-119. doi.org/10.1007/s10886-
545 012-0234-3
546 Kowalchuk GA, Hol WHG, van Veen JA (2006) Rhizosphere fungal communities are
547 influenced by *Senecio jacobaea* pyrrolizidine alkaloid content and composition. *Soil*
548 *Biology & Biochemistry* 38: 2852–2859 doi: 10.1016/j.soilbio.2006.04.043
549 Leigh J, Hodge A, Fitter AH (2009) Arbuscular mycorrhizal fungi can transfer substantial
550 amounts of nitrogen to their host plant from organic material. *New Phytol* 181:199-
551 207. doi:10.1111/j.1469-8137.2008.02630.x
552 Li AR, Smith SE, Smith FA, Guan KY (2012) Inoculation with arbuscular mycorrhizal fungi
553 suppresses initiation of haustoria in the root hemiparasite *Pedicularis tricolor*. *Ann*
554 *Bot* 109:1075-1080. doi:10.1093/aob/mcs028
555 Li HY, Yang GD, Shu HR, Yang YT, Ye BX, Nishida I, Zheng CC (2006) Colonization by
556 the arbuscular mycorrhizal fungus *Glomus versiforme* induces a defense response
557 against the root-knot nematode *Meloidogyne incognita* in the grapevine (*Vitis*
558 *amurensis* Rupr.), which includes transcriptional activation of the class III chitinase
559 gene VCH3. *Plant Cell Physiol* 47:154-163. doi:10.1093/pcp/pci231
560 Liland KH (2011) Multivariate methods in metabolomics - from pre-processing to dimension
561 reduction and statistical analysis. *Trends Anal Chem* 30:827-841.
562 doi:10.1016/j.trac.2011.02.007

- 563 Macel M, Bruinsma M, Dijkstra SM, Ooijendijk T, Niemeyer HM, Klinkhamer PGL (2005)
1
2 564 Differences in effects of pyrrolizidine alkaloids on five generalist insect herbivore
3
4
5 565 species. *J Chem Ecol* 31:1493-1508. doi:10.1007/s10886-005-5793-0
6
- 7 566 Maier W, Peipp H, Schmidt J, Wray V, Strack D (1995) Levels of a terpenoid glycoside
8
9 567 (blumenin) and cell wall-bound phenolics in some cereal mycorrhizas. *Plant Physiol*
10
11 568 109:465-470. doi:10.1104/pp.109.2.465
13
- 14 569 Maier W, Schmidt J, Nimtz M, Wray V, Strack D (2000) Secondary products in mycorrhizal
15
16 570 roots of tobacco and tomato. *Phytochem* 54:473-479. doi:10.1016/s0031-
17
18 571 9422(00)00047-9
20
- 21 572 Minton MM, Barber NA, Gordon LL (2016) Effects of arbuscular mycorrhizal fungi on
22
23 573 herbivory defense in two *Solanum* (Solanaceae) species. *Plant Ecol Evol* 149:157-
24
25 574 164. doi:10.5091/plecevo.2016.1176
27
- 28 575 Narberhaus I, Theuring C, Hartmann T, Dobler S (2003) Uptake and metabolism of
29
30 576 pyrrolizidine alkaloids in *Longitarsus* flea beetles (Coleoptera : Chrysomelidae)
31
32 577 adapted and non-adapted to alkaloid-containing host plants. *J Comp Physiol B-
33
34 578 Biochem Syst Environ Physiol* 173:483-491. doi:10.1007/s00360-003-0356-6
35
36
- 37 579 Narberhaus I, Zintgraf V, Dobler S (2005) Pyrrolizidine alkaloids on three trophic levels -
38
39 580 evidence for toxic and deterrent effects on phytophages and predators. *Chemoecology*
40
41 581 15:121-125. doi:10.1007/s00049-005-0302-z
42
43
- 44 582 Peipp H, Maier W, Schmidt J, Wray V, Strack D (1997) Arbuscular mycorrhizal fungus-
45
46 583 induced changes in the accumulation of secondary compounds in barley roots.
47
48 584 *Phytochem* 44:581-587. doi:10.1016/s0031-9422(96)00561-4
49
50
- 51 585 Redecker D, Schussler A, Stockinger H, Sturmer SL, Morton JB, Walker C (2013) An
52
53 586 evidence-based consensus for the classification of arbuscular mycorrhizal fungi
54
55 587 (Glomeromycota). *Mycorrhiza* 23:515-531. doi:10.1007/s00572-013-0486-y
56
57
58
59
60
61
62
63
64
65

- 588 Reidinger S, Eschen R, Gange AC, Finch P, Bezemer TM (2012) Arbuscular mycorrhizal
1
2 589 colonization, plant chemistry, and aboveground herbivory on *Senecio jacobaea*. *Acta*
3
4
5 590 *Oecol* 38:8-16. doi:10.1016/j.actao.2011.08.003
6
- 7 591 Rivero J, Gamir J, Aroca R, Pozo MJ, Flors V (2015) Metabolic transition in mycorrhizal
8
9 592 tomato roots. *Front Microbiol* 6: 598. doi: 10.3389/fmicb.2015.00598
10
11
- 12 593 Rodriguez-Echeverria S, de la Pena E, Moens M, Freitas H, van der Putten WH (2009) Can
13
14 594 root-feeders alter the composition of AMF communities? Experimental evidence from
15
16 595 the dune grass *Ammophila arenaria*. *Basic Appl Ecol* 10:131-140.
17
18 596 doi:10.1016/j.baae.2008.01.004
19
20
- 21 597 Saia S, Ruisi P, Fileccia V, Di Miceli G, Amato G, Martinelli F (2015) Metabolomics
22
23 598 suggests that soil inoculation with arbuscular mycorrhizal fungi decreased free amino
24
25 599 acid content in roots of durum wheat grown under N-limited, P-rich field conditions.
26
27 600 *PLoS One* 10:15. doi:10.1371/journal.pone.0129591
28
29
30
- 31 601 Sana TR, Fischer S, Wohlgemuth G, Katrekar A, Jung KH, Ronald PC, Fiehn O (2010)
32
33 602 Metabolomic and transcriptomic analysis of the rice response to the bacterial blight
34
35 603 pathogen *Xanthomonas oryzae* pv. *oryzae*. *Metabolomics* 6:451-465.
36
37 604 doi:10.1007/s11306-010-0218-7
38
39
40
- 41 605 Schliemann W, Ammer C, Strack D (2008a) Metabolite profiling of mycorrhizal roots of
42
43 606 *Medicago truncatula*. *Phytochem* 69:112-146. doi:10.1016/j.phytochem.2007.06.032
44
45
- 46 607 Schliemann W, Ammer C, Strack D (2008b) Metabolite profiling of mycorrhizal roots of
47
48 608 *Medicago truncatula*. *Phytochem* 69:1446-1447.
49
50 609 doi:10.1016/j.phytochem.2008.02.004
51
52
- 53 610 Schliemann W, Schmidt J, Nimtz M, Wray V, Thomas F, Strack D (2006) Accumulation of
54
55 611 apocarotenoids in mycorrhizal roots of *Ornithogalum umbellatum*. *Phytochem*
56
57 612 67:1196-1205. doi:10.1016/j.phytochem.2006.05.005
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 613 Schweiger R, Baier MC, Persicke M, Muller C (2014) High specificity in plant leaf metabolic
614 responses to arbuscular mycorrhiza. *Nat Commun* 5:11. doi:10.1038/ncomms4886
- 615 Schweiger R, Muller C (2015) Leaf metabolome in arbuscular mycorrhizal symbiosis. *Curr*
616 *Opin Plant Biol* 26:120-126. doi:10.1016/j.pbi.2015.06.009
- 617 Smith SE, Read DJ (2008) *Mycorrhizal symbiosis*. Academic Press, San Diego
- 618 Strack D, Fester T (2006) Isoprenoid metabolism and plastid reorganization in arbuscular
619 mycorrhizal roots. *New Phytol* 172:22-34. doi:10.1111/j.1469-8137.2006.01837.x
- 620 Taylor J, Harrier LA (2003) Expression studies of plant genes differentially expressed in leaf
621 and root tissues of tomato colonised by the arbuscular mycorrhizal fungus *Glomus*
622 *mosseae*. *Plant Mol Biol* 51:619-629. doi:10.1023/a:1022341422133
- 623 Thirkell TJ, Cameron DD, Hodge A (2016) Resolving the 'nitrogen paradox' of arbuscular
624 mycorrhizas: fertilization with organic matter brings considerable benefits for plant
625 nutrition and growth. *Plant Cell Environ* 39:1683-1690. doi:10.1111/pce.12667
- 626 Thoden TC, Boppre M, Hallmann J (2009) Effects of pyrrolizidine alkaloids on the
627 performance of plant-parasitic and free-living nematodes. *Pest Manag Sci* 65:823-
628 830. doi:10.1002/ps.1764
- 629 Troelstra SR, Wagenaar R, Smant W, Peters BAM (2001) Interpretation of bioassays in the
630 study of interactions between soil organisms and plants: involvement of nutrient
631 factors. *New Phytol* 150:697-706. doi:10.1046/j.1469-8137.2001.00133.x
- 632 Trygg J, Wold S (2002) Orthogonal projections to latent structures (O-PLS). *J Chemometr*
633 16:119-128. doi:10.1002/cem.695
- 634 van der Putten WH, Vet LEM, Harvey JA, Wackers FL (2001) Linking above- and
635 belowground multitrophic interactions of plants, herbivores, pathogens, and their
636 antagonists. *Trends Ecol Evol* 16:547-554. doi:10.1016/s0169-5347(01)02265-0

- 637 Van Wees SCM, Van der Ent S, Pieterse CMJ (2008) Plant immune responses triggered by
1
2 638 beneficial microbes. *Curr Opin Plant Biol* 11:443-448. doi:10.1016/j.pbi.2008.05.005
3
4 639 Vannette RL, Rasmann S (2012) Arbuscular mycorrhizal fungi mediate below-ground plant-
5
6 herbivore interactions: a phylogenetic study. *Funct Ecol* 26:1033-1042.
7 640
8
9 641 doi:10.1111/j.1365-2435.2012.02046.x
10
11 642 Vierheilig H, Coughlan AP, Wyss U, Piche Y (1998) Ink and vinegar, a simple staining
12
13 technique for arbuscular-mycorrhizal fungi. *Appl Environ Microbiol* 64:5004-5007.
14 643
15
16 644 Walter MH, Floss DS, Strack D (2010) Apocarotenoids: hormones, mycorrhizal metabolites
17
18 and aroma volatiles. *Planta* 232:1-17. doi:10.1007/s00425-010-1156-3
19 645
20
21 646 Wardle DA, Bardgett RD, Klironomos JN, Setälä H, van der Putten WH, Wall DH (2004)
22
23 Ecological linkages between aboveground and belowground biota. *Science* 304:1629-
24 647
25 1633. doi:10.1126/science.1094875
26 648
27
28 649 Wei X, Vrieling K, Mulder PPJ, Klinkhamer PGL (2015) Testing the generalist-specialist
29
30 dilemma: the role of pyrrolizidine alkaloids in resistance to invertebrate herbivores in
31 650
32 *Jacobaea* species. *J Chem Ecol* 41:159- 167. doi: 10.1007/s10886-015-0551-4
33 651
34
35 652 Wiklund S, Johansson E, Sjöström L, Mellerowicz EJ, Edlund U, Shockcor JP, Gottfries J,
36
37 Moritz T, Trygg J (2008) Visualization of GC/TOF-MS-based metabolomics data for
38 653
39 identification of biochemically interesting compounds using OPLS class models. *Anal*
40 654
41
42 *Chem* 80:115-122. doi:10.1021/ac0713510
43 655
44
45
46
47 656
48 657
49 658
50
51 659
52
53
54
55
56
57
58
59
60
61
62
63
64
65

TABLE 1 METABOLITES IDENTIFIED IN RAGWORT ROOTS THAT WERE SIGNIFICANTLY INCREASED AFTER COLONISATION BY RHIZOPHAGUS IRREGULARIS

Metabolite number [¶]	Observed ion (m/z)	UPLC-TOFMS r.t.	Putative formula	Theoretical mass of ion	Putative identity	Fold change ^a	P-value ^b
1	389.2177	6.37	C ₁₉ H ₃₃ O ₈	389.2175	13-hydroxyblumenol C glycoside [M+H] ⁺	1151.2*	6.65 x 10 ⁻⁹
1	387.2018	6.37	C ₁₉ H ₃₃ O ₈	387.2018	13-hydroxyblumenol C glycoside [M-H] ⁻	746.6*	6.65 x 10 ⁻⁹
2	373.2228	7.23	C ₁₉ H ₃₃ O ₇	373.2226	Blumenol C glycoside [M+H] ⁺	8065.8	6.65 x 10 ⁻⁹
3	547.2389	7.30	C ₂₅ H ₃₉ O ₁₃	547.2391	Blumenol C glycosyl-glucuronide [M-H] ⁻	1553.3	6.65 x 10 ⁻⁹
4	475.2182	7.55	C ₂₂ H ₃₅ O ₁₁	475.2179	13-hydroxyblumenol C malonylglycoside [M+H] ⁺	489.7	2.99 x 10 ⁻⁷
5	373.2222	8.20	C ₁₉ H ₃₃ O ₇	373.2226	Blumenol C glycoside [M+H] ⁺	5032.0	6.65 x 10 ⁻⁹
6	635.2551	8.24	C ₂₈ H ₄₃ O ₁₆	635.2551	Blumenol C malonylglycosyl-glucuronide [M+H] ⁺	3101.3*	6.65 x 10 ⁻⁹
6	633.2396	8.24	C ₂₈ H ₄₁ O ₁₆	633.2395	Blumenol C malonylglycosyl-glucuronide [M-H] ⁻	250.4*	6.65 x 10 ⁻⁹
7	459.2232	9.38	C ₂₂ H ₃₅ O ₁₀	459.2230	Blumenol C malonylglycoside [M+H] ⁺	1109.2	6.65 x 10 ⁻⁹
8	494.3247	14.20	C ₂₄ H ₄₉ NO ₇ P	494.3247	Hexadecenoyl-glycero-phosphocholine [M+H] ⁺	7.86	2.00 x 10 ^{-7c}

[¶]Structures given in Figure 2.

r.t.= retention time

^a Fold change indicates the concentration increase in roots colonised with AMF when compared to the concentrations observed in control plants.

^b Significance determined using t-tests (°) or Mann-Whitney U tests (unmarked) after Bonferroni adjustments.

* Differences in estimation of fold change between [M+H]⁺ and [M-H]⁻ signals for the same metabolite were due to presence of a Na adduct competing with the [M+H]⁺ ion in positive ESI mode.

TABLE 2 SPEARMAN RANK-ORDER CORRELATIONS (N = 16) BETWEEN THE CONCENTRATION OF THE BLUMENOL METABOLITES MEASURED IN RAGWORT ROOTS AND RHIZOPHAGUS IRREGULARIS COLONISATION IN THE AMF-TREATED PLANTS

Metabolite structure (see Figure 3)	Putative identity	Correlation coefficient	P-value [‡]
1	13-hydroxyblumenol C glycoside	0.852	1.16 x 10 ⁻⁹
2	Blumenol C glycoside	0.902	4.33 x 10 ⁻¹²
3*	Blumenol C glycosyl-glucuronide	0.828	9.37 x 10 ⁻⁹
4	13-hydroxyblumenol C malonylglycoside	0.841	3.21 x 10 ⁻⁹
5	Blumenol C glycoside	0.844	2.41 x 10 ⁻⁹
6	Blumenol C malonylglycosyl-glucuronide	0.832	6.39 x 10 ⁻⁹
7	Blumenol C malonylglycoside	0.855	9.36 x 10 ⁻¹⁰

* Result from MS analysis in -ESI mode (required to detect metabolite 3); all other results from +ESI mode

[‡]All P-values significant after Bonferroni adjustment

TABLE 3 AVERAGE RELATIVE CONCENTRATIONS (\pm S.E.) OF THE PYRROLIZIDINE ALKALOID (PA) SIGNALS MEASURED IN THE POSITIVE ESI UPLC-TOFMS PROFILES OF CONTROL (N=15) AND RHIZOPHAGUS IRREGULARIS COLONISED (N=16) RAGWORT ROOTS

Theoretical mass of ion	Putative formula	UPLC-TOFMS r.t.	Control	G. intraradices colonised	P-value ^a	Fold change ^{b,c}	Putative identity ^c
334.1654	C ₁₈ H ₂₃ NO ₅	6.00	49.7 (\pm 5.3)	59.2 (\pm 5.3)	0.188		
336.1811	C₁₈H₂₅NO₅	6.86	211.7 (\pm17.7)	289.8 (\pm21.6)	3.63 x 10⁻³	1.37	Senecionine
350.1604	C ₁₈ H ₂₃ NO ₆	6.34	220.5 (\pm 27.7)	223.2 (\pm 21.6)	0.571		
	C ₁₈ H ₂₃ NO ₆	9.01	24.5 (\pm 3.2)	38.6 (\pm 6.2)	0.054		
352.1760	C ₁₈ H ₂₅ NO ₆	4.89	23.7 (\pm 8.7)	52.0 (\pm 14.2)	0.033		
	C ₁₈ H ₂₅ NO ₆	5.82	76.1 (\pm 11.3)	55.3 (\pm 5.4)	0.216		
	C ₁₈ H ₂₅ NO ₆	6.13	327.9 (\pm 23.9)	303.8 (\pm 17.9)	0.711		
	C ₁₈ H ₂₅ NO ₆	6.99	48.8 (\pm 2.5)	53.0 (\pm 2.4)	0.029		
	C ₁₈ H ₂₅ NO ₆	7.16	108.8 (\pm 4.2)	113.0 (\pm 3.5)	0.110		
366.1553	C ₁₈ H ₂₃ NO ₇	4.29	193.2 (\pm 46.1)	174.4 (\pm 31.4)	0.598		
	C ₁₈ H ₂₃ NO ₇	4.98	126.0 (\pm 33.8)	141.9 (\pm 39.5)	0.891		
368.1709	C₁₈H₂₅NO₇	5.24	45.9 (\pm6.7)	65.2 (\pm10.2)	4.81 x 10⁻³	1.42	Usaramine N-oxide
	C ₁₈ H ₂₅ NO ₇	5.62	93.8 (\pm 20.2)	119.3 (\pm 18.8)	0.231		
	C ₁₈ H ₂₅ NO ₇	6.39	550.5 (\pm 54.4)	530.8 (\pm 38.6)	0.953		
370.1866	C ₁₈ H ₂₇ NO ₇	5.94	24.0 (\pm 2.2)	25.8 (\pm 2.8)	0.379		
376.1760	C ₂₀ H ₂₅ NO ₆	9.55	118.5 (\pm 12.1)	118.4 (\pm 14.3)	1.000		
386.1815	C₁₈H₂₇NO₈	3.68	34.6 (\pm9.9)	105.6 (\pm16.0)	2.01 x 10⁻³	3.05	Jacoline N-oxide
392.1709	C ₂₀ H ₂₅ NO ₇	9.77	196.5 (\pm 26.17)	194.0 (\pm 31.4)	0.740		
404.1476	C₁₈H₂₆NO₇Cl	5.22	39.3 (\pm9.73)	122.2 (\pm13.5)	5.87 x 10⁻⁵	3.11	Jaconine N-oxide

Relative concentrations were quantified as the ratio of the analyte signal: internal standard signal per 0.3 mg plant mass.

^a Results that were significant after Benjamini and Hochberg corrections (Benjamini et al. 2001) are displayed in bold.


^b Fold change indicates the concentration increase in roots colonised with AMF when compared to the concentrations observed in control plants.


^c Only given for those PAs that differed significantly between treatment groups.

FIGURE LEGENDS

FIG. 1 a) Orthogonal partial least square-discriminate analysis (OPLS-DA) scores plot of the chemical profiles of root extracts from control and *Rhizophagus irregularis* treated ragwort, b) Partial least square-discriminate analysis (PLS-DA) scores plot of the chemical profiles of leaves from control and *Rhizophagus irregularis* treated ragwort. Open and closed circles represent control and *R. irregularis* treated ragwort plants, respectively. In b), blue symbols represent samples from new leaves and orange symbols those from old leaves and the percentages of explained variation (R^2Y) modelled by the first two latent variables are displayed on the axes. Both plots show datasets from samples profiled in positive ESI MS mode; data from negative ESI mode revealed similar model characteristics (Figure S1).

FIG. 2 Schematic structure of the identified blumenol apocarotenoids associated with *Rhizophagus irregularis* colonisation of ragwort roots.





Metabolite	R1	R2
1	-CH ₂ OH	-OH
2	-CH ₃	-OH
3	-CH ₃	-C ₆ H ₉ O ₇
4	-CH ₂ OH	-OCOCH ₂ CO ₂ H
5	-CH ₃	-OH
6	-CH ₃	-OCOCH ₂ CO-C ₆ H ₉ O ₇
7	-CH ₃	-OCOCH ₂ CO ₂ H



Click here to access/download

Supplementary Material

Hill Hartley et al J Chem Ecol REVISED Supp Ino
Resub.docx