

This is a repository copy of *Observed NO/NO₂ Ratios in the Upper Troposphere Imply Errors in NO-NO₂-O₃ Cycling Kinetics or an Unaccounted NO_x Reservoir.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/130516/>

Version: Accepted Version

Article:

Silvern, R. F., Jacob, D J, Travis, K. R. et al. (10 more authors) (2018) Observed NO/NO₂ Ratios in the Upper Troposphere Imply Errors in NO-NO₂-O₃ Cycling Kinetics or an Unaccounted NO_x Reservoir. *Geophysical Research Letters*. ISSN 0094-8276

<https://doi.org/10.1029/2018GL077728>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Observed NO/NO₂ ratios in the upper troposphere imply errors in NO-NO₂-O₃ cycling kinetics or an unaccounted NO_x reservoir

R. F. Silvern¹, D. J. Jacob^{1,2}, K. R. Travis³, T. Sherwen^{4,5}, M. J. Evans^{4,5}, R. C. Cohen^{6,7}, J. L. Laughner⁶, S. R. Hall⁸, K. Ullmann⁸, J. D. Crouse⁹, P. O. Wennberg^{9,10}, J. Peischl^{11,12}, and I. B. Pollack¹³

¹Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA.

²John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA.

³Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA.

⁴Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, UK.

⁵National Centre for Atmospheric Science, Department of Chemistry, University of York, York, UK.

⁶Department of Chemistry, University of California, Berkeley, CA, USA.

⁷Department of Earth and Planetary Science, University of California, Berkeley, CA, USA.

⁸Atmospheric Chemistry Division, National Center for Atmospheric Research, Boulder, CO, USA.

⁹Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA.

¹⁰Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA, USA.

¹¹Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA.

¹²Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA.

¹³Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado, USA.

Corresponding author: Rachel Silvern (rsilvern@g.harvard.edu)

Key Points:

- Large model overestimates of NO/NO₂ concentration ratios in the upper troposphere imply errors in NO-NO₂-O₃ cycling kinetics or the presence of an unaccounted labile NO_x reservoir.
- The presence of an unaccounted labile NO_x reservoir would affect the NO_x lifetime in the upper troposphere and would suggest unrecognized, likely organic, chemistry.
- Possible error in NO-NO₂-O₃ cycling kinetics would have large implications for global simulations of tropospheric ozone and for satellite retrievals of tropospheric NO₂.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1029/2018GL077728

Abstract

Observations from the SEAC⁴RS aircraft campaign over the Southeast US in August-September 2013 show NO/NO₂ concentration ratios in the upper troposphere that are approximately half of photochemical equilibrium values computed from JPL kinetic data. One possible explanation is the presence of labile NO_x reservoir species, presumably organic, decomposing thermally to NO₂ in the instrument. The NO₂ instrument corrects for this artifact from known labile HNO₄ and CH₃O₂NO₂ NO_x reservoirs. To bridge the gap between measured and simulated NO₂, additional unaccounted labile NO_x reservoir species would have to be present at a mean concentration of ~40 ppt for the SEAC⁴RS conditions (compared with 197 ppt for NO_x). An alternative explanation is error in the low-temperature rate constant for the NO+O₃ reaction (30% 1-σ uncertainty in JPL at 240 K) and/or in the spectroscopic data for NO₂ photolysis (20% 1-σ uncertainty). Resolving this discrepancy is important for understanding global budgets of tropospheric oxidants and for interpreting satellite observations of tropospheric NO₂ columns.

Plain Language Summary

We identify large discrepancies between observed NO/NO₂ ratios and models representing our best understanding of the chemistry controlling NO and NO₂ in the upper troposphere in the Southeast US during August‐September 2013. We suggest that either unrecognized chemistry or errors in modeled cycling between NO, NO₂ and O₃ could explain this discrepancy. Either explanation will have important implications for global tropospheric chemistry and for the interpretation of satellite observations of NO₂.

1. Introduction

Nitrogen oxide radicals (NO_x = NO + NO₂) are emitted by anthropogenic sources (fuel combustion) and natural sources (lightning, soils, fires). Anthropogenic emissions degrade surface air quality by producing ozone and nitrate particulate matter, and also affect ecosystems through nitrogen deposition. On a global scale, NO_x increases the concentration of tropospheric oxidants (ozone and OH) with complicated implications for climate forcing (Wild et al., 2001). NO_x in the upper troposphere and the associated cycling between NO and NO₂ is of particular importance for production of tropospheric ozone and OH (Murray et al., 2013; Newsome & Evans, 2017; Ridley et al., 2017). Recent observations from the SEAC⁴RS

aircraft campaign over the Southeast US in August-September 2013 show much lower NO/NO₂ ratios in the upper troposphere than expected from models (Travis et al., 2016). Here we suggest possible explanations for this discrepancy and discuss the implications for global tropospheric chemistry and for the interpretation of satellite NO₂ data.

The SEAC⁴RS observations over the Southeast US show NO_x concentrations averaging 0.20 ppb in the upper troposphere above 8 km (0.11 ppb as NO and 0.09 ppb as NO₂), as compared to 0.37 ppb in the boundary layer below 2 km (0.06 ppb NO and 0.31 ppb NO₂), and much lower concentrations (averaging less than 0.07 ppb) in the middle troposphere between 2 and 8 km (Travis et al., 2016). Such a “C-shaped” profile reflects influences from fuel combustion in the boundary layer and lightning in the upper troposphere (Bertram et al., 2007; Hudman et al., 2007; Huntrieser et al., 2002; Pickering et al., 1998).

The mean observed daytime NO/NO₂ ratios in SEAC⁴RS were 0.44 mol mol⁻¹ in the boundary layer and 1.4 mol mol⁻¹ in the upper troposphere, while the corresponding ratios in the GEOS-Chem chemical transport model sampled along the flight tracks were 0.33 mol mol⁻¹ in the boundary layer and 3.3 mol mol⁻¹ in the upper troposphere (Travis et al., 2016).

The NO/NO₂ ratio in the model increases rapidly with altitude because of the strong temperature dependence of the NO+O₃ reaction (Burkholder et al., 2015), but in the observations this increase is much less.

NO₂ measurements in the upper troposphere are prone to positive interferences from inlet decomposition of thermally unstable compounds including HNO₄, CH₃O₂NO₂, and other organic nitrates (Bradshaw et al., 1999; Browne et al., 2011; Murphy et al., 2004; Nault et al., 2015; Reed et al., 2016). The Berkeley thermal-dissociation laser-induced fluorescence (TD-LIF) instrument used in SEAC⁴RS (Thornton et al., 2000; Day et al., 2002; Wooldridge et al., 2010) was specifically designed to minimize and correct for these interferences. Inlet

residence time is only 0.23 seconds and the NO_2 measurement is corrected for partial thermal dissociation of HNO_4 (0-11%) and $\text{CH}_3\text{O}_2\text{NO}_2$ (0-21%) with a calibration accuracy of 5% (Nault et al., 2015). HNO_4 and $\text{CH}_3\text{O}_2\text{NO}_2$ are independently measured with calibration accuracies of 15%, and the $\text{CH}_3\text{O}_2\text{NO}_2$ measurement has an overall uncertainty of 40%, mainly due to uncertainty in the thermal decomposition rate constant (Nault et al., 2015). An independent NO_2 measurement made by chemiluminescence (Pollack et al., 2010) during SEAC⁴RS was biased high compared to the TD-LIF measurement, likely due to interferences from $\text{CH}_3\text{O}_2\text{NO}_2$ and HNO_4 (Travis et al., 2016). The Berkeley TD-LIF NO_2 measurements in the upper troposphere have been used in previous work to interpret NO_x chemistry (Nault et al., 2016), lightning NO_x emissions (Nault et al., 2017), and satellite observations of NO_2 columns (Choi et al., 2014; Laughner and Cohen, 2017).

The model-measurement discrepancy in NO/NO_2 partitioning in the upper troposphere has consequences not only for tropospheric chemistry but also for interpreting solar backscatter NO_2 observations from satellites. It is generally assumed that the tropospheric NO_2 column retrieved from satellites is mainly contributed by the boundary layer (Lamsal et al., 2014; Laughner et al., 2016; Martin et al., 2002) and can therefore be related to local NO_x emissions (Martin et al., 2003). However, the NO_2 vertical profiles from SEAC⁴RS imply a 35-50% contribution of the upper troposphere to the NO_2 tropospheric column observed from satellite (Travis et al., 2016), because sensitivity of backscattered solar radiation to NO_2 increases by a factor of 3 from the surface to the upper troposphere (Martin et al., 2002). Better understanding of this upper tropospheric NO_2 is crucial to the use of satellite data for estimating surface NO_x emissions.

Previous work starting in the 1990s has found varied levels of agreement between NO/NO₂ ratios observed from aircraft and model photochemical equilibrium computed from local conditions. Early work in the lower stratosphere found models to be too high by 20-30% (Del Negro et al., 1999; Jaeglé et al., 1994; Sen et al., 1998; Cohen et al., 2000). Observed NO/NO₂ ratios in the upper troposphere during the PEM-West A campaign over the tropical Pacific were three times lower than model predictions, which was attributed to NO₂ measurement interferences (Crawford et al., 1996), although Davis et al. (1996) also hypothesized a role of halogen chemistry. Bradshaw et al. (1999) found model agreement with observations to within 30% in the upper troposphere over the tropical Pacific using an improved NO₂ instrument that avoided positive interferences through short inlet residence time. More recent model studies of the upper troposphere have again found an overestimate of observed NO/NO₂ ratios (Travis et al., 2016; Williams et al., 2017), and attributed it to underestimate of peroxy radicals converting NO to NO₂.

2. NO-NO₂ cycling in the upper troposphere during SEAC⁴RS

According to current understanding, the NO/NO₂ ratio in the daytime upper troposphere is determined by rapid chemical cycling through the reactions in Table 1. Here we calculated the mean rates of individual reactions along the SEAC⁴RS flight tracks over the Southeast US by applying the recommended JPL rate constants (Burkholder et al., 2015) to aircraft measurements of species concentrations (NO, NO₂, O₃), NO₂ photolysis rate constant (J_{NO_2}), temperature, and pressure, together with peroxy and halogen radical concentrations computed by the GEOS-Chem model (Sherwen et al., 2016, 2017; Travis et al., 2016) along the flight tracks. We exclude data outside the 9-15 hour solar time window and in stratospheric air ($[\text{O}_3]/[\text{CO}] > 1.25 \text{ mol mol}^{-1}$). We focus on the Southeast US because it

accounts for most of the SEAC⁴RS flights and represents a relatively homogeneous environment.

The calculated rates in Table 1 indicate that conversion of NO to NO₂ in the upper troposphere is mainly by reaction with ozone (75%). Reactions with peroxy radicals contribute 18%, and reactions with halogen radicals (BrO, IO, and ClO) contribute 7%. The total calculated rate of conversion of NO to NO₂ has an aggregated uncertainty of 32% and balances only half of the NO₂ photolysis rate, which has an uncertainty of 21%. This represents a significant discrepancy, such that a model using JPL kinetics would overestimate the NO/NO₂ ratio observed in SEAC⁴RS.

Figure 1 shows the median observed vertical profiles of the principal variables relevant to NO/NO₂ cycling, along with the corresponding values simulated by the standard GEOS-Chem model along the aircraft flight tracks as in Travis et al. (2016). The NO/NO₂ ratio in the model is over twice that measured in the upper troposphere above 8 km. The bias is systematic over the frequency distribution of the observations (Figure 2) and also extends to stratospherically influenced air ($[O_3]/[CO] > 1.25 \text{ mol mol}^{-1}$), although SEAC⁴RS did not sample the actual stratosphere (maximum ozone concentration was 125 ppb). Travis et al. (2016) attributed the model bias in the NO/NO₂ ratio to an underestimate of peroxy radicals, but that underestimate would have to be a factor of 5 in order to close the budget of Table 1. This is incompatible with the SEAC⁴RS observations for H₂O₂, which is produced by self-reaction of HO₂ and is thus a sensitive proxy of HO₂ concentrations. Observed H₂O₂ concentrations are 30% higher than GEOS-Chem (Figure 1) but would be grossly overestimated if HO₂ concentrations were increased by a factor of 5. Similarly, simulated BrO concentrations would have to be underestimated by a factor of 20 in order to correct the model bias in the NO/NO₂ ratio. This would be grossly inconsistent with observations (Schmidt et al., 2016; Shah et al., 2016; Sherwen et al., 2016).

We see from Figure 1 that there is no systematic model bias in ozone, temperature, or J_{NO_2} that would explain an error in the NO/NO₂ ratio. The J_{NO_2} observations during SEAC⁴RS are from spectrally resolved actinic flux measurements (Shetter et al., 2003), converted to photolysis frequencies using the same temperature-dependent JPL spectroscopic data (absorption cross-sections and quantum yields) as in GEOS-Chem. These spectroscopic data are tabulated by JPL as a function of wavelength at 220 and 294 K, and are interpolated linearly for intermediate temperatures. 90% of J_{NO_2} photolysis is contributed by wavelengths shorter than 398 nm for which the quantum yield is unity. The estimated JPL uncertainty on J_{NO_2} is 20% with no temperature dependence (Sander et al., 2011), though laboratory studies show better agreement at surface temperatures (Orphal, 2003; Shetter et al., 2003). Observed photolysis frequencies of other relevant species (O₃, HCHO, H₂O₂, HNO₃, PAN, CH₃OOH) also agree with GEOS-Chem values to within 1-15% throughout the troposphere.

One possible explanation for the apparent departure of the NO/NO₂ ratio from photochemical equilibrium would be a positive bias in the NO₂ measurement. This could occur if there was an unrecognized labile reservoir of NO₂ (other than HNO₄ or CH₃O₂NO₂) decomposing in the instrument inlet, or if the correction for HNO₄ or CH₃O₂NO₂ was inadequate. We find that that this missing reservoir (likely organic) would need to be present at a mean concentration of at least 40 ppt in the upper troposphere in order to fit the model NO/NO₂ ratios, assuming 100% decomposition to NO₂ inside the instrument. For comparison, the correction to the NO₂ measurement from the decomposition of CH₃O₂NO₂ was 0-23 ppt (0-21% of NO₂ at temperatures less than 240 K) and 0-20 ppt HNO₄ (0-11%) during SEAC⁴RS (Nault et al., 2015). The high-flow pump to minimize the influence from these reservoirs malfunctioned in the first five SEAC⁴RS flights (Aug 6-16) but these flights were either not over the Southeast US or not in the upper troposphere and are not included in our analysis.

A significant uncertainty in the $\text{CH}_3\text{O}_2\text{NO}_2$ correction is the thermal decomposition rate constant, which has a JPL 1- σ uncertainty of 30%. Considering a cabin air temperature of 300 K, an exterior pressure of 230 hPa, and an inlet residence time of 0.23 s, a 30% increase in the $\text{CH}_3\text{O}_2\text{NO}_2$ thermal decomposition rate would double the corresponding correction to the NO_2 measurement from 0-23 ppt to 0-46 ppt. The effect would be at most 23 ppt at the upper end of the range and is not sufficient to correct the NO/NO_2 ratio. GEOS-Chem underestimates $\text{CH}_3\text{O}_2\text{NO}_2$ in the upper troposphere (18 ± 19 ppt modeled, 124 ± 98 ppt observed), reflecting in part the model underestimate of NO_2 but also suggesting missing organic chemistry. A faster thermal decomposition rate for $\text{CH}_3\text{O}_2\text{NO}_2$ would exacerbate the model underestimate.

Previous work has postulated missing organic chemistry in the upper troposphere to explain observations of volatile organic compounds (VOCs) including methanol (Jacob et al., 2005), acetaldehyde (Millet et al., 2010) and glyoxal (Volkamer et al., 2015). Aumont et al. (2005) and Mouchel-Vallon et al. (2013) showed how explicit VOC mechanisms produce a cascade of oxidation products globally that are not tracked in models. Bradshaw et al. (1999) did not need to invoke an unknown NO_x reservoir to reconcile their model with observations over the tropical Pacific, but the Southeast US may be a more propitious environment for VOC oxidation products to be lifted to the upper troposphere by deep convection (Li et al., 2005).

An alternative explanation for the apparent NO/NO_2 departure from photochemical equilibrium would be error in the kinetic data used to compute that equilibrium. Figure 3 shows the IUPAC and JPL recommendations for the temperature dependence of the $\text{NO}+\text{O}_3$ rate constant k_1 , along with the individual laboratory data that went into these recommendations. IUPAC recommends $k_1 = 2.07 \times 10^{-12} \exp[-1400/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ with a 1- σ uncertainty of ± 200 K for E/R and a 1- σ uncertainty of 8% for k_1 at 298 K (Atkinson et

al., 2004). JPL recommends $k_1 = 3.00 \times 10^{-12} \exp[-1500/T]$, again with 1- σ uncertainty of ± 200 K for E/R and a 1- σ uncertainty of 10% for k_1 at 298 K. The IUPAC and JPL rate expressions agree to within 4% over the 220-300 K temperature range. For a typical upper tropospheric temperature of 220-240 K the implied 1- σ uncertainty for the JPL rate is 30-40%. However, several studies have suggested a departure from Arrhenius behavior at low temperatures (Figure 3; Birks et al., 1976; Borders & Birks, 1982; Cohen et al., 2000; Michael et al., 1981).

The purple curves in Figures 1 and 2 show the effects in the GEOS-Chem simulation of decreasing J_{NO_2} by 20%, and decreasing the activation energy for k_1 by 400 K (2σ) relative to the JPL recommendation so that k_1 increases by a factor 1.4 on average in the upper troposphere, while remaining at the JPL recommended value at 298 K through adjustment of the pre-exponential factor. The resulting NO/NO₂ ratio in the upper troposphere decreases by 40% from the standard simulation and Figure 2 shows that the variance in the modeled NO/NO₂ ratio decreases by half, becoming more consistent with the observations. One could match the observations if there was in addition a 15 ppt positive bias in the NO₂ measurement due to an unaccounted labile NO_x reservoir.

We examined whether the variability of the observed NO/NO₂ ratio in the SEAC⁴RS data set could test the above corrections. For this purpose we used 10-minute observations of the ratio along the flight tracks at 8-12 km altitude and correlated them to the local values of the photoequilibrium constant $J_{\text{NO}_2}(h\nu)/k_1(T)[\text{O}_3]$ for NO-NO₂-O₃ cycling where the UV actinic fluxes ($h\nu$), temperatures (T), and [O₃] are taken from the observations. Results are shown in Figure 4. When using JPL values for $J_{\text{NO}_2}(h\nu)$ and $k_1(T)$, we find a significant ($p < 0.01$) correlation coefficient $r = 0.37$ and a reduced-major-axis (RMA) regression slope $S = 4.0$. The relatively low correlation coefficient can be attributed to noise and high-frequency variability in the observations. Reducing J_{NO_2} by 20% and increasing the low-temperature k_1

as described above improves the slope ($S = 2.3$) while not affecting the correlation coefficient ($r = 0.38$). Excluding the 15% of the data with $[O_3] < 40$ ppb increases the correlation coefficient to $r = 0.48-0.49$. Those conditions were associated with particularly high J_{NO_2} (reflecting clouds below) and enhancements in CO and HCHO indicative of recent convective influence (Barth et al., 2015; Fried et al., 2016; Snow et al., 2007).

3. Implications

The apparent departure of the NO/NO₂ concentration ratio from photochemical equilibrium in upper tropospheric observations cannot be explained by missing radicals converting NO to NO₂, as proposed in previous work, because the required radical concentrations would be far in excess of observational constraints. It must be due either to an unaccounted labile NO_x reservoir acting as positive interference on the NO₂ measurement, and/or to significant errors in the kinetic data for NO-NO₂-O₃ photochemical cycling at low temperatures. Either of these possibilities have important implications for upper tropospheric chemistry.

The NO₂ measurement specifically excludes interferences from HNO₄ and CH₃O₂NO₂, but other labile NO_x reservoirs could potentially be measured as NO₂ following thermal decomposition in the instrument. The presence of such a reservoir at a concentration of 40 ppt, as needed to explain the observed NO/NO₂ ratios, would increase the effective lifetime of NO_x by 20% under the SEAC⁴RS conditions. More importantly, it would likely imply organic chemistry missing from the models, as also suggested by observations of acetaldehyde and glyoxal in the upper troposphere (Millet et al., 2010; Volkamer et al., 2015) and by the large model underestimate of CH₃O₂NO₂ in SEAC⁴RS. .

The rate constants J_{NO_2} and k_1 involved in NO-NO₂-O₃ photochemical cycling have

relatively small uncertainties in kinetic assessments, and even then are found to be major sources of uncertainty in model simulations of tropospheric oxidants (Bergin et al., 1999; Newsome and Evans, 2017; Ridley et al., 2017; Vuilleumier et al., 1997). When the low-temperature $\text{NO} + \text{O}_3$ reaction rate constant ($1.4k_1$) and NO_2 photolysis frequency ($J_{\text{NO}_2} -20\%$) are adjusted in GEOS-Chem within these uncertainties to improve the simulation of the NO/NO_2 ratio in the SEAC⁴RS upper tropospheric data, we find that simulated ozone decreases by 7 ppb at 8-12 km altitude. This degrades the previously successful simulation of upper tropospheric ozone in the standard model (Travis et al., 2016), however that simulation overestimated the NO concentration (Figure 1).

Improved understanding of the contribution of the upper troposphere to the tropospheric NO_2 column also has implications for retrieving and interpreting NO_2 observations from satellites. Spectral fitting of the satellite data in and around the NO_2 absorption bands measures the slant column of NO_2 along the light path. Conversion of this slant column to the actual vertical tropospheric column requires removal of the stratospheric contribution, followed by division by an air mass factor (AMF) dependent on the vertical distribution of tropospheric NO_2 (Martin et al., 2002). The NASA operational retrieval for the OMI satellite instrument (Krotkov et al., 2017) assumes NO_2 vertical profiles from the GMI model (Lamsal et al., 2014) that greatly underestimate NO_2 concentrations in the upper troposphere as observed by SEAC⁴RS. The mean AMF over the Southeast US in August-September 2013 is 1.28 using vertical distributions from the NASA operational retrieval but 1.67 when using the median observed profile in Figure 1. This implies that the OMI operational retrieval for NO_2 may be 30% too high. Laughner and Cohen (2017) show that inclusion of lightning NO_x in the upper troposphere to match DC3 observations of NO_2 increases the OMI AMF by 34% for summertime, further demonstrating the importance of accurately modeling and measuring NO_x in the upper troposphere for the interpretation of

satellite NO₂ measurements.

In conclusion, models significantly overestimate recent observations of the NO/NO₂ ratio in the upper troposphere. This cannot be easily explained by known labile NO_x reservoirs (HNO₄, CH₃O₂NO₂) interfering with the NO₂ measurement. It implies either error in current recommendations for NO-NO₂-O₃ cycling kinetics or the presence of a missing labile NO_x reservoir, likely organic. Either explanation would have important implications for our understanding of tropospheric oxidants and/or the interpretation of satellite NO₂ measurements.

Acknowledgments. We thank Tom Ryerson for his measurements of NO and O₃ from the NOAA NO_yO₃ instrument. This work was supported by the NASA Earth Science Division. SEAC⁴RS airborne measurements are available from the NASA LaRC Airborne Science Data for Atmospheric Composition (<http://www-air.larc.nasa.gov/cgi-bin/ArcView/seac4rs>). OMI NO₂ observations are available from the NASA Goddard Earth Sciences Data and Information Services Center (https://aura.gesdisc.eosdis.nasa.gov/data/Aura_OMI_Level2/OMNO2.003/).

References

- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., . . . Troe, J. (2004). IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation. *Atmospheric Chemistry and Physics*, 4, 1461.
- Aumont, B., Szopa, S., & Madronich, S. (2005). Modelling the evolution of organic carbon during its gas-phase tropospheric oxidation: development of an explicit model based on a self generating approach. *Atmospheric Chemistry and Physics*, 5, 2497-2517. doi:10.5194/acp-5-2497-2005
- Barth, M. C., Cantrell, C. A., Brune, W. H., Rutledge, S. A., Crawford, J. H., Huntrieser, H., . . . Ziegler, C. (2015). The Deep Convective Clouds and Chemistry (DC3) field campaign. *Bulletin of the American Meteorological Society*, 96(8), 1281-1309. doi:10.1175/bams-d-13-00290.1
- Bergin, M. S., Noblet, G. S., Petrin, K., Dhieux, J. R., Milford, J. B., & Harley, R. A. (1999). Formal uncertainty analysis of a Lagrangian photochemical air pollution model. *Environmental Science & Technology*, 33(7), 1116-1126. doi:10.1021/es980749y

- Bertram, T. H., Perring, A. E., Wooldridge, P. J., Crounse, J. D., Kwan, A. J., Wennberg, P. O., . . . Cohen, R. C. (2007). Direct measurements of the convective recycling of the upper troposphere. *Science*, 315(5813), 816-820. doi:10.1126/science.1134548
- Birks, J. W., Shoemaker, B., Leck, T. J., & Hinton, D. M. (1976). Studies of reactions of importance in the stratosphere. 1. Reaction of nitric oxide with ozone. *Journal of Chemical Physics*, 65(12), 5181-5185. doi:10.1063/1.433059
- Borders, R. A., & Birks, J. W. (1982). High-Precision Measurements of Activation Energies over Small Temperature Intervals: Curvature in the Arrhenius Plot for the Reaction $\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$. *Journal of Physical Chemistry*, 86(17), 3295-3302. doi:10.1021/j100214a007
- Bradshaw, J., Davis, D., Crawford, J., Chen, G., Shetter, R., Muller, M., . . . Sandholm, S. (1999). Photofragmentation two-photon laser-induced fluorescence detection of NO_2 and NO : Comparison of measurements with model results based on airborne observations during PEM-Tropics A. *Geophysical Research Letters*, 26(4), 471-474. doi:10.1029/1999gl900015
- Browne, E. C., Perring, A. E., Wooldridge, P. J., Apel, E., Hall, S. R., Huey, L. G., . . . Cohen, R. C. (2011). Global and regional effects of the photochemistry of $\text{CH}_3\text{O}_2\text{NO}_2$: evidence from ARCTAS. *Atmospheric Chemistry and Physics*, 11(9), 4209-4219. doi:10.5194/acp-11-4209-2011
- Burkholder, J. B., Sander, S. P., Abbatt, J., Barker, J. R., Huie, R. E., Kolb, C. E., . . . Wine, P. H. (2015). Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 18. <http://jpldataeval.jpl.nasa.gov/>
- Choi, S., Joiner, J., Choi, Y., Duncan, B. N., Vasilkov, A., Krotkov, N., & Bucsela, E. (2014). First estimates of global free-tropospheric NO_2 abundances derived using a cloud-slicing technique applied to satellite observations from the Aura Ozone Monitoring Instrument (OMI). *Atmospheric Chemistry and Physics*, 14(19), 10565-10588. doi:10.5194/acp-14-10565-2014
- Cohen, R. C., Perkins, K. K., Koch, L. C., Stimpfle, R. M., Wennberg, P. O., Hanisco, T. F., . . . Bui, T. P. (2000). Quantitative constraints on the atmospheric chemistry of nitrogen oxides: An analysis along chemical coordinates. *Journal of Geophysical Research-Atmospheres*, 105(D19), 24283-24304. doi:10.1029/2000jd900290
- Crawford, J., Davis, D., Chen, G., Bradshaw, J., Sandholm, S., Gregory, G., . . . Rodriguez, J. (1996). Photostationary state analysis of the NO_2 - NO system based on airborne observations from the western and central North Pacific. *Journal of Geophysical Research-Atmospheres*, 101(D1), 2053-2072. doi:10.1029/95jd02201
- Crounse, J. D., McKinney, K. A., Kwan, A. J., & Wennberg, P. O. (2006). Measurement of gas-phase hydroperoxides by chemical ionization mass spectrometry. *Analytical Chemistry*, 78(19), 6726-6732. doi:10.1021/ac0604235
- Davis, D., Crawford, J., Liu, S., McKeen, S., Bandy, A., Thornton, D., . . . Blake, D. (1996). Potential impact of iodine on tropospheric levels of ozone and other critical oxidants. *Journal of Geophysical Research-Atmospheres*, 101(D1), 2135-2147. doi:10.1029/95jd02727
- Day, D. A., Wooldridge, P. J., Dillon, M. B., Thornton, J. A., & Cohen, R. C. (2002). A thermal dissociation laser-induced fluorescence instrument for in situ detection of NO_2 , peroxy nitrates, alkyl nitrates, and HNO_3 . *Journal of Geophysical Research-Atmospheres*, 107(D5-6), 14. doi:10.1029/2001jd000779
- Del Negro, L. A., Fahey, D. W., Gao, R. S., Donnelly, S. G., Keim, E. R., Neuman, J. A., . . . Bui, T. P. (1999). Comparison of modeled and observed values of NO_2 and JNO_2 during the Photochemistry of Ozone Loss in the Arctic Region in Summer

- (POLARIS) mission. *Journal of Geophysical Research-Atmospheres*, 104(D21), 26687-26703. doi:10.1029/1999jd900246
- Fried, A., Barth, M. C., Bela, M., Weibring, P., Richter, D., Walega, J., . . . Woods, S. (2016). Convective transport of formaldehyde to the upper troposphere and lower stratosphere and associated scavenging in thunderstorms over the central United States during the 2012DC3 study. *Journal of Geophysical Research-Atmospheres*, 121(12), 7430-7460. doi:10.1002/2015jd024477
- Hudman, R. C., Jacob, D. J., Turquety, S., Leibensperger, E. M., Murray, L. T., Wu, S., . . . Wooldridge, P. J. (2007). Surface and lightning sources of nitrogen oxides over the United States: Magnitudes, chemical evolution, and outflow. *Journal of Geophysical Research-Atmospheres*, 112(D12), 14. doi:10.1029/2006jd007912
- Huntrieser, H., Feigl, C., Schlager, H., Schroder, F., Gerbig, C., van Velthoven, P., . . . Schumann, U. (2002). Airborne measurements of NO_x, tracer species, and small particles during the European lightning nitrogen oxides experiment. *Journal of Geophysical Research-Atmospheres*, 107(D11), 26. doi:10.1029/2000jd000209
- Jacob, D. J., Field, B. D., Li, Q. B., Blake, D. R., de Gouw, J., Warneke, C., . . . Guenther, A. (2005). Global budget of methanol: Constraints from atmospheric observations. *Journal of Geophysical Research-Atmospheres*, 110(D8), 17. doi:10.1029/2004jd005172
- Jaeglé, L., Webster, C. R., May, R. D., Fahey, D. W., Woodbridge, E. L., Keim, E. R., . . . Pfister, L. (1994). In-situ measurements of the NO₂/NO ratio for testing atmospheric photochemical models. *Geophysical Research Letters*, 21(23), 2555-2558. doi:10.1029/94gl02717
- Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., Marchenko, S. V., Bucsela, E. J., . . . Zara, M. (2017). The version 3 OMI NO₂ standard product. *Atmospheric Measurement Techniques*, 10(9), 3133-3149. doi:10.5194/amt-10-3133-2017
- Lamsal, L. N., Krotkov, N. A., Celarier, E. A., Swartz, W. H., Pickering, K. E., Bucsela, E. J., . . . Knepp, T. N. (2014). Evaluation of OMI operational standard NO₂ column retrievals using in situ and surface-based NO₂ observations. *Atmospheric Chemistry and Physics*, 14(21), 11587-11609. doi:10.5194/acp-14-11587-2014
- Laughner, J. L., & Cohen, R. C. (2017). Quantification of the effect of modeled lightning NO₂ on UV-visible air mass factors. *Atmospheric Measurement Techniques*, 10(11), 4403-4419. doi:10.5194/amt-10-4403-2017
- Laughner, J. L., Zare, A., & Cohen, R. C. (2016). Effects of daily meteorology on the interpretation of space-based remote sensing of NO₂. *Atmospheric Chemistry and Physics*, 16(23), 15247-15264. doi:10.5194/acp-16-15247-2016
- Li, Q. B., Jacob, D. J., Park, R., Wang, Y. X., Heald, C. L., Hudman, R., . . . Evans, M. (2005). North American pollution outflow and the trapping of convectively lifted pollution by upper-level anticyclone. *Journal of Geophysical Research-Atmospheres*, 110(D10), 18. doi:10.1029/2004jd005039
- Lippmann, H. H., Jesser, B., & Schurath, U. (1980). The rate constant of NO+O₃ --> NO₂+O₂ in the temperature range of 283-443 K. *International Journal of Chemical Kinetics*, 12(8), 547-554. doi:10.1002/kin.550120805
- Martin, R. V., Chance, K., Jacob, D. J., Kurosu, T. P., Spurr, R. J. D., Bucsela, E., . . . Koelemeijer, R. B. A. (2002). An improved retrieval of tropospheric nitrogen dioxide from GOME. *Journal of Geophysical Research-Atmospheres*, 107(D20), 21. doi:10.1029/2001jd001027
- Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Palmer, P. I., & Evans, M. J. (2003). Global inventory of nitrogen oxide emissions constrained by space-based

- observations of NO₂ columns. *Journal of Geophysical Research-Atmospheres*, 108(D17), 12. doi:10.1029/2003jd003453
- Michael, J. V., Allen, J. E., & Brobst, W. D. (1981). Temperature dependence of the NO + O₃ reaction rate from 195 to 369 K. *Journal of Physical Chemistry*, 85(26), 4109-4117. doi:10.1021/j150626a032
- Millet, D. B., Guenther, A., Siegel, D. A., Nelson, N. B., Singh, H. B., de Gouw, J. A., . . . Barkley, M. (2010). Global atmospheric budget of acetaldehyde: 3-D model analysis and constraints from in-situ and satellite observations. *Atmospheric Chemistry and Physics*, 10(7), 3405-3425. doi:10.5194/acp-10-3405-2010
- Moonen, P. C., Cape, J. N., Storeton-West, R. L., & McCOLM, R. (1998). Measurement of the NO + O₃ Reaction Rate at Atmospheric Pressure Using Realistic Mixing Ratios. *Journal of Atmospheric Chemistry*, 29(3), 299-314. doi:10.1023/a:1005936016311
- Mouchel-Vallon, C., Brauer, P., Camredon, M., Valorso, R., Madronich, S., Herrmann, H., & Aumont, B. (2013). Explicit modeling of volatile organic compounds partitioning in the atmospheric aqueous phase. *Atmospheric Chemistry and Physics*, 13(2), 1023-1037. doi:10.5194/acp-13-1023-2013
- Murphy, J. G., Thornton, J. A., Wooldridge, P. J., Day, D. A., Rosen, R. S., Cantrell, C., . . . Cohen, R. C. (2004). Measurements of the sum of HO₂NO₂ and CH₃O₂NO₂ in the remote troposphere. *Atmospheric Chemistry and Physics*, 4, 377-384.
- Murray, L. T., Logan, J. A., & Jacob, D. J. (2013). Interannual variability in tropical tropospheric ozone and OH: The role of lightning. *Journal of Geophysical Research-Atmospheres*, 118(19), 11468-11480. doi:10.1002/jgrd.50857
- Nault, B. A., Garland, C., Pusede, S. E., Wooldridge, P. J., Ullmann, K., Hall, S. R., & Cohen, R. C. (2015). Measurements of CH₃O₂NO₂ in the upper troposphere. *Atmospheric Measurement Techniques*, 8(2), 987-997. doi:10.5194/amt-8-987-2015
- Nault, B. A., Garland, C., Wooldridge, P. J., Brune, W. H., Campuzano-Jost, P., Crouse, J. D., . . . Cohen, R. C. (2016). Observational Constraints on the Oxidation of NO_x in the Upper Troposphere. *Journal of Physical Chemistry A*, 120(9), 1468-1478. doi:10.1021/acs.jpca.5b07824
- Nault, B. A., Laughner, J. L., Wooldridge, P. J., Crouse, J. D., Dibb, J., Diskin, G., . . . Cohen, R. C. (2017). Lightning NO_x Emissions: Reconciling Measured and Modeled Estimates With Updated NO_x Chemistry. *Geophysical Research Letters*, 2017GL074436. doi:10.1002/2017GL074436
- Newsome, B., & Evans, M. (2017). Impact of uncertainties in inorganic chemical rate constants on tropospheric composition and ozone radiative forcing. *Atmospheric Chemistry and Physics*, 17, 14333-14352. doi:10.5194/acp-17-14333-2017
- Orphal, J. (2003). A critical review of the absorption cross-sections of O₃ and NO₂ in the ultraviolet and visible. *Journal of Photochemistry and Photobiology a-Chemistry*, 157(2-3), 185-209. doi:10.1016/s1010-6030(03)00061-3
- Pickering, K. E., Wang, Y. S., Tao, W. K., Price, C., & Muller, J. F. (1998). Vertical distributions of lightning NO_x for use in regional and global chemical transport models. *Journal of Geophysical Research-Atmospheres*, 103(D23), 31203-31216. doi:10.1029/98jd02651
- Pollack, I. B., Lerner, B. M., & Ryerson, T. B. (2010). Evaluation of ultraviolet light-emitting diodes for detection of atmospheric NO₂ by photolysis - chemiluminescence. *Journal of Atmospheric Chemistry*, 65(2-3), 111-125. doi:10.1007/s10874-011-9184-3
- Ray, G. W., & Watson, R. T. (1981). Kinetics of the reaction NO + O₃ --> NO₂ + O₂ from 212 to 422 K. *The Journal of Physical Chemistry*, 85(12), 1673-1676.
- Reed, C., Evans, M. J., Di Carlo, P., Lee, J. D., & Carpenter, L. J. (2016). Interferences in photolytic NO₂ measurements: explanation for an apparent missing oxidant?

Atmospheric Chemistry and Physics, 16(7), 4707-4724. doi:10.5194/acp-16-4707-2016

- Ridley, D., Cain, M., Methven, J., & Arnold, S. (2017). Sensitivity of tropospheric ozone to chemical kinetic uncertainties in air masses influenced by anthropogenic and biomass burning emissions. *Geophysical Research Letters*.
- Ryerson, T. B., Buhr, M. P., Frost, G. J., Goldan, P. D., Holloway, J. S., Hubler, G., . . . Fehsenfeld, F. C. (1998). Emissions lifetimes and ozone formation in power plant plumes. *Journal of Geophysical Research-Atmospheres*, 103(D17), 22569-22583. doi:10.1029/98jd01620
- Ryerson, T. B., Williams, E. J., & Fehsenfeld, F. C. (2000). An efficient photolysis system for fast-response NO₂ measurements. *Journal of Geophysical Research-Atmospheres*, 105(D21), 26447-26461. doi:10.1029/2000jd900389
- Sander, S. P., Abbatt, J., Barker, J. R., Burkholder, J. B., Friedl, R. R., G., D. M., Huie, R. E., Kolb, C. E., Kurylo, M. J., & Moortgat, G. K., Orkin, V. L., and Wine, P. H. (2011). *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies*, Evaluation No. 17.
- Schmidt, J. A., Jacob, D. J., Horowitz, H. M., Hu, L., Sherwen, T., Evans, M. J., . . . Volkamer, R. (2016). Modeling the observed tropospheric BrO background: Importance of multiphase chemistry and implications for ozone, OH, and mercury. *Journal of Geophysical Research-Atmospheres*, 121(19), 11819-11835. doi:10.1002/2015jd024229
- Sen, B., Toon, G. C., Osterman, G. B., Blavier, J. F., Margitan, J. J., Salawitch, R. J., & Yue, G. K. (1998). Measurements of reactive nitrogen in the stratosphere. *Journal of Geophysical Research-Atmospheres*, 103(D3), 3571-3585. doi:10.1029/97jd02468
- Shah, V., Jaegle, L., Gratz, L. E., Ambrose, J. L., Jaffe, D. A., Selin, N. E., . . . Mauldin, R. L. (2016). Origin of oxidized mercury in the summertime free troposphere over the southeastern US. *Atmospheric Chemistry and Physics*, 16(3), 1511-1530. doi:10.5194/acp-16-1511-2016s
- Sherwen, T., Evans, M. J., Sommariva, R., Hollis, L. D. J., Ball, S. M., Monks, P. S., . . . Bloss, W. J. (2017). Effects of halogens on European air-quality. *Faraday Discussions*, 200, 75-100. doi:10.1039/c7fd00026j
- Sherwen, T., Schmidt, J. A., Evans, M. J., Carpenter, L. J., Grossmann, K., Eastham, S. D., . . . Ordonez, C. (2016). Global impacts of tropospheric halogens (Cl, Br, I) on oxidants and composition in GEOS-Chem. *Atmospheric Chemistry and Physics*, 16(18), 12239-12271. doi:10.5194/acp-16-12239-2016
- Shetter, R. E., Junkermann, W., Swartz, W. H., Frost, G. J., Crawford, J. H., Lefer, B. L., . . . van Weele, M. (2003). Photolysis frequency of NO₂: Measurement and modeling during the International Photolysis Frequency Measurement and Modeling Intercomparison (IPMMI). *Journal of Geophysical Research-Atmospheres*, 108(D16), 15. doi:10.1029/2002jd002932
- Snow, J. A., Heikes, B. G., Shen, H. W., O'Sullivan, D. W., Fried, A., & Walega, J. (2007). Hydrogen peroxide, methyl hydroperoxide, and formaldehyde over North America and the North Atlantic. *Journal of Geophysical Research-Atmospheres*, 112(D12), 17. doi:10.1029/2006jd007746
- Thornton, J. A., Wooldridge, P. J., & Cohen, R. C. (2000). Atmospheric NO₂: In situ laser-induced fluorescence detection at parts per trillion mixing ratios. *Analytical Chemistry*, 72(3), 528-539. doi:10.1021/ac9908905
- Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., . . . Zhou, X. L. (2016). Why do models overestimate surface ozone in the Southeast United States?

Atmospheric Chemistry and Physics, 16(21), 13561-13577. doi:10.5194/acp-16-13561-2016

Volkamer, R., Baidar, S., Campos, T. L., Coburn, S., DiGangi, J. P., Dix, B., . . . Romashkin, P. A. (2015). Aircraft measurements of BrO, IO, glyoxal, NO₂, H₂O, O-2-O-2 and aerosol extinction profiles in the tropics: comparison with aircraft-/ship-based in situ and lidar measurements. *Atmospheric Measurement Techniques*, 8(5), 2121-2148. doi:10.5194/amt-8-2121-2015

Vuilleumier, L., Harley, R. A., & Brown, N. J. (1997). First- and second-order sensitivity analysis of a photochemically reactive system (a Green's function approach). *Environmental Science & Technology*, 31(4), 1206-1217. doi:10.1021/es960727g

Wild, O., Prather, M. J., & Akimoto, H. (2001). Indirect long-term global radiative cooling from NO_x emissions. *Geophysical Research Letters*, 28(9), 1719-1722. doi:10.1029/2000gl012573

Williams, J. E., Boersma, K. F., Le Sager, P., & Verstraeten, W. W. (2017). The high-resolution version of TM5-MP for optimized satellite retrievals: description and validation. *Geoscientific Model Development*, 10(2), 721-750. doi:10.5194/gmd-10-721-2017

Wooldridge, P. J., Perring, A. E., Bertram, T. H., Flocke, F. M., Roberts, J. M., Singh, H. B., . . . Cohen, R. C. (2010). Total Peroxy Nitrates (Sigma PNs) in the atmosphere: the Thermal Dissociation-Laser Induced Fluorescence (TD-LIF) technique and comparisons to speciated PAN measurements. *Atmospheric Measurement Techniques*, 3(3), 593-607. doi:10.5194/amt-3-593-2010

Accepted

Table 1. NO-NO₂ cycling in the upper troposphere during SEAC⁴RS^a

Reaction	Mean Rate (10 ⁶ molecules cm ⁻³ s ⁻¹)
Conversion of NO to NO ₂	
R1. NO + O ₃ → NO ₂ + O ₂	3.42 ± 1.04 ^{b,c,d}
R2. NO + HO ₂ → NO ₂ + OH	0.68 ± 0.23 ^{b,d,e}
R3. NO + CH ₃ O ₂ → NO ₂ + CH ₃ O	0.13 ± 0.04 ^{b,d,e}
R4. NO + BrO → NO ₂ + Br	0.18 ± 0.09 ^{b,d,f}
R5. NO + IO → NO ₂ + I	0.10 ± 0.05 ^{b,d,f}
R6. NO + ClO → NO ₂ + Cl	0.02 ± 0.01 ^{b,d,f}
Total	4.53 ± 1.46
Conversion of NO ₂ to NO	
R7. NO ₂ + hν → NO + O	8.20 ± 1.74 ^g

^aMain reactions cycling NO and NO₂ in the daytime upper troposphere (8-12 km) over the Southeast US. Mean rates are calculated using JPL kinetic data (Burkholder et al., 2015) applied to SEAC⁴RS aircraft observations (NO, NO₂, O₃, J_{NO2}, temperature, pressure) over the Southeast US (94.5-76°W, 30-37°N) in August-September 2013 and with radical concentrations (RO₂, halogens) computed by the GEOS-Chem model along the flight tracks (Sherwen et al., 2016, 2017; Travis et al., 2016). Only reactions with rates above 1 × 10⁴ molecules cm⁻³ s⁻¹ are listed. Data outside the 9-15 solar time window and in stratospheric air ([O₃]/[CO] > 1.25 mol mol⁻¹) have been excluded. Error standard deviations are calculated by propagation of measurement, rate constant, and GEOS-Chem radical concentration errors (precision).

^bPrecision of the NO measurement (4%; Ryerson et al., 2000).

^cPrecision of the O₃ measurement (3%; Ryerson et al., 1998).

^dPrecision of the kinetic rate constants (30% for NO+O₃; 15% for NO+HO₂, NO+CH₃O₂, NO+BrO, and NO+ClO; 20% for NO+IO; Burkholder et al., 2015).

^eUncertainty in the GEOS-Chem HO₂ concentration (30%) estimated from the H₂O₂ measurements in SEAC⁴RS (Crouse et al., 2006). The same relative error is assumed for CH₃O₂.

^fUncertainty in the GEOS-Chem BrO, IO, ClO concentrations (50%; Sherwen et al., 2016).

^gPrecision of the NO₂ measurement (5%; Nault et al., 2015), the measured UV-A actinic flux (5%; Shetter et al., 2003), and the NO₂ cross section and quantum yield (20%; Sander et al., 2011).




Figure 1. Median vertical tropospheric profiles of the NO/NO₂ concentration ratio and related quantities on SEAC⁴RS flights (9-15 solar time) during August-September 2013 over the Southeast US (94.5-76°W, 30-37°N). Data from urban plumes ([NO₂] > 4 ppb), open fire plumes ([CH₃CN] > 200 ppt), and stratospheric air ([O₃]/[CO] > 1.25 mol mol⁻¹) are excluded. Observations are compared to GEOS-Chem model results sampled along the flight tracks, for the standard model (Travis et al., 2016) and a sensitivity simulation with reduced J_{NO₂} and increased low-temperature NO+O₃ rate constant (see bottom right panel). Observed NO₂ is from the Berkeley TD-LIF measurement (Nault et al., 2015). The bottom right panel shows the k₁ (NO+O₃) rate constant versus temperature from JPL in red with 2σ uncertainty in gray shading (Burkholder et al., 2015), and sensitivity simulation values resulting in 1.4k₁ (purple) in the upper troposphere. The J_{NO₂} observations apply JPL spectroscopic data to actinic fluxes measured aboard the aircraft, and would be reduced similarly to the model if the spectroscopic data are corrected downward (black dashed line). The dashed purple line in the top left panel shows the NO-NO₂-O₃ photochemical equilibrium values J_{NO₂}(hν)/k₁(T)[O₃] calculated from observed actinic fluxes (hν), temperature (T), and [O₃] with 1.4k₁ and the 20% reduction in J_{NO₂} applied.




Figure 2. Probability density function of the NO/NO₂ concentration ratio under midday conditions (9-15 local time) in the upper troposphere (8-12 km) during SEAC⁴RS in August-September 2013. Observations are compared to the standard GEOS-Chem model and the model with J_{NO_2} reduced by 20% and the activation energy for the NO+O₃ reaction increased so that k_1 increases on average by 1.4 in the upper troposphere (purple). The same data criteria as stated for Table 1 are applied.




Figure 3. Temperature dependence of the $\text{NO} + \text{O}_3$ rate constant k_1 . The y-axis is a log scale and the x-axis is an inverse scale ($1/T$), so that an Arrhenius dependence plots as a straight line. Recommended rates from JPL (black; Burkholder et al., 2015) and IUPAC (light gray; Atkinson et al., 2004) are shown as solid lines with the $2\text{-}\sigma$ uncertainty in shading. Laboratory measurements used in the JPL and IUPAC recommendations are shown in circles with their respective uncertainties. The sensitivity simulation resulting in $1.4k_1$ in the upper troposphere (230-250 K) is shown as the purple line.




Figure 4. Correlation of the observed NO/NO₂ concentration ratio with the local photochemical equilibrium constant $J_{\text{NO}_2}(h\nu)/k_1(T)[\text{O}_3]$ under midday conditions (9-15 local time) in the upper troposphere (8-12 km) during SEAC⁴RS in August-September 2013. Data are 10-minute averages along the aircraft flight tracks. The photochemical equilibrium constant is calculated from local aircraft measurements of actinic fluxes ($h\nu$), temperature (T), and ozone concentrations. Calculations using the JPL recommendations for $J_{\text{NO}_2}(h\nu)$ and $k_1(T)$ (in red) are compared to calculations reducing J_{NO_2} by 20% and increasing k_1 on average by 1.4 in the upper troposphere (purple). Solid lines show reduced major axis regressions and the 1:1 line is dashed. Correlation coefficients (r) and regression slopes (S) are given inset.