



UNIVERSITY OF LEEDS

This is a repository copy of *The effects of environmental enrichment on white matter pathology in a mouse model of chronic cerebral hypoperfusion*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/131119/>

Version: Published Version

---

**Article:**

Hase, Y, Craggs, L [orcid.org/0000-0001-9192-1072](https://orcid.org/0000-0001-9192-1072), Hase, M et al. (9 more authors) (2018) The effects of environmental enrichment on white matter pathology in a mouse model of chronic cerebral hypoperfusion. *Journal of Cerebral Blood Flow & Metabolism*, 38 (1). pp. 151-165. ISSN 0271-678X

<https://doi.org/10.1177/0271678X17694904>

---

(c) Author(s) 2017, Hase, Y, Craggs, L , Hase, M et al. The effects of environmental enrichment on white matter pathology in a mouse model of chronic cerebral hypoperfusion. *Journal of Cerebral Blood Flow & Metabolism*, 38 (1). pp. 151-165. Reprinted by permission of SAGE Publications.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## **Effects of Environmental Enrichment on White Matter Pathology in a Mouse Model of Chronic Cerebral Hypoperfusion**

Yoshiki Hase<sup>1</sup>, Lucinda Craggs<sup>1\*</sup>, Mai Hase<sup>1\*</sup>, William Stevenson<sup>1</sup>, Janet Slade<sup>1</sup>, Aiqing Chen<sup>1</sup>, Di Liang<sup>1</sup>, Abdel Ennaceur<sup>2</sup>, Arthur Oakley<sup>1</sup>, Masafumi Ihara<sup>3</sup>, Karen Horsburgh<sup>4</sup>, and Raj N Kalaria<sup>1</sup>

<sup>1</sup> Neurovascular Research Group, Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, UK

<sup>2</sup> Department of Pharmacy, Sunderland Pharmacy School, University of Sunderland, Sunderland, UK

<sup>3</sup> Department of Stroke and Cerebrovascular Diseases, National Cerebral and Cardiovascular Centre, Osaka, Japan

<sup>4</sup> Centre for Neuroregeneration, University of Edinburgh, Edinburgh, UK

\*These authors contributed equally to this work.

Running headline: White Matter Changes and Environmental Enrichment

Corresponding author:

Professor Raj N. Kalaria

Institute of Neuroscience, Newcastle University, Campus for Ageing and Vitality, Newcastle upon Tyne, NE4 5PL, United Kingdom

Tel: +44-(0)191-208-1352

E-mail: raj.kalaria@newcastle.ac.uk

Article: Total text 5,262 words. Tables/Figures: 7 (plus two supplementary figures)

## **Abstract**

White matter (WM) disintegration is common in the older population and associated with vascular cognitive impairment (VCI). This study explored the effects of environmental enrichment (EE) on pathological sequelae in a mouse model of chronic cerebral hypoperfusion induced by bilateral common carotid artery stenosis (BCAS). Male C57BL/6J mice underwent BCAS or sham surgery. One-week after surgery, mice were exposed to three different degrees of EE; either standard housing conditions (std), limited 3 hours exposure to EE per day (3hrs) or full-time exposure to EE (full) for 12 weeks. At 13-weeks after surgery, cognitive testing was performed using a 3-dimensional 9-arm radial maze. At 16-weeks after surgery, nesting ability was assessed in each mouse immediately before euthanasia. Brains retrieved after perfusion fixation were examined for WM pathology. BCAS caused WM changes, as demonstrated by corpus callosum atrophy and greater WM disintegrity. BCAS also caused impaired nesting ability and cognitive function. These pathological changes and working memory deficits were attenuated, more so by limited rather than full-time exposure to EE regime. Our results suggest that limited exposure to EE delays the onset of WM degeneration. Therefore, implementation of even limited EE may be beneficial for patients diagnosed with VCI.

## **Keywords**

Animal model, Chronic cerebral hypoperfusion, Environmental enrichment, Vascular dementia, White matter

## **Introduction**

There is currently no cure for vascular dementia (VaD), which is the second most common type of dementia diagnosed amongst the elderly population.<sup>1</sup> Safe and effective interventional strategies, which delay the onset or slow down the progression of cognitive decline, are urgently required. Subcortical ischaemic vascular dementia (SIVD), the most prevalent subtype of VaD is a common feature of vascular cognitive impairment (VCI).<sup>2,3</sup> SIVD is characterised by lacunar infarcts and white matter (WM) lesions,<sup>4</sup> and is responsible for VCI in the ageing population.<sup>5</sup> Chronic cerebral hypoperfusion, a major component of SIVD and diffuse WM changes caused by small-vessel disease (SVD) are strongly correlated to VCI and depression, as well as gait disturbance as evident in SIVD patients.<sup>6,7</sup> However, little is known about the neuropathological processes that lead to the progression of VCI in SIVD patients. To understand the biological mechanisms that occur in the progression of SVD, animal models of chronic cerebral hypoperfusion have been developed in rats<sup>8-12</sup> and gerbils.<sup>13</sup> Our research group has been working at the forefront to establish several novel rodent models of VCI or VaD by assessing the effects of reduced brain perfusion on the dynamics of the cerebral circulation and brain tissue changes.<sup>14-19</sup> Remarkably, the bilateral carotid artery stenosis (BCAS) mouse model has been evaluated as one of the most germane rodent models of VaD.<sup>20-23</sup>

Environmental enrichment (EE) has been suggested as a strategy to improve cognitive impairment in humans, and possibly reverse WM and neuronal damage. There is some evidence to show that physical and cognitive stimulations, which are incorporated in EE and physical exercise training, could enhance synaptic plasticity and attenuate cognitive deficits in rodents<sup>24-27</sup> and humans.<sup>28</sup> These studies reported that EE increased brain plasticity, enhanced neurogenesis, and increased synaptogenesis as well as preserved motor and cognitive function. However, although several previous studies reported the beneficial effects of EE in rodents and humans, few studies explored the protective effects of differing degrees of EE in VaD. Here, we investigated the effects of different degrees of EE on the pathological sequelae in mice subjected to BCAS, with the aim of assessing the morphology of the WM and cognitive dysfunction as it relates to VCI and VaD.<sup>4</sup>

## **Material and Methods**

### *Animals and surgical procedures*

Male C57BL/6J mice (9 weeks old, 23.1–25.3g) were purchased from the Jackson Laboratory, USA. The mice were housed in group cages with a 12 hrs day and 12 hrs night cycle (6am–6pm, day; 6pm–6am, night) and were given access to food and water *ad libitum*. After one-week acclimatisation, a total of 74 mice were randomly selected for either bilateral common carotid artery stenosis (BCAS, n=41) or sham (n=33) surgery. BCAS surgery was performed as described previously.<sup>14</sup> Briefly, mice were anaesthetised by 1.5% isoflurane in oxygen and air. Middle neck incision was made, bilateral common carotid arteries (CCAs) were exposed and isolated from the vagus nerves. Microcoils of inner diameter of 0.18mm (Sawane Spring, Japan) were applied to the both CCAs. Sham animals were exposed to the same operative procedures as BCAS mice, except for the application of microcoils. Body temperature was monitored and maintained between 36.5°C and 37.5°C with the aid of a feedback warming pad and a blanket during the operative procedure. Animals were appropriately identified with coded numbers. All of the experiments including behavioural test and data analyses in this study were performed under investigator-blinded conditions. Animals dying during the experimental period before the planned euthanasia at 16 weeks after surgery were excluded from further analyses, other than assessment of survival rate. All procedures were pre-approved by the Home Office, UK based upon ASPA: The Animals (Scientific Procedures) Act 1986, UK and performed in accordance with the guidelines stipulated by the ethical committee of Newcastle University and adhering to ARRIVE guidelines. Figure 1 outlines the entire experimental protocol.

#### *Measurement of body weight and analysis of survival rate*

Body weight (BW) of each mouse was determined before surgery and followed up to 3 times a week as necessary for a period of 12 weeks and prior to cognitive testing (Supplementary Figures 1a and b). During the cognitive testing period, BW was assessed daily in order to monitor consistency in weight. Survival rates were determined until 16 weeks after surgery and before euthanasia. These results were expressed as Kaplan-Meier survival curves (Supplemental Figures 1c and d).

#### *Environmental Enrichment (EE) paradigm and animal groups*

Animals were initially divided into two main groups, sham and BCAS (Figure 1b). One week after surgery, mice were randomly assigned to six subgroups, three per group: three different

EE conditions per main group for 12 weeks either standard housing, limited exposure to EE or full-time exposure to EE. Standard housing denotes a normal housing condition with no EE and it incorporated a paper house and shredded tissue. EE cages had extra toys in addition to the standard housing e.g. running wheels, hanging chains, igloos and a paper tunnel. Limited exposure to EE was performed as described previously.<sup>29</sup> Briefly, for the first four weeks, mice were transferred to the EE cages for 3 hours every day in the morning from 9am to noon. From the 5th week to the 12th week, mice placed in EE cages for 3 hours, 3 days a week. Full-time EE group was exposed to EE every day for 24 hours over the entire 12 weeks. Thus, altogether 6 experimental groups were compared: Sham-std, sham operated mice with standard housing; Sham-3hrs, sham mice with limited exposure to EE; Sham-full, sham operated mice with full-time exposure to EE; BCAS-std, BCAS operated mice with standard housing; BCAS-3hrs, BCAS operated mice with limited exposure to EE; BCAS-full, BCAS operated mice with full-time exposure to EE.

#### *Cognitive testing using an innovative three dimensional 9-arm radial maze (3D-RAM)*

Cognitive function (mainly working memory) was assessed in each mouse from 13 weeks post BCAS surgery during a period of 20 days, using a novel 3-dimensional (3D) 9-arm radial maze (3D-RAM).<sup>30</sup> This was a modified version of the conventional 8-arm radial maze. Behavioural tests were performed in a dimly lit behavioural testing room. The 3D maze (Grey PVC, 5 mm thick) consisted of nine arms connected to bridges (slope part) radiating from a central platform. Each arm (35 cm x 11.2 cm) was attached to a bridge (slope part at 40 degrees, 15.2 cm x 11.2 cm). The surface of the bridge was made of metal mesh, which enabled mice to maintain grip. A small transparent wall panel (9 cm x 6 cm) was randomly placed at each entrance of a bridge in order to narrow entry, which avoids a continuous sequential entry from one bridge to the next. In order to enter each arm, mice paced in the centre platform and needed to cross a bridge (elevated ramp).<sup>30</sup> At the end of each arm, a small pellet (Dustless Precision Pellets<sup>®</sup> Rodent, Purified, Bio Serv, USA) was placed in order to entice food restricted mice to enter the arm. Different colours/shapes of pictures were placed vertically at the end of each arm as visual cues, enabling mice to distinguish individual arms. The maze was placed at the centre of a behavioural test room affixed during the entire testing period. All of the sessions for each mouse were streamed and recorded using a camcorder (LEGRIA HF R56, Canon), suspended directly over the

maze, via a Wi-Fi network to an iPad (Apple Inc., USA), which was sited outside the behavioural test room.

A day before the first session, each mouse was weighed to ascertain BW and food deprivation was induced to achieve 10% reduction in their body weight. Mice were weighed immediately prior to each testing session to ensure their BW was maintained at 90% of baseline level. Mice were randomly tested for 20 consecutive days without prior habituation to the maze.<sup>31,32</sup> Mice placed into a transparent plastic beaker (7cm diameter, 18.5cm height) were gently introduced on the central platform at the start of each session. The session was terminated when each mouse completed 9 arm entries or when 10 minutes had elapsed. After each session, the mouse was allowed to return to the central platform before removing it from the maze in a similarly gentle manner. Between each session, the maze was cleaned with hypochlorous acid soaked tissue followed by distilled water to remove any traces smells left by urine or faeces.

The number of arm entries, number of arm repeats and number of arm entries before first repeat (NABFR) were recorded. The probability of each pair of “arm entries and arm repeats (top of each cell)” and “arm repeat score (bottom of each cell)” (Figure 7a), probability of each “NABFR (top of each cell)” and “NABFR score (bottom of each cell)” (Figure 7d) was then calculated. Arm repeat scores for each pair were defined as the difference in percentage compared to the most probable pair. For example, the most probable arm repeats in nine arm entries can be 3, and its probability is calculated as 41.0%. Arm repeat score of this pair was set as zero (Figure 7a). The probability of 1 arm repeat in 9 arm entries was theoretically 3.37%. Therefore, the arm repeat score of this pair was calculated as:  $41.0 - 3.37 = 37.63\%$ . However, the probability of 5 arm repeats in 9 arm entries is 6.0%. In this case, the arm repeat score was calculated as a negative value:  $6.0 - 41.0 = -32.0\%$ , as more than 3 arm repeats are counted as a negative score. NABFR scores were similarly calculated in the same manner as the arm repeat score. For example, the most probable NABFR is 3 and its probability was calculated as 23.05%. NABFR score of this was set as zero (Figure 7d). The probability of NABFR =5 is theoretically 14.23%. Therefore, the NABFR score of this was calculated as:  $23.05 - 14.23 = 8.82\%$ . On the other hand, the probability of NABFR=1 is 11.11%. In this case, NABFR score was calculated as a negative value:  $11.11 - 23.05 = -11.93\%$ , as less than 3 NABFR is counted as a negative score. The final scores represent the average of the scores from all sessions.

### *Nesting behaviour*

The NESTLETS™ (Ancare, USA) test was used to assess the nesting ability of mice<sup>33</sup> before BCAS or sham surgery (as baseline) as well as 16 weeks after surgery before euthanasia. Briefly, one NESTLET, composed of a pressed square of cotton material (2 x 2 inches) was placed at one corner of each cage at 4pm and left there overnight. First, mice shredded the tightly packed cotton material and then arranged it into a nest. Next morning at 10am, each cage was assessed in terms of the appearance of the nest (NESTLET score), height of the nest (cm) and the percentage NESTLET used (%). Scoring was defined as follows:<sup>33</sup> 1, Nestlet not noticeably touched (more than 90% intact); 2, Nestlet partially torn (50–90% remaining intact); 3, Nestlet mostly shredded but often no identifiable nest site: less than 50% of the Nestlet remains intact, but less than 90% is within a quarter of the cage floor area; i.e., the cotton is not gathered into a nest but is spread around the cage. The material may sometimes be in a broadly defined nest area, but the critical definition is that 50–90% is shredded; 4, An identifiable but flat nest: >90% of the Nestlet is torn and the material is gathered into a nest within a quarter of the cage floor area, but the nest is flat, with walls higher than the average mouse BW for less than 50% of its circumference; 5, A (near) perfect nest: more than 90% of the Nestlet is torn and the nest is a crater, with walls higher than the average mouse BW for >50% of its circumference.

### *Histopathological analysis*

Mice were anaesthetised by intraperitoneal injection of sodium pentobarbital (50 mg/kg) and perfused transcardially at 20 ml/min with 0.01M phosphate buffered saline (PBS), pH 7.4. Brains were removed and divided into two hemispheres. The right or left hemisphere of each brain was randomly assigned for histological analysis and post fixed in 4% paraformaldehyde (PFA) in 0.01M PBS (pH 7.4) for 48 hours. Each hemisphere was cut into 5 blocks at different coronal levels after fixation: Block OB, coronal level of olfactory bulb (OB); Block 1, coronal level of bregma +0.5 mm; Block 2, coronal level of bregma -1.0 mm; Block 3, coronal level of bregma -2.0 mm; Block 4, level of cerebellum and brain stem. Each sub-dissected block was dehydrated and embedded in paraffin to produce 5-µm thick sections for histological staining and analysis.



**Volume of the corpus callosum:** Luxol-fast blue and cresyl fast violet (LFB+CFV): Klüver-Barrera (KB) stain was used to assess area and volume of the corpus callosum (CC). KB-stained sections obtained from Blocks 1, 2, and 3: coronal level of bregma +0.5 mm, -1.0 mm and -2.0 mm respectively were used so that the whole CC from rostral to caudal segment was analysed. Images of the CC from paramedian to the lateral edge were captured using a bright field microscope (Leitz DIALUX 20, Leica) with a 5x objective lens coupled to a lumenera infinity digital camera (Lumenera Corporation, Canada). The entire CC was also traced using imageJ software and the area of CC (mm<sup>2</sup>) for each block was calculated. To assess the volume, we assumed each CC area was of similar shape. Areas of the CC were defined as: CC area obtained from block 1 (bregma +0.5 mm), S1 (mm<sup>2</sup>); from block 2 (bregma -1.0 mm), S2 (mm<sup>2</sup>); from block 3 (bregma -2.0 mm), S3 (mm<sup>2</sup>). Volume of the rostral and caudal side of CC were defined as: V1 (mm<sup>3</sup>), volume of the CC between block 1 and block 2 (distance=1.5 mm); V2 (mm<sup>3</sup>), volume of the CC between block 2 and block 3 (distance = 1.0 mm). Using the following formula, each volume was calculated as follows: V1 (mm<sup>3</sup>) =  $\frac{1.5}{3} (S1 + \sqrt{S1 * S2} + S2)$ ; V2 (mm<sup>3</sup>) =  $\frac{1}{3} (S2 + \sqrt{S2 * S3} + S3)$ . Total estimated volume of CC between coronal level of bregma + 0.5 mm and bregma – 2.0 mm (mm<sup>3</sup>) = V1 + V2 (mm<sup>3</sup>).

**White matter severity scores:** WM severity score was assessed in KB-stained coronal sections obtained from Blocks 1, 2, and 3. We analysed all distinct regions of the WM including the CC, anterior commissure, hippocampal fimbria, internal capsule, optic nerve and optic tract. For analysis of the CC, images were captured using a bright field microscope (Leitz DIALUX 20, Leica) with a 20x objective lens coupled to a lumenera infinity digital camera (Lumenera Corporation, Canada). The severity of the WM pathology was graded as follows: Grade 0, normal; Grade 1, disarrangement of the nerve fibres; Grade 2, the formation of marked vacuoles; Grade 3, and the disappearance of myelinated fibres as described previously.<sup>8</sup>

**Oligodendrocytes:** Haematoxylin-stained sections obtained from Block 3, coronal level of bregma +2.0 mm were used for the analysis of total number of oligodendrocytes in the CC. Oligodendrocytes were identified based upon its size and morphology: size of its nucleus around 6-µm, rounding shape with a dense-stained nucleus.<sup>4</sup> The whole CC from

paramedian to lateral edge of each section was imaged and analysed. All oligodendrocytes within each image were counted to assess the total number of oligodendrocytes.

### *Statistical analysis*

Data were expressed mean  $\pm$  SEM. Using IBM SPSS statistics 22 software, non-paired t-test or one-way analysis of variance (ANOVA) followed by post-hoc Tukey's test were performed for multiple comparison of each group. Mann-Whitney U test or Kruskal-Wallis H Test was performed for non-parametric analysis. Two-way repeated measures ANOVA was performed for assessment of body weight change and behavioural data. Log-rank test was performed for Kaplan-Meier survival analysis.  $P < 0.05$  was defined as statistically significant in all analyses.

## **Results**

### *BW compared with sham animals after BCAS*

Pre-operative BWs of each group were: Sham-std,  $23.6 \pm 0.3$  g; Sham-3hrs,  $23.6 \pm 0.5$  g; Sham-full,  $24.0 \pm 0.7$  g; BCAS-std,  $24.6 \pm 0.3$  g; BCAS-3hrs,  $24.4 \pm 0.6$  g; BCAS-full,  $24.9 \pm 0.4$  g respectively ( $P > 0.05$ ). At one week after surgery, all BCAS groups had lost some BW compared with the pre-operative value and began to recover until 13 weeks post-operation. However, weight gain in all BCAS groups was slower compared with sham animals at each indicated time points ( $P < 0.01$ ) (Supplemental Figures 1a, 1b), although the amount of food intake after surgery was not different between the sham and BCAS animals (data not shown). There were no significant differences in BW at any time point between sham and BCAS subgroups (Supplementary Figure 1B). However, limited exposure to EE (BCAS-3hrs) tended to gain more BW compared to other BCAS subgroups, especially from 7 weeks after surgery.

### *BCAS animals tended to survive less compared to sham animals*

All sham-operated mice ( $n=33$ ) survived until 16 weeks after surgery (Supplementary figure 1c). However, survival rate of the BCAS mice was 85.4% (35 of original 41 mice) at 16

weeks after surgery (Kaplan-Meier survival analysis, log-rank  $P=0.01$ ) (Supplemental Figure 1c). The survival rates of the BCAS mice were as follows: BCAS-std 92.9% (13 of original 14), BCAS-3hrs 91.0% (10 of 11); BCAS-full 75% (12 of 16) (Supplemental Figure 1d). There were no significant differences between BCAS groups by Kaplan-Meier survival analysis. However, BCAS-full showed the lowest survival rate during 16 weeks observation period post-surgery. The causes of death in the BCAS group were: cerebral haemorrhage (n=1), severe enterocolitis (n=1), kidney anomaly (n=1), acute renal failure due to dehydration (n=3). The final number of survivors in each group was Sham-std, n=11; Sham-3hrs, n=11; Sham-full, n=11; BCAS-std, n=13, BCAS-3hrs, n=10; BCAS-full, n=12 (Figure 1b).

#### *Limited versus full-time exposure to EE and corpus callosum (CC) atrophy*

The CC volume in each group was calculated to assess atrophy. Compared with all the sham animals (Figure 2 b-d). BCAS-std and BCAS-full subgroups showed significantly smaller volumes of CC. This was apparent in the rostral as well as caudal segments and the total volume of the CC. The BCAS-3hrs subgroup showed significantly larger volumes of CC compared with BCAS-std subgroups ( $*P<0.05$ ).

#### *Limited versus full-time exposure to EE on WM integrity*

BCAS resulted in higher WM severity scores compared with sham animals. In the paramedian area of the CC (Figure 3 a (A-F)), BCAS-3hrs mice showed reduced WM score (preserved WM integrity) compared to BCAS-std at all coronal levels comprising blocks 1, 2 and 3 at coronal levels relative to bregma +0.5 mm, -1.0 mm and -2.0 mm respectively ( $*P<0.05$ ) (Figure 4 a, d and g). In the most caudal part, BCAS-3hrs mice also had reduced WM severity score compared to BCAS-full ( $*P<0.05$ ) (Figure 4g). In turn, BCAS-full mice had more preserved WM integrity compared to BCAS-std ( $*P<0.05$ ) (Figure 4i). In both the median (Figure 3 b (A-F)) and lateral (Figure 3 c (A-F)) regions of the CC, BCAS-3hrs and BCAS-full subgroups exhibited reduced WM scores compared to BCAS-std at all coronal levels ( $*P<0.05$ ) (Figure 4 b-f and i-h). There was also a reduction in the WM score of the anterior commissure of BCAS-3hrs and BCAS-full animals compared to BCAS-std ( $*P<0.05$ ) (Figure 4j). Similarly, there was less WM damage in the hippocampal fimbria and internal capsule in the BCAS-3hrs animals compared to BCAS-std ( $*P<0.05$ ) (Figure 4 k and

l). In the optic nerve and optic tract, BCAS subgroups showed greater WM scores compared to sham subgroups.

*Status of oligodendrocytes during limited versus full-time exposure to EE*

All sham subgroups showed normal alignment and appearance of oligodendrocytes in the caudal segment of the CC (Figure 5 a (A-C)). On the other hand, BCAS-std mice had lower numbers and disrupted cells (Figure 5 a (D)). EE, especially BCAS-3hrs, preserved the number and alignment of oligodendrocytes (Figure 5 a (E)). The total number of oligodendrocytes in the entire caudal segment was greater in BCAS-3hrs mice compared with BCAS-std and BCAS-full subgroups ( $P<0.01$ ) (Figure 5b). BCAS-full also showed increased numbers of cells compared with BCAS-std ( $P<0.05$ ) (Figure 5b).

*Impaired nesting ability induced by BCAS was reversed by EE*

Sham subgroups had almost the same level of nesting ability (Nestlet score) compared with preoperative baseline level at 16 weeks after surgery. However, BCAS-std subgroup exhibited lower Nestlet scores (impaired nesting ability) compared to baseline level ( $*P<0.05$ ). EE reversed impaired nesting ability, and limited exposure to EE (BCAS-3hrs) tended to improve nesting ability compared with BCAS-std and BCAS-full subgroups (Figure 6c). BCAS plus EE subgroups made higher nests, used more Nestlet to make nests compared with BCAS-std (data not shown).

*Impaired working memory partially reversed by EE*

Working memory was significantly impaired in BCAS compared to sham animals.

**Arm repeat scores:** Average arm repeat scores in sessions 1-10, sessions 11-20 and sessions 1-20, all showed significantly lower scores in BCAS compared to sham animals ( $P<0.05$  in sessions 1-10;  $P<0.01$  in sessions 11-20 and sessions 1-20) (Figure 7b). Sham group dramatically improved arm repeat score by 47.4% in sessions 11-20 compared to sessions 1-10, whereas BCAS group reduced score by 6.6% (Figure 7b). All the sham animals improved arm repeat scores (average 49%) in sessions 11-20 compared to sessions 1-10. The BCAS-3hrs mice improved their score by 37.3%, whereas BCAS-std only improved by 13.0% and BCAS-full even reduced the average score by 45.7% (Figure 7c). For the arm

repeat score averaged from sessions 1-20, the BCAS-3hrs subgroup showed the highest mean score between BCAS subgroups, followed by BCAS-full and BCAS-std. In sessions 11-20 and sessions 1-20, the BCAS-3hrs mice showed no differences compared to sham-std. In sessions 11-20, BCAS-3hrs also showed no significant change in lower arm repeat scores compared with those on sham-3hrs ( $P=0.08$ ).

**NABFR scores:** the number of arm entries before first repeat (NABFR) was lower in BCAS mice compared to the sham ( $P<0.01$ ) (Figure 7e). All sham subgroups also completed more arm entries before first repeat compared with all BCAS subgroups. Between BCAS subgroups, the BCAS-3hrs mice subgroup completed more arm entries before first repeat, in sessions 1-5 and 11-15 (Figure 7f). The BCAS animals also showed lower NABFR scores compared with sham in sessions 1-10, 11-20 and 1-20, ( $P<0.01$ ) (Figure 7g). The NABFR scores averaged from sessions 1-20, were highest for the BCAS-3hrs mice followed by BCAS-full and BCAS-std (Figure 7h). BCAS-3hrs mice also showed a trend towards lower scores compared with sham-3hrs in sessions 11-20 ( $P=0.06$ ).

## Discussion

Our findings in this study suggest beneficial effects of limited EE against WM changes after chronic BCAS as related to the features of VCI and VaD. We found that BCAS caused: 1) WM damage demonstrated by CC atrophy as well as WM disintegration and loss of oligodendrocytes; 2) impaired nesting ability as well as cognitive function. These unfavourable changes were attenuated by environmental enrichment (EE) intervention, by limited exposure to EE rather than full-time exposure to EE.

Few previous studies reported overall BW change after long-term BCAS. We found BCAS animals did not gain weight compared to sham, despite lack of differences in the amount of food consumption between BCAS and sham animals. The reduced trailing effects on BW in the BCAS mice may be explained by systemic metabolic changes. Overall survival rate after surgery was significantly lower in BCAS compared to sham animals. This is compatible with previous reports indicating that survival rates after BCAS was about 80%.<sup>14, 34</sup> However, in our study, an overall 4-month survival rate after BCAS was >90% in BCAS-std and BCAS-3hrs subgroups. We also suggest the higher mortality in the BCAS-full subgroup could be explained by dehydration and acute renal failure possibly due to increased physical activity promoted by the full-time exposure to EE. Although EE exposes the animals to a number of

different features including physical exercise and social interaction, we noted the average number of wheel rotations in BCAS-full subgroup was 5-fold greater compared with BCAS-3hrs, Sham-3hrs and Sham-full subgroups (data not shown). Limited exposure to EE could have also initiated and activated the fronto-subcortical circuits more often when the animals experienced limited exposure to EE every time. Repeated activation of fronto-subcortical circuits could enhance cognitive function and prevent cognitive decline after BCAS.

It has been reported that nesting is important for heat conservation, reproduction and shelter and shown to relate to brain lesions, pharmacological agents and genetic mutations.<sup>33</sup> We found that BCAS caused impaired nesting ability, and it was attenuated by EE, which is a new finding in the BCAS mouse model. BCAS plus limited exposure to EE mice tended to make better nests compared to other BCAS subgroups. These results suggest that EE after chronic cerebral hypoperfusion could protect changes in different cognitive domains or could attenuate mild undetectable sensory-motor deficit involved in nesting ability.

Previous reports suggest that BCAS causes impaired working memory.<sup>15, 35, 36</sup> These studies used the conventional 8-arm radial maze or the Y-maze. We utilised an innovative 3D 9-arm radial arm maze for the assessment of cognitive function, which is a modified version of the conventional radial arm maze and designed for assessing working memory.<sup>31</sup> This maze allowed us to measure more indices compared with the conventional 8-arm radial arm and Y-mazes.<sup>32, 37</sup> The arm repeat scores allowed us to assess probability as well as number of arm repeats. The NABFR scores allowed us to detect differences in cognitive function of each group with greater sensitivity. We found that all BCAS showed significantly lower arm repeat scores and NABFR scores compared with the sham animals, suggesting that BCAS impaired working memory. This is consistent with the heterogeneity observed in VaD and normally ageing individuals. We also found that limited exposure to EE ameliorated cognitive decline after BCAS. The average arm repeat score and NABFR score in sessions 11-20, BCAS plus limited exposure to EE subgroup (BCAS-3hrs) showed no significant difference compared with sham subgroups. This result suggests that moderate, rather than full-time exposure to EE could be an effective intervention against impaired working memory as evident in VCI.<sup>3</sup> Overall, our observations are consistent with previous evidence indicating that EE elicits beneficial effects on neuronal plasticity and cognitive functioning in rats,<sup>26</sup> mice<sup>27, 38-40</sup> and humans.<sup>27, 38-40</sup> However, the underlying neurophysiological mechanisms of EE remains poorly understood. It is proposed that the structural and

biomechanical changes in the WM induced by EE, is due to the upregulation of genetic expressions associated with neurogenesis, synaptic plasticity, neuroprotection and intact memory function.<sup>41</sup>

Carotid artery stenosis is an important risk factor for cerebral WM disease<sup>42, 43</sup> and relates to WM damage and cognitive dysfunction<sup>44-46</sup> apparent in SIVD.<sup>14</sup> We propose that chronic cerebral hypoperfusion in mice or rats<sup>47</sup> leads to reduced CBF. Thus, we noted that all BCAS subgroups showed thickened carotid arterial wall and prolonged reduction of CBF as evident by laser speckled flowmetry at 16 weeks after surgery.<sup>48</sup> This suggested that compensatory mechanisms such as collateral flow, were not able to prevent CBF reduction due to severely stenosed carotid arteries but the reduced CBF in turn facilitated damage in the deep WM, particularly affecting the cellular structure including oligodendrocytes in the CC.<sup>14, 34-36</sup> However, the WM pathology is indicative of overall subcortical damage and neuronal disconnectivity between cerebral cortices that result in cognitive and motor deficits observed in SIVD.<sup>4, 45, 46, 49</sup> As BCAS-std and BCAS-full mice both showed more severe WM damage i.e. CC atrophy than in the BCAS-3hrs subgroup, this suggests that continuous but moderate levels of EE is more beneficial to sustaining WM integrity than no enrichment or full-time exposure to EE.

WM in the aged mouse is vulnerable to prolonged cerebral hypoperfusion and hypoxic stress, and loses its ability to recruit cyclic AMP response element-binding protein (CREB)-mediated oligodendrogenesis for responding to WM injury and stress.<sup>50</sup> In our study, limited exposure to EE significantly attenuated WM pathology and preserved oligodendrocytes, suggesting that EE can restore WM integrity and result in better cognitive function, possibly via CREB-mediated oligodendrogenesis. Assessment of CREB-mediated oligodendrogenesis in the WM could enable elucidation of the mechanisms of limited exposure to EE against WM damage.

There are some limitations in our study. First, we assessed cognitive function and brain histological changes in brain only at one time point, 16 weeks after surgery before euthanasia. Physiological and histological data obtained from several more time points would be useful for greater understanding of the pathological mechanisms as well as effects of EE in BCAS. Secondly, we set limited exposure to EE regime as 3 hours a day. Testing other durations of exposure to EE, such as shorter than 3 hours or longer than 3 hours could more precisely elucidate the effects of limited EE against BCAS. Thirdly, as VCI or VaD also involves changes in the medial temporal lobe,<sup>51</sup> we have not presented data on the status of the

hippocampus along with our present findings. This requires an in depth labour intensive analysis, which is currently in progress. Finally, we deem that further in depth analysis of the fibre tracts of the CC would allow us to identify the nature of the damage in specific fronto-subcortical circuits that are affected after long-term cerebral hypoperfusion.

## **Conclusions**

In summary, long-term chronic cerebral hypoperfusion induced by BCAS produced similar neuropathological changes to those seen in VaD indicative the relevance of the model. This study also demonstrated that limited exposure to EE protects the WM and elicits a beneficial effect on oligodendrocyte loss after chronic cerebral hypoperfusion-induced vascular injury. Although further experimental studies are needed, moderate EE appears a safe and effective future interventional strategy for cerebrovascular diseases, especially for patients with VCI.

## **Funding**

Our study was supported by a grant (ARUK PG2013-22) from Alzheimer's Research UK. Y.H. was supported by SENSHIN Medical Research Foundation, Osaka, Japan and The Great Britain Sasakawa Foundation, London, UK.

## **Acknowledgements**

We specially thank Neil Hamilton and Sandra Hogg for excellent technical assistance with respect to the daily care of mice and animal husbandry. We thank Mary Johnson and Lynn Ramsey (Newcastle University) for excellent technical assistance in processing brain tissue.

## **Declaration of conflicting interests**

The authors declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

## **Author's contributions**

YH, LC, MH, WS and JS performed or contributed to different aspects of the experiments in the study. YH wrote the first draft of the manuscript. AC, DL, AE, AO, MI, KH and RNK



contributed to critically revising the manuscript for important intellectual content, and all approved the final version of the manuscript for submission.

**Supplementary material**

Supplementary Material for this paper can be found at <http://jcbfm.sagepub.com/content/by/supplemental-data>.

## References

1. Kalaria RN. Neuropathological diagnosis of vascular cognitive impairment and vascular dementia with implications for Alzheimer's disease. *Acta Neuropathol.* 2016; 131: 659-85.
2. Roman GC, Erkinjuntti T, Wallin A, Pantoni L and Chui HC. Subcortical ischaemic vascular dementia. *Lancet Neurol.* 2002; 1: 426-36.
3. Hachinski V, Iadecola C, Petersen RC, et al. National Institute of Neurological Disorders and Stroke-Canadian Stroke Network vascular cognitive impairment harmonization standards. *Stroke; a journal of cerebral circulation.* 2006; 37: 2220-41.
4. Ihara M, Polvikoski TM, Hall R, et al. Quantification of myelin loss in frontal lobe white matter in vascular dementia, Alzheimer's disease, and dementia with Lewy bodies. *Acta Neuropathol.* 2010; 119: 579-89.
5. Pantoni L. Cerebral small vessel disease: from pathogenesis and clinical characteristics to therapeutic challenges. *Lancet Neurol.* 2010; 9: 689-701.
6. Poggesi A, Pantoni L, Inzitari D, et al. 2001-2011: A Decade of the LADIS (Leukoaraiosis And DISability) Study: What Have We Learned about White Matter Changes and Small-Vessel Disease? *Cerebrovasc Dis.* 2011; 32: 577-88.
7. Longstreth WT, Jr., Manolio TA, Arnold A, et al. Clinical correlates of white matter findings on cranial magnetic resonance imaging of 3301 elderly people. The Cardiovascular Health Study. *Stroke; a journal of cerebral circulation.* 1996; 27: 1274-82.
8. Wakita H, Tomimoto H, Akiguchi I and Kimura J. Glial activation and white matter changes in the rat brain induced by chronic cerebral hypoperfusion: an immunohistochemical study. *Acta Neuropathol.* 1994; 87: 484-92.
9. Wakita H, Tomimoto H, Akiguchi I and Kimura J. Protective effect of cyclosporin A on white matter changes in the rat brain after chronic cerebral hypoperfusion. *Stroke; a journal of cerebral circulation.* 1995; 26: 1415-22.
10. Farkas E, Donka G, de Vos RA, Mihaly A, Bari F and Luiten PG. Experimental cerebral hypoperfusion induces white matter injury and microglial activation in the rat brain. *Acta Neuropathol.* 2004; 108: 57-64.
11. Bueche CZ, Hawkes C, Garz C, et al. Hypertension drives parenchymal beta-amyloid accumulation in the brain parenchyma. *Ann Clin Transl Neurol.* 2014; 1: 124-9.
12. Kalaria RN, Bhatti SU, Palatinsky EA, et al. Accumulation of the beta amyloid precursor protein at sites of ischemic injury in rat brain. *Neuroreport.* 1993; 4: 211-4.

13. Kudo T, Tada K, Takeda M and Nishimura T. Learning impairment and microtubule-associated protein 2 decrease in gerbils under chronic cerebral hypoperfusion. *Stroke; a journal of cerebral circulation*. 1990; 21: 1205-9.
14. Shibata M, Ohtani R, Ihara M and Tomimoto H. White matter lesions and glial activation in a novel mouse model of chronic cerebral hypoperfusion. *Stroke; a journal of cerebral circulation*. 2004; 35: 2598-603.
15. Shibata M, Yamasaki N, Miyakawa T, et al. Selective impairment of working memory in a mouse model of chronic cerebral hypoperfusion. *Stroke; a journal of cerebral circulation*. 2007; 38: 2826-32.
16. Kitamura A, Fujita Y, Oishi N, et al. Selective white matter abnormalities in a novel rat model of vascular dementia. *Neurobiology of aging*. 2012; 33: 1012.e25-35.
17. Kitamura A, Saito S, Maki T, et al. Gradual cerebral hypoperfusion in spontaneously hypertensive rats induces slowly evolving white matter abnormalities and impairs working memory. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2015.
18. Hattori Y, Enmi J, Kitamura A, et al. A novel mouse model of subcortical infarcts with dementia. *J Neurosci*. 2015; 35: 3915-28.
19. Hattori Y, Enmi J, Iguchi S, et al. Gradual Carotid Artery Stenosis in Mice Closely Replicates Hypoperfusive Vascular Dementia in Humans. *J Am Heart Assoc*. 2016; 5.
20. Bink DI, Ritz K, Aronica E, van der Weerd L and Daemen MJ. Mouse models to study the effect of cardiovascular risk factors on brain structure and cognition. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2013; 33: 1666-84.
21. Ihara M and Tomimoto H. Lessons from a mouse model characterizing features of vascular cognitive impairment with white matter changes. *J Aging Res*. 2011; 2011: 978761.
22. Madigan JB, Wilcock DM and Hainsworth AH. Vascular Contributions to Cognitive Impairment and Dementia: Topical Review of Animal Models. *Stroke; a journal of cerebral circulation*. 2016.
23. Holland PR, Searcy JL, Salvadores N, et al. Gliovascular disruption and cognitive deficits in a mouse model with features of small vessel disease. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2015; 35: 1005-14.

24. Sun H, Zhang J, Zhang L, Liu H, Zhu H and Yang Y. Environmental enrichment influences BDNF and NR1 levels in the hippocampus and restores cognitive impairment in chronic cerebral hypoperfused rats. *Curr Neurovasc Res.* 2010; 7: 268-80.
25. Cechetti F, Worm PV, Lovatel G, Moyses F, Siqueira IR and Netto CA. Environmental enrichment prevents behavioral deficits and oxidative stress caused by chronic cerebral hypoperfusion in the rat. *Life Sci.* 2012; 91: 29-36.
26. Bayat M, Sharifi MD, Haghani M and Shabani M. Enriched environment improves synaptic plasticity and cognitive deficiency in chronic cerebral hypoperfused rats. *Brain Res Bull.* 2015; 119: 34-40.
27. Fischer A. Environmental enrichment as a method to improve cognitive function. What can we learn from animal models? *Neuroimage.* 2016; 131: 42-7.
28. Ihara M, Okamoto Y, Hase Y and Takahashi R. Association of physical activity with the visuospatial/executive functions of the montreal cognitive assessment in patients with vascular cognitive impairment. *J Stroke Cerebrovasc Dis.* 2013; 22: e146-51.
29. Lazarov O, Robinson J, Tang YP, et al. Environmental enrichment reduces Abeta levels and amyloid deposition in transgenic mice. *Cell.* 2005; 120: 701-13.
30. Ennaceur A, Michalikova S, van Rensburg R and Chazot PL. Detailed analysis of the behavior and memory performance of middle-aged male and female CD-1 mice in a 3D maze. *Behav Brain Res.* 2008; 187: 312-26.
31. Ennaceur A. Omission of the habituation procedure in the acquisition of a working memory task - evidence from Balb/c, C57/BL6J, and CD-1 mice. *Behav Brain Res.* 2011; 223: 203-10.
32. Ennaceur A, Michalikova S, van Rensburg R and Chazot PL. MK-801 increases the baseline level of anxiety in mice introduced to a spatial memory task without prior habituation. *Neuropharmacology.* 2011; 61: 981-91.
33. Deacon RM. Assessing nest building in mice. *Nat Protoc.* 2006; 1: 1117-9.
34. Nishio K, Ihara M, Yamasaki N, et al. A mouse model characterizing features of vascular dementia with hippocampal atrophy. *Stroke; a journal of cerebral circulation.* 2010; 41: 1278-84.
35. Washida K, Ihara M, Nishio K, et al. Nonhypotensive dose of telmisartan attenuates cognitive impairment partially due to peroxisome proliferator-activated receptor-gamma activation in mice with chronic cerebral hypoperfusion. *Stroke; a journal of cerebral circulation.* 2010; 41: 1798-806.

36. Maki T, Ihara M, Fujita Y, et al. Angiogenic and vasoprotective effects of adrenomedullin on prevention of cognitive decline after chronic cerebral hypoperfusion in mice. *Stroke; a journal of cerebral circulation*. 2011; 42: 1122-8.
37. Abuhamdah RM, Hussain MD, Chazot PL and Ennaceur A. Effects of chronic fluoxetine treatment on anxious behaviour of BALB/c mice in a 3-dimensional maze. *Stress*. 2015; 18: 677-85.
38. Nie J and Yang X. Modulation of Synaptic Plasticity by Exercise Training as a Basis for Ischemic Stroke Rehabilitation. *Cell Mol Neurobiol*. 2016.
39. McKee AC, Daneshvar DH, Alvarez VE and Stein TD. The neuropathology of sport. *Acta Neuropathol*. 2014; 127: 29-51.
40. Zou C, Shi Y, Ohli J, Schuller U, Dorostkar MM and Herms J. Neuroinflammation impairs adaptive structural plasticity of dendritic spines in a preclinical model of Alzheimer's disease. *Acta Neuropathol*. 2016; 131: 235-46.
41. Rampon C, Jiang CH, Dong H, et al. Effects of environmental enrichment on gene expression in the brain. *Proc Natl Acad Sci U S A*. 2000; 97: 12880-4.
42. Kandiah N, Goh O, Mak E, Marmin M and Ng A. Carotid stenosis: a risk factor for cerebral white-matter disease. *J Stroke Cerebrovasc Dis*. 2014; 23: 136-9.
43. Baradaran H, Mtui EE, Richardson JE, et al. White Matter Diffusion Abnormalities in Carotid Artery Disease: A Systematic Review and Meta-Analysis. *J Neuroimaging*. 2016.
44. Tomimoto H, Lin JX, Matsuo A, et al. Different mechanisms of corpus callosum atrophy in Alzheimer's disease and vascular dementia. *J Neurol*. 2004; 251: 398-406.
45. Jokinen H, Ryberg C, Kalska H, et al. Corpus callosum atrophy is associated with mental slowing and executive deficits in subjects with age-related white matter hyperintensities: the LADIS Study. *J Neurol Neurosurg Psychiatry*. 2007; 78: 491-6.
46. Ryberg C, Rostrup E, Paulson OB, et al. Corpus callosum atrophy as a predictor of age-related cognitive and motor impairment: a 3-year follow-up of the LADIS study cohort. *J Neurol Sci*. 2011; 307: 100-5.
47. Tomimoto H, Ihara M, Wakita H, et al. Chronic cerebral hypoperfusion induces white matter lesions and loss of oligodendroglia with DNA fragmentation in the rat. *Acta Neuropathol*. 2003; 106: 527-34.
48. Hase Y, Craggs L, Stevenson W, et al. Effects of Environmental Enrichment/Physical Exercise on White Matter Integrity and Stroke injury in a Mouse Model of Chronic Cerebral Hypoperfusion. *Cerebrovascular Diseases*. 2016; 41, Suppl 1: 199.

49. Yamauchi H, Fukuyama H and Shio H. Corpus callosum atrophy in patients with leukoaraiosis may indicate global cognitive impairment. *Stroke; a journal of cerebral circulation*. 2000; 31: 1515-20.
50. Miyamoto N, Pham LD, Hayakawa K, et al. Age-related decline in oligodendrogenesis retards white matter repair in mice. *Stroke; a journal of cerebral circulation*. 2013; 44: 2573-8.
51. Gemmell E, Bosomworth H, Allan L, et al. Hippocampal neuronal atrophy and cognitive function in delayed poststroke and aging-related dementias. *Stroke; a journal of cerebral circulation*. 2012; 43: 808-14.

## Figure legends

**Figure 1. Experimental protocol and animal groups.** **a**, Experimental protocol – BCAS, bilateral common carotid artery stenosis; EE, environmental enrichment **b**, Six different subgroups of mice in this study.

**Figure 2. Volume of the corpus callosum (CC) a(A-F)**, Representative images of LFB + CFV stained sections from block 3 (coronal level of bregma -2.0 mm) in each group. **a(A)**, sham-std; **a(B)**, sham-3hrs; **a(C)**, sham-full; **a(D)**, BCAS-std; **a(E)**, BCAS-3hrs; **a(F)**, BCAS-full. Scale bar represents 200  $\mu$ m. Arrows indicate the peripheral edge of the CC. **b-d**, Histograms showing CC volume of different levels from rostral (**b**), caudal side (**c**) as well as total CC volume (**d**). All BCAS-std and BCAS-full subgroups showed smaller CC volume compared with all sham subgroups ( $P<0.01$ ) (**b-d**). BCAS-3hrs showed larger total CC volume compared with BCAS-std (\*\* $P<0.01$ ) (**c**) ( $*P<0.05$ ) (**d**). Lack of differences were apparent between BCAS-std and BCAS-full (**b, c and d**) or between sham-std and BCAS-3hrs (**b, c and d**). There were no difference between the sham subgroups (**b, c and d**).

**Figure 3. Representative images of LFB + CFV stained corpus callosum a**, Representative images of LFB + CFV stained paramedian part of the CC in each group. **a(A)**, sham-std; **a(B)**, sham-3hrs; **a(C)**, sham-full; **a(D)**, BCAS-std; **a(E)**, BCAS-3hrs; **a(F)**, BCAS-full. Scale bar represents 50  $\mu$ m. **b**, Representative images of LFB + CFV stained median part of the CC in each group. **b(A)**, sham-std; **b(B)**, sham-3hrs; **b(C)**, sham-full; **b(D)**, BCAS-std; **b(E)**, BCAS-3hrs; **b(F)**, BCAS-full. Scale bar represents 50  $\mu$ m. **c**, Representative images of LFB + CFV stained lateral part of the CC in each group. **c(A)**, sham-std; **c(B)**, sham-3hrs; **c(C)**, sham-full; **c(D)**, BCAS-std; **c(E)**, BCAS-3hrs; **c(F)**, BCAS-full. Scale bar represents 50  $\mu$ m.

**Figure 4. White matter (WM) severity score in WM a-i**, Histograms showing WM severity score in the paramedian (**a, d and g**), median (**b, e and h**) and lateral (**c, f and i**) side of corpus callosum (CC) of all different coronal levels. **j-l**, Histograms showing WM severity score in the anterior commissure (**j**), hippocampal fimbria (**k**) and internal capsule (**l**). In all of these WM, BCAS plus limited exposure to EE (BCAS-3hrs) subgroup showed lower WM scores (preserved WM integrity) compared with BCAS-std ( $*P<0.05$ ) (**a-i**) and BCAS-full ( $*P<0.05$ ) (**g**). BCAS-full also attenuated WM damage compared with BCAS-std ( $*P<0.05$ ) (**b, e, h, i and j**) especially in the caudal part of the CC ( $*P<0.05$ ) (**g, h and i**).

**Figure 5. Total number of oligodendrocytes in the corpus callosum (CC).** **a (A-F)**, Representative images of Haematoxylin stained corpus callosum (CC) in each group. **a(A)**, sham-std; **a(B)**, sham-3hrs; **a(C)**, sham-full; **a(D)**, BCAS-std; **a(E)**, BCAS-3hrs; **a(F)**, BCAS-full. Scale bar represents 25  $\mu$ m. **b**, Histogram showing total number of oligodendrocytes in the entire CC. BCAS-3hrs showed increased number of oligodendrocytes in the entire CC compared with BCAS-std and BCAS-full (both  $**P<0.01$ ). BCAS-full showed increased number of oligodendrocytes compared with BCAS-std ( $*P<0.05$ ).

**Figure 6. Assessment of nesting ability a and b(A-F)**, Representative images of nest created from the Nestlet in each group. **a**, A nest seen before surgery (baseline); **b(A-F)**, nests seen 16 weeks post-operation. **b(A)**, sham-std; **b(B)**, sham-3hrs; **b(C)**, sham-full; **b(D)**, BCAS-std; **b(E)**, BCAS-3hrs; **b(F)**, BCAS-full. **c**, Boxplot showing Nestlet score at baseline level and post operation in each group. BCAS-std subgroup had impaired nesting ability compared to baseline level ( $*P<0.05$ ). EE reversed impaired nesting ability induced by BCAS, especially by limited EE (BCAS-3hrs) (**c**).

**Figure 7. Cognitive function assessed in the 9-arm radial maze a**, Table showing probability of each pair of “arm entries and arm repeats (top of each cell)” and “arm repeat score (bottom of each cell)”. Pairs of N/A did not occur in this experiment. **b-c**, Histograms showing arm repeat score of all sham and all BCAS (**b**) and score of each subgroup (**c**). **b**, Average arm repeat scores of all BCAS in sessions 1-10, sessions 11-20 and sessions 1-20, were lower compared with sham ( $*P<0.05$  vs sham;  $\dagger P<0.01$  vs sham;  $\ddagger P<0.01$  vs sham). sham group had dramatically increased arm repeat scores, by average 47.4% in sessions 11-20 compared to sessions 1-10. BCAS group reduced their score by 6.6%. **c**, All sham subgroups had increased arm repeat score (average 49%) in sessions 11-20 compared to sessions 1-10. Limited exposure to EE (BCAS-3hrs) increased their score by 37.3%. BCAS-std only increased by 13.0% and BCAS-full reduced the score by 45.7%. Limited exposure to EE (BCAS-3hrs) subgroup showed the best score between BCAS subgroups. **d**, Table showing probability of each “number of arm entries before first repeat, NABFR (top of each cell)” and “NABFR score (bottom of each cell)”. **e-f**, Graphs showing number of arm entries before first repeat (NABFR) in all sham and all BCAS (**e**) as well as NABFR of each subgroups (**f**). Every consecutive 5 sessions were averaged. **e**, All sham completed more arm entries before first repeat, compared with all BCAS ( $P<0.01$ ). **f**, Between BCAS subgroup, BCAS-3hrs tends to complete more arm entries, especially in sessions 1-5 as well as sessions 11-15. **g-h**, Histograms showing NABFR score of all sham and all BCAS (**g**) and subgroups



**(h)**. Average NABFR scores for all BCAS in sessions 1-10, sessions 11-20 and all sessions 1-20, were lower compared with all sham animals (\*\* $P < 0.01$  vs sham; † $P < 0.01$  vs sham; ‡ $P < 0.01$  vs sham) **(g)** and limited exposure to EE (BCAS-3hrs) subgroup showed the best NABFR score between BCAS subgroups **(h)**.