This is a repository copy of *Metabolic regulation of photosynthetic membrane structure tunes electron transfer function*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/131797/>

Version: Accepted Version

Article:

Johnson, M.P. orcid.org/0000-0002-1663-0205 (2018) Metabolic regulation of photosynthetic membrane structure tunes electron transfer function. *Biochemical Journal*, 475 (7). pp. 1225-1233.

<https://doi.org/10.1042/BCJ20170526>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Metabolic regulation of photosynthetic membrane structure tunes electron transfer function.

Matthew P. Johnson

Department of Molecular Biology and Biotechnology, University of Sheffield, Firth Court, Western Bank, Sheffield, S10 2TN, United Kingdom.

The photosynthetic chloroplast thylakoid membrane of higher plants is a complex 3-dimensional structure that is morphologically dynamic on a timescale of just a few minutes. The membrane dynamics are driven by the phosphorylation of light harvesting complex II (LHCII) by the STN7 kinase, which controls the size of the stacked grana region relative to the unstacked stromal lamellae region. Here I hypothesise that the functional significance of these membrane dynamics is in controlling the partition of electrons between photosynthetic linear and cyclic electron transfer (LET and CET), which determines the ratio of NADPH/ATP produced. The STN7 kinase responds to the metabolic state of the chloroplast by sensing the stromal redox state. High NADPH/ATP ratio leads to reduction of thioredoxin f (TRXf), which reduces a CxxxC motif in the stromal domain of STN7 leading to its inactivation, whereas low NADPH/ATP ratio leads to oxidation of TRXf and STN7 activation. Phosphorylation of LHCII leads to smaller grana, which favour LET by speeding up diffusion of electron carriers plastoquinone (PQ) and plastocyanin (PC) between the domains. In contrast, dephosphorylation of LHCII leads to larger grana that slow the diffusion of PQ and PC, leaving the PQ pool in the stroma more oxidised, thus enhancing the efficiency of CET. The feedback regulation of electron transfer by the downstream metabolism is crucial to plant fitness, since perturbations in the NADPH/ATP ratio can rapidly lead to the inhibition of photosynthesis and photo-oxidative stress.

Introduction

The plant chloroplast thylakoid membrane houses the photosynthetic electron transfer reactions (PET) that provide the NADPH and ATP required for CO₂ fixation by the Calvin-Benson-Bassham (CBB) cycle. Generally, PET and the CBB cycle are conceptualised and studied separately, even though the two processes are intimately coupled. Interactions take place not only at the level of shared turnover of the NADP⁺/NADPH and ADP/ATP pools in the chloroplast, but also in the feedforward and feedback regulation the two processes impart on one another [1-3]. Feedforward regulation of the CBB by PET involves the alteration of the stromal environment in terms of redox potential, pH and ionic status, which modulates the activity of numerous enzymes involved in CO₂ assimilation [3]. In turn, feedback regulation of PET by the CBB cycle influences the redox state of the electron carriers and the size of the proton motive force (Δp) of the thylakoid membrane [1,2]. These factors control the balance between light energy utilisation in the photochemical reactions and the dissipation of this energy as heat by non-photochemical quenching (NPQ) through regulation of the LHCII antenna system and the ATP synthase [4-8]. In principle, the CBB cycle in C₃ plants consumes ATP to NADPH in the strict ratio of 1.5:1, yet the linear electron transfer pathway (LET) from H₂O to NADP⁺ during photosynthesis only produces 1.28 ATP per NADPH if the H⁺/ATP ratio is 4.67, as suggested by early biochemical studies and later the structure of the c-ring of the spinach ATP synthase [9,10]. In addition to production of glyceraldehyde 3-phosphate by the CBB cycle, a range of other metabolic processes also make varying demands on the chloroplast NADPH and ATP pools, including photorespiration, nitrogen and sulphur reduction, and protein, lipid and carbohydrate biosynthesis [1,11,12] (Fig. 1A). The activity of these pathways varies depending on environmental conditions and developmental state of the leaf, thus affecting the ratio of ATP to NADPH required by the cell [1,11,12]. Since the ATP and NADPH pools turnover so rapidly any mismatch in the ATP/NADPH ratio can rapidly inhibit photosynthesis [12-14]. Therefore, several regulatory mechanisms exist in the plant cell for the dissipation of the metabolite in excess or the augmentation of the metabolite in deficit. These mechanisms

include the water-water cycle (WWC or Mehler reaction) [15], the malate valve [16], chlororespiration via plastoquinol oxidase (PTOX) [17] and cyclic electron transfer (CET) [18,19]. In plants CET appears to be the dominant pathway for ATP augmentation and has been shown to be crucial to plant fitness, particularly in fluctuating light conditions [20-22]. Whereas LET involves photosystem II (PSII), plastoquinone (PQ), cytochrome *b₆f* (*cytb₆f*), plastocyanin (PC), photosystem I (PSI), ferredoxin (Fd) and ferredoxin-NADP⁺ reductase (FNR) complexes acting in series and generates NADPH and a proton gradient that is utilised by ATP synthase to make ATP, CET transfers electrons from Fd back to PQ, forming a cycle around PSI. This allows proton transfer and thus synthesis of ~~allowing~~ ATP ~~to be produced~~ without net NADPH formation (Fig. 1B) [18,19]. Therefore, by controlling the balance between LET and CET plants can adjust the ATP/NADPH ratio and contribute to the fulfilment of metabolic demand [1,4,11]. A greatly lowered capacity for electron transfer and photoprotection is observed when CET is knocked out by genetic mutation, suggesting that the balance between LET and CET is central to the proper regulation of photosynthesis *in vivo* [20-22].

In plants two major routes of CET have been discovered to date, the antimycin A-(AA) sensitive Proton Gradient Regulation complex (PGRL1/PGR5)-dependent pathway and AA-insensitive NADPH-like dehydrogenase (NDH)-dependent pathway [18,19] (Fig. 1B). There is evidence that PGRL1/PGR5 acts as an Fd-PQ oxidoreductase that can transiently interact with both PSI and *cytb₆f* [23,24]. Alternatively, it has been suggested that PGR5 may be a regulator of the *cytb₆f* complex [21] and that CET may involve transfer of electrons from a Fd-FNR complex to the Q_n site of *cytb₆f* [25,26], a pathway which may involve the unusual haem c [27,28] (Fig. 1B). The NDH (NADPH-like dehydrogenase complex) is an Fd-PQ oxidoreductase that forms a supercomplex with PSI and shows similarity to Complex I in mitochondria [29-31]. The NDH complex has been shown to act as a proton pump, thus increasing the H⁺/e⁻ ratio of this CET pathway compared to the PGR5/PGRL1 and Fd/FNR/*cytb₆f* routes, which only translocate protons at *cytb₆f* [32].

B

Figure 1. Role of linear and cyclic electron transfer (LET and CET) in photosynthesis. (A) Sunlight initiates electron transfer in the chloroplast thylakoid membrane that leads to the synthesis of NADPH and ATP by LET, with ATP production further augmented by CET. The thylakoid membrane must continually adjust the LET / CET balance to provide the correct ratio of NADPH/ ATP to meet changeable metabolic demand that depends on the relative activity of the Calvin cycle, N and S fixation, photorespiration and biosynthesis in the chloroplast stroma, which vary according to the environmental and developmental state of the plant. **(B)** Lateral heterogeneity in organisation of photosynthetic complexes in the thylakoid membrane. PSII is located in the grana stacks, PSI, ATP synthase and PGRL1/PGR5 and NDH are located in the stromal lamellae and cytb₆f is present in both domains. LET involves electron transfer from water to NADP⁺ via both PSII and PSI, with water oxidation at PSII and PQH₂ oxidation at cytb₆f leading to proton accumulation in the lumen, which is utilised by ATP synthase to drive ATP synthesis. CET involves recycling electrons from Fd to the stromal PQ pool and therefore produces ATP without net NADPH synthesis. Three possible CET pathways based on the NDH, PGRL1/PGR5 and FNR/cytb₆f are shown.

Current models for regulation of the LET/CET balance.

Since the CBB cycle uses 1.5 ATP/NADPH and LET only produces a ratio of 1.28, a constant portion of CET can be expected even in the steady state, estimated at ~15% of the total LET flux [4]. Indeed, different isoforms of Fd and FNR genes, which encode proteins with activities devoted to LET or CET [33] could provide the basis for a constant portion of CET regulated by protein copy number. However, as the developmental state of the leaf and its environment can change with time so the relative demand for ATP/NADPH and so the ratio of LET/CET can be expected to dynamically vary [1,4,18,19]. Several metabolic signals, indicative of an imbalance in the NADPH/ ATP ratio, have been suggested to regulate the amount of CET including the PQ redox state [1, 34], ADP/ATP ratio [35], stromal redox state [36-38], reactive oxygen species [39,40], phosphorylation of NDH [41] and PGRL1 [42] and calcium signalling [43].

The suggested regulation of the LET/CET balance via the PQ redox state is based on the state transition [1, 34]. State transitions in plants involve the dynamic reallocation of LHCII between PSI and PSII, provoked by preferential excitation of PSI (far-red light) or PSII (red/ blue light). PQ reduction in red/blue light activates STN7 kinase [44,45] mediated phosphorylation of LHCII, favouring its association with PSI (State II). PQ oxidation in far-red light inactivates STN7 allowing the constitutively active TAP38 [46,47] to dephosphorylate LHCII, favouring its association with PSII (State I). Reduction of PQ was suggested to

parallel PSI acceptor side limitation due to a high NADPH/ATP ratio [1,34]. The transition to State II would then lead to an increased excitation of PSI relative to PSII, thus increasing CET and making good the ATP deficit. In contrast, a low NADPH/ATP ratio would lead to PQ oxidation via removal of the acceptor side limitation of PSI and a transition to State I and thus enhanced LET to augment NADPH production [1,34]. In line with this, anoxic conditions that lead to induction of State II in the green alga *Chlamydomonas*, also cause increased CET flux and accumulation of a supercomplex containing PGRL1, PSI and *cytb₆f* [37, 48]. However, the CET supercomplex formation and enhanced CET flux still occurred in the absence of state transitions in *Chlamydomonas* [37], similarly in *Arabidopsis* CET activity was unaffected by the absence of the STN7 kinase [49].

An alternative means through which metabolic signals may regulate the LET/CET balance is via control of the partition of electrons between the pathways. CET is sensitive to the 'redox poise' of the electron transfer chain, thus it is inhibited by over-reduction or over-oxidation [50]. Therefore, when high NADPH levels inhibit LET, PQ will tend to be over-reduced thus potentially curtailing the ability of CET to operate. For this reason, LET and CET may require some degree of structural compartmentalisation. One possibility is through the formation of supercomplexes dedicated to CET [29-31,37,48] with a sequestered PQ pool. Yet to date no counterpart to the PGRL1/PSI/*cytb₆f* supercomplex in *Chlamydomonas* [37,48] has been found in plants, although associations between PGRL1 and PSI, PGRL1 and *cytb₆f*, NDH and PSI [29-31], *cytb₆f* and FNR [51], NDH and FNR [52], PSI and FNR [53], and more recently between PSI and *cytb₆f* [31] have been described, although none of these has yet been shown to be more abundant in conditions favouring CET. Another possibility is the maintenance of a separate PQ pools within membrane domains [54, 55]. The thylakoid membrane in plants is divided into two domains; the stacked grana and the interconnecting stromal lamellae with the various electron transfer components heterogeneously distributed among them [56]. PSII is found mainly within the grana stacks while PSI and ATP synthase is found mainly within the stromal lamellae, and *cytb₆f* is found within both domains [56, 57]. There is evidence for compartmentation of PQ in dark adapted thylakoids, with ~70% of the PQ pool available for rapid reduction by PSII ($t_{1/2}$ =25 ms) and the remaining 30% only slowly reduced ($t_{1/2}$ =6 s) [54]. These fast and slow PQ pools were suggested to arise from the grana and stromal lamellae respectively, with diffusion of PQ confined by the extreme protein crowding within the grana [54, 58]. Joliot suggested that the presence of a distinct PQ pool associated with *cytb₆f* in the stromal lamellae, isolated by diffusion limitation from PSII in the grana, could create the conditions necessary for CET to co-exist with LET [54]. In this view, CET and LET would share a common pool of PC and PSI, with injection of electrons into the CET pathways determined by competition at the level of Fd [18]. Importantly the restricted diffusion of PQ is largely abolished upon either unstacking of thylakoids [58] or light adaptation [54], suggesting that changes in membrane organisation can significantly influence the rate of diffusion and potentially therefore the balance between CET and LET.

Figure 2. Variable grana stacking alters the partition of the thylakoid PQ pool and so LET/CET balance. (A) Smaller grana, create a larger area of contact between the grana and stromal lamellae membranes. Under these conditions the diffusion of PC and PQ between grana and stromal lamellae is relatively fast, engaging *cytb₆f* in both domains efficiently in LET which predominates over CET. **(B)** Larger grana reduce the area of contact between the grana and stromal lamellae membranes. Under these conditions the diffusion of PC and PQ between grana and stromal lamellae is relatively slow, leaving the stromal lamellae PQ pool in a more oxidised state that promotes more efficient CET.

A **B**

A new model for the regulation of the LET/CET balance based on dynamic thylakoid stacking.

Here I propose that dynamic changes in thylakoid stacking regulate the LET/ CET balance by altering the degree of partition of the PQ pool. Rapid changes in thylakoid stacking have been observed in spinach upon transition from darkness to low light and from low light to high light [59-61]. Darkness and high light led to larger grana stacks, whereas low light led to smaller grana stacks. These changes parallel the trends in LHCII phosphorylation, which peak at low light and are decreased in high light and darkness [62, 63]. Phosphorylation regulates the surface charge on LHCII and so the cation dependency of stacking [64,65].

Smaller grana have been shown to facilitate faster diffusion of PQH₂ and PC from the grana to stromal lamellae thylakoids and more rapid reduction of PSI, i.e. reduced partition of electron carriers between domains [61] (Fig. 2A). Since the rate-limiting step of LET is the diffusion of PQH₂ from PSII to *cytb₆f* and its oxidation therein, any change in the diffusion time has the potential to affect the overall rate considerably [66]. Indeed, more efficient LET has also been observed in Δ TAP38 *Arabidopsis* plants [46], which possess smaller grana, whereas LET is compromised in the Δ CURT mutant with larger grana [67]. In contrast to enhanced LET, smaller grana and reduced stacking negatively impact on the ability to induce CET [61]. The rate of PSI reduction following far-red illumination was slower when the grana size was smaller indicating reduced CET and transient NPQ generation upon moderate light challenge, which is a proxy for Δ pH generation by CET [22], was also compromised [61]. Thus, larger grana may benefit CET through increased partition of electron carriers between the domains, leaving the stromal PQ pool in a relatively more oxidised state, that is more competitive for electrons from Fd (Fig. 2B). Larger grana are also favoured in high light and darkness conditions [59], where there is evidence CET is upregulated [20,35,68].

How might the STN7 kinase be regulated to allow these dynamics? STN7 is activated when PQH₂ binds to the Q_o site of the *cytb₆f* complex [69], wherein it is suggested to reduce a luminal facing disulphide linkage between two cysteines in the kinase [70]. Therefore, increasing reduction of the PQ pool with light intensity should upregulate STN7. However, a second negative feedback loop exists that sees STN7 inactivated by thioredoxin *f* (TRXf) upon a build-up of reducing power on the PSI acceptor side with increasing light intensity [71, 72]. TRXf is suggested to reduce a buried disulphide linkage between cysteines near the ATP-binding motif on the stromal side of STN7 [70, 72]. STN7 is therefore maximally active at around 100-200 μ mol photons m⁻² s⁻¹ in white light and inactivated in darkness (by PQ oxidation) and in high light (by TRXf) [44, 62, 71, 72]. Interestingly, changes in white light intensity cause no alterations in relative PSI and PSII antenna sizes in contrast to the preferential excitation of the photosystems described above [8, 63]. Thus, under natural white light conditions the hyper-phosphorylation of LHCII, and thus the state transition, is prevented by the TRXf regulation of STN7. Thus, white light driven LHCII phosphorylation is sufficient for changes in grana size, but insufficient to provoke the state transition in plants. This situation contrasts with that in *Chlamydomonas* where the STT7 LHCII kinase (counterpart of STN7) is not inhibited by TRXf and therefore state transitions and stromal over-reduction (and therefore increased CET) co-exist [37, 48].

I suggest that the prevailing NADPH/ ATP ratio in the chloroplast controls the activity of STN7 and thus the size of the grana. In this view, PQ reduction in the light activates STN7 and its activity is then largely regulated by the stromal redox state and therefore the extent of TRXf reduction. Thus, when the NADPH/ ATP ratio is low, TRXf is oxidised, allowing ATP binding and phosphorylation of LHCII (Fig. 3A). In contrast, when ATP is in deficit (high NADPH/ ATP ratio), STN7 is inactivated by reduced TRXf leading to larger grana stacks and increased CET. The amount thylakoid membrane stacking and thus LET/ CET balance is therefore regulated by the metabolic state of the chloroplast. Evidence for metabolic regulation of STN7 already exists in the literature. For instance, treatment of thylakoids with reduced dithiothreitol (DTT) resulted in decreased LHCII phosphorylation and increased CET activation [38, 71], while oxidised DTT stimulated LHCII phosphorylation [62, 71]. In turn, addition of ADP or an ATP sink such as ribulose-5-phosphate, which would elevate the NADPH/ATP ratio, also provokes dephosphorylation of LHCII [73, 74], while addition of ATP or increased adenylate pool energy charge stimulates LHCII phosphorylation through consumption of NADPH and therefore oxidation of TRXf [73]. It is also possible that the PGR5/PGRL1 complex may play a crucial role in sensing the stromal redox state [26] and directly or indirectly regulating STN7 [63]. Such a role for PGR5 would be consistent with two other observations; the lack of LHCII dephosphorylation upon shift from low to high light in the Δ PGR5 mutant of *Arabidopsis* and the stimulation of LHCII phosphorylation in the presence of antimycin A [63, 75]. If grana size remains small in the Δ PGR5 mutant in high light this may partly explain the inefficient CET in this mutant.

Figure 3. Metabolic regulation of the Stn7 kinase. In the light PQ reduction leads to STN7 activation. Two conditions can then ensue **(A)** When the NADPH/ ATP ratio is low, the STN7 kinase is active and LHCII phosphorylation is increased leading to smaller grana that favour LET augmenting NADPH production (dashed arrow) **(B)** When the NADPH/ ATP ratio is high, reduced TRX *f* inactivates the STN7 kinase and the constitutively active TAP38 phosphatase dephosphorylates LHCII leading to larger grana that favour CET, augmenting ATP production (dashed arrow). The PGR5/PGRL1 complex appears to have some as yet undefined regulatory role in sensing the stromal redox poise and its absence [63] or inhibition by antimycin A [76] prevents the inactivation of STN7 by TRX *f*.

Concluding remarks

I have suggested a novel hypothesis that links grana dynamics with LHCII phosphorylation and the regulation of the LET/ CET balance. The key player in this regulatory model is the effect of TRX *f* on STN7 activity, which provides information on the NADPH/ATP ratio in the stroma. There are several important implications of this hypothesis both in terms of future experiments and biological function. The hypothesis predicts a central role for grana stacking in controlling the degree of partition of the PQ pool and thus control of the relative efficiencies of CET versus LET. Therefore, CET should be less competitive for electrons in the Δ TAP38 and *oe*CURT mutants of *Arabidopsis*, which have smaller grana. It is equally important to establish whether the absence of STN7 and /or TAP38 prevent dynamic changes in stacking and understand whether changing PSII (and possibly CURT) phosphorylation play any role. In principle, by manipulating the metabolic state of the chloroplast, it should be possible to manipulate the extent of grana stacking, while keeping the light intensity constant. Moreover, with modern metabolomic approaches it should now be possible to accurately quantify the dynamic transients in metabolite pools that elicit changes in LHCII phosphorylation upon fluctuations in light intensity.

Acknowledgments

I wish to thank the late Professor Jan Anderson FRS for inspiration in writing this hypothesis and dedicate it to her memory. I also wish to thank Professor Peter Horton FRS (University of Sheffield), Professor Neil Hunter FRS (University of Sheffield) and Professor Alexander Ruban (Queen Mary University of London) for useful discussions and critiquing the work. I acknowledge funding from the Biotechnology and Biological Sciences Research Council (U.K.), Leverhulme Trust, Human Frontiers Science Programme and the University of Sheffield Krebs Institute and Grantham Centre for Sustainable Futures.

References

- [1] Horton, P. (1985) Interactions between electron transfer and carbon assimilation. In: Barber, J. and Baker, N.R (eds) *Photosynthetic Mechanisms and the Environment*, pp 135- 187. Elsevier, Amsterdam, New York
- [2] Foyer, C., Furbank, R., Harbinson, J. and Horton, P. (1990) The mechanisms contributing to photosynthetic control of electron transport by carbon assimilation in leaves. *Photosyn. Res.* **25**, 83-100.
- [3] Walker, D. A. and Robinson, S. P. (1978) Regulation of photosynthetic carbon assimilation. In: Siegelman, H.W. and Hind, G. *Photosynthetic carbon assimilation*, pp 43-59. Plenum Press, New York.
- [4] Kramer, D. M., Avenson, T. J. and Edwards, G. E. (2004) Dynamic flexibility in the light reactions of photosynthesis governed by both electron and proton transfer reactions. *Trends Plant Sci.* **9**, 349–357.
- [5] Li, Z., Wakao, S., Fischer, B. B., and Niyogi, K. K. (2009) Sensing and Responding to Excess Light. *Annu. Rev. Plant Biol.* **60**, 239-260.
- [6] Horton, P. (2012) Optimization of light harvesting and photoprotection: molecular mechanisms and physiological consequences. *Phil. Trans. R. Soc. B* **367**, 3455-3465.
- [7] Ruban, A. V., Johnson, M. P. and Duffy, C. D. P (2012) The photoprotective molecular switch in the Photosystem II antenna. *Biochim. Biophys. Acta*, **1817**, 167-181.
- [8] Tikkanen, M. and Aro. E. M. (2014) Integrative regulatory network of plant thylakoid energy transduction. *Trends in Plant Sci.* **19**, 10-17.
- [9] Ort, D. R. and Izawa, S. (1974) Studies on the energy coupling sites of photosynthesis. *Plant Physiol.* **53**, 370-376.
- [10] Seelert, H., Poetsch, A., Dencher, N. A., Engel, A., Stahlberg, H. and Muller, D. J. (2000) Structural biology: proton-powered turbine of a plant motor. *Nature* **405**, 418–419.
- [11] Kramer, D. M. and Evans, J. R. (2011) The importance of energy balance in improving photosynthetic productivity. *Plant Physiol.* **155**, 70–78.
- [12] Noctor, G., and Foyer, C. H. (2000) Homeostasis of adenylate status during photosynthesis in a fluctuating environment. *J. Exp. Bot.* **51**, 347–56.
- [13] Foyer, C. H., Furbank, R. T. and Walker, D. A. (1989) Co-regulation of electron transport and Benson-Calvin cycle activity in isolated spinach chloroplasts. Studies on glycerate 3-phosphate reduction. *Arch Biochem Biophys* **268**, 687-697.
- [14] Slabas, A. R. and Walker D. A. (1976) Transient inhibition by ribose 5-phosphate of photosynthetic O₂ evolution in a reconstituted chloroplast system. *Biochim. Biophys. Acta* **430**, 154-164.
- [15] Miyake, C. (2010) Alternative electron flows (water-water cycle and cyclic electron flow around PSI) in photosynthesis: molecular mechanisms and physiological functions. *Plant Cell Physiol.* **51**, 1951–1963.
- [16] Scheibe, R. (2004) Malate valves to balance cellular energy supply. *Physiol. Plantar.* **120**, 21-26.
- [17] Nawrocki, W. J., Tourasse, N. J., Taly, A., Rappaport, F. and Wollman, F. A. (2015) The Plastid Terminal Oxidase: Its Elusive Function Points to Multiple Contributions to Plastid Physiology. *Annu. Rev. Plant Biol.* **66**, 49–74.
- [18] Johnson, G. N. (2011) Physiology of PSI cyclic electron transport in higher plants *Biochim. Biophys. Acta* **1807**, 384-389.
- [19] Yamori, W. and Shikanai, T. (2016) Physiological functions of cyclic electron transport around photosystem I in sustaining photosynthesis and plant growth. *Annu. Rev. Plant Biol.* **67**, 81-106.
- [20] Munekage, Y., Hashimoto, M., Miyake, C., Tomizawa, K., Endo, T., Tasaka, M., *et al.* (2004) Cyclic electron flow around photosystem I is essential for photosynthesis. *Nature* **429**, 579-582.

- [21] Soursa, M., Jarvi, S., Grieco, M., Nurmi, M., Pietrzykowska, M., Rantala, M., *et al* (2012) Proton gradient regulation 5 is essential for proper acclimation of *Arabidopsis* photosystem I to naturally and artificially fluctuating light conditions. *Plant Cell* **24**, 2394-2948.
- [22] Munekage, Y., Hojo, M., Meurer, J. Endo, T., Tasaka, M. and Shikanai, T. (2002) PGR5 Is Involved in Cyclic Electron Flow around Photosystem I and Is Essential for Photoprotection in *Arabidopsis*. *Cell*. **110**, 361-371.
- [23] DalCorso, G., Pesaresi, P., Masiero, S., Aseeva, E., Schunemann, D., Finazzi, G., *et al* (2008) A complex containing PGRL1 and PGR5 is involved in the switch between linear and cyclic electron flow in *Arabidopsis*. *Cell* **132**, 273–285.
- [24] Hertle, A. P., Blunder, T., Wunder, T., Pesaresi, P., Pribil, M., Armbruster, U., *et al.* (2013) PGRL1 is the elusive ferredoxin- plastoquinone reductase in photosynthetic cyclic electron flow. *Mol. Cell* **49**, 511–23
- [25] Joliot, P., and Johnson, G. (2011) Regulation of cyclic and linear electron flow in higher plants. *Proc. Nat. Acad. Sci. USA*. **108**, 13317-13322.
- [26] Nandha, B., Finazzi, G., Joliot, P., Hald, S. and Johnson, G. N. (2007) The role of PGR5 in the redox poisoning of photosynthetic electron transport. *Biochim. Biophys. Acta* **1767**, 1252–1259.
- [27] Stroebel, D., Choquet, Y., Popot, J.-L., and Picot, D. (2003) An Atypical Haem in the Cytochrome B6F Complex. *Nature* **426**, 413-418.
- [28] Kurisu, G., Zhang, H., Smith, J. L. and Cramer, W. A. (2003) Structure of the cytochrome *b6f* complex of oxygenic photosynthesis: tuning the cavity. *Science* **302**, 1009–14.
- [29] Kouřil, R., Strouhal, O., Nosek, L., Lenobel, R., Chamrad, I., Boekema, E., *et al.* (2014) Structural characterization of a plant photosystem I and NAD(P)H dehydrogenase supercomplex, *Plant J.* **77**, 568–576.
- [30] Peng, L. W., Fukao, Y., Fujiwara, M., Takami, T. and Shikanai, T. (2009) Efficient operation of NAD (P)H dehydrogenase requires supercomplex formation with photosystem I via minor LHCl in *Arabidopsis*, *Plant Cell* **21**, 623–3640.
- [31] Sathish Yadav, K. N., Semchonok, D. A., Nosek, L., Kouril, R., Fucile, G., Boekema, E., *gret al.* (2016) Supercomplexes of plant photosystem I with cytochrome b6f, light-harvesting complex II and NDH. *Biochim. Biophys. Acta* **1858**, 12-20.
- [32] Strand, D., Fisher, N. and Kramer, D. A. (2017) The higher plant plastid NAD(P)H dehydrogenase-like complex (NDH) is a high efficiency proton pump that increases ATP production by cyclic electron flow. *J. Biol. Chem.* **292**, 11850-11860.
- [33] Goss, T. and Hanke, G. (2014) The End of the Line: Can Ferredoxin and Ferredoxin NADP(H) Oxidoreductase Determine the Fate of Photosynthetic Electrons? *Curr. Prot. Pep. Res.* **15**, 384-393.
- [34] Allen, J. F. (1984) Protein Phosphorylation and Optimal Production of ATP in Photosynthesis. *Biochem. Soc. Trans.* **12**, 774-775.
- [35] Joliot, P., and Joliot, A. (2002) Cyclic electron transfer in plant leaf. *Proc. Nat. Acad. Sci. USA* **99**, 10209–10214.
- [36] Breyton, C., Nandha, B., Johnson, G. N., Joliot, P. and Finazzi, G. (2006) Redox modulation of cyclic electron flow around PSI in C3 plants. *Biochemistry*, **45**, 13465-13475.
- [37] Takahashi, H., Clowez, S., Wollman, F. A., Vallon, O. and Rappaport, F. (2013) Cyclic electron flow is redox-controlled but independent of state transition. *Nature Comm.* **4**, 1954.
- [38] Strand, D., Fisher, N., Davis, G. A. and Kramer, D. M. (2016) Redox regulation of the antimycin A sensitive pathway of cyclic electron flow around photosystem I in higher plant thylakoids. *Biochim. Biophys. Acta* **1857**, 1-6.
- [39] Strand, D. D., Livingston, A. K., Satoh-Cruz, M., Froehlich, J. E., Maurino, V. G. and Kramer, D. M. (2015) Activation of cyclic electron flow by hydrogen peroxide *in vivo*. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 5539 – 5544.
- [40] Casano, L.M., Martín, M. and Sabater, B. (2001) Hydrogen peroxide mediates the induction of chloroplastic Ndh complex under photooxidative stress in barley. *Plant Physiol.* **125**, 1450–1458.

- [41] Lascano, H.R., Casano, L.M., Martín, M. and Sabater, B. (2003) The activity of the chloroplastic Ndh complex is regulated by phosphorylation of the NDH-F subunit. *Plant Physiol.* **132**, 256–262.
- [42] Reiland, S., Finazzi, G., Endler, A., Willig, A., Baerenfaller, K., Grossman, J. *et al.* (2011) Comparative phosphoproteome profiling reveals a function of the STN8 kinase in fine-tuning of cyclic electron flow (CEF). *Proc. Nat. Acad. Sci. USA* **108**, 12955–12960.
- [43] Terashima, M., Petroutsos, D., Hudig, M., Tolsygina, I., Trompelt, K., Gabelein, P. *et al.* (2012) Calcium-dependent regulation of cyclic photosynthetic electron transfer by a CAS, ANR1, and PGRL1 complex. *Proc. Nat. Acad. Sci. USA* **109**, 17717–17722.
- [44] Horton, P. and Black, M.T. (1980) Activation of adenosine 5-triphosphate induced quenching of chlorophyll fluorescence by reduced plastoquinone. The basis of State I–State II transitions in chloroplasts. *FEBS Lett* **119**, 141–144.
- [45] Bellafore, S., Barneche, F., Peltier, G. and Rochaix, J. D. (2005) State transitions and light adaptation require chloroplast thylakoid protein kinase STN7. *Nature* **433**, 892–895
- [46] Pribil, M., Pesaresi, P., Hertle, A., Barbato, R. and Leister, D. (2010) Role of plastid protein phosphatase TAP38 in LHCII dephosphorylation and thylakoid electron flow. *PLoS Biol.* **8**, e1000288.
- [47] Shapiguzov, A., Ingelsson, B., Samol, I., Andres, C., Kessler, F. Rochaix, J. D. *et al.* (2010) PPH1 phosphatase is specifically involved in LHCII dephosphorylation and state transitions in *Arabidopsis*. *Proc. Nat. Acad. Sci. USA*. **107**, 4782–4787.
- [48] Iwai, M., Takizawa, K., Tokutsu, R., Okamuro, A., Takahashi, Y. and Minagawa, J. (2010) Isolation of the elusive supercomplex that drives cyclic electron flow in photosynthesis. *Nature* **464**, 1210–1213.
- [49] Pesaresi, P., Hertle, A., Pribil, M., Kleine, T., Wagner, R., Strissel, H. *et al.* (2009) *Arabidopsis* STN7 Kinase Provides a Link between Short- and Long-Term Photosynthetic Acclimation. *Plant Cell* **21**, 2402–2423.
- [50] Grant, B.R. and Whatley, F.R. (1967) Some factors affecting the onset of cyclic photophosphorylation. In: Goodwin TW (ed) *Biochemistry of Chloroplasts*, pp 505–521. Academic Press.
- [51] Zhang, H. M., Whitelegge, J. P. and Cramer, W. A. Ferredoxin : NADP(+) oxidoreductase is a subunit of the chloroplast cytochrome b(6)f complex. *J. Biol. Chem.* **276**, 38159–38165.
- [52] Jose Quiles, M. and Cuello, J. (1998) Association of ferredoxin-NADP oxidoreductase with the chloroplastic pyridine nucleotide dehydrogenase complex in barley leaves. *Plant Physiol.*, **117**, 235–244.
- [53] Andersen, B., Scheller, H.V. and Moller, B.L. (1992) The PSI-E subunit of photosystem I binds ferredoxin:NADP+ oxidoreductase. *FEBS Lett.* **311**, 169–173.
- [54] Joliot, P., Lavergne, J. and Beal, D. (1992) Plastoquinone compartmentation in chloroplasts. I. Evidence for domains with different rates of photo-reduction. *Biochim. Biophys. Acta* **1101**, 1–12.
- [55] Dumas, L., Chazaux, M., Peltier, G., Johnson, X. and Alric, J. (2016) Cytochrome b6f function and localization, phosphorylation state of thylakoid membrane proteins and consequences on cyclic electron flow. *Photosyn. Res.* **29**, 307–320.
- [56] Andersson, B. and Anderson, J. M. (1980) Lateral heterogeneity in the distribution of chlorophyll-protein complexes of the thylakoid membranes of spinach chloroplasts. *Biochim. Biophys. Acta* **593**, 427–440.
- [57] Cox, R.P., and Andersson, B. (1981). Lateral and transverse organisation of cytochromes in the chloroplast thylakoid membrane. *Biochem. Biophys. Res. Commun.* **103**, 1336–1342.
- [58] Kirchhoff, H., Horstmann, S., and Weis, E. (2000). Control of photosynthetic electron transport by PQ diffusion microdomains in thylakoids of higher plants. *Biochim. Biophys. Acta* **1459**, 148–168.
- [59] Rozak, P. R., Seiser, R. M., Wacholtz, W. F. and Wise, R. R. (2002) Rapid, reversible alterations in spinach thylakoid appression upon changes in light intensity. *Plant Cell Environ.* **25**, 421–429.
- [60] Anderson, J. M., Horton, P., Kim, E.-H. and Chow, W. S. (2012) Towards elucidation of dynamic structural changes of plant thylakoid architecture. *Phil. Trans. Roy. Soc. London* **367**, 3515–3524.

- [61] Wood, W. H. J., MacGregor-Chatwin, C., Barnett, S., Mayneord, G., Huang, X., Hobbs, J., *et al.* (2018) Dynamic thylakoid stacking regulates the balance between linear and cyclic photosynthetic electron transfer. *Nature Plants*. doi:10.1038/s41477-017-0092-7
- [62] Rintamaki, E., Salonen, M., Suoranta, U. M., Carlberg, I., Andersson, B. and Aro, E. M. (1997) Phosphorylation of Light-harvesting Complex II and Photosystem II Core Proteins Shows Different Irradiance-dependent Regulation *in Vivo*. *J. Biol. Chem.* **272**, 30476-30482.
- [63] Mekala, N. R., Soursa, M., Rantala, M., Aro, E.M. and Tikkanen, M. (2015) Plants Actively Avoid State Transitions upon Changes in Light Intensity: Role of Light-Harvesting Complex II Protein Dephosphorylation in High Light. *Plant Phys.* **168**, 721-734.
- [64] Barber, J. (1982) Influence of surface charges on thylakoid structure and function. *Annu. Rev. Plant Physiol.*, **33**, 261-295.
- [65] Puthiyaveeti, S., van Oort, B. and Kirchhoff, H. (2017) Surface charge dynamics in photosynthetic membranes and the structural consequences. *Nature Plants* **3**, 17020.
- [66] Haehnel, W. (1984) Photosynthetic electron transport in higher plants. *Ann. Rev. Plant Physiol.* **35**, 659-693.
- [67] Armbruster, U., Labs, M., Pribil, M., Viola, S., Xu, W., Scharfenberg, M., *et al.* (2013) *Arabidopsis* CURVATURE THYLAKOID1 proteins modify thylakoid architecture by inducing membrane curvature. *Plant Cell* **25**, 2661–2678.
- [68] Miyake, C., Miyata, M., Shinzaki, Y. and Tomizawa, K. (2005) CO₂ response of cyclic electron flow around PSI (CEF-PSI) in tobacco leaves—relative electron fluxes through PSI and PSII determine the magnitude of non-photochemical quenching (NPQ) of Chl fluorescence. *Plant Cell Physiol.* **46**, 629–3.
- [69] Vener, A.V., van Kan, P. J. M., Rich, P. R., Ohad, I. and Andersson, B. (1997) Plastoquinol at the quinol oxidation site of reduced cytochrome *bf* mediates signal transduction between light and protein phosphorylation: thylakoid protein kinase deactivation by a single-turnover flash. *Proc. Natl. Acad. Sci. USA* **94**, 1585–1590.
- [70] Puthiaveetil, S. (2011) A mechanism for regulation of chloroplast LHC II kinase by plastoquinol and thioredoxin. *FEBS Lett.* **585**, 1717-1721.
- [71] Rintamaki, E, Martinsuo, P, Pursihemo, S. and Aro, E.M. (2000) Cooperative regulation of light-harvesting complex II phosphorylation via the plastoquinol and ferredoxin-thioredoxin system in chloroplasts. *Proc. Natl Acad. Sci. USA* **97**, 11644-11649.
- [72] Wunder, T., Xu, W., Liu, Q., Wanner, G., Leiaster, D. and Pribil, M. (2013) The major thylakoid protein kinases STN7 and STN8 revisited: effects of altered STN8 levels and regulatory specificities of the STN kinases. *Front. Plant Sci.* **4**, 417.
- [73] Markwell, J. P, Baker, N. R. and Thornber, J. P. (1982) Metabolic regulation of the thylakoid protein kinase. *FEBS Lett.* **142**, 171-174.
- [74] Horton, P. and Foyer, C. (1983) Relationships between protein phosphorylation and electron transport in the reconstituted chloroplast system. *Biochem J.* **210**, 517-521.
- [75] Oxborough, K., Lee, P. and Horton, P. (1987) Regulation of thylakoid protein phosphorylation by high-energy-state quenching. *FEBS Lett.* **221**, 211-214.