



This is a repository copy of *Longitudinal bioimpedance assessments to evaluate hydration in POEMS syndrome*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/131908/>

Version: Published Version

Article:

Nwosu, A., Morris, L., Mayland, C.R. orcid.org/0000-0002-1440-9953 et al. (3 more authors) (2016) Longitudinal bioimpedance assessments to evaluate hydration in POEMS syndrome. *BMJ Supportive and Palliative Care*, 6 (3). pp. 369-372. ISSN 2045-435X

<https://doi.org/10.1136/bmjspcare-2015-000991>

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://www.bmj.com/company/products-services/rights-and-licensing/> This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



OPEN ACCESS

Longitudinal bioimpedance assessments to evaluate hydration in POEMS syndrome

Amara Callistus Nwosu,¹ Lauren Morris,² Catriona Mayland,¹ Stephen Mason,¹ Andrew Pettitt,² John Ellershaw¹

¹Marie Curie Palliative Care Institute Liverpool (MCPCIL), University of Liverpool, Liverpool, UK
²Royal Liverpool and Broadgreen University Hospitals NHS Trust, Merseyside, UK

Correspondence to

Dr Amara Nwosu, Department of Molecular & Clinical Cancer Medicine, Marie Curie Palliative Care Institute Liverpool (MCPCIL), University of Liverpool, Cancer Research Centre, 200 London Rd, Liverpool L3 9TA, UK; anwosu@liverpool.ac.uk

Received 29 July 2015
Revised 27 January 2016
Accepted 4 April 2016
Published Online First
28 April 2016

ABSTRACT

Polyneuropathy, organomegaly, endocrinopathy, M-protein and skin changes (POEMS) syndrome is a rare paraneoplastic disorder associated with an underlying plasma cell dyscrasia and multiorgan failure. POEMS syndrome is potentially fatal and adversely affects quality of life. Oedema is common with many patients affected by pleural effusions, ascites and lower limb oedema. Bioelectrical impedance vector analysis (BIVA) is a non-invasive assessment tool, which enables rapid bedside assessments of nutrition and hydration. This paper describes the use of sequential BIVA assessments to evaluate the response to diuretic therapy in a woman aged 52 years with POEMS syndrome. This case illustrates the potential to use BIVA to conduct longitudinal assessments of hydration status. This provides opportunities for further research using BIVA to monitor hydration and response to interventions. This may be useful in specific situations, for example at the end of life.

INTRODUCTION

Polyneuropathy, organomegaly, endocrinopathy, M-protein and skin changes (POEMS) syndrome is a rare paraneoplastic disorder associated with an underlying plasma cell dyscrasia and multiorgan failure.¹ The pathogenesis of POEMS syndrome is likely caused by overproduction of vascular endothelial growth factor (VEGF).² POEMS syndrome is potentially fatal and adversely affects quality of life. Oedema is common with many patients affected by pleural effusions and ascites.^{1 2}

There is limited evidence to determine the association between hydration and symptoms in advanced cancer.³ Physical examination has a low sensitivity and specificity for identifying fluid deficit and, biochemical methods show little association with hydration status.^{3 4} There is

no routine hydration assessment method for patients with advanced cancer. The evidence for the efficacy of clinically assisted hydration (CAH) in advanced cancer is limited, conflicting and inconclusive. The subject of hydration is extremely important to patients and caregivers; there is concern about the risk of harm to patients through the use or non-use of CAH.³

Bioelectrical impedance vector analysis (BIVA) is a non-invasive, validated body composition assessment method, which may be useful in the assessment of hydration.⁵ To date, there are no published reports about the utility of BIVA in POEMS syndrome.

CASE HISTORY

This case describes the use of BIVA in a woman aged 52 years with POEMS syndrome. She had a history of two autologous stem cell transplants, renal impairment and recurrent lower limb and abdominal oedema. Oedema was a cause of great discomfort and had adversely affected her mobility. She was referred to the specialist palliative care team for symptom management and fluid assessment using BIVA. Following the baseline assessment, she received 40 mg of oral furosemide daily in combination with advice about fluid restriction. Two further BIVA assessments were conducted at weekly intervals following the commencement of diuretic therapy to assess the response to diuretic therapy. A clinical evaluation of peripheral oedema (upper and lower limbs) was also conducted during these assessments.

BIOELECTRICAL IMPEDANCE VECTOR ANALYSIS (BIVA)

Bioimpedance analysis involves a tetrapolar technique to deliver a single-frequency



CrossMark

To cite: Nwosu AC, Morris L, Mayland C, et al. *BMJ Supportive & Palliative Care* 2016;**6**:369–372.

electrical current of 50 kHz. The technique works on the principle that fluid and cellular structures will provide different levels of resistance to an electrical current as it passes through the human body. Bioimpedance provides the following direct measurements: resistance (R) assessing cellular hydration, reactance (Xc) assessing tissue integrity and phase angle (PA), which is reported to be a useful indicator of health and prognosis.⁵ Bioimpedance analysis was conducted using the EFG-3 ElectroFluidGraph Vector Impedance Analyser (Akern) in line with methods and recommendations described elsewhere.⁵ Regression equations of the manufacturer (Akern BodyGram Pro 3.0) were used to calculate total body water (TBW), intracellular water (ICW) and extracellular water (ECW).⁶ These validated equations were derived from previous research.⁷

BIVA enables interpretation of bioimpedance data, which is independent of regression equations and body weight. To establish BIVA, the direct impedance measurements (R and Xc) were plotted as a point (bivariate random vector) on a probability graph (RXc graph); this represented the sex-specific and race-specific tolerance intervals of a non-cancer reference population used for the analysis (figure 1).⁸

RESULTS

At baseline, BIVA demonstrated the participant's overall body composition was just outside the normal ellipse 50th centile and did not suggest fluid overload (figure 1). Following diuretic therapy, the subsequent




Figure 1 Longitudinal change of hydration represented by the BIVA RXc graph. The RXc graph method allows statistical analysis of bivariate distributions of successive impedance vectors of an individual relative to the 50%, 75% and 95% tolerance ellipses of a non-cancer reference population. The black dots represent the study assessments. The black arrow demonstrates the linear trend of the assessments. BIVA, bioelectrical impedance vector analysis; R, resistance; Xc, reactance.

assessments demonstrated a reduction in hydration volume. This corresponded with weight loss and a reduction in clinically detectable oedema. Throughout the assessments, TBW was low relative to body weight (table 1) and ECW was high relative to TBW.

DISCUSSION

Main findings

BIVA demonstrated a reduction in hydration volume following the intervention (diuretic therapy and fluid restriction). This corresponded clinically with weight loss and a reduction in clinically detectable oedema.

Strengths

To our knowledge, this is the first study to use BIVA to evaluate hydration in POEMS syndrome. Furthermore, the evaluation of longitudinal change in hydration in POEMS syndrome using BIVA, following intervention with diuretics, is novel. The advantage of BIVA is that it allows information to be obtained simultaneously about changes in tissue hydration or soft tissue mass, independent of regression equations, or body weight. This allows for accurate interpretation of BIVA readings even if patients are at extremes of weight or volume distribution.

Significance of the findings and comparison with previous studies

The correlations between hydration change, body weight and observable oedema are consistent with non-cancer studies using BIVA to measure hydration.⁹ The participant commented on the simplicity of the BIVA method. Her experience is consistent with the findings of our previous research using BIVA to evaluate hydration in advanced cancer, which adds to the evidence that the BIVA method is not burdensome in palliative care.¹⁰

TBW was low relative to body weight; however, ECW was high compared with TBW. This suggests an imbalance of ECW:ICW ratio, which may be the potential cause of the oedema. Although there was a clinical reduction in weight and oedema, this was not reflected by change in the ECW volume. The reason for this is not clear; however, a potential explanation may be that the bioimpedance regression equations lack the sensitivity to detect small changes in fluid volume.

Limitations

Bioimpedance regression equations are limited in cancer owing to their reliance on physiological assumptions (which include requirements about body shape, the relationship between trunk and leg lengths, soft tissue hydration level and fat fraction).⁵ Although the equations have been validated against reference methods with good accuracy, they may lack accuracy in situations where basic assumptions of the bioimpedance method are not met (eg, in cancer or at

Table 1 Results of bioelectrical impedance analysis

	Baseline	Follow-up 1	Follow-up 2	
Bioimpedance variables N (ohms)				
R (Ω)	679	715	696	
Xc (Ω)	57	57	63	
PA (Ω)	4.8	4.6	5.2	
Weight (kg)	69.5	67.7	66.4	
Height (cm)	157.9	157.9	157.9	
BMI (kg/m ²)	27.9	27.1	27.0	
Hydration variables N (L)				Normal values in an adult woman
TBW	29.6 (43% of body weight)	28.6 (42% of body weight)	28.9 (44% of body weight)	45–50% of body weight
ICW	14.2 (48% of TBW)	13.3 (47% of TBW)	13.3 (47% of TBW)	55% of TBW
ECW	15.4 (52% of TBW)	15.3 (53% of TBW)	15.3 (53% of TBW)	45% of TBW
Clinical examination of oedema				
	Pitting oedema evident around ankles and shin	Small non-pitting oedema to ankle	No oedema detectable	

BMI, body mass index; ECW, extracellular water; ICW, intracellular water; PA, phase angle; R, resistance; TBW, total body water; Xc, reactance.

extremes of body mass and hydration).⁵ The limitations of regression equations are overcome by BIVA, which is independent of the regression analysis and body weight. We acknowledge that we are unable to account for potential confounding factors, which may have influenced hydration between assessments. Furthermore, the relative under-representation of the trunk by whole-body impedance limits our ability to quantify or localise the oedema.⁵ Consequently, BIVA should be interpreted with acknowledgement of the clinical presentation of the individual.

Implications for practice

The properties of BIVA (non-invasive, safe, accurate and painless) highlight its potential to assess hydration in palliative care. BIVA could potentially be used to measure hydration in specific clinical scenarios (eg, bowel obstruction, vomiting and diarrhoea) or to monitor the response to interventions (eg, CAH).⁹ In cancer, BIVA has advantages over regression equations; however, the results should be interpreted carefully with knowledge of the clinical presentation of the patient.

Research opportunities

In research, BIVA can be used to monitor hydration in studies that evaluate the efficacy of CAH. Furthermore, longitudinal hydration assessments could be conducted at specific phases of illness (eg, to examine the relationship between symptomatic burden and clinical hydration states in the last hours and days of life).

CONCLUSION

This case report highlights the potential to use BIVA to monitor hydration states over time in response to

interventions. BIVA overcomes the limitations of bioimpedance regression equations. More research is needed to determine the potential of BIVA to improve the evaluation and management of hydration states in advanced cancer.

Twitter Follow Amara Nwosu at @amaranwosu

Contributors ACN designed the study, conducted the analysis, interpreted the results and wrote the paper. LM, CM, SM, AP and JE assisted with the data analysis and provided critical review of the final manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The results of this analysis will be available upon request.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- 1 Ji ZF, Zhang DY, Weng SQ, *et al.* POEMS syndrome: a report of 14 cases and review of the literature. *ISRN Gastroenterol* 2012;2012:584287.
- 2 Kuwabara S, Dispenzieri A, Arimura K, *et al.* Treatment for POEMS (polyneuropathy, organomegaly, endocrinopathy, M-protein, and skin changes) syndrome. *Cochrane Database Syst Rev* 2012;6:CD006828.
- 3 Nwosu AC, Mayland CR, Mason SR, *et al.* Hydration in advanced cancer: can bioelectrical impedance analysis improve the evidence base? A systematic review of the literature. *J Pain Symptom Manage* 2013;46:433–46.e6.
- 4 Nwosu A, Mayland C, Varro A, *et al.* The assessment of hydration in advanced cancer patients using bioelectrical

- impedance analysis. *BMJ Support Palliat Care* 2014;4 (Suppl 1):A3.
- 5 Barbosa-Silva MC, Barros AJ. Bioelectrical impedance analysis in clinical practice: a new perspective on its use beyond body composition equations. *Curr Opin Clin Nutr Metab Care*. 2005;8:311–17.
 - 6 Akern. Information about the equations used BodyGram Pro Version 3.0 Handbook 2011. 8]. http://www.akern.com/images/stories/Manuals/ENG/BGPRO_ENGManualrev4.pdf
 - 7 Sun SS, Chumlea WC, Heymsfield SB, *et al*. Development of bioelectrical impedance analysis prediction equations for body composition with the use of a multicomponent model for use in epidemiologic surveys. *Am J Clin Nutr* 2003;77:331–40.
 - 8 Piccoli A, Nigrelli S, Caberlotto A, *et al*. Bivariate normal values of the bioelectrical impedance vector in adult and elderly populations. *Am J Clin Nutr* 1995;61:269–70.
 - 9 Jones SL, Tanaka A, Eastwood GM, *et al*. Bioelectrical impedance vector analysis in critically ill patients: a prospective, clinician-blinded investigation. *Crit Care* 2015;19:290.
 - 10 Nwosu AC, Mayland CR, Mason S, *et al*. Patients want to be involved in end-of-life care research. *BMJ Support Palliat Care* 2013;3:457.