



**UNIVERSITY OF LEEDS**

This is a repository copy of *Global Assessment of Agricultural System Redesign for Sustainable Intensification*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/133326/>

Version: Accepted Version

---

**Article:**

Pretty, J, Benton, TG [orcid.org/0000-0002-7448-1973](https://orcid.org/0000-0002-7448-1973), Bharucha, ZP et al. (14 more authors) (2018) *Global Assessment of Agricultural System Redesign for Sustainable Intensification*. *Nature Sustainability*, 1 (8). pp. 441-446. ISSN 2398-9629

<https://doi.org/10.1038/s41893-018-0114-0>

---

© 2018, Springer Nature. This is a post-peer-review, pre-copyedit version of an article published in *Nature Sustainability*. The final authenticated version is available online at: <https://doi.org/10.1038/s41893-018-0114-0>. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

1 **Global Assessment of Agricultural System Redesign for Sustainable**  
2 **Intensification**

3  
4 Jules Pretty\*, School of Biological Sciences, University of Essex, UK  
5 Tim G. Benton, Faculty of Biological Sciences, University of Leeds, UK  
6 Zareen Pervez Bharucha, Global Sustainability Institute, Anglia Ruskin University, UK  
7 Lynn V. Dicks, School of Biological Sciences, University of East Anglia, UK  
8 Cornelia Butler Flora, Kansas State University and Iowa State University, USA  
9 H. Charles J. Godfray, Oxford Martin School, University of Oxford, UK  
10 Dave Goulson, School of Life Sciences, University of Sussex, UK  
11 Sue Hartley, York Environmental Sustainability Institute, University of York, UK  
12 Nic Lampkin, Organic Research Centre, UK  
13 Carol Morris, School of Geography, University of Nottingham, UK  
14 Gary Pierzynski, Ohio Agricultural Experiment Station, Ohio State University, USA  
15 P. V. Vara Prasad, Sustainable Intensification Innovation Lab, Kansas State University, USA  
16 John Reganold, Department of Crop and Soil Sciences, Washington State University, USA  
17 Johan Rockström, Stockholm Resilience Centre, Sweden (from Oct 2018: Potsdam Institute for  
18 Climate Impact Research, Germany)  
19 Pete Smith, Institute of Biological and Environmental Sciences, University of Aberdeen, UK  
20 Peter Thorne, Sustainable Livestock Systems, International Livestock Research Institute, Ethiopia  
21 Steve Wratten, Bio-Protection Research Centre, Lincoln University, New Zealand

22  
23 [4250 words and 50 references, word count includes the two tables, but not supplementary  
24 information and table; abstract is exactly 150 words]

25  
26 \* corresponding author: [jpretty@essex.ac.uk](mailto:jpretty@essex.ac.uk)

27  
28 **Abstract**

29  
30 The sustainable intensification (SI) of agricultural systems offers synergistic opportunities for the co-  
31 production of agricultural and natural capital outcomes. Efficiency and Substitution are steps  
32 towards SI, but system Redesign is essential to deliver optimum outcomes as ecological and  
33 economic conditions change. We show global progress towards SI by farms and hectares, using  
34 seven SI sub-types: integrated pest management, conservation agriculture, integrated crop and  
35 biodiversity, pasture and forage, trees, irrigation management, and small/patch systems. From 47 SI  
36 initiatives at scale (each >10<sup>4</sup> farms or hectares), we estimate 163M farms (29% of all worldwide)  
37 have crossed a redesign threshold, practising forms of SI on 453Mha of agricultural land (9% of  
38 worldwide total). Key challenges include investing to integrate more forms of SI in farming systems,  
39 creating agricultural knowledge economies, and establishing policy measures to scale SI further. We  
40 conclude that SI may be approaching a tipping point where it could be transformative.

41  
42  
43

44 Here we show that the sustainable intensification (SI) of agricultural systems offers synergistic  
45 opportunities for the co-production of agricultural and environmental outcomes. Efficiency and  
46 Substitution are steps towards SI, but system Redesign is essential to deliver optimum outcomes as  
47 ecological and economic conditions change. This global assessment of SI by farms and hectares  
48 categorises SI by seven sub-types: integrated pest management, conservation agriculture, integrated  
49 crop and biodiversity, pasture and forage, trees, irrigation management, and small and patch  
50 systems. From 47 SI initiatives at scale (each >10<sup>4</sup> farms or hectares), we estimate 163M farms (29%  
51 of all worldwide) have crossed a redesign threshold, practising forms of SI on 453 Mha of agricultural  
52 cropped and pasture land (9% of worldwide total). The key challenges centre now on creating  
53 agricultural knowledge economies and establishing policy measures to scale SI further. We conclude  
54 that SI may be at a tipping point where it could be transformative.

55

56

57 The past half century has seen substantial increases in global food production. World population has  
58 risen 2.5 fold since 1960 and yet per-capita food production has grown by 50% over the same period  
59 (1). At the same time, evidence shows that agriculture is the single largest cause of biodiversity loss,  
60 greenhouse gas emissions, consumptive use of freshwater, loading of nutrients into the biosphere  
61 (nitrogen and phosphorus), and a major cause of pollution due to pesticides (2). This is manifested in  
62 soil erosion and degradation, pollution of rivers and seas, depletion of aquifers, and climate forcing  
63 (3). As a consequence, efforts have advanced to develop production systems that at least reduce the  
64 damage footprint per unit produced (4).

65

66 This desire for agricultural systems to produce sufficient and nutritious food without environmental  
67 harm, and going further to produce positive contributions to natural, social and human capital, has  
68 been reflected in calls for a wide range of different types of more sustainable agriculture (5-7). The  
69 dominant paradigm for agricultural development centres on intensification (productivity  
70 enhancement) without integrating sustainability. When the environment is considered, the  
71 conventional focus is on reducing negative impacts rather than exploring synergies between  
72 intensification and sustainability. There is increasing evidence that sustainability frameworks can  
73 improve intensity through shifts in the factors of agricultural production: such as shifts from  
74 fertilizers to nitrogen-fixing legumes as part of rotations or intercropping, from pesticides to natural  
75 enemies, and from ploughing to reduced-intensity tillage.

76

## 77 **Sustainable Intensification**

78

79 Compatibility of *sustainability* and *intensification* was hinted at in the 1980s, then first used in  
80 conjunction with an examination of African agriculture (8). Intensification had previously become  
81 synonymous with types of agriculture that resulted in environmental harm (9). The combination of  
82 the two terms was an attempt to indicate that desirable outcomes, such as more food and better  
83 ecosystem services, need not be mutually exclusive. Both could be achieved by making better use of  
84 land, water, biodiversity, labour, knowledge and technologies. SI was further proposed in a number  
85 of key commissions, its adoption since increasing from about ten papers annually before 2010 to  
86 over 100 per year by 2015 (10). SI is now central to both the UN's Sustainable Development Goals  
87 and wider efforts to improve global food and nutritional security (11).

88

89 Sustainable intensification (SI) is defined as an agricultural process or system where valued  
90 outcomes are maintained or increased while at least maintaining and progressing to substantial  
91 enhancement of environmental outcomes. It incorporates the principles of doing this without the  
92 cultivation of more land (and thus loss of non-farmed habitats), in which increases in overall system  
93 performance incur no net environmental cost (12-15). The concept is open, emphasising outcomes  
94 rather than means, applying to any size of enterprise, and not predetermining technologies,  
95 production type, or particular design components. SI seeks synergies between agricultural and  
96 landscape-wide system components, and can be distinguished from earlier manifestations of  
97 intensification because of the explicit emphasis on a wider set of environmental as well as socially-  
98 progressive outcomes. Central to the concept of SI is an acceptance that there will be no perfect end  
99 point due to the multi-objective nature of sustainability. Thus, no designed system is expected to  
100 succeed forever, with no package of practices fitting the shifting dynamics of every location.

101

102 SI is a necessary but not sufficient component of transformation in the wider food system. Changes  
103 in consumption behaviours (e.g., in animal products), as well as reductions in food waste, may make  
104 greater contributions to the overall sustainability of food and agriculture systems (7), as well as  
105 helping to address the challenge of over-consumption of calorie-dense food, which has become a  
106 global threat to health. System level changes will be necessary from production to consumption, and  
107 eating better is now a priority for affluent countries. At the farm and landscape level, the need for  
108 effective SI is nonetheless urgent. Pressure continues to grow on existing agricultural lands.

109 Environmental degradation reduces the asset base (4, 16), expansion of urban and road  
110 infrastructure captures agricultural land (in the EU28, agricultural land area fell by 31Mha over 50  
111 years from 1961; in the USA and Canada, 0.5Mha are lost annually (17-18)); and climate change and  
112 associated extreme weather create new stresses, testing the resilience of the global food system  
113 (19).

114

115 Attempts to implement SI can result in beneficial outcomes for both agricultural output and natural  
116 capital (14, 20-21). The largest increases in food productivity have occurred in less developed  
117 countries, mostly starting from a lower output base. In industrialised countries, systems have tended  
118 to see increases in efficiency (lower costs), minimizing harm to ecosystem services, and often some  
119 reductions in crop and livestock yields (22). However, the global challenge is significant: planetary  
120 boundaries are under threat or have been exceeded, world population will continue to grow from  
121 7.6 billion (2018) to 10 billion by 2050 (23), and consumption patterns are converging on those  
122 typical in affluent countries for some sections of populations, yet still leaving some 800 million  
123 people hungry worldwide. One question centres on scale: can agriculture still provide sufficient  
124 nutritious food whilst improving natural capital and not compromising other aspects of well-being;  
125 and can this occur at a scale to benefit millions of lives, reverse biodiversity loss and environmental  
126 contamination, and limit greenhouse gas emissions? A further question centres on how much wider  
127 food system changes towards healthier diets could shape the requirements for agricultural  
128 production to focus on both food and environmental outcomes: healthier diets tend to be higher in  
129 fruit, pulse and nut content, therefore more dependent on pollination services (24). Healthier diets  
130 could also generate enhanced consumer demand for lower pesticide residues.

131

132 As SI is an umbrella term that includes a wide range of different agricultural practices and  
133 technologies, the precise extent of existing SI practice has been largely unknown. We use an

134 analytical framework developed for this global assessment data sets of large-scale changes (by  
135 numbers of farms and hectares) that have been made towards SI in this millennium.

136

137

### 138 **Beyond Improved Efficiency and Substitution to Redesign**

139

140 Hill (25) proposed three non-linear stages in transitions towards sustainability: i) efficiency; ii)  
141 substitution; and iii) redesign. While both efficiency and substitution are valuable stages towards  
142 system sustainability, they are not sufficient for ensuring greatest co-production of both favourable  
143 agricultural and environmental outcomes at regional and continental scales (26).

144

145 The first stage: *Efficiency* focuses on making better use of on-farm and imported resources within  
146 existing system configurations. Many agricultural systems are wasteful, permitting natural capital  
147 degradation within the farm or the escape of inputs across system boundaries to cause external  
148 costs on-farm and beyond. Post-harvest losses reduce food availability: tackling them contributes  
149 directly to efficiency gains and amplifies the benefits of yield increases generated by other means.  
150 On-farm efficiency gains can arise from targeting and rationalizing inputs of fertilizer (such as  
151 through deep-fertilizer placement: in Bangladesh used by 1M farmers on 2Mha (27), pesticide, and  
152 water to focus impact, reduce use, and cause less damage to natural capital and human health. Such  
153 precision farming can incorporate sensors, detailed soil mapping, GPS and drone mapping, scouting  
154 for pests, weather and satellite data, information technology, robotics, improved diagnostics and  
155 delivery systems to ensure inputs (e.g., pesticide, fertilizer, water) are applied at the rate and time to  
156 the right place, and only when needed (17, 28-29). Automatic control and satellite navigation of  
157 agricultural vehicles and machinery can enhance energy efficiency and limit soil compaction.

158

159 The second stage: *Substitution* focuses on the replacement of technologies and practices. The  
160 development of new crop varieties and livestock breeds deploys substitution to replace less efficient  
161 system components with alternatives, such as plant varieties better at converting nutrients to  
162 biomass, tolerating drought and/or increases in salinity, and with resistance to specific pests and  
163 diseases. Other forms of Substitution include the release of biological control agents to substitute  
164 for inputs); the use of RNA-based gene silencing pesticides; water-based architecture replacing the  
165 use of soil in hydroponics; and in no-tillage systems new forms of direct seeding and weed  
166 management replacing inversion tillage (14).

167

168 The third stage is a fundamental prerequisite for SI to achieve impact at scale. *Redesign* centres on  
169 the composition and structure of agro-ecosystems to deliver sustainability across all dimensions to  
170 facilitate food, fibre and fuel production at increased rates. Redesign harnesses predation,  
171 parasitism, allelopathy, herbivory, nitrogen fixation, pollination, trophic dependencies and other  
172 agro-ecological processes to develop components that deliver beneficial services for the production  
173 of crops and livestock (30-31). A prime aim is to influence the impacts of agroecosystem management  
174 on externalities (negative and positive), such as greenhouse gas emissions, clean water, carbon  
175 sequestration, biodiversity, and dispersal of pests, pathogens and weeds. While Efficiency and  
176 Substitution tend to be additive and incremental within current production systems, Redesign brings  
177 the most transformative changes across systems.

178

179 *Redesign* is, however, a social and institutional as well as agricultural challenge (31-32), as there is a  
180 need to create and make productive use of human capital in the form of knowledge and capacity to  
181 adapt and innovate, and social capital to promote common landscape-scale change, such as for  
182 positive biodiversity, water quantity and quality, pest management, and soil health outcomes (33-  
183 34). Negative unintended consequences for human, social and economic capital associated with the  
184 system must also be identified and mitigated as part of the redesign process.

185

186 Redesign is critical as ecological, economic, social and political conditions change across whole  
187 landscapes. The changing nature of pest, disease and weed threats illustrates the continuing  
188 challenge (35). New pests and diseases can suddenly emerge in different ways: development of  
189 resistance to pesticides; secondary pests outbreaks due to pesticide overuse; climate change  
190 facilitating new invasions; and accidental long-distance organism transfer. Recent appearances  
191 include wheat blast (*Myrnoportha oryzae*) in Bangladesh (2016), and Fall Army Worm (*Spodoptera*  
192 *fruigiperda*) in sub-Saharan Africa (2017). The papaya mealybug (*Paracoccus marginatus*) is native to  
193 Mexico, but spread to the Caribbean in 1994 then to Pacific islands by 2002, was reported in  
194 Indonesia, India and Sri Lanka by 2008, then to West Africa; the preferred host is papaya, but it has  
195 now colonised mulberry, cassava, tomato and eggplant. Each geographic spread, each shift of host,  
196 requires redesigns of local agricultural systems, and rapid responses from research and extension.  
197 Such new pests and diseases may also impact crop pollinators, as illustrated by host shifts and the  
198 accidental anthropogenic spread of bee parasites (e.g., *Varroa* mites) and pathogens (e.g., *Nosema*  
199 *ceranae*) (36).

200

201

## 202 **Redesign Typology and Methods**

203

204 We analysed transitions towards redesign in agricultural systems worldwide. We reviewed literature  
205 on SI, including meta-analyses and practices, to produce a typology of seven system types that we  
206 classify as redesign: (i) integrated pest management, (ii) conservation agriculture, (iii) integrated  
207 crop and biodiversity, (iv) pasture and forage, (v) trees in agricultural systems, (vi) irrigation water  
208 management and (vii) intensive small and patch systems (Table 1). These seven systems and  
209 illustrative sub-types are discussed in more detail in Supplementary Section 1.

210

211 The seven system types span both industrialised and less-developed countries, and zones from  
212 temperate to tropical. Progress towards SI in developing countries is occurring in the context of the  
213 pressing need to implement sustainable development goals for poverty reduction, improved  
214 livelihoods and better nutrition by building more productive and sustainable systems of smallholder  
215 agriculture. There are some 570 million farms worldwide, 84% of which are landholdings of less than  
216 2 ha (37). These small farms make up 12% of total agricultural area, yet produce 70% of food in  
217 Africa and Asia. Sustainable intensification will have to be effective worldwide, yet will have to reach  
218 larger numbers of farms in less developed countries: 74% of all farms are in Asia (of which 35% are in  
219 China and 24% in India), 9% in Sub-Saharan Africa, 7% in Central Europe and Central Asia, 3% in Latin  
220 America and the Caribbean, and 3% in Middle East and North Africa. Owing to the average size of  
221 the 4% of farms in industrialised countries, the choices made by a single farmer can have landscape-  
222 wide consequences.

223

224  
225

**Table 1. Redesign typology and examples of sub-types of intervention**

<b>Redesign type</b>	<b>Illustrative redesign sub-types of intervention</b>
1. Integrated pest management (IPM)	IPM through farmer field schools Integrated plant and pest management Push-pull systems
2. Conservation agriculture (CA)	Conservation agriculture practices Zero- and low-tillage Soil conservation and soil erosion prevention Enhancement of soil health
3. Integrated crop and biodiversity redesign	Organic agriculture Rice-fish systems Systems of crop and rice intensification (SCI, SRI) Zero-budget natural farming (ZBNF) Science and technology backyard platforms Farmer wisdom networks Landcare and watershed management groups
4. Pasture and forage redesign	Mixed forage-crop systems Management intensive rotational grazing systems (MIRGs) Agropastoral field schools
5. Trees in agricultural systems	Agroforestry Joint and collective forest management Leguminous fertilizer trees and shrubs
6. Irrigation water management	Water user associations Participatory irrigation management Watershed management Micro-irrigation technologies
7. Intensive small and patch scale systems	Community farms, allotments, backyard gardens, raised beds Vertical farms Group purchasing associations and artisanal small producers (in Community Supported Agriculture, tekei groups, guilds) Micro-credit groups for small-scale intensification Integrated aquaculture

226 Note: i) This is an illustrative list of sub-types; ii) Some sub-types span a number of types (e.g., organic agriculture also  
227 appears in elements of 4 and 7); iii) Community Supported Agriculture operations (CSAs) are group purchasing associations  
228 in North America and the UK, tekei groups are in Japan, guilds in France, Belgium and Switzerland.

229  
230

231 We have screened 400 SI projects, programmes and initiatives worldwide (drawn from literature or  
232 existing data sets (20-21, 35) and selected those implemented to a scale greater than 10<sup>4</sup> farms or  
233 hectares. Our intention is not to map all innovation for SI worldwide, but to assess where innovation  
234 has scaled to have potentially positive outcomes on ecosystem services as well as agricultural  
235 objectives across landscapes.

236  
237

## 238 **Results**

239

240 Forty-seven SI initiatives have exceeded the 10<sup>4</sup> scale, of which 17 exceed the 10<sup>5</sup> threshold, and 14  
241 the 10<sup>6</sup> scale (Supplementary Table 1; Figures 1 and 2). Many SI initiatives worldwide show promise  
242 but remain limited in scale (either demonstrating locally-dependent conditioning, or the lack of  
243 attention to scalar mechanisms). We estimate from these projects-initiatives in some 100 countries  
244 that 163 million farms have crossed an important substitution-redesign threshold, and are using SI


245 methods, in at least one farm enterprise, on an area approaching 453 million ha of agricultural land  
 246 (not counting the SI initiatives in home and urban gardens and on field boundaries). This comprises  
 247 29% of all farms worldwide; and 9% of agricultural land (total worldwide crop and pasture land is 4.9  
 248 x 10<sup>9</sup> hectares).

249

250 We note that this global assessment might imply numbers of farms and hectares are fixed: on the  
 251 ground, there will be a flux in numbers as a result of both adoption and dis-adoption. This may arise  
 252 from farmer choice and agency, but equally from the actions of vested interests, agricultural input  
 253 companies, consolidation of small farms into larger operations, changes in agricultural policy or  
 254 shifts in market demand, and discrepancies between on-paper claims and what farmers have  
 255 implemented. We have also not included apparent adoption in this assessment: for example, EU  
 256 regulations require all farms to use IPM, but this has not yet led to significant uptake of agricultural  
 257 practices that significantly benefit ecosystem services (21).

258


259



260

261





262

263

264

265

## 266 The Co-creation of Agricultural Knowledge Economies

267

268 For SI to have a transformative impact on whole landscapes, it requires cooperation, or at least  
 269 individual actions that collectively result in additive or synergistic benefits. For farmers to be able to  
 270 adapt their agroecosystems in the face of stresses, they will need to have the confidence to  
 271 innovate. As ecological, climatic, and economic conditions change, and as knowledge evolves, so  
 272 must the capacity of farmers and communities to allow them to drive transitions through processes  
 273 of collective social learning. This suggests a valued property of intrinsic adaptability, whereby  
 274 interventions that can be adapted by users to evolve with changing environmental, economic and  
 275 social conditions are likely to be more sustainable than those requiring a rigid set of conditions to  
 276 function. Every example of successful redesign for SI at scale has involved the prior building of social  
 277 capital (32), in which emphasis is paid to: i) relations of trust, ii) reciprocity and exchange, iii)  
 278 common rules, norms and sanctions, and iv) connectedness in groups. As social capital lowers the  
 279 costs of working together, it facilitates co-operation, and people have the confidence to invest in  
 280 collective activities, knowing that others will do so too. They are also less likely to engage in free-  
 281 rider actions that result in resource degradation.

282

283 This suggests the need for new knowledge economies for agriculture (38). The technologies and  
 284 practices increasingly exist to provide both positive food and ecosystem outcomes: new knowledge  
 285 needs to be co-created and deployed in an interconnected fashion, with an emphasis on ecological  
 286 as well as technological innovation. This includes the need to rebuild extension systems and extend  
 287 them to environmental as well as agronomic skills, with farmer field schools already dense enough in  
 288 some locations that they have transformed knowledge co-creation and behavioural change (34).  
 289 Important examples in industrialised countries include the Landcare movement in Australia with

290 6000 groups, farmer-led watershed councils and the long-term agroecosystem research network in  
291 the USA, the French network of agroecology farms, and the 49 Farmer Cluster Initiatives in the UK  
292 (39-40). These have created platforms for creation of practices to address locally specific problems  
293 of erosion, nutrient loss, pathogen escape and waterlogging. In Cuba, the *Campesino-a-Campesino*  
294 movement integrates agroecology into redesign, with knowledge and technologies spread through  
295 exchange and cooperatives: productivity of 100,000 farmers increased by 150% over ten years, and  
296 pesticide use fell to 15% of former levels (41). In West Africa, innovation platforms have increased  
297 yield in maize and cassava systems (42), and in Bangladesh have resulted in the development and  
298 spread of direct seeded and early-maturing rice (43). In China, Science and Technology Backyard  
299 (STB) platforms operate in 21 provinces covering many crops: wheat, maize, rice, soybean, potato,  
300 mango and lychee (44). STB platforms bring agricultural scientists to live in villages, and use field  
301 demonstrations and farm schools to engage farmers in developing innovations: reasons for success  
302 centre on in-person communication, socio-cultural bonding, and the trust developed among farmer  
303 groups of 30-40 individuals.

304  
305

### 306 **Next Steps: A Tipping Point**

307

308 This analysis shows that the expansion of SI has begun to occur at scale across a wide range of  
309 agroecosystems. The benefits of both scientific and farmer input into technologies and practices that  
310 combine crops and animals with appropriate agro-ecological and agronomic management are  
311 increasingly evident. The associated creation of novel social infrastructure results in both flows of  
312 information and builds trust among individuals and agencies. This should result in the improvement  
313 of farmer knowledge and capacity through the use of platforms for cooperation together with digital  
314 communication technologies.

315

316 The key question thus centres on what could happen next. SI has been shown to increase  
317 productivity (4-5), raise system diversity (3), reduce farmer costs (20, 22, 30), reduce negative  
318 externalities (12-13, 30), and improve ecosystem services (26, 30). There are thus a range of  
319 potential motivations for farmers to adopt SI approaches, and for policy support to be provided by  
320 national government, third sector and international organisations. SI requires investments, though,  
321 to build natural, social and human capital, so is not costless (6-7). In all 47 initiatives, there are  
322 differences in SI adoption by types of farm, farmers, and SI sub-type. All innovations begin on a small  
323 scale, yet here expanded to exceed the 10<sup>4</sup> scale for farm numbers and/or hectares. But several  
324 hundred more projects remain small in scale or are at early stages of development. In some cases,  
325 innovations started with efficiency or substitution interventions, and then spread to redesign (31). In  
326 every case, social capital formation leading to knowledge co-creation has been a critical pre-  
327 requisite. In every case, too, farmer benefit (e.g. food output, income, health) will have been  
328 demonstrated and understood.

329

330 In most contexts, though, state policies for SI remain poorly developed or counter-productive. In the  
331 EU, farm subsidies have increasingly been shifting towards targeted environmental outcomes rather  
332 than payments for production, a process the UK Government has plans to accelerate (45-46), but  
333 this seldom guarantees synergistic benefits across whole landscapes. Several countries have offered  
334 explicit public policy support to social group formation, such as for Landcare (Australia), watershed

335 management (India), joint forest management (India, Nepal, DR Congo), irrigation user groups  
 336 (Mexico) and farmer field schools (Indonesia, Burkina Faso). In India's state of Andhra Pradesh, the  
 337 state government has made explicit its support to zero-budget natural farming (local form of  
 338 uncertified organic farming), aiming to reach 6 million farmers by 2027 (47); in Bhutan and the  
 339 Indian states of Kerala and Sikkim, policy commitments have been made to convert all land to  
 340 organic agriculture; the greening of the Sahel through agroforestry began when national tree  
 341 ownership regulations were changed to favour local people (12). In China, the 2016 No 1 Central  
 342 Document emphasises innovation, coordination, greening and sharing as key parts of a new strategy  
 343 for SI (48). At the same time, consumers are increasingly playing a role in connecting directly with  
 344 farmers in affluent countries, such as through group purchasing schemes, farmers' markets and  
 345 certification schemes, which may in turn change consumption choices (49).

346  
 347 With this growing understanding of the positive roles governments can play in structuring incentives  
 348 and policies, as well as supporting agricultural knowledge economies, we anticipate that SI may be at  
 349 a tipping point (2, 4). A further small increase in the number of farms successfully operating re-  
 350 designed agricultural systems could lead rapidly to re-design of agriculture on a global scale. To  
 351 transform agriculture to provide comprehensive sustainably intensified systems that can deliver  
 352 adequate, healthy food for all people, will require the integration of different redesign types to  
 353 create system-wide transitions, and the internalisation of agricultural externalities into prices or  
 354 through consumer demand. Our hypothesis is that important synergies are occurring, where  
 355 redesigned systems will deliver more than the sum of the parts, and that when more than one SI  
 356 sub-type is combined, the likelihood will increase that redesigned systems will be better fitted to  
 357 local circumstances and thus be more resilient. In the 47 initiatives analysed here, we scored for the  
 358 number of types used in each initiative (Table 2). Most initiatives are deploying one (25% of farms,  
 359 37% of hectares) or two (66% of farms, 52% of hectares) types. The most common paired  
 360 combinations were integrated crop and biodiversity redesign with either IPM, CA and soil health,  
 361 agroforestry and irrigation management. The most common deployment of only one sub-type was  
 362 trees in agricultural systems. This suggests a clear challenge centres on further integration: this  
 363 might include, for example, combining conservation agriculture for soil health with integrated  
 364 watershed management, nutrient recycling and integrated pest management.

365  
 366  
 367 **Table 2. Number of redesign types of SI deployed in each of 47 initiatives, by farm and hectare numbers and**  
 368 **proportions**

	Number of redesign types deployed				
	1	2	3	4	5-7
Farms (M)	50.7	132.5	16.1	1.0	0.0
Proportion of farms in each redesign type	25.3%	66.1%	8.0%	0.5%	0.0%
Hectares (Mha)	170.2	240.5	32.8	19.5	0.0
Proportion of hectares in each redesign type	36.8%	51.9%	7.1%	4.2%	0.0%

369  
 370  
 371 There is much to be done to ensure agricultural and food systems worldwide increase the  
 372 production of nutritious food whilst ensuring positive impacts on natural and social capital. Some

373 efficiency-based initiatives are reaching large numbers of farmers, such as the 21M reducing  
374 fertilizer use in China (50). We conclude that a transition from efficiency through substitution to  
375 redesign will be essential, suggesting that the concept and practice of SI of agriculture will be a  
376 process of adaptation, driven by a wide range of actors cooperating in new agricultural knowledge  
377 economies. This will still need farmers and society to invest in SI, not just for the sake of  
378 sustainability, but for livelihoods and profitability. There are risks: technologies could be dis-  
379 adopted, advances lost, and competing interests could co-opt and dilute innovations. Positive  
380 changes towards consuming healthier food and reductions in food waste may also not occur, putting  
381 more pressure on farmers to produce more food at any cost.

382

383 We conclude by recommending that three key questions will need addressing for SI to fulfil its  
384 potential across agro-ecosystems worldwide:

385

- 386 1. What further evidence is needed to spread SI innovations as options of choice and best  
387 practice globally, thus contributing to further progress towards global food security and  
388 landscape-wide benefits for natural capital?
- 389 2. How can agricultural systems be redesigned to ensure it is more profitable to maintain,  
390 rather than erode, natural capital?
- 391 3. How can national policy support for the mainstreaming of SI be strengthened and  
392 implemented within and across all countries?

393

394

395

396

397 **A Note on Terminology**

398 There is no single accepted terminology for grouping of types of countries. Terms relate to past  
399 stages of development (developed, developing, less developed), state of economy or wealth  
400 (industrialised, affluent), geographic location (global south or north), or membership (OECD, non-  
401 OECD). None are perfect: China has the second largest economy measured by GDP (which does not  
402 measure all aspects of economies, environments and societies well), yet might be considered still  
403 developing or less-developed. The USA has the largest economy by GDP, yet has nearly 50M hungry  
404 people. Here we have simply used *industrialised* and *less-developed*, and acknowledge the  
405 shortcomings. We also use the term pesticide to incorporate all synthesised pest, disease, weed and  
406 other control compounds.

407

408 **Acknowledgements**

409 We are grateful to a number of people for their guidance and updates on numbers of farmers and  
410 hectares for some of the illustrative sub-types: Henk van den Berg, Roland Bunch, Kevin Gallagher  
411 and Vijay Kumar.

412

413 **Authors**

414 The author to whom correspondence and requests for materials should be addressed is JP. The  
415 design of this study was conducted by JP and ZB; all authors (JP, TB, CBF, LD, CG, DG, SH, NL, CM, GP,  
416 VP, JR, JR, PS, PT, SW, ZB) were equally engaged in data gathering, analysis and assessment, and  
417 writing of the paper and supplementary file.

418

419 **Data Statement and Availability**

420 The data that support the findings of this study are available from the corresponding author upon  
421 request. The supplemental file contains detail of each of the initiatives (farmers, hectares), and all  
422 references to the data are provided in both the paper and supplementary information.

423

424 **Competing Interests**

425 The authors declare there are no competing interests in this paper, as defined as financial and non-  
426 financial interests that could directly undermine, or be perceived to undermine the objectivity,  
427 integrity and value of a publication, through a potential influence on the judgements and actions of  
428 authors with regard to objective data presentation, analysis and interpretation.

429

430 **References**

431

- 432 1. FAO. 2017. FAOSTAT Online database. Rome
- 433 2. West P C et al. 2014. Leverage points for improving global food security and the environment. *Science* 345 (6194),  
434 325-328
- 435 3. Pywell R F et al. 2015. Wildlife friendly farming increases crop yield: evidence for ecological intensification. *Proc Royal*  
436 *Society Lond. B* 282: 20151740
- 437 4. Rockström J et al. 2017. Sustainable intensification of agriculture for human prosperity and global sustainability.  
438 *Ambio* 46, 4–17
- 439 5. Foresight. 2011. *The Future of Global Food and Farming*. London: Government Office for Science London
- 440 6. FAO 2016. *Save and Grow: Maize, Rice and Wheat – A Guide to Sustainable Crop Production*. Rome
- 441 7. Benton T. 2015. Sustainable intensification. In Pritchard B, Ortiz R and Shekar M (eds). *Routledge Handbook of Food*  
442 *and Nutrition Security*. Routledge, Abingdon
- 443 8. Pretty J. 1997. The sustainable intensification of agriculture. *Natural Resources Forum* 21(4) 247–256
- 444 9. Collier W L et al. 1973. Recent changes in rice harvesting methods. Some serious social implications. *Bulletin of*  
445 *Indonesian Economic Studies* 9(2): 36-45

- 446 10. Gunton R M et al. 2016. How scalable is sustainable intensification? *Nature Plants* 2: 1-4
- 447 11. UN Sustainable Development Platform. 2017. Sustainable Development Goals.
- 448 <http://www.un.org/sustainabledevelopment/sustainable-development-goals/>
- 449 12. Godfray H C J et al. 2010. Food security: the challenge of feeding 9 billion people. *Science* 327: 812–818
- 450 13. Smith P. 2013. Delivering food security without increasing pressure on land. *Global Food Security* 2(1), 18-23
- 451 14. Pretty J and Bharucha Z P. 2014. Sustainable intensification in agricultural systems. *Annals of Botany* 205, 1-26.
- 452 15. Geertsema W et al. 2016. Actionable knowledge for ecological intensification of agriculture. *Frontiers in Ecology and*
- 453 *the Environment*, 14(4), 209-216
- 454 16. Hallman, C A et al. 2017. More than 75 percent decline over 27 years in total flying insect biomass in protected
- 455 areas. *PLOS ONE* 12 (10). e0185809
- 456 17. Buckwell A et al. 2014. *The Sustainable Intensification of European Agriculture*. Brussels: RISE Foundation
- 457 18. Francis C A et al. 2012. Farmland conversion to non-agricultural uses in the US and Canada: current impacts and
- 458 concerns for the future. *Int J Agric Sust* 10(1), 8-24
- 459 19. IPCC. 2014. *Fifth Assessment Report (AR5)*. Geneva
- 460 20. Pretty J et al. 2006. Resource-conserving agriculture increases yields in developing countries. *Environmental Science*
- 461 *and Technology* 40: 1114-19
- 462 21. Pretty J et al. 2011. Sustainable intensification in African agriculture. *Internat. J Agric Sustainability* 9(1): 5-24
- 463 22. Reganold J P and Wachter J M. 2016. Organic agriculture in the 21<sup>st</sup> century. *Nature Plants* 2(2): 15221
- 464 23. UN. 2017. *World Population Prospects: 2017 Revision*. Department of Economic and Social Affairs, NY
- 465 24. Smith M R et al. 2015. Effects of decreases of animal pollinators on human nutrition and global health: a modelling
- 466 analysis. *Lancet* 386, 1964–1972
- 467 25. Hill S. 1985. Redesigning the food system for sustainability. *Alternatives* 12, 32-36
- 468 26. Sandhu H et al. 2015. Significance and value of non-traded ecosystem services on farmland. *PeerJ*, 3, p.e762
- 469 27. Mulligan K. 2016. *Fertilizer Deep Placement*. Feed the Future, USAID, Washington DC
- 470 28. Garbach K et al. 2017. Examining multi-functionality for crop yield and ecosystem services in five systems of
- 471 agroecological intensification. *Internat J Agric Sust* 15(1), 11-28
- 472 29. Lampkin N H et al. 2015. The role of agroecology in sustainable intensification. Report for the Land Use Policy Group.
- 473 Organic Research Centre, Elm Farm and Game & Wildlife Conservation Trust
- 474 30. Gurr G M et al. 2016. Multi-country evidence that crop diversification promotes ecological intensification of
- 475 agriculture. *Nature Plants* 2(3), 16014
- 476 31. Gliessman S R and Rosemeyer M (eds.) 2009. *The Conversion to Sustainable Agriculture: Principles, Processes, and*
- 477 *Practices*. CRC Press, Boca Raton
- 478 32. Hartley S E et al. 2015. Defending the leaf surface: intra- and inter-specific differences in silicon deposition in grasses
- 479 in response to damage and silicon supply. *Frontiers in Plant Science* 6: 35
- 480 33. Pretty J. 2003. Social capital and the collective management of resources. *Science* 302: 1912-1915
- 481 34. FAO. 2016. *Farmer Field School Guidance Document*. Rome.
- 482 35. Pretty J and Bharucha Z P. 2015. Integrated pest management for sustainable intensification of agriculture in Asia and
- 483 Africa. *Insects* 6, 152-82
- 484 36. Goulson D et al. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers.
- 485 *Science* 347(6229), p.1255957
- 486 37. Lowder S K et al. 2016. The number, size, and distribution of farms, smallholder farms, and family farms worldwide.
- 487 *World Development* 87, 16-29
- 488 38. MacMillan T and Benton T. 2014 Engage farmers in research. *Nature* 509, 25-27
- 489 39. Spiegall S et al. 2018. Evaluating strategies for sustainable intensification of US agriculture through the Long-Term
- 490 Agroecosystem Research network. *Environ. Res. Lett.* 13 (2018) 034031
- 491 40. Campbell A et al. 2017. Reflections on four decades of land restoration in Australia. *The Rangeland Journal*
- 492 doi.org/10.1071/RJ17056
- 493 41. Rosset P M et al. 2011. The Campesino-to-Campesino agroecology movement of ANAP in Cuba: social process
- 494 methodology in the construction of sustainable peasant agriculture and food sovereignty. *The Journal of peasant*
- 495 *studies*, 38(1), 161-191
- 496 42. Jatoe J P D et al. 2015. Does sustainable agricultural growth require a system of innovation? Evidence from Ghana and
- 497 Burkina Faso. *Int J Agric Sust* 13(2): 104-119
- 498 43. Malabayabas A J B et al. 2014. Impacts of direct-seeded and early-maturing varieties of rice on mitigating seasonal
- 499 hunger for farming communities in northwest Bangladesh. *Internat J Agric Sust*, 12(4), 459-470
- 500 44. Zhang W et al. 2016. Closing yield gaps in China by empowering smallholder farmers. *Nature* 537(7622): 671-674
- 501 45. Defra. 2018. *A Green Future: Our 25 Year Plan to Improve the Environment*. London
- 502 46. Morris C et al. 2017. Sustainable Intensification: the view from the farm. *Aspects of Applied Biology* 136, 19-26
- 503 47. Kumar V. T. 2017. Zero-Budget Nature Farming. Department of Agriculture , Government of Andhra Pradesh,
- 504 Hyderabad
- 505 48. Xinhua 2016. CPC and State Council Guide Opinion on Using New Development Concepts to Accelerate Agricultural
- 506 Modernisation and Realise Moderate Prosperity Society. URL [http://news.xinhuanet.com/fortune/2016-](http://news.xinhuanet.com/fortune/2016-01/27/c_1117916568.htm)
- 507 [01/27/c\\_1117916568.htm](http://news.xinhuanet.com/fortune/2016-01/27/c_1117916568.htm)

- 508 49. Allen J E et al. 2017. Do Community Supported Agriculture programmes encourage change to food lifestyle behaviours  
509 and health outcomes? New evidence from shareholders. *Internat J Agric Sust* 15 (1), 70-82  
510 50. Cui Z L et al. 2018. Pursuing sustainable productivity with millions of smallholder farmers. *Nature*  
511 doi:10.1038/nature25785  
512  
513  
514

515 **Figure legends**

516

517

518 Figure 1. Farm numbers and hectares under seven types of Sustainable Intensification (47  
519 initiatives)

520

521 Figure 2: Seven types of sustainable intensification (47 initiatives): disc area (Mha)

522

523