UNIVERSITY OF LEEDS

This is a repository copy of *Tackling burnout in UK trainee doctors is vital for a sustainable, safe, high quality NHS*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/135842/>

Version: Accepted Version

Article:

Johnson, J orcid.org/0000-0003-0431-013X, Bu, C and Panagioti, M (2018) Tackling burnout in UK trainee doctors is vital for a sustainable, safe, high quality NHS. BMJ, 362. k3705. ISSN 0959-8138

<https://doi.org/10.1136/bmj.k3705>

Protected by copyright. BMJ Publishing Group Limited. "This article has been accepted for publication in BMJ 2018 following peer review, and the Version of Record can be accessed online at <https://doi.org/10.1136/bmj.k3705>.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Tackling burnout in UK trainee doctors is vital for a sustainable, safe, high quality NHS

Judith Johnson, Christopher Bu and Maria Panagioti

Letter to the BMJ; published September 2018

We read the GMC survey findings on burnout in trainee and trainer doctors with great interest. Increasing evidence suggests doctor wellbeing is a marker of the quality of care delivered to patients. These results therefore raise cause for concern in relation to patient care. Our previous studies have shown a consistent link between healthcare staff burnout and patient safety[1-3]. In a recent meta-comprehensive analysis accepted for publication in JAMA Internal Medicine[4], we examined the association between burnout in doctors and patient safety incidents, low professionalism and patient satisfaction. Of particular relevance is our finding that the association between burnout and low professionalism was two times higher in trainee and early career doctors compared with more experienced doctors. As such, addressing burnout particularly in trainee doctors has strategic importance for the provision of safe, high quality patient care.

Two recent meta-analytic reviews have established that interventions to reduce burnout in doctors are effective [5,6]. Unfortunately, these interventions have been developed and evaluated mostly in the US. National research investments are needed to develop and evaluate interventions for enhancing the wellbeing of trainee UK doctors which take into consideration the distinctive characteristics of the UK health care system. We have recently piloted the delivery of i) a resilience training intervention in obstetric and gynaecological registrar doctors as part of the Yorkshire and Humber Patient Safety Translational Research Centre and ii) mindfulness courses in foundation year doctors in the Royal Liverpool Hospital following the award of successful institutional funding. Both courses have been well-received by trainee doctors. Although such programmes should be accompanied by organisational improvements, these early encouraging findings suggest that the wellbeing of trainee UK doctors is subject to establishing a flourishing training environment and an engaging career culture.

Judith Johnson; Lecturer; University of Leeds and Bradford Institute for Health Research

Christopher Bu, Foundation Year 2 Doctor, Royal Liverpool & Broadgreen University Hospitals NHS Trust

Maria Panagioti; Senior Research Fellow; NIHR School for Primary Care Research, NIHR Greater Manchester Patient Safety Translational Research Centre, Manchester Academic Health Science Centre, University of Manchester

1. Johnson J, Louch G, Dunning A, et al. Burnout mediates the association between symptoms of depression and patient safety perceptions: A cross-sectional study in hospital nurses Journal of Advanced Nursing. 2017;73(7):1667-1680.

2. Hall LH, Johnson J, Heyhoe J, Watt I, Anderson K, O'Connor DB. Exploring the impact of primary care physician burnout and wellbeing on patient care: A focus group study. Journal of Patient Safety. 2018;in press.

3. Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare Staff Wellbeing, Burnout, and Patient Safety: A Systematic Review. *PloS One*. 2016;11(7):e0159015.
4. Panagioti M, Geraghty K, Johnson J, et al. Association between physician burnout and patient safety, professionalism and patient satisfaction: a sytematic review and meta-analysis. *JAMA internal medicine*. 2018;in press.
5. Panagioti M, Panagopoulou E, Bower P, et al. Controlled Interventions to Reduce Burnout in Physicians: A Systematic Review and Meta-analysis. *JAMA Internal Medicine*. 2017;177(2):195-205.
6. West CP, Dyrbye LN, Erwin PJ, Shanafelt TD. Interventions to prevent and reduce physician burnout: a systematic review and meta-analysis. *The Lancet*. 2016;388(10057):2272-2281.