UNIVERSITY OF LEEDS

This is a repository copy of *Breastfeeding is associated with the risk of ovarian cancer in the UK Women's Cohort Study*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/137241/>

Version: Accepted Version

Proceedings Paper:

Mohammed Danial, FN, Cade, JE, Greenwood, DC et al. (1 more author) (2018)
Breastfeeding is associated with the risk of ovarian cancer in the UK Women's Cohort Study. In: Proceedings of the Nutrition Society. Nutrition Society Summer Meeting, 10-12 Jul 2018, Leeds. Cambridge University Press , E224.

<https://doi.org/10.1017/S0029665118002306>

© The Authors 2018. This article has been published in a revised form in Proceedings of the Nutrition Society <https://doi.org/10.1017/S0029665118002306>. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Breastfeeding is associated with the risk of ovarian cancer in the UK Women's Cohort Study

F.N. Mohammed Danial^{1,3}, J.E. Cade¹, D.C. Greenwood² and V.J. Burley¹

¹Nutritional Epidemiology Group, School of Food Science and Nutrition, University of Leeds, LS2 9JT United Kingdom,

²Centre for Epidemiology and Biostatistics, University of Leeds, LS2 9JT United Kingdom

³Faculty of Hospitality, Tourism and Wellness, Universiti Malaysia Kelantan, Malaysia.

Studies have found that longer breastfeeding duration is associated with the risk of having ovarian cancer and other hormonal related cancers(1). However, the evidence is still inconsistent with a limited number of prospective studies providing evidence(2). Therefore, this analysis aimed to explore the association between breastfeeding duration and the risks of ovarian cancer in a large prospective cohort of middle-aged women, the UK Women's Cohort Study (UKWCS).

The UKWCS participants were recruited in 1995–1998 at the age of 35–69 years(3). Cancer incidence was obtained from the Office of National Statistics (ONS). A baseline questionnaire was used to estimate breastfeeding duration, ranging from >0 months to >6 months. Patients with prevalent cancers were excluded. The relationship between breast feeding duration and incidence of ovarian cancer was tested using survival analysis adjusting for age, children, socio-economic class, menopausal status, smoking status, body mass index, energy intake and age at menarche.

The total number of incident ovarian cancer cases was 224 over 16 years follow up. Mean age at recruitment was 54 years. Breastfeeding duration for each child shows that ever breastfeeding was associated with 6 % risk reduction of ovarian cancer, after adjustment for confounders (HR: 0.94; 95 % CI: 0.90 to 0.99). Further analysis on breastfeeding duration per child shows that the risk reduces by 43 % among those who breastfed for more than 6 months per child (HR: 0.57; 95 % CI: 0.35 to 0.95). Additional analysis was done on cumulative breastfeeding duration across all children, with ovarian cancer risk, which shows a 2 % risk reduction for every 1 month of breastfeeding (HR: 0.98; 95 % CI: 0.97 to 1.00).

In conclusion, ever breastfeeding per child was associated with 6 % risk reduction of ovarian cancer in a fully adjusted model and those who were breastfeeding for more than 6-month benefit more with a high-risk reduction of ovarian cancer by 43 %. This finding supports the World Health Organisation (WHO) recommendation for women

to breastfeed for at least 6 months for each child. Future studies are warranted to determine if the duration of breastfeeding per child provides more meaningful data than cumulative breastfeeding duration. The average duration of breastfeeding per child might have more public health applicability than total duration and is a better reflection of lactation amenorrhea(1).

1. Luan NN, Wu QJ, Gong TT et al. (2013) Breastfeeding and ovarian cancer risk: a meta-analysis of epidemiologic studies. *AmJ Clin Nutr* 98, 1020–31.
2. Feng LP, Chen HL & Shen MY (2014) Breastfeeding and the risk of ovarian cancer: a meta-analysis. *J Midwifery Womens Health* 59, 428–37.
3. Cade JE, Burley VJ, Alwan NA et al. (2017) Cohort Profile: The UK Women's Cohort Study (UKWCS). *Int J Epidemiol* 46, e11.