



**UNIVERSITY OF LEEDS**

This is a repository copy of *Coloring the rationals in reverse mathematics*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/138942/>

Version: Accepted Version

---

**Article:**

Frittaion, E [orcid.org/0000-0003-4965-9271](https://orcid.org/0000-0003-4965-9271) and Patey, L (2017) Coloring the rationals in reverse mathematics. *Computability*, 6 (4). pp. 319-331. ISSN 2211-3568

<https://doi.org/10.3233/COM-160067>

---

© 2017 – IOS Press and the author. This is an author produced version of a paper published in *Computability*. Uploaded in accordance with the publisher's self-archiving policy. The final publication is available at IOS Press through <https://doi.org/10.3233/COM-160067>

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# COLORING THE RATIONALS IN REVERSE MATHEMATICS

EMANUELE FRITTAION AND LUDOVIC PATEY

ABSTRACT. Ramsey’s theorem for pairs asserts that every 2-coloring of the pairs of integers has an infinite monochromatic subset. In this paper, we study a strengthening of Ramsey’s theorem for pairs due to Erdős and Rado, which states that every 2-coloring of the pairs of rationals has either an infinite 0-homogeneous set or a 1-homogeneous set of order type  $\eta$ , where  $\eta$  is the order type of the rationals. This theorem is a natural candidate to lie strictly between the arithmetic comprehension axiom and Ramsey’s theorem for pairs. This Erdős-Rado theorem, like the tree theorem for pairs, belongs to a family of Ramsey-type statements whose logical strength remains a challenge.

## CONTENTS

1. Introduction	1
1.1. Definitions and notation	4
2. The Erdős-Rado theorem in reverse mathematics	4
3. Pigeonhole principle on $\mathbb{Q}$	6
4. $ER_2^2$ does not computably reduce to $RT_2^2$	8
5. Discussion and questions	16
References	18

## 1. INTRODUCTION

In this paper, we investigate the reverse mathematics of a well-known theorem due to Erdős and Rado about 2-colorings of pairs of rationals. This theorem is a natural strengthening of Ramsey’s theorem for pairs and two colors. We say that an order type  $\alpha$  is *Ramsey*, and write  $\alpha \rightarrow (\alpha)_2^2$ , if for every coloring  $f: [L]^2 \rightarrow 2$ , where  $L$  is a linear order of order type  $\alpha$ , there is a homogeneous set  $H$  such that  $(H, \leq_L)$  has order type  $\alpha$ . Ramsey’s theorem for pairs and two colors asserts that  $\omega$  is Ramsey.

---

Emanuele Frittaion’s research is supported by the Japan Society for the Promotion of Science.

Ludovic Patey is funded by the John Templeton Foundation (‘Structure and Randomness in the Theory of Computation’ project). The opinions expressed in this publication are those of the author(s) and do not necessarily reflect the views of the John Templeton Foundation.

It turns out that  $\omega$  and  $\omega^*$  are the only countable Ramsey order types. In particular,  $\eta \rightarrow (\eta)_2^2$  does not hold, where  $\eta$  is the order type of the rationals. A standard counterexample is as follows. Fix a one-to-one map  $i: \mathbb{Q} \rightarrow \mathbb{N}$ . Define  $f: [\mathbb{Q}]^2 \rightarrow 2$  by letting

$$f(x, y) = \begin{cases} 0 & \text{if } x <_{\mathbb{Q}} y \wedge i(x) < i(y); \\ 1 & \text{if } x <_{\mathbb{Q}} y \wedge i(x) > i(y). \end{cases}$$

A homogeneous set of order type  $\eta$  would give an embedding of  $\mathbb{Q}$  into  $\omega$  (with color 0) or  $\omega^*$  (with color 1), which is impossible. Even though Ramsey's theorem for rationals fails, Erdős and Rado [6, Theorem 4, p. 427] proved the following Ramsey-type theorem (see also Rosenstein [17, Theorem 11.7, p. 207]).

**Theorem 1.1** (Erdős-Rado theorem) The partition relation  $\eta \rightarrow (\aleph_0, \eta)^2$  holds.

The relation  $\eta \rightarrow (\aleph_0, \eta)^2$  asserts that for every coloring  $f: [L]^2 \rightarrow 2$ , where  $L$  is a linear order of order type  $\eta$ , there is either an infinite 0-homogeneous set or a 1-homogeneous set  $H$  such that  $(H, \leq_{\mathbb{Q}})$  has order type  $\eta$ .

We study Theorem 1.1 within the framework of reverse mathematics (see Simpson [21]). Reverse mathematics is a vast mathematical program whose goal is to study the logical strength of ordinary theorems in terms of set existence axioms. It uses the framework of subsystems of second-order arithmetic, with the base theory  $\text{RCA}_0$  (recursive comprehension axiom).  $\text{RCA}_0$  is composed of  $P^-$ , that is, the basic first-order Peano axioms for  $0, 1, +, \times, <$ , together with  $\Delta_1^0$ -comprehension and  $\Sigma_1^0$ -induction with number and set parameters.  $\text{RCA}_0$  is usually thought of as capturing *computable mathematics*. It turns out that the large majority of countable mathematics can be proven in  $\text{ACA}_0$ , where  $\text{ACA}_0$  is  $\text{RCA}_0$  together with arithmetic comprehension. See Hirschfeldt [8] for a gentle presentation of the reverse mathematics below  $\text{ACA}_0$ .

We formalize Theorem 1.1 in  $\text{RCA}_0$  as follows.

$\text{ER}_2^2$  For every coloring  $f: [\mathbb{Q}]^2 \rightarrow 2$  there exists either an infinite 0-homogeneous set or a 1-homogeneous set  $H$  such that  $(H, \leq_{\mathbb{Q}})$  is dense.

Here  $\mathbb{Q}$  is any fixed primitive recursive presentation of the rationals. We may safely assume that the domain of  $\mathbb{Q}$  is  $\mathbb{N}$ . Note that provably in  $\text{RCA}_0$  every two (countable) linear orders of order type  $\eta$  are isomorphic and any dense linear order obviously contains a linear order of order type  $\eta$ . Therefore  $\text{ER}_2^2$  is provably equivalent over  $\text{RCA}_0$  to the statement of Theorem 1.1.

In order to study  $\text{ER}_2^2$  we also consider a version of the infinite pigeonhole principle over the rationals, namely the statement:

$ER^1$  For every  $n$  and for every  $n$ -coloring  $f: \mathbb{Q} \rightarrow n$  there exists a dense homogeneous set.

The early study of reverse mathematics has led to the observation that most of the theorems happen to be equivalent to five main subsystems of second-order arithmetic that Montalban [13] called the “Big Five”. However, Ramsey’s theory provides many statements escaping this observation. Perhaps the most well-known example is Ramsey’s theorem for pairs ( $RT_2^2$ ). The effective analysis of Ramsey’s theorem was started by Jockusch [10]. In the framework of reverse mathematics, Simpson (see [21]), building on Jockusch results, proved that whenever  $n \geq 3$ ,  $RT_k^n$  is equivalent to  $ACA_0$  over  $RCA_0$ . The case of  $RT_2^2$  has been a long-standing open problem until Seetapun [19] proved that  $RT_2^2$  is strictly weaker than  $ACA_0$  over  $RCA_0$ . Cholak, Jockusch and Slaman [1] paved the way to the reverse mathematics analysis of Ramsey’s theorem for pairs. Since then, many consequences of Ramsey’s theorem for pairs have been studied, leading to a whole zoo of independent statements. However, no natural statement is known to be strictly between  $ACA_0$  and  $RT_2^2$  over  $RCA_0$ . The only known candidate is the tree theorem for pairs ( $TT_2^2$ ) studied in [2, 3, 5, 15]. We show that  $ER_2^2$  also lies between  $ACA_0$  and  $RT_2^2$ , and so represents another candidate, arguably more natural than  $TT_2^2$ .

Although no relation is known between them,  $TT_2^2$  and  $ER_2^2$  share some essential combinatorial features and put the emphasis on a new family of Ramsey-type theorems, characterized by what we call a *disjoint extension commitment*. See section 5 for a discussion on this notion. Some separations known for variants of  $TT_2^2$  are essentially due to this common feature, which enables us to prove the same separations for variants of  $ER_2^2$ . In particular, we prove that  $ER_2^2$  does not computably reduce to Ramsey’s theorem for pairs *with an arbitrary number of colors* ( $RT_{<\infty}^2$ ). However, we cannot simply adapt this “one-step separation” to a separation over  $\omega$ -models, and in particular over  $RCA_0$ , as in the case of  $TT_2^2$  [15]. This is the first known example of such an inability. Indeed, a diagonalization against an  $RT_4^2$ -instance is similar to a diagonalization against two  $RT_2^2$ -instances. Therefore, diagonalizing against  $RT_{<\infty}^2$  has some common flavor with a separation over standard models.

Among the consequences of Ramsey’s theorem for pairs, Ramsey’s theorem for singletons ( $RT_{<\infty}^1$ ), also known as the infinite pigeonhole principle, is of particular interest.  $RT_{<\infty}^1$  happens to be equivalent to the  $\Sigma_2^0$  bounding scheme (see Hirst [9]). The  $\Sigma_2^0$  bounding scheme ( $B\Sigma_2^0$ ) is formally defined as

$$(\forall x < a)\exists y\varphi(x, y, a) \implies \exists b(\forall x < a)(\exists y < b)\varphi(x, y, a)$$

where  $\varphi$  is any  $\Sigma_2^0$  formula. One may think of  $B\Sigma_2^0$  as asserting that the finite union of finite sets is finite (see for instance [7]). We show that

$\text{ER}^1$ , the corresponding pigeonhole principle for rationals, is strictly stronger than  $\text{B}\Sigma_2^0$ , and hence has the same reverse mathematics status as the tree theorem for singletons ( $\text{TT}^1$ ) [3].

For the purpose of separating  $\text{ER}_2^2$  from  $\text{RT}_{<\infty}^2$  over computable reducibility, we also introduce the asymmetric version of  $\text{ER}^1$  for two colors, namely  $\text{a-ER}_2^1$ , stating that for every partition  $A_0 \cup A_1 = \mathbb{Q}$  of the rationals there exists either an infinite subset of  $A_0$  or a dense subset of  $A_1$ . Indeed, we show the existence of a  $\Delta_2^0$ -instance of  $\text{a-ER}_2^1$ , and hence of a computable instance of  $\text{ER}_2^2$ , which does not reduce to any computable instance of  $\text{RT}_{<\infty}^2$ .

**1.1. Definitions and notation.** *String.* A *string* is an ordered tuple of bits  $b_0, \dots, b_{n-1}$ , that is, such that  $b_i < 2$  for every  $i < n$ . The empty string is written  $\langle \rangle$ . A *real* is an infinite listing of bits  $b_0, b_1, \dots$ . Given  $s \in \omega$ ,  $2^s$  is the set of strings of length  $s$  and  $2^{<s}$  is the set of strings of length  $< s$ . Similarly,  $2^{<\omega}$  is the set of finite strings and  $2^\omega$  is the set of reals. Given a string  $\sigma \in 2^{<\omega}$ , we denote by  $|\sigma|$  its length. Given two strings  $\sigma, \tau \in 2^{<\omega}$ , we write  $\sigma \hat{\ } \tau$  for the concatenation of  $\sigma$  and  $\tau$ , and we say that  $\sigma$  is a *prefix* of  $\tau$  (written  $\sigma \preceq \tau$ ) if there exists a string  $\rho \in 2^{<\omega}$  such that  $\sigma \hat{\ } \rho = \tau$ . Given a real  $X$ , we write  $\sigma \prec X$  if  $\sigma = X \upharpoonright n$  for some  $n \in \omega$ , where  $X \upharpoonright n$  denotes the restriction of  $X$  to its first  $n$  elements. We may identify a real with a set of integers by considering that the real is its characteristic function.

*Tree, path.* A *binary tree*  $T \subseteq 2^{<\omega}$  is a set downward-closed under the prefix relation. A real  $P$  is a *path* through  $T$  if for every  $\sigma \prec P$ ,  $\sigma \in T$ .

*Sets, partitions.* Given two sets  $A$  and  $B$ , we denote by  $A < B$  the formula  $(\forall x \in A)(\forall y \in B)[x < y]$  and by  $A \subseteq^* B$  the formula  $(\forall^\infty x \in A)[x \in B]$ , meaning that  $A$  is included in  $B$  up to finitely many elements. Given a set  $X$  and some integer  $k$ , a *k-partition* of  $X$  is a  $k$ -tuple of pairwise disjoint sets  $A_0, \dots, A_{k-1}$  such that  $A_0 \cup \dots \cup A_{k-1} = X$ . A *Mathias condition* is a pair  $(F, X)$  where  $F$  is a finite set,  $X$  is an infinite set and  $F < X$ . A condition  $(F_1, X_1)$  *extends*  $(F, X)$  (written  $(F_1, X_1) \leq (F, X)$ ) if  $F \subseteq F_1$ ,  $X_1 \subseteq X$  and  $F_1 \setminus F \subseteq X$ . A set  $G$  *satisfies* a Mathias condition  $(F, X)$  if  $F \subseteq G$  and  $G \setminus F \subseteq X$ .

## 2. THE ERDŐS-RADO THEOREM IN REVERSE MATHEMATICS

We start off the analysis of the Erdős-Rado theorem by proving that the statement  $\text{ER}_2^2$  lies between  $\text{ACA}_0$  and  $\text{RT}_2^2$ . On the lower bound hand,  $\text{ER}_2^2$  can be seen as an immediate strengthening of  $\text{RT}_2^2$ . The upper bound is an effectivization of the original proof of  $\text{ER}_2^2$  by Erdős and Rado in [6].

**Lemma 2.1**  $(\text{RCA}_0) \text{ER}_2^2 \rightarrow \text{RT}_2^2$ .

*Proof.* An instance of  $\text{RT}_2^2$  can be regarded as an instance of  $\text{ER}_2^2$ . Moreover, provably in  $\text{RCA}_0$ , a dense set is infinite.  $\square$

The rest of this section is devoted to show that  $\text{ER}_2^2$  is provable in  $\text{ACA}_0$ . For this purpose, we give the following definition.

**Definition 2.2** ( $\text{RCA}_0$ ) By *interval* we mean a set of the form  $I = (x, y)_{\mathbb{Q}}$  for  $x, y \in \mathbb{Q}$ . We say that  $A \subseteq \mathbb{Q}$  is *somewhere dense* if  $A$  is dense in some interval of  $\mathbb{Q}$ , i.e., there exists an interval  $I$  such that for all intervals  $J \subseteq I$  we have that  $A \cap J \neq \emptyset$ . We call  $A$  *nowhere dense* otherwise.

Notice that the above notion is the usual notion of topological density with respect to the order topology of  $\mathbb{Q}$ . In general, the nowhere dense sets of a topological space form an ideal. This is crucial in the proof by Erdős and Rado. For this reason, we also use the terminology *positive* and *small* for somewhere dense and nowhere dense respectively. In  $\text{RCA}_0$  we can show that nowhere dense subsets of  $\mathbb{Q}$  are small, meaning that:

- (1) If  $A \subseteq \mathbb{Q}$  is small and  $B \subseteq A$ , then  $B$  is small;
- (2) If  $A, B \subseteq \mathbb{Q}$  are small, then  $A \cup B$  is small.

With enough induction, it is possible to generalize (2) to finitely many sets.

**Lemma 2.3** ( $\text{RCA}_0 + \text{I}\Sigma_2^0$ ) If  $A_i$  is a small subset of  $\mathbb{Q}$  for all  $i < n$ , then  $\bigcup_{i < n} A_i$  is small.

*Proof.* Suppose that  $A_i$  is small for every  $i < n$ . Fix an interval  $I$ . We aim to show that  $A^n = \bigcup_{i < n} A_i$  is not dense in  $I$ . By bounded  $\Sigma_2^0$ -comprehension we can form the set

$$C = \{i \leq n : (\exists J \subseteq I \text{ interval}) A^i \cap J = \emptyset\}.$$

where  $A^i = \bigcup_{j < i} A_j$ . The set  $C$  is nonempty as  $0 \in C$ . Let  $i$  be the maximum of  $C$ . We claim that  $i = n$ . Suppose not. By definition there exists an interval  $J \subseteq I$  such that  $A^i \cap J = \emptyset$ . By the assumption  $A_i$  is small and so there exists an interval  $K \subseteq J$  such that  $A_i \cap K = \emptyset$ . It follows that  $A^{i+1} \cap K = (A^i \cup A_i) \cap K = \emptyset$ , and hence  $i + 1 \in C$ , for the desired contradiction.  $\square$

**Theorem 2.4**  $\text{ER}_2^2$  is provable in  $\text{ACA}_0$ .

*Proof.* Let  $f: [\mathbb{Q}]^2 \rightarrow 2$  be given. For any  $x \in \mathbb{Q}$ , let  $\text{Red}(x) = \{y \in \mathbb{Q} \setminus \{x\} : f(x, y) = 0\}$ . Define  $\text{Blue}(x)$  accordingly. We say that  $A \subseteq \mathbb{Q}$  is *red-admissible* if there exists some  $x \in A$  such that  $A \cap \text{Red}(x)$  is positive.

Case I. Every positive subset of  $\mathbb{Q}$  is red-admissible. We aim to show that there exists an infinite 0-homogeneous set. We define by

arithmetical recursion a sequence  $(x_n)_{n \in \mathbb{N}}$  as follows. Suppose we have defined  $x_i$  for all  $i < n$ , and assume by arithmetical induction that  $A_n = \bigcap_{i < n} \text{Red}(x_i)$  is positive, and hence 0-admissible (where  $\bigcap_{i < 0} \text{Red}(x_i) = \mathbb{Q}$ ). Search for the  $\omega$ -least  $x_n \in A_n$  such that  $A_n \cap \text{Red}(x_n) = \bigcap_{i < n+1} \text{Red}(x_i)$  is positive. By definition, the set  $\{x_n : n \in \mathbb{N}\}$  is infinite and 0-homogeneous.

Case II. There is a positive subset  $A$  of  $\mathbb{Q}$  which is not red-admissible. In this case, we show that there exists a dense 1-homogeneous set. Let  $I$  be a witness of  $A$  being positive. Fix an enumeration  $(I_n)_{n \in \mathbb{N}}$  of all subintervals of  $I$ . Notice that by definition  $A$  intersects every  $I_n$ .

We define by arithmetical recursion a sequence  $(x_n)_{n \in \mathbb{N}}$  as follows. Let  $x_0 \in A \cap I_0$ . Suppose we have defined  $x_i \in A \cap I_i$  for all  $i < n$ . By Lemma 2.3, since every  $A \cap \text{Red}(x_i)$  with  $i < n$  is small, it follows that  $E = \bigcup_{i < n} (A \cap \text{Red}(x_i))$  is small. Let  $J \subseteq I_n$  be such that  $E \cap J = \emptyset$ . We may safely assume that no  $x_i$  with  $i < n$  belongs to  $J$ . Since  $A$  is dense in  $I$  and  $J \subseteq I$ , we can find  $x_n \in A \cap J$ . In particular,  $x_n \in \bigcap_{i < n} \text{Blue}(x_i)$ . Therefore  $\{x_n : n \in \mathbb{N}\}$  is dense and 1-homogeneous.  $\square$

**Remark 2.5** A similar proof shows that  $\text{RT}_2^2$  is provable in  $\text{ACA}_0$ . In fact, we can consider the ideal of finite sets of  $\mathbb{N}$  so that a positive set is just an infinite set and a red-admissible set is a set  $A \subseteq \mathbb{N}$  such that  $A \cap \text{Red}(x)$  is infinite for some  $x \in A$ .

### 3. PIGEONHOLE PRINCIPLE ON $\mathbb{Q}$

We next consider the statement  $\text{ER}^1$  asserting that every finite coloring of rationals has a dense homogeneous set. The main result is that  $\text{ER}^1$  is stronger than  $\text{B}\Sigma_2^0$  over  $\text{RCA}_0$ . We achieve this by adapting the model-theoretic proof of Corduan, Groszek, and Mileti [3] that separates  $\text{TT}^1$  from  $\text{B}\Sigma_2^0$ . Basically, in a model of  $\text{RCA}_0 + \neg \text{I}\Sigma_2^0$ , there are a real  $X$  and an  $X$ -recursive instance of  $\text{ER}^1$  with no  $X$ -recursive solutions. Before going into the details of this proof, we establish the following simple reverse mathematics facts.

**Lemma 3.1** Over  $\text{RCA}_0$ ,

- 1)  $\text{ER}_2^2 \vee \text{I}\Sigma_2^0 \rightarrow \text{ER}^1$
- 2)  $\text{ER}^1 \rightarrow \text{RT}_{< \infty}^1$ .

*Proof.* 1) Let  $f: \mathbb{Q} \rightarrow n$  be a given coloring. Assume  $\text{ER}_2^2$ , and let  $g: [\mathbb{Q}]^2 \rightarrow 2$  be defined by  $g(x, y) = 1$  if and only if  $f(x) = f(y)$ . Provably in  $\text{RCA}_0$  every one-to-one function from an infinite set is unbounded. Then by  $\text{ER}_2^2$  there exists a dense 1-homogeneous set for  $g$ , which is homogeneous for  $f$ . Assume  $\text{I}\Sigma_2^0$  and define  $A_i = f^{-1}(i)$  for  $i < n$ . As  $\mathbb{Q} = \bigcup_{i < n} A_i$  is positive, by lemma 2.3, there exists  $i < n$  such

that  $A_i$  is positive. From  $A_i$  we can compute a dense  $i$ -homogeneous set.

2) is trivial.  $\square$

As in [3], the proof of our separation result consists of a few lemmas. We start by first adapting [3, Lemma 3.4] (see Lemma 3.3 below). The combinatorial core of the proof is based on the following.

**Lemma 3.2** ( $P^- + \text{I}\Sigma_1$ ) For each  $e < n$ , let  $\Gamma_e$  consist of  $4n$  pairwise disjoint intervals. Then there exist  $2n$  pairwise disjoint intervals  $\langle I_{e,i} : e < n, i < 2 \rangle$  such that  $I_{e,i} \in \Gamma_e$  for all  $e < n$  and  $i < 2$ .

*Proof.* Let  $\Gamma_e$ ,  $e < n$ , be given. Consider the following recursive procedure. At each stage we define  $\Gamma_{e,s}$  for  $e < n$  and  $\Delta_s$  as follows. At stage 0,  $\Gamma_{e,0} = \Gamma_e$  and  $\Delta_0 = \langle \rangle$ . At stage  $s + 1$ , if  $|\Delta_s| = 2n$  or  $\Gamma^s = \bigcup_{e < n} \Gamma_{e,s}$  is empty, we are done. Otherwise search for  $I \in \Gamma^s$  minimal with respect to inclusion (such an interval exists by  $\text{I}\Sigma_1$ ). Add  $I$  to  $\Delta_s$ , that is,  $\Delta_{s+1} = \Delta_s \hat{\cup} I$ . Let  $e$  be such that  $I \in \Gamma_{e,s}$ . If  $\Delta_s$  already contains an interval in  $\Gamma_e$ , let  $\Gamma_{e,s+1} = \emptyset$ , otherwise let  $\Gamma_{e,s+1} = \Gamma_{e,s} \setminus \{I\}$ . For all  $j \neq e$ , let  $\Gamma_{j,s+1} = \{J \in \Gamma_{j,s} : I \cap J = \emptyset\}$ . Notice that by the choice of  $I$  as minimal, at most two intervals from each  $\Gamma_{j,s}$  with  $j \neq e$  have nonempty intersection with  $I$ .

By  $\text{I}\Sigma_1$  (indeed  $\text{I}\Sigma_0$ ) it is easy to show that, for all  $s < 2n + 1$ ,  $\Delta_s$  consists of  $s$  disjoint intervals from  $\bigcup_{e < n} \Gamma_e$  with at most two intervals from the same  $\Gamma_e$ , that every interval in  $\Delta_s$  is disjoint from any interval in  $\Gamma^s$ , and that if  $\Delta_s$  does not contain 2 intervals from  $\Gamma_e$ , then  $\Gamma_{e,s}$  contains at least  $4n - 2s$  intervals. In particular,  $\Delta_{2n}$  is as desired.  $\square$

**Lemma 3.3** ( $P^- + \text{I}\Sigma_1^X$ ) There exists an  $X$ -recursive function  $d : \mathbb{N} \times \mathbb{Q} \rightarrow 2$  such that for all  $n$  and  $e < n$ , if  $W_e^X$  is a dense set of  $\mathbb{Q}$ , then there exist two disjoint intervals  $I_0, I_1$  such that  $W_e^X \cap I_i$  is infinite for all  $i < 2$  and  $d(n, x) = i$  for all  $i < 2$  and for almost every  $x \in I_i$ .

*Proof sketch.* Our strategy to defeat  $n$ -many dense sets  $\{A_e : e < n\}$  is to choose  $2n$  pairwise disjoint intervals  $I_{e,0}, I_{e,1}$  for  $e < n$  so that each  $I_{e,i}$  has end-points in  $A_e$ , and assign color  $i$  to the interval  $I_{e,i}$  for all  $e < n$  and  $i < 2$ . As we want to diagonalize against  $n$ -many potential dense sets of the form  $W_e^X$  for  $e < n$  and we cannot decide uniformly in  $n$  which ones are dense, we act only when some  $W_e^X$  outputs  $4n + 1$  points. We then specify a set  $\Gamma_e$  of  $4n$  disjoint intervals with end-points in  $W_e^X$  and from each  $\Gamma_e$  currently defined we choose intervals  $I_{e,0}$  and  $I_{e,1}$  as in Lemma 3.2. Every time we act, our choice of  $I_{e,0}$  and  $I_{e,1}$  might change, but this happens at most  $n$ -many times. As the actual construction is essentially the one in the proof of Lemma [3, Lemma 3.4], we leave the details to the reader.  $\square$

The next lemma is the key part of the whole argument (see [3, Proposition 3.5]).

**Lemma 3.4** Let  $M$  be a first-order model of  $P^-$  and suppose that  $M$  satisfies  $\text{I}\Sigma_1^X$  but not  $\text{I}\Sigma_2^X$  for some  $X \subseteq M$ . Then there is an  $X$ -recursive (in the sense of  $M$ ) coloring  $f$  of  $\mathbb{Q}$  into  $M$ -finitely many colors such that no  $X$ -recursive dense set is homogeneous for  $f$ .

*Proof.* Since  $\text{I}\Sigma_2^X$  fails, there is an  $X$ -recursive function  $h: \mathbb{N}^2 \rightarrow \mathbb{N}$  such that for some number  $a$ , the range of the partial function  $h(y) = \lim_{s \rightarrow \infty} h(y, s)$  is unbounded on  $\{y: y < a\}$  (see [3, Lemma 3.6]). Define  $f: \mathbb{Q} \rightarrow 2^a$  by

$$f(x) = \langle d(h(y, x), x) : y < a \rangle,$$

where  $d(n, x)$  is the function of Lemma 3.3. Let  $W_e^X$  be a dense set of  $\mathbb{Q}$ . We aim to show that  $W_e^X$  is not homogeneous for  $f$ . Let  $y < a$  such that  $h(y) > e$ . Observe that for almost every  $x \in \mathbb{Q}$  the  $y$ th bit of  $f(x)$  is  $d(h(y), x)$ . As  $e < h(y)$ , let  $I_0$  and  $I_1$  be two intervals as in Lemma 3.3. Now for sufficiently large  $x_0 \in W_e^X \cap I_0$  and  $x_1 \in W_e^X \cap I_1$  we have  $d(h(y, x_i), x_i) = d(h(y), x_i) = i$ , and hence  $f(x_0) \neq f(x_1)$ .  $\square$

We can finally prove the analogue of [3, Corollary 3.8], which is the main result.

**Theorem 3.5** Let  $P$  be a  $\Pi_1^1$  sentence. Then  $\text{RCA}_0 + P \vdash \text{ER}^1$  if and only if  $\text{RCA}_0 + P \vdash \text{I}\Sigma_2^0$ . In particular,  $\text{RCA}_0 + \text{B}\Sigma_2^0 \not\vdash \text{ER}^1$ .

*Proof.* The argument is the same as in the proof of [3, Theorem 3.7]. As  $\text{RCA}_0 + \text{I}\Sigma_2^0 \vdash \text{ER}^1$ , we just need to prove one implication. Suppose that  $\text{RCA}_0 + P \not\vdash \text{I}\Sigma_2^0$ , and let  $M$  be a model of  $\text{RCA}_0 + P$  where  $\text{I}\Sigma_2^0$  fails. Then there is a real  $X$  of  $M$  witnessing  $\neg \text{I}\Sigma_2^0$ . Let  $M'$  be the submodel of  $M$  with the same first-order part as  $M$  and second-order part consisting of the reals recursive in  $X$  (in the sense of  $M$ ). Since  $M'$  has same first-order part as  $M$ ,  $M'$  satisfies the  $\Pi_1^1$  sentence  $P$ . Also, as the reals of  $M'$  are the ones recursive in a given real of  $M$ ,  $M$  satisfies  $\text{RCA}_0$ . Now, by Lemma 3.4, there exists an  $X$ -recursive instance of  $\text{ER}^1$  with no  $X$ -recursive solutions, and so  $\text{ER}^1$  fails in  $M'$ . Therefore,  $\text{RCA}_0 + P \not\vdash \text{ER}^1$ .  $\square$

#### 4. $\text{ER}_2^2$ DOES NOT COMPUTABLY REDUCE TO $\text{RT}_2^2$

Many proofs of  $\mathbf{Q} \rightarrow \mathbf{P}$  over  $\text{RCA}_0$  make use only of one  $\mathbf{Q}$ -instance to solve a  $\mathbf{P}$ -instance. This is the notion of computable reducibility.

**Definition 4.1** (Computable reducibility) Fix two  $\Pi_2^1$  statements  $\mathbf{P}$  and  $\mathbf{Q}$ .  $\mathbf{P}$  is *computably reducible* to  $\mathbf{Q}$  (written  $\mathbf{P} \leq_c \mathbf{Q}$ ) if every  $\mathbf{P}$ -instance  $X_0$  computes a  $\mathbf{Q}$ -instance  $X_1$  such that for every solution  $Y$  to  $X_1$ ,  $Y \oplus X_0$  computes a solution to  $X_0$ .

Proving that  $P \leq_c Q$  is not sufficient to deduce that  $\text{RCA}_0 \vdash Q \rightarrow P$ . One needs to prove that this reducibility can be formalized within  $\text{RCA}_0$ , and in particular that  $\Sigma_1^0$ -induction is sufficient to prove its validity. The fine-grained nature of computable reducibility enables one to exhibit distinctions between statements which would not have been revealed in reverse mathematics. For example,  $\text{RT}_k^2$  and  $\text{RT}_{k+1}^2$  are equivalent over  $\text{RCA}_0$  whereas  $\text{RT}_{k+1}^2 \not\leq_c \text{RT}_k^2$  [16].

This notion of reducibility can be also seen as an intermediary step to tackle difficult separations [4]. Proving that  $P \not\leq_c Q$  is simpler than separating  $Q$  from  $P$  over  $\omega$ -models. Lerman, Solomon and Towsner [12] introduced a framework to separate Ramsey-type statements over  $\omega$ -models, in which they transform a one-step diagonalization, that is, computable non-reducibility, into a separation in the sense of reverse mathematics. In this section, we prove that the Erdős-Rado theorem for pairs does not reduce to Ramsey's theorem for pairs in one step.

**Theorem 4.2**  $\text{ER}_2^2 \not\leq_c \text{RT}_{<\infty}^2$ .

Interestingly, this diagonalization does not seem to be easily generalizable to a separation over  $\omega$ -models. A reason is that the fairness property ensured by the  $\text{ER}_2^2$ -instance does not seem to be preserved by weak König's lemma. This is hitherto the first example of a computable non-reducibility of a principle  $P$  to  $\text{RT}_{<\infty}^2$  which is not generalizable to a proof that  $\text{RT}_2^2$  does not imply  $P$  over  $\text{RCA}_0$ .

The remainder of this section is devoted to a proof of Theorem 4.2. The notion of fairness presented below may have some ad-hoc flavor. It has been obtained by applying the main ideas of the framework of Lerman, Solomon and Towsner [12, 14]. Thanks to an analysis of the combinatorics of Ramsey's theorem for pairs and the Erdős-Rado theorem for pairs, we prove our computable non-reducibility result by constructing an instance of  $\text{ER}_2^2$  ensuring the density of the diagonalizing conditions in the forcing notion of  $\text{RT}_2^2$ . Then we abstract the diagonalization to any  $\Sigma_1^0$  formula, to get rid of the specificities of the forcing notion of  $\text{RT}_2^2$  in the notion of fairness preservation. See [15] for a detailed example of the various steps of this framework, leading to a separation of  $\text{RT}_2^2$  from the tree theorem for pairs over  $\text{RCA}_0$ .

**Definition 4.3** (Simple partition) A *simple partition*  $\text{int}_{\mathbb{Q}}(S)$  is a finite sequence of open intervals  $(-\infty, x_0), (x_0, x_1), \dots, (x_{n-1}, +\infty)$  for some set of rationals  $S = \{x_0 <_{\mathbb{Q}} \dots <_{\mathbb{Q}} x_{n-1}\}$ . We set  $\text{int}_{\mathbb{Q}}(\emptyset) = \{\mathbb{Q}\}$ . A simple partition  $I_0, \dots, I_{n-1}$  *refines* another simple partition  $J_0, \dots, J_{m-1}$  if for every  $i < n$ , there is some  $j < m$  such that  $I_i \subseteq J_j$ . Given two simple partitions  $I_0, \dots, I_{n-1}$  and  $J_0, \dots, J_{m-1}$ , the product  $\vec{I} \otimes \vec{J}$  is the simple partition

$$\{I \cap J : I \in \vec{I} \wedge J \in \vec{J}\}$$

One can easily see that  $\text{int}_{\mathbb{Q}}(S)$  refines  $\text{int}_{\mathbb{Q}}(T)$  if  $T \subseteq S$  and that  $\text{int}_{\mathbb{Q}}(S \cup T) = \text{int}_{\mathbb{Q}}(S) \otimes \text{int}_{\mathbb{Q}}(T)$ . Note that every simple partition has a finite description, since the set  $S$  and each rational has a finite description. Also note that a simple partition is not a true partition of  $\mathbb{Q}$  since the endpoints do not belong to any interval. However, we have  $S \cup \bigcup \text{int}_{\mathbb{Q}}(S) = \mathbb{Q}$ .

**Definition 4.4** (Matrix) An  $m$ -by- $n$  matrix  $M$  is a rectangular array of rationals  $x_{i,j} \in \mathbb{Q}$  such that  $x_{i,j} <_{\mathbb{Q}} x_{i,k}$  for each  $i < m$  and  $j < k < n$ . The  $i$ th row  $M(i)$  of the matrix  $M$  is the  $n$ -tuple of rationals  $x_{i,0} < \dots < x_{i,n-1}$ . The simple partition  $\text{int}_{\mathbb{Q}}(M)$  is defined by  $\bigotimes_{i < m} \text{int}_{\mathbb{Q}}(M(i))$ . In particular,  $\bigotimes_{i < m} \text{int}_{\mathbb{Q}}(M(i))$  refines the simple partition  $\text{int}_{\mathbb{Q}}(M(i))$  for each  $i < m$ .

It is important to notice that an  $m$ -by- $n$  matrix is formally a 3-tuple  $\langle m, n, M \rangle$  and not only the matrix itself  $M$ . This distinction becomes important when dealing with the degenerate cases. An  $m$ -by-0 matrix  $M$  and a 0-by- $n$  matrix  $N$  are both empty. However, they have different sizes. In particular, we shall define the notion of  $M$ -type for a matrix, and this definition will depend on the number of columns of the matrix  $M$ , which is 0 for  $M$ , and  $n$  for  $N$ . Notice also that, for a degenerate matrix  $M$ , the simple partition  $\text{int}_{\mathbb{Q}}(M)$  is the singleton  $\{\mathbb{Q}\}$ .

Given a simple partition  $\vec{I}$ , we want to classify the  $k$ -tuples of rationals according to which interval of  $\vec{I}$  they belong to. This leads to the notion of  $(\vec{I}, k)$ -type.

**Definition 4.5** (Type) Given a simple partition  $I_0, \dots, I_{n-1}$  and some  $k \in \omega$ , an  $(\vec{I}, k)$ -type is a tuple  $T_0, \dots, T_{k-1}$  such that  $T_i \in \vec{I}$  for each  $i < k$ . Given an  $m$ -by- $n$  matrix  $M$ , an  $M$ -type is an  $(\text{int}_{\mathbb{Q}}(M), n)$ -type.

We now state two simple combinatorial lemmas which will be useful later. The first trivial lemma simply states that each  $m$ -tuple of rational (different from the endpoints of a simple partition) belongs to a type.

**Lemma 4.6** For every simple partition  $I_0, \dots, I_{n-1}$  and every  $k$ -tuple of rationals  $x_0, \dots, x_{k-1} \in \bigcup_{i < n} I_i$ , there is an  $(\vec{I}, k)$ -type  $T_0, \dots, T_{k-1}$  such that  $x_j \in T_j$  for each  $j < k$ .

*Proof.* Fix  $k$  rationals  $x_0, \dots, x_{k-1}$ . For each  $i < k$ , there is some interval  $T_i \in \vec{I}$  such that  $x_i \in T_i$  since  $x_i \in \bigcup_{j < n} I_j$ . The sequence  $T_0, \dots, T_{k-1}$  is the desired  $(\vec{I}, k)$ -type.  $\square$

The next lemma is a consequence of the pigeonhole principle.

**Lemma 4.7** For every  $m$ -by- $n$  matrix  $M$  and every  $M$ -type  $T_0, \dots, T_{n-1}$ , there is an  $m$ -tuple of intervals  $J_0, \dots, J_{m-1}$  with  $J_i \in \text{int}_{\mathbb{Q}}(M(i))$  such

that

$$\left(\bigcup_{j < n} T_j\right) \cap \left(\bigcup_{i < m} J_i\right) = \emptyset$$

*Proof.* Let  $T_0, \dots, T_{n-1}$  be an  $M$ -type. For every  $i < m$  and  $j < n$ , there is some  $J \in \text{int}_{\mathbb{Q}}(M(i))$  such that  $T_j \subseteq J$ . Since  $|\text{int}_{\mathbb{Q}}(M(i))| = n + 1$ , there is an interval  $J_i \in \text{int}_{\mathbb{Q}}(M(i))$  such that  $(\bigcup_{j < n} T_j) \cap J_i = \emptyset$ .  $\square$

**Definition 4.8** (Formula, valuation) Given an  $m$ -by- $n$  matrix  $M$ , an  $M$ -formula is a formula  $\varphi(\vec{U}, \vec{V})$  with distinguished set variables  $U_j$  for each  $j < n$  and  $V_{i,I}$  for each  $i < m$  and  $I \in \text{int}_{\mathbb{Q}}(M(i))$ . An  $M$ -valuation  $(\vec{R}, \vec{S})$  is a tuple of finite sets  $R_j \subseteq \mathbb{Q}$  for each  $j < n$  and  $S_{i,I} \subseteq I$  for each  $i < m$  and  $I \in \text{int}_{\mathbb{Q}}(M(i))$ . The  $M$ -valuation  $(\vec{R}, \vec{S})$  is of type  $\vec{T}$  for some  $M$ -type  $T_0, \dots, T_{n-1}$  if moreover  $R_j \subseteq T_j$  for each  $j < n$ . The  $M$ -valuation  $(\vec{R}, \vec{S})$  satisfies  $\varphi$  if  $\varphi(\vec{R}, \vec{S})$  holds.

Given some valuation  $(\vec{R}, \vec{S})$  and some integer  $s$ , we write  $(\vec{R}, \vec{S}) > s$  to say that for every  $x \in (\bigcup \vec{R}) \cup (\bigcup \vec{S})$ ,  $x > s$ . Following the terminology of [12], we define the notion of essentiality for a formula (an abstract requirement), which corresponds to the idea that there is room for diagonalization since the formula is satisfied for arbitrarily far valuations.

**Definition 4.9** (Essential formula) Given an  $m$ -by- $n$  matrix  $M$ , and  $M$ -formula  $\varphi$  is *essential* if for every  $s \in \omega$ , there are an  $M$ -type  $\vec{T}$  and an  $M$ -valuation  $(\vec{R}, \vec{S}) > s$  of type  $\vec{T}$  such that  $\varphi(\vec{R}, \vec{S})$  holds.

The notion of fairness is defined accordingly. If some formula is essential, that is, gives enough room for diagonalization, then there is an actual valuation which will diagonalize against the  $\text{ER}_2^2$ -instance.

**Definition 4.10** (Fairness) Fix two sets  $A_0, A_1 \subseteq \mathbb{Q}$ . Given an  $m$ -by- $n$  matrix  $M$ , an  $M$ -valuation  $(\vec{R}, \vec{S})$  diagonalizes against  $A_0, A_1$  if  $\bigcup \vec{R} \subseteq A_1$  and for every  $i < m$ , there is some  $I \in \text{int}_{\mathbb{Q}}(M(i))$  such that  $S_{i,I} \subseteq A_0$ . A set  $X$  is *fair* for  $A_0, A_1$  if for every  $m, n \in \omega$ , every  $m$ -by- $n$  matrix  $M$  and every  $\Sigma_1^{0,X}$  essential  $M$ -formula, there is an  $M$ -valuation  $(\vec{R}, \vec{S})$  diagonalizing against  $A_0, A_1$  such that  $\varphi(\vec{R}, \vec{S})$  holds.

Of course, if  $Y \leq_T X$ , then every  $\Sigma_1^{0,Y}$  formula is  $\Sigma_1^{0,X}$ . As an immediate consequence, if  $X$  is fair for some  $A_0, A_1$  and  $Y \leq_T X$ , then  $Y$  is fair for  $A_0, A_1$ .

Now we have introduced the necessary terminology, we create a non-effective instance of  $\mathbf{a}\text{-ER}_2^1$  which will serve as a bootstrap for fairness preservation.

**Lemma 4.11** For every set  $C$ , there exists a  $\Delta_2^{0,C}$  partition  $A_0 \cup A_1 = \mathbb{Q}$  such that  $C$  is fair for  $A_0, A_1$ .

*Proof.* The proof is done by a no-injury priority construction. Let  $M_0, M_1, \dots$  be an enumeration of all  $m$ -by- $n$  matrices and  $\varphi_0, \varphi_1, \dots$  be an effective enumeration of all  $\Sigma_1^{0,C}$   $M_k$ -formulas for every  $m, n \in \omega$ . We want to satisfy the following requirements for each pair of integers  $e, k$ .

$\mathcal{R}_{e,k}$ : If the  $M_k$ -formula  $\varphi_e$  is essential, then  $\varphi_e(\vec{R}, \vec{S})$  holds for some  $M_k$ -valuation  $(\vec{R}, \vec{S})$  diagonalizing against  $A_0, A_1$ .

The requirements are ordered via the standard pairing function  $\langle \cdot, \cdot \rangle$ . The sets  $A_0$  and  $A_1$  are constructed by a  $C'$ -computable list of finite approximations  $A_{i,0} \subseteq A_{i,1} \subseteq \dots$  such that all elements added to  $A_{i,s+1}$  from  $A_{i,s}$  are strictly greater than the maximum of  $A_{i,s}$  (in the  $\mathbb{N}$  order) for each  $i < 2$ . We then let  $A_i = \bigcup_s A_{i,s}$  which will be a  $\Delta_2^{0,C}$  set. At stage 0, set  $A_{0,0} = A_{1,0} = \emptyset$ . Suppose that at stage  $s$ , we have defined two disjoint finite sets  $A_{0,s}$  and  $A_{1,s}$  such that

- (i)  $A_{0,s} \cup A_{1,s} = [0, b]_{\mathbb{N}}$  for some integer  $b \geq s$
- (ii)  $\mathcal{R}_{e',k'}$  is satisfied for every  $\langle e', k' \rangle < s$

Let  $\mathcal{R}_{e,k}$  be the requirement such that  $\langle e, k \rangle = s$ . Decide  $C'$ -computably whether there are some  $M_k$ -type  $\vec{T}$  and some  $M_k$ -valuation  $V = (\vec{R}, \vec{S}) > b$  of type  $\vec{T}$  such that  $\varphi_e(V)$  holds. If so,  $C$ -effectively fetch  $\vec{T} = T_0, \dots, T_{n-1}$  and such a  $(\vec{R}, \vec{S}) > b$ . Let  $d$  be an upper bound (in the  $\mathbb{N}$  order) on the rationals in  $(\vec{R}, \vec{S})$ . By Lemma 4.7, for each  $i < m$ , there is some  $J_i \in \text{int}_{\mathbb{Q}}(M(i))$  such that

$$\left( \bigcup_{j < n} T_j \right) \cap \left( \bigcup_{i < m} J_i \right) = \emptyset$$

Set  $A_{0,s+1} = A_{0,s} \cup \bigcup_{i < m} J_i \cap (b, d]_{\mathbb{N}}$  and  $A_{1,s+1} = [0, d]_{\mathbb{N}} \setminus A_{0,s+1}$ . This way,  $A_{0,s+1} \cup A_{1,s+1} = [0, d]_{\mathbb{N}}$ . By the previous equation,  $\bigcup_{j < n} T_j \cap (b, d]_{\mathbb{N}} \subseteq [0, d]_{\mathbb{N}} \setminus A_{0,s+1}$  and the requirement  $\mathcal{R}_{e,k}$  is satisfied. If no such  $M_k$ -valuation is found, the requirement  $\mathcal{R}_{e,k}$  is vacuously satisfied. Set  $A_{0,s+1} = A_{0,s} \cup \{b+1\}$  and  $A_{1,s+1} = A_{1,s}$ . This way,  $A_{0,s+1} \cup A_{1,s+1} = [0, b+1]_{\mathbb{N}}$ . In any case, go to the next stage. This finishes the construction.  $\square$

**Lemma 4.12** If  $X$  is fair for some sets  $A_0, A_1 \subseteq \mathbb{Q}$ , then  $X$  computes neither an infinite subset of  $A_0$ , nor a dense subset of  $A_1$ .

*Proof.* Since fairness is downward-closed under the Turing reducibility, it suffices to prove that if  $X$  is infinite and fair for  $A_0, A_1$ , then it intersects both  $A_0$  and  $A_1$ .

We first prove that  $X$  intersects  $A_1$ . Let  $M$  be the 0-by-1 matrix and  $\varphi(U)$  be the  $\Sigma_1^{0,X}$   $M$ -formula which holds if  $U \cap X \neq \emptyset$ . The

only  $M$ -type is  $\mathbb{Q}$  and since  $X$  is infinite,  $\varphi$  is essential. By fairness of  $X$ , there is an  $M$ -valuation  $R$  diagonalizing against  $A_0, A_1$  such that  $\varphi(R)$  holds. By definition of diagonalization,  $R \subseteq A_1$ . Since  $R \cap X \neq \emptyset$ , this shows that  $X \cap A_1 \neq \emptyset$ .

We now prove that  $X$  intersects  $A_0$ . Let  $M$  be the 1-by-0 matrix and  $\varphi(V)$  be the  $\Sigma_1^{0,X}$   $M$ -formula which holds if  $V \cap X \neq \emptyset$ . The  $M$ -formula  $\varphi$  is essential since  $X$  is infinite. By fairness of  $X$ , there is an  $M$ -valuation  $S$  diagonalizing against  $A_0, A_1$  such that  $\varphi(S)$  holds. By definition of diagonalization,  $S \subseteq A_0$ . Since  $R \cap X \neq \emptyset$ , this shows that  $X \cap A_0 \neq \emptyset$ .  $\square$

Note that we did not use the fact that  $X$  is dense to make it intersect  $A_0$ . Density will be useful in the proof of Theorem 4.14.

**Definition 4.13** A *Scott set* is a set  $\mathcal{S} \subseteq 2^\omega$  such that

- (i)  $(\forall X \in \mathcal{S})(\forall Y \leq_T X)[Y \in \mathcal{S}]$
- (ii)  $(\forall X, Y \in \mathcal{S})[X \oplus Y \in \mathcal{S}]$
- (iii) Every infinite, binary tree in  $\mathcal{S}$  has an infinite path in  $\mathcal{S}$ .

**Theorem 4.14** Let  $A_0, A_1 \subseteq \mathbb{Q}$  and  $\mathcal{S}$  be a Scott set whose members are all fair for  $A_0, A_1$ . For every set  $C \in \mathcal{S}$ , every  $C$ -computable coloring  $f : [\omega]^2 \rightarrow k$ , there is an infinite  $f$ -homogeneous set  $H$  such that  $H \oplus C$  computes neither an infinite subset of  $A_0$ , nor a dense subset of  $A_1$ .

*Proof.* The proof is done by induction over the number of colors  $k$ . The case  $k = 1$  is ensured by Lemma 4.12. Fix a set  $C \in \mathcal{S}$  and let  $f : [\omega]^2 \rightarrow k$  be a  $C$ -computable coloring. If  $f$  has an infinite  $f$ -thin set  $H \in \mathcal{S}$ , that is, an infinite set over which  $f$  avoids at least one color, then  $H \oplus C$  computes a coloring  $g : [\omega]^2 \rightarrow k - 1$  such that every infinite  $g$ -homogeneous set computes relative to  $H \oplus C$  an infinite  $f$ -homogeneous set. Since  $H \oplus C \in \mathcal{S}$ , by induction hypothesis, there is an infinite  $g$ -homogeneous set  $H_1$  such that  $H_1 \oplus H \oplus C$  computes neither an infinite subset of  $A_0$ , nor a dense subset of  $A_1$ . So suppose that  $f$  has no infinite  $f$ -thin set in  $\mathcal{S}$ .

We construct  $k$  infinite sets  $G_0, \dots, G_{k-1}$ . We need therefore to satisfy the following requirements for each  $p \in \omega$ .

$$\mathcal{N}_p : \quad (\exists q_0 > p)[q_0 \in G_0] \quad \wedge \cdots \wedge \quad (\exists q_{k-1} > p)[q_{k-1} \in G_{k-1}]$$

Furthermore, we want to ensure that one of the  $G$ 's computes neither an infinite subset of  $A_0$ , nor a dense subset of  $A_1$ . To do this, we will satisfy the following requirements for every  $k$ -tuple of integers  $e_0, \dots, e_{k-1}$ .

$$\mathcal{Q}_{\vec{e}} : \quad \mathcal{R}_{e_0}^{G_0} \quad \vee \cdots \vee \quad \mathcal{R}_{e_{k-1}}^{G_{k-1}}$$

where  $\mathcal{R}_e^H$  holds if  $W_e^{H \oplus C}$  is neither an infinite subset of  $A_0$ , nor a dense subset of  $A_1$ .

We construct our sets  $G_0, \dots, G_{k-1}$  by forcing. Our conditions are variants of Mathias conditions  $(F_0, \dots, F_{k-1}, X)$  such that  $X$  is an infinite set in  $\mathcal{S}$  and the following property holds:

(P)  $(\forall i < k)(\forall x \in X)[F_i \cup \{x\}$  is  $f$ -homogeneous with color  $i]$

A condition  $d = (E_0, \dots, E_{k-1}, Y)$  extends  $c = (F_0, \dots, F_{k-1}, X)$  if  $(E_i, Y)$  Mathias extends  $(F_i, X)$  for every  $i < k$ . We now prove the progress lemma, stating that we can force the  $G$ 's to be infinite. This is where we use the fact that there is no infinite  $f$ -thin set in  $\mathcal{S}$ .

**Lemma 4.15** For every condition  $c = (F_0, \dots, F_{k-1}, X)$ , every  $i < k$  and every  $p \in \omega$  there is some extension  $d = (E_0, \dots, E_{k-1}, Y)$  such that  $E_k \cap (p, +\infty)_{\mathbb{N}} \neq \emptyset$ .

*Proof.* Fix  $c$ ,  $i$  and  $p$ . If for every  $x \in X \cap (p, +\infty)_{\mathbb{N}}$  and almost every  $y \in X$ ,  $f(x, y) \neq i$ , then  $X$  computes an infinite  $f$ -thin set, contradicting our hypothesis. Therefore, there is some  $x \in X \cap (p, +\infty)_{\mathbb{N}}$  such that  $f(x, y) = i$  for infinitely many  $y \in X$ . Let  $Y$  be the collection of such  $y$ 's. The condition  $(F_0, \dots, F_{i-1}, F \cup \{x\}, F_{i+1}, \dots, F_k, Y)$  is the desired extension.  $\square$

We now prove the core lemma stating that we can satisfy each  $\mathcal{Q}$ -requirement. A condition  $c$  forces a requirement  $\mathcal{Q}$  if  $\mathcal{Q}$  holds for every set  $G$  satisfying  $c$ .

**Lemma 4.16** For every condition  $c = (F_0, \dots, F_{k-1}, X)$  and every  $k$ -tuple of indices  $\vec{e}$ , there is an extension  $d = (E_0, \dots, E_{k-1}, Y)$  forcing  $\mathcal{Q}_{\vec{e}}$ .

*Proof.* We can assume that  $W_{e_i}^{F_i \oplus C}$  has already outputted at least  $k$  elements and is either included in  $A_0$  or in  $A_1$  for each  $i < k$ . Indeed, if  $c$  has no such extension, then  $c$  forces  $W_{e_i}^{G_i \oplus C}$  to be finite or not to be a valid solution for some  $i < k$  and therefore forces  $\mathcal{Q}_{\vec{e}}$ . For each  $i < k$ , we associate the label  $\ell_i < 2$  and the number  $p_i$  such that  $W_{e_i}^{F_i \oplus C}$  is the  $(p_i + 1)$ th set included in  $A_{\ell_i}$ .

Let  $n$  be the number of sets  $W_{e_i}^{F_i \oplus C}$  which are included in  $A_0$ , and let  $M$  be the  $(k - n)$ -by- $n$  matrix such that the  $j$ th row is composed of the  $n$  first elements already outputted by the set  $W_{e_i}^{F_i \oplus C}$  where  $p_i = j$  and  $\ell_i = 1$ . In other words,  $M(j)$  are the  $n$  first elements outputted by the  $j$ th set  $W_{e_i}^{F_i \oplus C}$  included in  $A_1$ .

Let  $\varphi(\vec{U}, \vec{V})$  be the  $\Sigma_1^{0, X \oplus C}$  formula which holds if for every  $k$ -partition  $Z_0 \cup \dots \cup Z_{k-1} = X$ , there are some  $i < k$  and some finite set  $E \subseteq Z_i$  which is  $f$ -homogeneous with color  $i$  and such that either  $\ell_i = 0$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap U_{p_i} \neq \emptyset$ , or  $\ell_i = 1$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap V_{p_i, I} \neq \emptyset$  for each  $I \in \text{int}_{\mathbb{Q}}(M(p_i))$ . We have two cases.

In the first case,  $\varphi(\vec{U}, \vec{V})$  is essential. Since  $X \oplus C$  is fair for  $A_0, A_1$ , there is an  $M$ -valuation  $(\vec{R}, \vec{S})$  diagonalizing against  $A_0, A_1$  such that  $\varphi(\vec{R}, \vec{S})$

holds. By compactness and definition of diagonalization against  $A_0, A_1$ , there is a finite subset  $D \subset X$  such that for every  $k$ -partition  $D_0 \cup \dots \cup D_{k-1} = D$ , there are some  $i < k$  and some finite set  $E \subseteq D_i$  which is  $f$ -homogeneous with color  $i$  and such that either  $\ell_i = 0$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap A_1 \neq \emptyset$ , or  $\ell_i = 1$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap A_0 \neq \emptyset$ .

Each  $y \in X \setminus D$  induces a  $k$ -partition  $D_0 \cup \dots \cup D_{k-1}$  of  $D$  by setting  $D_i = \{x \in D : f(x, y) = i\}$ . Since there are finitely many possible  $k$ -partitions of  $D$ , there are a  $k$ -partition  $D_0 \cup \dots \cup D_{k-1} = D$  and an infinite  $X$ -computable set  $Y \subseteq X$  such that

$$(\forall i < k)(\forall x \in D_i)(\forall y \in Y)[f(x, y) = i]$$

We furthermore assume that  $\min(Y)$  is larger than the use of the computations. Let  $i < k$  and  $E \subseteq D_i$  be the  $f$ -homogeneous set with color  $i$  such that either  $\ell_i = 0$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap A_1 \neq \emptyset$ , or  $\ell_i = 1$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap A_0 \neq \emptyset$ . The condition  $(F_0, \dots, F_{i-1}, F_i \cup E, F_{i+1}, \dots, F_{k-1}, Y)$  is an extension of  $c$  forcing  $\mathcal{Q}_{\vec{e}}$  by the  $i$ th side.

In the second case, there is some threshold  $s \in \omega$  such that for every  $M$ -type  $\vec{T}$ , there is no  $M$ -valuation  $(\vec{R}, \vec{S}) > s$  of type  $\vec{T}$  such that  $\varphi(\vec{R}, \vec{S})$  holds. By compactness, it follows that for every  $M$ -type  $\vec{T}$ , the  $\Pi_1^{0, X \oplus C}$  class  $\mathcal{C}_{\vec{T}}$  of all  $k$ -partitions  $Z_0 \cup \dots \cup Z_{k-1} = X$  such that for every  $i < k$  and every finite set  $E \subseteq Z_i$  which is  $f$ -homogeneous with color  $i$ , either  $\ell_i = 0$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap T_{p_i} \cap (s, +\infty)_{\mathbb{N}} = \emptyset$ , or  $\ell_i = 1$  and  $W_{e_i}^{(F_i \cup E) \oplus C} \cap I \cap (s, +\infty)_{\mathbb{N}} = \emptyset$  for some  $I \in \text{int}_{\mathbb{Q}}(M(p_i))$  is non-empty. Since  $\mathcal{S}$  is a Scott set, for each  $M$ -type  $\vec{T}$ , there is a  $k$ -partition  $\vec{Z}^{\vec{T}} \in \mathcal{C}_{\vec{T}}$  such that  $\bigoplus_{\vec{T}} \vec{Z}^{\vec{T}} \oplus X \oplus C \in \mathcal{S}$ .

If there are some  $M$ -type  $\vec{T}$  and some  $i < k$  such that  $\ell_i = 1$  and  $Z_i^{\vec{T}}$  is infinite, then the condition  $(F_0, \dots, F_{k-1}, Z_i^{\vec{T}})$  extends  $X$  and forces  $W_{e_i}^{G_i \oplus C}$  not to be dense. So suppose that it is not the case. Let  $Y \in \mathcal{S}$  be an infinite subset of  $X$  such that for each  $M$ -type  $\vec{T}$ , there is some  $i < k$  such that  $Y \subseteq Z_i^{\vec{T}}$ . Note that by the previous assumption,  $\ell_i = 0$  for every such  $i$ . We claim that the condition  $(F_0, \dots, F_{k-1}, Y)$  forces  $W_{e_i}^{G_i \oplus C}$  to be finite for some  $i < k$  such that  $\ell_i = 0$ . Suppose for the sake of contradiction that there are some rationals  $x_0, \dots, x_{n-1} > s$  such that  $x_{p_i} \in W_{e_i}^{G_i \oplus C}$  for each  $i < k$  where  $\ell_i = 0$ . Since  $x_0, \dots, x_{n-1} > s$ ,  $x_0, \dots, x_{n-1} \in \bigcup \text{int}_{\mathbb{Q}}(M)$ . Therefore, by Lemma 4.6, let  $\vec{T}$  be the unique  $M$ -type such that  $x_j \in T_j$  for each  $j < n$ . By assumption, there is some  $i < k$  such that  $Y \subseteq Z_i^{\vec{T}}$  and  $\ell_i = 0$ . By definition of  $Z_i^{\vec{T}}$ ,  $W_{e_i}^{G_i \oplus C} \cap T_{p_i} \cap (s, +\infty)_{\mathbb{N}} = \emptyset$ , contradicting  $x_{p_i} \in W_{e_i}^{G_i \oplus C}$ .  $\square$

Using Lemma 4.15 and Lemma 4.16, define an infinite descending sequence of conditions  $c_0 = (\emptyset, \dots, \emptyset, \omega) \geq c_1 \geq \dots$  such that for each  $s \in \omega$

- (i)  $|F_{i,s}| \geq s$  for each  $i < k$

(ii)  $c_{s+1}$  forces  $\mathcal{Q}_{\bar{e}}$  if  $s = \langle e_0, \dots, e_{k-1} \rangle$

where  $c_s = (F_{0,s}, \dots, F_{k-1,s}, X_s)$ . Let  $G_i = \bigcup_s F_{i,s}$  for each  $i < k$ . The  $G$ 's are all infinite by (i) and  $G_i$  does not compute an  $\mathbf{a}\text{-ER}_2^1$ -solution to the  $A$ 's for some  $i < k$  by (ii). This finishes the proof of Theorem 4.14.  $\square$

We are now ready to prove the main theorem.

*Proof of Theorem 4.2.* By the low basis theorem [11], there is a low set  $P$  of PA degree. By Scott [18], every PA degree bounds a Scott set. Let  $\mathcal{S}$  be a Scott set such that  $X \leq_T P$  for every  $X \in \mathcal{S}$ . By Lemma 4.11, there is a  $\Delta_2^{0,P}$  (hence  $\Delta_2^0$ ) partition  $A_0 \cup A_1 = \mathbb{Q}$  such that  $P$  is fair for  $A_0, A_1$ . In particular, every set  $X \in \mathcal{S}$  is fair for  $A_0, A_1$  since fairness is downward-closed under the Turing reducibility.

By Schoenfield's limit lemma [20], there is a computable function  $h : [\mathbb{Q}]^2 \rightarrow 2$  such that for each  $x \in \mathbb{Q}$ ,  $\lim_s h(x, s)$  exists and  $x \in A_{\lim_s h(x,s)}$ . Note that for every infinite set  $D$  0-homogeneous for  $h$ ,  $D \subseteq A_0$ , and for every dense set  $D$  1-homogeneous for  $h$ ,  $D \subseteq A_1$ .

Fix a computable  $\text{RT}_{<\infty}^2$ -instance  $f : [\omega]^2 \rightarrow k$ . In particular,  $f \in \mathcal{S}$ . By Theorem 4.14, there is an infinite  $f$ -homogeneous set  $H$  such that  $H$  computes neither an infinite subset of  $A_0$ , nor a dense subset of  $A_1$ . Therefore,  $H$  computes no  $\text{ER}_2^2$ -solution to  $h$ .  $\square$

## 5. DISCUSSION AND QUESTIONS

This Erdős-Rado theorem shares an essential feature with another strengthening of Ramsey's theorem for pairs already studied in reverse mathematics: the tree theorem for pairs [2, 3, 5, 15].

**Definition 5.1** (Tree theorem) We denote by  $[2^{<\mathbb{N}}]^n$  the collection of *linearly ordered* subsets of  $2^{<\mathbb{N}}$  of size  $n$ . A tree  $S \subseteq 2^{<\mathbb{N}}$  is *order isomorphic* to  $2^{<\mathbb{N}}$  (written  $S \cong 2^{<\mathbb{N}}$ ) if there is a bijection  $g : 2^{<\mathbb{N}} \rightarrow S$  such that for all  $\sigma, \tau \in 2^{<\mathbb{N}}$ ,  $\sigma \preceq \tau$  if and only if  $g(\sigma) \preceq g(\tau)$ . Given a coloring  $f : [2^{<\mathbb{N}}]^n \rightarrow k$ , a tree  $S$  is  $f$ -homogeneous if  $S \cong 2^{<\mathbb{N}}$  and  $f \upharpoonright [S]^n$  is monochromatic.  $\text{TT}_k^n$  is the statement "Every coloring  $f : [2^{<\mathbb{N}}]^n \rightarrow k$  has an  $f$ -homogeneous tree."

Both  $\text{TT}_2^2$  and  $\text{ER}_2^2$  lie between the arithmetic comprehension axiom and  $\text{RT}_2^2$ , but more than that, they share a *disjoint extension commitment*. Let us try to explain this informal notion with a case analysis.

Suppose we want to construct a computable  $\text{RT}_2^1$ -instance  $f : \mathbb{N} \rightarrow 2$  which diagonalizes against two opponents  $W_0^f$  and  $W_1^f$ . After some finite amount of time, each opponent  $W_i^f$  will have outputted a finite approximation of a solution to  $f$ , that is, a finite  $f$ -homogeneous set  $F_i$ . The two opponents share a common strategy.  $W_0^f$  tries to build an infinite  $f$ -homogeneous set  $H_0$  for color 0, and  $W_1^f$  tries to build an infinite  $f$ -homogeneous set  $H_1$  for color 1. It is therefore difficult to defeat both

opponents at the same time, since if from now on we set  $f(x) = 1$ ,  $W_1^f$  will succeed in extending  $F_1$  to an infinite  $f$ -homogeneous set, and if we set  $f(x) = 0$ ,  $W_0^f$  will succeed with its dual strategy.

Consider now the same situation, where we want to construct a computable  $\text{TT}_2^1$ -instance  $f : 2^{<\mathbb{N}} \rightarrow 2$ . After some time, the opponent  $W_0^f$  will have outputted a finite tree  $S_0 \cong 2^{<b}$  which is  $f$ -homogeneous for color 0, and the opponent  $W_1^f$  will have done the same with a finite tree  $S_1 \cong 2^{<b}$   $f$ -homogeneous for color 1. The main difference with the  $\text{RT}_2^1$  case is that each opponent will *commit to extend* each leaf of his finite tree  $S_i$  into an infinite tree isomorphic to  $2^{<\mathbb{N}}$ . In particular, for each tree  $S_i$ , the sets  $X_\sigma$  of nodes extending the leaf  $\sigma \in S_i$  are *pairwise disjoint*. Therefore, each opponent commits to extend its partial solution to disjoint sets. Moreover, by asking  $b$  to be large enough, each opponent will commit to extend enough pairwise disjoint sets so that we can choose two of them for each opponent and operate the diagonalization without any conflict.

This combinatorial property works in the same way for  $\text{ER}^1$ -instances. Indeed, in this case, each opponent will commit to extend its partial solution to pairwise disjoint intervals due to the density requirement of an  $\text{ER}^1$ -solution. Since the combinatorial arguments of the Erdős-Rado theorem and the tree theorem for pairs are very similar, one may wonder whether they are equivalent in reverse mathematics.

**Question 5.2** How do  $\text{ER}_2^2$  and  $\text{TT}_2^2$  compare over  $\text{RCA}_0$ ?

The failure of Seetapun's argument for  $\text{ER}_2^2$  comes from this disjoint extension commitment feature. In particular, there it is hard to find a forcing notion for  $\text{ER}_2^2$  whose conditions are extendible.

**Question 5.3** Does  $\text{ER}_2^2$  imply  $\text{ACA}_0$  over  $\text{RCA}_0$ ?

$\text{ER}^1$  and  $\text{TT}^1$  have the same state of the art due to their common combinatorial flavor. However, when looking at their statements for pairs,  $\text{ER}_2^2$  and  $\text{TT}_2^2$  have a fundamental difference:  $\text{ER}_2^2$  has only a half disjoint extension commitment feature. This weaker property prevents one from separating  $\text{RT}_2^2$  from  $\text{ER}_2^2$  over  $\text{RCA}_0$  by adapting the argument of  $\text{TT}_2^2$  in [15].

**Question 5.4** Does  $\text{RT}_2^2$  imply  $\text{ER}_2^2$  over  $\text{RCA}_0$ ?

We have seen in section 3 that the separation of  $\text{B}\Sigma_2^0$  from  $\text{ER}^1$  is directly adaptable from the separation of  $\text{B}\Sigma_2^0$  from  $\text{TT}^1$  from Corduan, Groszek, and Mileti [3], since the combinatorial core of this separation comes from this shared disjoint extension commitment. It is natural to conjecture that the status of  $\text{ER}^1$  with respect to  $\text{I}\Sigma_2^0$  will be the same as  $\text{TT}^1$ .

**Question 5.5** Does  $ER^1$  imply  $I\Sigma_2^0$  over  $RCA_0$ ?

It is worth mentioning that, over  $RCA_0 + I\Sigma_2^0$  proves a strengthening of both  $TT^1$  and  $ER^1$ , namely the statement “For every  $n$  and every  $f: 2^{<\mathbb{N}} \rightarrow n$  there exists a strong copy  $S$  of the full binary tree such that  $f$  is constant on  $S$ ”, where by *strong copy* we mean an isomorphic copy of  $2^{<\mathbb{N}}$  with respect to order and minima. It is easy to see that a strong copy computes a dense set of  $2^{<\mathbb{N}}$ , when  $2^{<\mathbb{N}}$  is equipped with the standard dense linear ordering on binary strings, i.e., the only linear order such that  $\{\tau: \tau \succeq \sigma \hat{\ } 0\} <_{\mathbb{Q}} \sigma <_{\mathbb{Q}} \{\tau: \tau \succeq \sigma \hat{\ } 1\}$  for all  $\sigma \in 2^{<\mathbb{N}}$ . It is likely that if we can separate  $TT^1$  or  $ER^1$  from  $I\Sigma_2^0$ , then we can already separate this stronger statement by essentially the same proof.

## REFERENCES

- [1] Peter A. Cholak, Carl G. Jockusch, and Theodore A. Slaman. On the strength of Ramsey’s theorem for pairs. *Journal of Symbolic Logic*, 66(01):1–55, 2001. 3
- [2] Jennifer Chubb, Jeffrey L Hirst, and Timothy H McNicholl. Reverse mathematics, computability, and partitions of trees. *The Journal of Symbolic Logic*, 74(01):201–215, 2009. 3, 16
- [3] Jared Corduan, Marcia J Groszek, and Joseph R Mileti. Reverse mathematics and Ramsey’s property for trees. *The Journal of Symbolic Logic*, 75(03):945–954, 2010. 3, 4, 6, 7, 8, 16, 17
- [4] Damir D Dzhafarov. Strong reductions between combinatorial principles. In preparation. 9
- [5] Damir D Dzhafarov, Jeffrey L Hirst, and Tamara J Lakins . Ramsey’s theorem for trees: the polarized tree theorem and notions of stability. *Archive for Mathematical Logic*, 49(3):399–415, 2010. 3, 16
- [6] Paul Erdos and Richard Rado. Combinatorial theorems on classifications of subsets of a given set. *Proceedings of the London mathematical Society*, 3(1):417–439, 1952. 2, 4
- [7] Emanuele Frittaion and Alberto Marcone. Linear extensions of partial orders and reverse mathematics. *MLQ Math. Log. Q.*, 58(6), 2012. 3
- [8] Denis R Hirschfeldt. Slicing the truth. *Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore*, 28, 2014. 2
- [9] Jeffrey L. Hirst. *Combinatorics in subsystems of second order arithmetic*. PhD thesis, Pennsylvania State University, August 1987. 3
- [10] Carl G Jockusch. Ramsey’s theorem and recursion theory. *Journal of Symbolic Logic*, 37(2):268–280, 1972. 3
- [11] Carl G Jockusch and Robert I Soare.  $\Pi_1^0$  classes and degrees of theories. *Transactions of the American Mathematical Society*, 173:33–56, 1972. 16
- [12] Manuel Lerman, Reed Solomon, and Henry Towsner. Separating principles below Ramsey’s theorem for pairs. *Journal of Mathematical Logic*, 13(02):1350007, 2013. 9, 11
- [13] Antonio Montalbán. Open questions in reverse mathematics. *Bulletin of Symbolic Logic*, 17(03):431–454, 2011. 3
- [14] Ludovic Patey. Iterative forcing and hyperimmunity in reverse mathematics. In Arnold Beckmann, Victor Mitran, and Mariya Soskova, editors, *Evolving Computability*, volume 9136 of *Lecture Notes in Computer Science*, pages 291–301. Springer International Publishing, 2015. 9

- [15] Ludovic Patey. The strength of the tree theorem for pairs in reverse mathematics. Submitted, 2015. 3, 9, 16, 17
- [16] Ludovic Patey. The weakness of being cohesive, thin or free in reverse mathematics. Submitted. Available at <http://arxiv.org/abs/1502.03709>, 2015. 9
- [17] Joseph G. Rosenstein. *Linear orderings*, volume 98 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1982. 2
- [18] Dana Scott. Algebras of sets binumerable in complete extensions of arithmetic. In *Proc. Sympos. Pure Math*, volume 5, pages 117–121, 1962. 16
- [19] David Seetapun and Theodore A. Slaman. On the strength of Ramsey’s theorem. *Notre Dame Journal of Formal Logic*, 36(4):570–582, 1995. 3
- [20] Joseph R. Shoenfield. On degrees of unsolvability. *Annals of Mathematics*, 69(03):644–653, May 1959. 16
- [21] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Cambridge University Press, 2009. 2, 3

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, JAPAN

*E-mail address:* [frittaion@math.tohoku.ac.jp](mailto:frittaion@math.tohoku.ac.jp)

*URL:* <http://www.math.tohoku.ac.jp/~frittaion/>

LABORATOIRE PPS, PARIS DIDEROT UNIVERSITY, FRANCE

*E-mail address:* [ludovic.patey@computability.fr](mailto:ludovic.patey@computability.fr)

*URL:* <http://ludovicpatey.com/>