

ORIGINAL ARTICLE

T lymphocyte phenotype of contact-allergic patients: experience with nickel and *p*-phenylenediamine

Kate Wicks¹  | Clare Stretton¹ | Amy Popple¹ | Lorna Beresford¹ | Jason Williams² | Gavin Maxwell³ | John Paul Gosling⁴ | Ian Kimber¹ | Rebecca J. Dearman¹

¹Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

²Contact Dermatitis Investigation Unit, Salford Royal NHS Foundation Trust, Salford, UK

³Unilever Safety and Environmental Assurance Centre, Colworth Science Park, Sharnbrook, UK

⁴School of Mathematics, University of Leeds, Leeds, UK

Correspondence

Dr Kate Wicks, University of Manchester, Faculty of Biology, Medicine and Health, Michael Smith Building, Oxford Road, Manchester M13 9PT, UK.

Tel: +44 1612755368.

Email: kate.wicks@manchester.ac.uk

Funding information

Unilever

Background: There is considerable interest in understanding the immunological variables that have the greatest influence on the effectiveness of sensitization by contact allergens, particularly in the context of developing new paradigms for risk assessment of novel compounds.

Objectives: To examine the relationship between patch test score for three different contact allergens and the characteristics of T cell responses.

Methods: A total of 192 patients with confirmed nickel, *p*-phenylenediamine (PPD) or methylisothiazolinone (MI) allergy were recruited from the Contact Dermatitis Investigation Unit at Salford Royal Hospital. Severity of allergy was scored by the use of patch testing, peripheral blood lymphocytes were characterized for T cell phenotype by flow cytometry, and proliferative activity was characterized by radiolabelled thymidine incorporation. Comparisons were drawn with buffy coat samples from healthy volunteers.

Results: Patch test positivity for nickel, PPD and MI was associated with changes in the phenotype of peripheral blood T cells: increases in naïve cells, decreases in regulatory T cell frequency and the CD4⁺/CD8^{hi} ratio, and increased expression of the skin-homing marker cutaneous lymphocyte antigen (CLA), particularly for those patients with a +++ patch test score.

Conclusions: This increased understanding of the characteristics of the T cell responses to contact allergens may provide parameters with which to better measure health risks associated with skin sensitization.

KEYWORDS

CD4, CD8, inverted CD4/CD8, skin-homing CLA, lymphocyte transformation test, T cell phenotype, RRID:SCR_001905

1 | INTRODUCTION

Skin sensitization resulting in allergic contact dermatitis is an important health issue, and it is now clear that many hundreds of chemicals, possibly in excess of several thousands, have some potential to cause skin sensitization.¹ There remains considerable interest in defining the cellular and molecular mechanisms through which skin sensitization is acquired and allergic contact dermatitis is elicited, particularly in the context of developing new paradigms for risk assessment of novel compounds. Of particular relevance are the nature and dynamics of T lymphocyte responses to contact allergens,

and contributions made by functional subpopulations and differentiated subsets of T cells.²⁻⁷

One important tool for the characterization of human T cell responses to contact allergens is the lymphocyte transformation test, in which the proliferation of cultured T lymphocytes induced by contact allergens, or by contact allergens conjugated with proteins, is measured.⁸⁻¹⁵ Experience with the lymphocyte transformation test has been predominantly with metals, and in particular nickel, although somewhat variable results have also been reported with certain organic contact allergens, including: 2,4-dinitrochlorobenzene,⁹ methylisothiazolinone (MI),⁷ and *p*-phenylenediamine (PPD).^{11,12}

In the present investigations, the lymphocyte transformation test was used to explore further the characteristics of T lymphocyte responses to nickel (as nickel sulfate), and to PPD, which is a proven contact allergen in hair dyes and henna tattoos^{1,16} (delivered in culture as a hapten-protein conjugate). Furthermore, the relative frequencies of discrete subpopulations of T lymphocytes were measured in subjects sensitized to PPD or nickel. In previous investigations of subjects with allergic contact dermatitis caused by MI, it was found that there was an association between skin sensitization and the relative frequencies of CD4⁺ and CD8⁺ naïve, memory and effector T lymphocytes.⁷ Comparable studies were conducted here with PPD and nickel to determine whether similar patterns are found in subjects sensitized to these contact allergens, and previously published analyses of MI have been extended. Finally, experiments were performed to determine whether sensitization to PPD or nickel (and also to MI) is associated with changes in the relative frequency in peripheral blood of regulatory T cells (Tregs)¹⁷ or skin-homing (cutaneous lymphocyte antigen [CLA]⁺) T lymphocytes.¹⁸

2 | MATERIALS AND METHODS

2.1 | Patient recruitment

A total of 192 individuals were recruited to this study (NRES Ethics Committee North West-Greater Manchester East [12/NW/0602]). All subjects were recruited from patients attending the Contact Dermatology Investigation Unit at Salford Royal Hospital for diagnosis of contact allergy who gave their informed consent to participate in this study. Of the 57 individuals recruited who were allergic to nickel, five were also allergic to PPD. Of the 53 individuals recruited who were allergic to PPD, five were also allergic to nickel. In addition to the 56 individuals with MI allergy who had been reported in the previous publication,⁷ a further 27 individuals were recruited, of whom eight also had an allergy to nickel or PPD.

2.2 | Healthy volunteers

Blood samples were obtained from generally healthy individuals who met the NHS Blood and Transplant Service screening criteria for blood donation ($n = 14$). Samples were obtained in the form of a "buffy coat" preparation, which was prepared by centrifugation of whole blood at 18°C within 24 hours of blood draw, and was received by our laboratory for processing within the following 48 hours.

2.3 | Patch testing and blood draw

Patch testing was performed on the skin of the upper back with Finn Chambers (inner diameter, 8 mm) that contained potential contact allergens at a standard concentration (nickel, 5% [as nickel sulfate]; PPD, 1%; MI, 0.2%). After 4 days, patients' reactions were scored as no reaction (–), weak positive (+), strong positive (++) or extreme positive (+++), according to standards set by the ICDRG, and up to 30 mL of peripheral blood was collected in a lithium heparin-coated collection tube (Sigma Aldrich, St Louis, Missouri). Of the patients recruited, six consented to an additional, pre-patch test blood draw of up

to 30 mL, which was taken immediately before application of the Finn Chambers. This was in addition to the post-patch test blood sample, which was taken 4 days after application of the Finn Chambers as standard.

2.4 | Peripheral blood mononuclear cell isolation

Peripheral blood mononuclear cells (PBMCs) were isolated by centrifugation (500g, 30 minutes, no brake) of a maximum of 30 mL of whole blood (diluted 1:1 in phosphate-buffered saline [PBS]), layered over an equal volume of Histopaque 1077 (Sigma Aldrich), according to the manufacturer's protocol. An aliquot of 1×10^6 cells was collected and stored in 10% dimethylsulfoxide/90% human AB serum (Sigma Aldrich) at -80°C for flow cytometric analysis, and the remaining cells were cultured for thymidine incorporation assays. Cells for thymidine incorporation assays were cultured as described previously² in the presence or absence of cognate contact allergen or carrier control (nickel sulfate, 0.5–50 $\mu\text{g}/\text{mL}$; free PPD, 0.01–1 $\mu\text{g}/\text{mL}$; PPD conjugated to human serum albumin [HSA], 10–100 $\mu\text{g}/\text{mL}$; HSA alone, 100 $\mu\text{g}/\text{mL}$), or in medium alone.

2.5 | Preparation of PPD conjugates

PPD conjugates were prepared as previously described.³ Briefly, four separate preparations of PPD-HSA were made by incubating 15 mM PPD (Sigma Aldrich; $\times 100$ molar excess) with 10 mg/mL (150 μM) HSA (Sigma Aldrich) in PBS for 2 days. After dialysis against PBS for 3 days, the solution was lyophilized for 24 hours and stored at -80°C until required. PPD-HSA was dissolved in PBS for cell activation. Only two of the batches of conjugate resulted in cell activation; these batches were used for subsequent experiments. New batches of PPD-HSA were tested in a proliferation assay alongside the existing batch with both patient samples and negative control samples. Only those new batches giving comparable results to the existing batch were used in subsequent assays.

2.6 | Thymidine incorporation assays

Thymidine incorporation assays were performed on freshly isolated cells cultured for 5 days as described above. For the final 16 to 20 hours of culture, cells were pulsed with 22.2 μBq of [³H]thymidine (specific activity, 2 Ci/mmol; Amersham International, Amersham, UK) per well. Cells were harvested into scintillation fluid, and incorporation of tritiated thymidine was measured by β -scintillation counting. A positive proliferative response was defined as one that resulted in a stimulation index (SI: mean disintegrations per minute of test sample divided by mean disintegrations per minute of medium-alone control sample) of ≥ 2.5 as compared with control (medium-only) cells.²

2.7 | Flow cytometric analyses

Flow cytometric analyses of T cell population composition were performed with a standard protocol and one of four antibody panels designed to detect CD4⁺ T cell subsets, CD8⁺ T cell subsets, Tregs, and expression of CLA by CD4⁺ and CD8⁺ cells, respectively. All cells used for flow cytometric analysis had been frozen upon receipt at the

laboratory, and were thawed prior to staining. Antibody panels were as follows: panel 1—allophycocyanin (APC)-anti-CD4, fluorescein isothiocyanate (FITC)-anti-CD45RA, and phycoerythrin (PE).Cy5-anti-CD27; panel 2—APC-anti-CD8, FITC-anti-CD45RA, and PE.Cy5-anti-CD27; panel 3—FITC-anti-CD4 and APC-anti-CD25 pool, and PE-anti-FoxP3; and panel 4—FITC-anti-CD4, PE.Cy5-anti-CD8, and eFluor660-anti-CLA. All antibodies were from eBioscience (San Diego, California), and were used at 1:20 dilution, with staining carried out in flow cytometry staining buffer for 30 minutes at 4°C. Single-stained controls were used to set compensation settings prior to acquisition, and to establish a compensation matrix for data analysis. Flow cytometry was performed on a minimum of 10 000 cells with a FACScalibur (Becton Dickinson, Mountain View, California). FLOWJO (TreeStar, Ashland, Oregon) was used for analysis. Gating was standardized across all samples. Non-viable cells and debris were excluded from analyses by gating on the viable lymphocyte population on the basis of forward and side scatter. Gates were then drawn to delineate the CD4⁺ population and the CD8^{hi} population. For T cell subset analysis, quadrants were drawn to divide both the CD4⁺ population and the CD8^{hi} population into CD27⁺ CD45RA⁺ (naïve), CD27⁺ CD45RA⁻ (central memory), CD27⁻ CD45RA⁻ (effector memory) and CD27⁻ CD45RA⁺ (EMRA) subsets.¹⁹ Representative flow plots showing the gating strategy for CD4⁺ and CD8^{hi} cells are shown in Supplementary Information Figure S51. CD4/CD8 ratios were then calculated from the percentage of CD4⁺ cells and the percentage of CD8^{hi} cells of the total viable lymphocytes. CLA⁺ cells were defined as those that expressed CD4 or high levels of CD8, and also expressed CLA. Tregs were defined as CD4⁺ CD25⁺ cells that also expressed FoxP3.

2.8 | Statistical analyses

Statistical analyses were performed with R (v3.2.5; R Core Team, 2016; RRID:SCR_001905). Unless otherwise stated, comparisons were performed between samples taken from the general population ("buffy coat" group) and each of the three sensitized subgroups (+, ++, and +++). For the ratios of CD4⁺ to CD8^{hi} cells for nickel-sensitive, PPD-sensitive and MI-sensitive patients, abundance of Tregs, and expression of CLA, the non-parametric Mann-Whitney *U* test was used. For comparisons of distribution of CD4⁺ and CD8⁺ T cells across the four T cell subsets, including in samples taken from the same individual before and after patch testing, these compositional data were first transformed by the use of an additive log ratio (as described in Ref. ²⁰) to remove linear dependence, and a paired Hotelling's *t*² test²¹ was then applied to test for significance of differences across the multivariate means.

3 | RESULTS

3.1 | Lymphocyte proliferative responses to contact allergen

The vigour of the T cell response to contact allergen was assessed in individuals identified by patch testing as being weakly (+), strongly (+) or extremely (+++) allergic to nickel or to PPD (a small number were

allergic to both, as detailed in Materials and Methods). PBMCs were cultured for 120 hours with various concentrations of cognate allergen, in the presence of radiolabelled thymidine for the final 16 to 20 hours of culture. An SI of ≥ 2.5 was regarded as being indicative of a biologically meaningful level of proliferation.⁷ For nickel-sensitive donors, the range of SI values recorded and the proportions of positive donors were similar for weakly (5/7) and strongly (7/12) allergic individuals (Figure 1A). However, only 5 of 12 extremely allergic (+++) donors achieved a positive SI. Previous reports have shown that, despite individuals being strongly allergic to PPD, proliferative responses to free organic PPD (rather than the known metabolite Brandowski's base) were either not detected or were detected in only a minority of patients unless they had extremely severe allergic reactions.^{11,22–24} Our approach was to prepare water-soluble conjugates of PPD with the carrier protein HSA; interestingly, although four batches of conjugate were prepared under identical conditions, only two of these conjugates elicited robust proliferative responses (data not shown). Whereas negative or a minority of borderline positive results were recorded in the presence of carrier protein alone, marked proliferative responses were observed when cells were stimulated with PPD-HSA conjugate, particularly in those individuals with +++ patch test scores (8/11) (Figure 1B). Fewer patients in the ++ and + patch test groups had SI values of >2.5 (3/8 in each group). Interestingly, this +++ cohort of patients also responded to free PPD in greater numbers (14/16) and with greater vigour than did weakly (+; 2/13) or strongly allergic (++; 4/17) patients.

3.2 | Characterization of T lymphocyte naïve and memory populations in peripheral blood of contact-allergic individuals

We showed recently that the composition of the T cell compartment in the peripheral blood of patients with allergy to MI was significantly altered as compared with healthy (buffy coat) controls, with increases in the proportions of naïve CD4⁺ and CD8⁺ T cells, and corresponding decreases in the proportions of central memory, effector memory and EMRA cells.⁷ For comparison, we have now characterized the T cell populations in nickel-allergic and PPD-allergic patients by flow cytometric analysis of the proportions of CD4⁺ and CD8⁺ T cells in each of four T cell subsets, that is, naïve, central memory, EMRA, and effector memory, defined on the basis of expression of CD27 and CD45RA.¹⁹ The CD4⁺ or the CD8^{hi} population was gated (Supporting Information Figure 1), and the proportion of cells in each of four subpopulations was determined in comparison with healthy (buffy coat) control samples. The patient cohort for each contact allergen was subdivided on the basis of degree of allergy (+, ++, and +++), as described previously. In nickel-allergic individuals, an increase in the proportion of naïve CD4⁺ T cells and corresponding decreases in the proportions of EMRA and effector memory cells were observed across all three patient cohorts as compared with healthy controls, but these differences only reached statistical significance in the severely allergic patients (+++) ($P < 0.01$; Figure 2A). The same analysis of the CD4⁺ T cell compartment of PBMCs from PPD-allergic patients showed the same trend of an increase in the proportion of the naïve cell subpopulation with corresponding decreases in the proportions of EMRA and

effector memory cells, but these differences did not reach statistical significance (Figure 2B). The composition of the CD8^{hi} T cell compartment was then analysed for the cohorts of nickel-allergic and PPD-allergic patients. The CD8^{hi} population in both cohorts showed a trend of an increased proportion of naïve cells, and reductions in the EMRA and effector memory populations, as compared with healthy controls. This change in population composition was statistically significant in strongly nickel-allergic individuals (++) ($P < 0.05$; Figure 2C) and in both the strongly (++) and extremely (+++) PPD-allergic cohorts ($P < 0.001$ and $P < 0.05$, respectively; Figure 2D).

Given that the blood sampling in these studies was performed 4 days after the application of diagnostic patch tests, a potential explanation for the expansion of the naïve T cell population and a contraction of memory populations in the allergic patients is that there is recruitment of allergen-specific memory T cells to the site of application, which decreases the numbers detectable in peripheral blood. To examine this possibility, blood samples were taken from individuals with either a weak (+) or a strong (++) allergy ($n = 3$ for both groups) immediately prior to application of the suspected contact allergen for patch testing, and at the standard time point (4 days after initiation of patch testing). The proportions of CD4⁺ and CD8^{hi} T cells in the naïve, central memory, EMRA and effector memory subsets were then analysed by flow cytometry as described previously (Figure 3). Although there was some interindividual variation in the relative contributions of the different subsets, it was notable that there were no significant changes in either CD4 or CD8 cell phenotype with time, even for those individuals who had the highest proportion of naïve cells at the day 4 time point. This supports the hypothesis that individuals with allergic contact dermatitis may have an expanded pool of naïve T cells, and that this is independent of whether there is an ongoing skin reaction.

Finally, the ratio of CD4⁺ to CD8^{hi} cells was also calculated for the same samples (Figure 4). For the buffy coat control donors, a mean CD4⁺/CD8^{hi} value of 2.6 was recorded, although there was

considerable interindividual variation. This is consistent with previous studies of healthy donors, in which ratios of 2.0 and 2.3 were observed.^{25,26} In the nickel-allergic cohort, a significant decrease in the CD4⁺/CD8^{hi} ratio was shown in strongly allergic individuals (++; $P < 0.05$), whereas, in the PPD-allergic donors, there was a marked decrease in the CD4⁺/CD8^{hi} ratio in the weakly, strongly and extremely allergic cohorts (combined population as compared with buffy coat samples; $P < 0.001$). To complement the previous studies of T cell dynamics in allergy to MI, the ratio of CD4⁺ to CD8^{hi} cells was measured for an additional cohort of 27 MI-allergic individuals (+, 14; ++, 7; +++, 6). Although the mean ratio of CD4⁺ to CD8^{hi} cells in the extremely allergic cohort was considerably lower than in healthy controls (approximately 1.5 as compared with 2.6), the difference was not statistically significant.

3.3 | Further phenotypic characterization of the T lymphocyte population: contribution of Tregs and CLA-positive cells

Additional flow cytometric analyses were then performed in order to investigate further the phenotype of CD4⁺ and CD8^{hi} T cells in contact-allergic individuals (Figure 5). These analyses also incorporated a number of newly recruited MI-allergic patients. Tregs constitute a subset of the CD4⁺ population that, in humans, can be defined as CD25⁺ FoxP3⁺ CD4⁺ cells.¹⁷ Their role is to dampen inflammation, and defective or reduced numbers of Tregs have been associated with inflammatory and autoimmune diseases, including multiple sclerosis, systemic lupus erythematosus, and atopic dermatitis.^{27,28} We therefore hypothesized that individuals with more severe nickel, PPD or MI allergy would have reduced numbers of Tregs in peripheral blood. Flow cytometry analysis was performed in which CD4⁺ cells from allergic individuals or buffy coat controls were then gated for expression of both CD25 and FoxP3 (Figure 5A representative gating strategy). The abundance of Tregs as a percentage of the CD4⁺ population was then assessed for weakly ($n = 22$), strongly ($n = 15$) and




FIGURE 1 Proliferative responses of peripheral blood mononuclear cells (PBMCs) isolated from nickel-allergic or *p*-phenylenediamine (PPD)-allergic individuals. PBMCs were isolated by density gradient centrifugation from nickel (A) or PPD (B) patch test-positive donors ($n = 31$ and $n = 46$), grouped according to allergy severity. Cells were cultured in 96-well flat-bottomed plates, at 1×10^5 cells/well, stimulated with allergen or cultured in the presence of medium alone, and incubated at 37°C (\geq triplicate wells) for 120 hours. Cells were pulsed with [³H]thymidine for the final 16 to 20 hours of culture, and then harvested and processed for β -scintillation counting. PBMCs from nickel-allergic patients (A) were stimulated with 0.5 to 50 μ g/mL nickel sulfate (+, $n = 7$; ++, $n = 12$; +++, $n = 12$). PBMCs from PPD-allergic patients (B) were incubated with 0.01 to 1 μ g/mL free PPD ((i) +, $n = 13$; (ii) ++, $n = 17$; (iii) +++, $n = 16$), and a subset was also stimulated with 10 to 100 μ g/mL of one of two PPD-human serum albumin (HSA) conjugates or with 100 μ g/mL HSA (+, $n = 8$; ++, $n = 8$; +++, $n = 11$). For the allergen stimulation, the data from the concentration that gave maximal proliferation are shown. Data are shown as a stimulation index (SI) (fold change in proliferation above baseline [medium alone]) for individual patients (●) and mean and standard error of the mean per group; an SI of ≥ 2.5 was regarded as being indicative of cellular proliferation (illustrated with horizontal dotted lines)




FIGURE 2 T lymphocyte phenotype of freshly isolated peripheral blood mononuclear cells (PBMCs) of nickel-allergic and *p*-phenylenediamine (PPD)-allergic patients: comparisons with buffy coat donors. PBMCs were isolated from nickel (A, C; $n = 56$) or PPD (B, D; $n = 30$) patch test-positive donors'. The A and C for nickel and the B and D for PPD (in parentheses) refer to panels 2A–2D. Cells were stained with anti-human CD4-allophycocyanin (APC), CD8-APC, CD45RA-fluorescein isothiocyanate and CD27-phycoerythrin-Cy5 antibodies. Data are shown as percentage of total viable CD4⁺ or CD8^{hi} cells in each phenotype (naïve [CD45RA⁺ CD27⁺] [i], central memory [CD45RA⁻ CD27⁺] [ii], effector memory RA [CD45RA⁺ CD27⁻] [iii], and effector memory [CD45RA⁻ CD27⁻] [iv]) within the CD4⁺ (A, B) or CD8^{hi} (C, D) T lymphocyte populations. Donors were segregated according to patch test status (buffy coat donors [$n = 14$], + donors [nickel, $n = 14$; PPD, $n = 8$], ++ donors [nickel, $n = 27$; PPD, $n = 11$], and +++ donors [nickel, $n = 15$; PPD, $n = 11$]), and data are shown as group mean (horizontal line) and standard error of the mean and for each individual donor (•). Statistically significant shifts (Hotelling's t^2 test after additive-log-ratio transformation) as compared with buffy coat controls were seen in CD4⁺ population composition in patients with severe nickel allergy ($P < 0.01$) and in CD8^{hi} population composition in patients with moderate nickel allergy ($P < 0.05$), and in patients with moderate or severe PPD allergy ($P < 0.001$ and $P < 0.05$, respectively)

extremely ($n = 12$) allergic individuals (pooled across the allergens nickel, PPD, and MI), as well as for nine buffy coat controls. For the buffy coat controls, Tregs constituted ~1.1% of the total CD4⁺ population, which is somewhat lower than in previous reports, in which 3.2% of the CD4 population was identified as Tregs²⁹ (Figure 4B). Despite the relatively low control levels, a significant decrease in the percentage of CD4⁺ cells that were Tregs was recorded for extremely allergic individuals (Figure 5B; $P < 0.05$).

Finally, expression of the skin-homing marker CLA¹⁸ by CD4⁺ and CD8^{hi} cells was measured by flow cytometry, to determine whether contact allergy was associated with an increase in CLA expression in peripheral blood T cells (Figure 6). Representative flow cytometry plots for (A) CD4⁺ cells and (B) CD8^{hi} cells are shown in Figure 5.

Approximately 14% of healthy control (buffy coat) CD4⁺ cells expressed CLA, consistent with previous publications, which also reported ~14% CLA⁺ CD4⁺ cells in healthy controls.³⁰ A marked increase in the percentage of CLA⁺ CD4⁺ cells was recorded for all three categories of patch test-positive individuals (+, $P < 0.05$; ++, $P < 0.001$; +++, $P < 0.05$; pooled for PPD-allergic, MI-allergic or nickel-allergic donors). For the CD8^{hi} population, there was little change in the number of CLA⁺ cells (~7%), but the mean level of CLA expression (mean fluorescence intensity) was significantly elevated in individuals strongly (++; $P < 0.001$) or extremely (+++; $P < 0.05$) allergic to PPD, MI, or nickel. This indicates that increased CLA expression may be a hallmark of allergic contact dermatitis, particularly in the case of severe allergies.




FIGURE 3 Influence of patch test application on CD4 and CD8 cell phenotype. Peripheral blood mononuclear cells were isolated and stained as described in the legend to Figure 2 for the identification of naïve, central memory, EMRA and effector memory CD4⁺ (A) and CD8^{hi} (B) subsets. Blood samples were drawn immediately before patch testing (black bars) and on day 4 after a positive patch test reaction (white bars) was recorded for nickel, methylisothiazolinone, or *p*-phenylenediamine (PPD) for three individuals with weak (+) allergy and three individuals with strong (++) allergy. The results are displayed as mean and standard error of the mean. There were no statistically significant changes in subset composition with time (Hotelling's one-sample t^2 test after additive-log-ratio transformation)




FIGURE 4 Influence of allergy on CD4/CD8 cell ratio. Peripheral blood mononuclear cells (PBMCs) were isolated and stained as described in the legend to Figure 2 for the identification of CD4⁺ and CD8^{hi} subsets. The percentage of CD4⁺ and CD8^{hi} cells in each PBMC sample was calculated as a function of the total viable cell population, and is displayed as a ratio of CD4⁺ to CD8^{hi} cells. PBMCs from methylisothiazolinone (MI)-allergic (n = 27), nickel-allergic (n = 56) and *p*-phenylenediamine (PPD)-allergic (n = 30) patients were compared with those from healthy (buffy coat) donors (n = 14). Patients were segregated according to patch test status, that is, + donors (MI, n = 14; nickel, n = 14; PPD, n = 9), ++ donors (MI, n = 7; nickel, n = 27; PPD, n = 10), and +++ donors (MI, n = 6; nickel, n = 15; PPD, n = 11), and data are shown as group mean (horizontal line) and standard error of the mean and for each individual donor (•). Statistically significant differences vs buffy coat donors were assessed with the Mann-Whitney U test. * $P < 0.05$; *** $P < 0.001$




FIGURE 5 Association of severe contact allergy with decreased regulatory T cell (Treg) cell numbers. (A) Peripheral blood mononuclear cells (PBMCs) were isolated from individuals allergic to nickel (n = 30), *p*-phenylenediamine (PPD) (n = 5), or methylisothiazolinone (MI) (n = 14), or from buffy coat samples (n = 9), and stained with anti-human CD4/CD25-fluorescein isothiocyanate (FITC)/allophycocyanin (APC) antibody cocktail, and anti-human FoxP3-phycoerythrin (PE) for flow cytometric analysis. Data are shown as the proportion of CD4⁺ PBMCs coexpressing (i) CD25 and (ii) FoxP3, and therefore classed as Tregs. (B) The individuals analysed constituted a subset of the total allergic population, and were mildly (+), moderately (++) or strongly (+++) allergic to nickel (closed circles: +, n = 11; ++, n = 10; +++, n = 9), PPD (open circles: +, n = 3; ++, n = 1; +++, n = 1), or MI (half-filled circles: +, n = 8; ++, n = 4; +++, n = 2). Statistically significant differences vs buffy coat donors (n = 9) were assessed with the Mann-Whitney U test; * $P < 0.05$. Line and error bars indicate mean and standard error of the mean




FIGURE 6 Expression of cutaneous lymphocyte antigen (CLA) on CD4 and CD8 cells: comparisons with buffy coat donors. Peripheral blood mononuclear cells (PBMCs) were isolated from individuals allergic to nickel ($n = 28$), *p*-phenylenediamine (PPD) ($n = 6$), methylisothiazolinone (MI) ($n = 9$), or from 'buffy coat' samples representative of the general population, and stained with anti-human CD4-fluorescein isothiocyanate, CD8-phycoerythrin-Cy5, and CLA-eFluor660. A representative gating strategy is shown in (A) and (B) for analysis by flow cytometry of CLA expression by (C) the CD4⁺ and (D) the CD8^{hi} PBMC subsets. For each of (A) and (B), panel (i) indicates expression of CLA by buffy coat samples, and panel (ii) indicates CLA expression by patient-derived cells. For the CD4⁺ subset, the proportion of CD4⁺ cells expressing CLA is shown. For CD8^{hi} cells, the mean fluorescence intensity of CLA-eFluor660 as a proxy for surface levels of CLA is shown. Data are from a pool of individuals who were weakly (+), strongly (++) or extremely (+++) allergic to nickel (half-filled circles: +, $n = 9$; ++, $n = 10$; +++, $n = 9$), PPD (filled circles: +, $n = 4$; ++, $n = 1$; +++, $n = 1$), or MI (open circles: +, $n = 3$; ++, $n = 4$; +++, $n = 2$), and buffy coat controls ($n = 10$). Significant differences vs the buffy coat population were analysed with the Mann-Whitney *U* test. * $P < 0.05$; *** $P < 0.001$. Error bars indicate standard error of the mean. APC, allophycocyanin

4 | DISCUSSION

In the current study, T lymphocyte responses have been characterized in individuals with contact allergy to nickel, which, despite Europe-wide attempts to reduce exposure, remains one of the most common skin sensitizers,^{31,32} and to PPD, which is a potent allergen and a major cause of adverse reactions to hair dyes.^{33,34} Consistent with

previous analyses of PBMCs derived from MI-sensitized individuals,⁷ patch test positivity for both nickel and PPD was associated with increases in the proportions of both naïve CD4⁺ and naïve CD8⁺ lymphocytes, with corresponding decreases in the EMRA and effector memory cell populations. It should, however, be noted that statistical significance was seen only in severely allergic patients. The healthy control comparator population of PBMCs utilized in the study

reported herein was isolated from buffy coat samples ($n = 14$). It could be argued that this was an inappropriate control, as there are differences in the storage and isolation procedure as compared with the patch test donor samples. Whereas patch test donor PBMCs are isolated from blood within hours of blood being drawn and then snap-frozen prior to phenotypic analyses, buffy coats are prepared within 24 hours of venepuncture, received in the laboratory for PBMC preparation within 48 hours, and then snap-frozen. However, there are several studies that have compared directly freshly isolated and buffy coat PBMCs, and these have concluded that there are no significant differences in cell viability and purity and CD4/CD8 phenotype between the two preparation methods.^{35,36} The parameter that is most likely to impact on cell phenotype is freeze/thawing³⁷; given that both patch test-positive patient samples and buffy coats are frozen once PBMCs have been isolated, this should not be a cause for concern. Reference data for the distribution of naïve, central memory, effector memory and EMRA T cell subsets in peripheral whole blood anticoagulated with EDTA are available from a cohort of 253 healthy adults aged between 19 and 67 years.³⁸ The distribution of subsets among CD4⁺ cells was similar to the buffy coat data reported herein (mean values: naïve, 43.1% [vs ~32%]; central memory, 32.8% [vs ~38%]; and effector memory, 16.7% [vs ~20%]), with the exception of the EMRA subset, which was somewhat larger in the buffy coat samples, although still the smallest subset (~10% vs 1.6%). For CD8⁺ cell subsets, taking into account the markers that were used in the two different studies, there were more differences between these control data and the buffy coat data, but values were broadly similar (mean values: naïve, 44.6% [vs ~35%]; effector memory, 4.7% [vs ~14%]; EMRA, 9.9% [vs ~16%]; and central memory, 28.5% [vs 35%]). Taken together, these data suggest not only that the buffy coat sample represents an appropriate control for the patch test-positive donor samples, but also that the patch test-positive donor samples show a different distribution from that in healthy control samples. Had it been possible to obtain such samples, a more appropriate negative control would have been individuals who were patch-test negative for all patch tests applied; however, during the course of the study, only one such individual was identified.

One additional technical issue regarding the study design was that, in order to be confident of the identity of the allergen to which the patch test-positive donors were allergic, blood samples were drawn 4 days after the initial application of the patch tests, on the day of patch test reading. In other studies using peripheral blood samples taken from patients with contact allergy, blood sampling has tended to be conducted before patch testing, as the authors were concerned that exposure to contact allergen might activate the cells.^{15,39} Alternatively, donors have been drawn from a pool of previously patch test-positive or patch test-negative individuals.⁴⁰ Given that significant differences were found between the phenotype of the healthy control population and that of the nickel-positive and PPD-positive donors in our study, particularly with respect to increases in the proportions of naïve CD4⁺ and CD8^{hi} cells, it is possible that the observed differences were attributable to cellular recruitment of skin-homing lymphocytes to the patch test site. For a subset of donors, we therefore took blood samples before and 4 days after patch test application. There were no significant differences observed in the phenotype of

CD4⁺ or CD8^{hi} T cells in samples taken prior to and after patch test readings, suggesting that individuals with allergic contact dermatitis do indeed have an expanded pool of naïve T cells and reduced effector memory and EMRA cell numbers, regardless of an ongoing skin patch test response. Because of low sample numbers, it was not possible to determine whether ongoing patch test inflammation was contributing to the alteration in Treg numbers or CLA expression.

It should be noted that peripheral blood T cell phenotypes show considerable interindividual (and intraindividual) variation, and can be influenced by factors such as age and disease, with, for example, cytomegalovirus infection resulting in decreasing numbers of naïve CD4⁺ T cells⁴¹ and decreased numbers of both CD4 and CD8 naïve cells being reported with age.^{42,43} However, it is of interest that an expanded pool of naïve T cells was recorded in patients with allergy to nickel and PPD, and, in a previous publication, to MI,⁷ suggesting that this might be a common feature of contact allergy. Increased levels of naïve T cells could result from the release from the bone marrow and thymus of increased numbers of naïve T cells and T cell precursors, or from increased survival of these cells. It is known that the half-life of naïve human T cells is ~50 days, and that survival of naïve T lymphocytes requires signalling from interleukin (IL)-7.⁴⁴ It is therefore possible that the general systemic cytokine milieu that is a result of contact allergy provides an environment for improved naïve T cell survival, through increased levels of cytokines such as IL-7.

The ability of PBMCs from allergic individuals to proliferate *in vitro* in response to challenge with contact allergen was also examined with a lymphocyte transformation test. There is considerable interest in developing such *in vitro* measures of allergen-specific responses in order to supplement, or perhaps even to replace, patch testing when there are concerns about severe reactions or unintended sensitization.^{8–15} In the absence of cytokine supplements (IL-2 or IL-7, for example)^{11,39} being used to enhance proliferative responses, the overall positive rate for nickel was 17 of 31, which is lower than the rate found in many other reported studies (16/18; 62/74; and 13/14),^{15,39,40} although it is similar to the frequency of 13 of 19 reported in 2013 by Pacheco et al for patients referred by local orthopaedists for evaluation for possible nickel sensitization who were patch test-positive.⁴⁵ The reasons for this lower share of positivity in our study are not clear, although it is noted that minor technical differences in the lymphocyte transformation test protocol can have marked effects, such as the use of autologous serum,⁴⁶ or the length of time in culture (5 days vs 6 or 7 days).^{39,40,47} The lowest number of responders (5/12) was recorded for the extremely (+++) sensitized group, although mean proliferation was similar between the ++ and +++ groups. This inverse dose-response relationship may indicate that the culture conditions were not optimal, rather than a true negative correlation of proliferation with patch test sensitivity. In general, concerns regarding nickel and the measurement of proliferative responses tend to be related to non-specific responses in negative control groups (false positives), so alternative endpoints are being investigated. Cytokine expression profiling, including the ratio of IL-5 to IL-8 production⁴⁷ or the secretion of interferon- γ , are being explored currently.⁴⁸ Experience with PPD is rather more variable, with some groups reporting vigorous proliferative responses to free PPD (6 of 9 +++ patch test donors; 7 of 8 ++ or +++ patch test donors; and

10 of 11 donors said to have a strong history of sensitization to PPD).^{11,12,24} However, other studies found that free PPD stimulated proliferative responses in a minority of donors (1/6), despite them having ++ or +++ scores in patch tests,⁴⁹ or that 0 of 9 sensitized donors reacted positively to free PPD, although 7 of 9 responded to the metabolite Brandowski's base.²² We previously observed, with a different cohort of PPD patch test-positive donors, that PPD alone gave variable responses in a minority of individuals, and that PPD-HSA conjugates were more successful at stimulating proliferative responses, with the most vigorous effects being observed among the +++ patch test group.⁶ In the current study, free PPD was more effective in the +++ patch test-positive group, but the conjugate was more effective at provoking significant levels of thymidine incorporation in the more weakly sensitized groups. Given the level of sensitivity, the number of false negatives, and interlaboratory variation, the lymphocyte transformation test seems very unlikely to be a replacement for patch testing, but does have the advantages that additional endpoints such as the identity of the proliferating cells (CD4 or CD8) or the nature of cytokine production can be incorporated to improve mechanistic understanding.^{7,47,48}

In addition to sensitization being associated with changes in the profile of naïve and effector/central memory and EMRA cells, we also observed that, particularly for extremely allergic individuals, the Treg frequency was decreased, as was the CD4⁺/CD8^{hi} ratio, whereas CLA expression was increased. These findings appeared to be a feature of sensitization to each of the three different allergens, despite the fact that they have very different chemistries. It has been shown previously that PBMCs from nickel-allergic individuals express lower levels of the Treg cytokines IL-10 and transforming growth factor- β ,³⁹ and that their Tregs had limited suppressive capacity as compared with those from nickel non-allergic individuals.⁴⁹ There has been one previous report that allergic contact dermatitis is associated with a decreased frequency of Tregs. As compared with healthy controls (4.06% Tregs), significantly decreased levels of Tregs were recorded (1.6%) in eight individuals with skin sensitization to unspecified allergens, despite the fact that the average length of time since diagnosis was only 54 days.²⁷ Perhaps the most interesting report came from a study of melanoma patients who received an *in vivo* infusion of a recombinant IL-2-diphtheria toxin fusion protein, which depleted Tregs from ~3% (3.5%-2%) to ~2% (2.5%-1%) of CD4⁺ cells. Prior to depletion, only weak contact allergic responses were provoked by sensitization with the potent contact allergen diphenylprone, whereas, after depletion, very vigorous elicitation reactions to this allergen were recorded.⁵⁰ If a reduced frequency of Tregs is a feature of contact allergy, particularly severe contact allergy, then this could be a factor in polysensitization, as those individuals with +++ patch test scores, and potentially low Treg numbers, have a higher risk of becoming polysensitized.⁵¹ In this study, individuals with +++ patch test scores for nickel and MI also tended to have positive patch test reactions to other allergens.

There are reports that suggest that CD4⁺ cells may be more prevalent in skin lesions⁵²; however, it is generally accepted that both CD4⁺ and CD8⁺ cells producing interferon- γ , IL-17 and IL-22 are involved in contact allergic responses.^{1,53-55} To our knowledge, there are no previous reports showing that contact allergy is associated with

increased numbers of total CD8⁺ cells and a decreased numbers of total CD4⁺ cells. This is also not a feature of other forms of allergic disease such as allergic asthma, as even severe asthmatics had CD4⁺/CD8⁺ ratios similar to those of healthy controls.⁵⁶ There are other situations in which decreased CD4/CD8 cell ratios have been recorded; in early menopause with inadequate oestrogen levels, for example, where they can be restored with hormone replacement therapy (from $1.4\% \pm 0.16$ to $2.4\% \pm 0.3\%$),⁵⁷ or with immune senescence resulting from ageing.⁵⁸ Viral infection, particularly cytomegalovirus infection, is also associated with a similar increase in the CD4/CD8 ratio through an expansion of CD8 cells, although these tend to be in the EM and EMRA T cell compartments rather than naïve T cells.⁵⁹ Other chronic infections, such as tuberculosis, dimorphic fungal infections, toxoplasmosis, and leishmaniasis, also lead to CD8 expansion with altered ratios.⁶⁰ It therefore seems likely that the inverted CD4/CD8 ratio observed in highly sensitized donors is attributable to an expanded CD8⁺ T cell population.

The final features of T cells isolated from sensitized individuals were increased numbers of skin-homing marker CLA⁺ CD4 cells and increased expression of this marker on CD8 cells. It is well accepted that skin-resident T cells are CLA⁺; indeed, it has been calculated, from comparisons of the frequency of these cells in skin and blood and the relative volume/surface area of blood and skin, that 98% of CLA effector memory T cells are found in the skin.⁶¹⁻⁶³ In peripheral blood, this marker has been reported to be expressed on 17% of CD4 cells and 11% of CD8 cells,⁶⁴ primarily on central and effector memory cells.⁶⁵ To our knowledge, this marker has not been examined systematically in peripheral blood of contact-allergic patients, or previously correlated with the severity of allergy. It may be that this increase in CLA expression reflects activation of T cells as a result of the patch test applied 4 days prior to the blood draw, or could represent a bystander effect, with cytokines produced by allergen-specific T cells non-specifically activating other T lymphocytes.⁶⁶

In conclusion, these studies have shown that patch test positivity for nickel, PPD and MI is associated with changes in the phenotype of peripheral blood T cells: increases in the numbers of naïve cells, decreases in Treg frequency and the CD4⁺/CD8^{hi} ratio, and increased expression of the skin-homing marker CLA. This increased understanding of the characteristics of the T cell response to contact allergens may provide insights regarding useful parameters to consider in the development of health risk assessment models for chemical skin sensitizers.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to report.

FUNDING SOURCE

Unilever.

ORCID

Kate Wicks  <https://orcid.org/0000-0002-7724-9678>

REFERENCES

- de Groot AC. Patch testing. *Test Concentrations and Vehicles for 4350 Chemicals*. 3rd ed. Wapserveen: Wapserveen:Acdegroot Publishing; 2008.
- Kimber I, Dearman RJ. Allergic contact dermatitis: the cellular effectors. *Contact Dermatitis*. 2002;46:1-5.
- Kimber I, Travis MA, Martin SF, Dearman RJ. Immunoregulation of skin sensitization and regulatory T cells. *Contact Dermatitis*. 2012;67:179-183.
- Kimber I, Maxwell G, Gilmour N, Dearman RJ, Friedmann PS, Martin SF. Allergic contact dermatitis: a commentary on the relationship between T lymphocytes and skin sensitising potency. *Toxicology*. 2012;291:18-24.
- Esser PR, Kimber I, Martin SF. Correlation of contact sensitizer potency with T cell frequency and TCR repertoire diversity. *EXS*. 2014;104:101-114.
- Oakes T, Popple AL, Williams J, et al. The T cell response to contact sensitizer paraphenylenediamine is characterized by a polyclonal diverse repertoire of antigen specific receptors. *Front Immunol*. 2017; 8:162.
- Popple A, Williams J, Maxwell G, Gellatly N, Dearman RJ, Kimber I. T lymphocyte dynamics in methylisothiazolinone-allergic patients. *Contact Dermatitis*. 2016;75:1-13.
- Nordlind K. Lymphocyte transformation test in diagnosis of nickel allergy. A comparison between the separation of peripheral blood lymphocytes on Ficoll-Paque, Percoll or by gravity sedimentation. *Int Arch Allergy Appl Immunol*. 1984;73:151-154.
- Kimber I, Quirke S, Beck MH. Attempts to identify the causative allergen in cases of allergic contact dermatitis using an in vitro lymphocyte transformation test. *Toxicol In Vitro*. 1990;4:302-306.
- Hallab NJ. Lymphocyte transformation testing for quantifying metal-implant-related hypersensitivity responses. *Dermatitis*. 2004;15:82-90.
- Kneilling M, Caroli U, Grimm C, et al. Para-phenylenediamine-specific lymphocyte activation test: a sensitive in vitro assay to detect para-phenylenediamine sensitization in patients with severe allergic reactions. *Exp Dermatol*. 2010;19:435-441.
- Coulter EM, Jenkinson C, Wu Y, et al. Activation of T-cells from allergic patients and volunteers by p-phenylenediamine and Bandrowski's base. *J Invest Dermatol*. 2008;128:897-905.
- Popple A, Williams J, Maxwell G, Gellatly N, Dearman RJ, Kimber I. The lymphocyte transformation test in allergic contact dermatitis: new opportunities. *J Immunotoxicol*. 2016;13:84-91.
- Spoerri I, Scherer K, Michel S, Link S, Bircher AJ, Heijnen IA. Detection of nickel and palladium contact hypersensitivity by a flow cytometric lymphocyte proliferation test. *Allergy*. 2015;70:323-327.
- Ständer S, Opiel E, Thomas P, Summer B. Evaluation of lymphocyte transformation tests as compared with patch tests in nickel allergy diagnosis. *Contact Dermatitis*. 2017;76:228-234.
- Encabo Durán B, Romero-Pérez D, Silvestre Salvador JF. Allergic contact dermatitis to paraphenylenediamine: an update. *Acras Dermosfilogr*. 2018;109(7):602-609.
- Walker MR, Kasproicz DJ, Gersuk VH, et al. Induction of FoxP3 and acquisition of T regulatory activity by stimulated human CD4+CD25-T cells. *J Clin Invest*. 2003;112:1437-1443.
- Berg EL, Yoshino T, Rott LS, et al. The cutaneous lymphocyte antigen is a skin lymphocyte homing receptor for the vascular lectin endothelial cell-leukocyte adhesion molecule 1. *J Exp Med*. 1991;174:1461-1466.
- Appay V, van Lier RA, Sallusto F, Roederer M. Phenotype and function of human T lymphocyte subsets: consensus and issues. *Cytometry A*. 2008;73:975-983.
- Aitchison J. *The Statistical Analysis of Compositional Data*. London: Chapman and Hall; 1986.
- Hotelling H. The generalization of Student's Ratio. *Ann Math Statist*. 1931;2:360-378.
- Krasteva M, Nicolas JF, Chabeau G, et al. Dissociation of allergenic and immunogenic functions in contact sensitivity to para-phenylenediamine. *Int Arch Allergy Immunol*. 1993;102:200-204.
- Moed H, von Blomberg M, Bruynzeel DP, Scheper R, Gibbs S, Rustemeyer T. Improved detection of allergen-specific T-cell responses in allergic contact dermatitis through the addition of 'cytokine cocktails'. *Exp Dermatol*. 2005;14:634-640.
- Sieben S, Kawakubo Y, Al Masaoudi T, Merk HF, Blömeke B. Delayed-type hypersensitivity reaction to paraphenylenediamine is mediated by 2 different pathways of antigen recognition by specific alphabeta human T-cell clones. *J Allergy Clin Immunol*. 2002;109:1005-1011.
- Elisia I, Lam V, Hofs E, et al. Effect of age on chronic inflammation and responsiveness to bacterial and viral challenges. *PLoS One*. 2017;12: e0188881.
- Matos LC, Batista P, Monteiro N, et al. Lymphocyte subsets in alcoholic liver disease. *World J Hepatol*. 2013;5:46-55.
- Ma L, Xue HB, Guan XH, et al. The imbalance of Th17 cells and CD4 (+) CD25(high) Foxp3(+) Treg cells in patients with atopic dermatitis. *J Eur Acad Dermatol Venereol*. 2014;28:1079-1086.
- Nie J, Li YY, Zheng SG, Tsun A, Li B. FOXP3(+) treg cells and gender bias in autoimmune diseases. *Front Immunol*. 2015;6:493.
- Xu WH, Zhang AM, Ren MS, et al. Changes of Treg-associated molecules on CD4+CD25 +Treg cells in myasthenia gravis and effects of immunosuppressants. *J Clin Immunol*. 2012;32:975-983.
- Teraki Y, Hotta T, Shiohara T. Increased circulating skin-homing cutaneous lymphocyte-associated antigen (CLA)+ type 2 cytokine-producing cells, and decreased CLA+ type 1 cytokine-producing cells in atopic dermatitis. *Br J Dermatol*. 2000;143:373-378.
- Duarte I, Mendonça RF, Korkeas KL, Lazzarini R, Hafner MFS. Nickel, chromium and cobalt: the relevant allergens in allergic contact dermatitis. Comparative study between two periods: 1995-2002 and 2003-2015. *An Bras Dermatol*. 2018;93:59-62.
- Uter W, Amario-Hita JC, Balato A, et al. European Surveillance System on Contact Allergies (ESSCA): results with the European baseline series, 2013/14. *J Eur Acad Dermatol Venereol*. 2017;31:1516-1525.
- DeLeo VA. p-Phenylenediamine. *Dermatitis*. 2016;7:53-55.
- Basketter DA, Alépée N, Ashikaga T, et al. Categorization of chemicals according to their relative human skin sensitizing potency. *Dermatitis*. 2014;25:11-21.
- Jia Y, Xu H, Li Y, et al. A modified Ficoll-Paque gradient method for isolating mononuclear cells from the peripheral and umbilical cord blood of humans for biobanks and clinical laboratories. *Biopreserv Biobank*. 2018;16:82-91.
- Adams M, Lee TH, Busch MP, et al. Rapid freezing of whole blood or buffy coat samples for polymerase chain reaction and cell culture analysis: application to detection of human immunodeficiency virus in blood donor and recipient repositories. The Transfusion Safety Study Group. *Transfusion*. 1993;33:504-508.
- Fowke KR, Behnke J, Hanson C, Shea K, Cosentino LM. Apoptosis: a method for evaluating the cryopreservation of whole blood and peripheral blood mononuclear cells. *J Immunol Methods*. 2000;244:139-144.
- Apoil PA, Puissant-Lubraro B, Congy-Jolivet N, et al. Reference values for T, B and NK human lymphocyte subpopulations in adults. *Data Brief*. 2017;12:400-404.
- Rustemeyer T, von Blomberg BM, van Hoogstraten IM, Bruynzeel DP, Scheper RJ. Analysis of effector and regulatory immune reactivity to nickel. *Clin Exp Allergy*. 2004;34:1458-1466.
- Spiewak R, Moed H, von Blomberg BM, et al. Allergic contact dermatitis to nickel: modified in vitro test protocols for better detection of allergen-specific response. *Contact Dermatitis*. 2007;56:63-69.
- van der Heiden M, van Zelm MC, Bartol SJ, et al. Differential effects of cytomegalovirus carriage on the immune phenotype of middle-aged males and females. *Sci Rep*. 2016;20:26892.
- Fagnoni FF, Vescovini R, Passeri G, et al. Shortage of circulating naive CD8(+) T cells provides new insights on immunodeficiency in aging. *Blood*. 2000;95:2860-2868.
- Yan J, Greer JM, Hull R, et al. The effect of ageing on human lymphocyte subsets: comparison of males and females. *Immun Ageing* 2010; 7: 4.
- Moro-García MA, Mayo JC, Sainz RM, Alonso-Arias R. Influence of inflammation in the process of T lymphocyte differentiation: proliferative, metabolic, and oxidative changes. *Front Immunol*. 2018;9:339.
- Pacheco K, Barker L, Maier L, Erb S, Sills M, Knight V. Development of a validated blood test for nickel sensitization. *J Allergy Clin Immunol*. 2013;132:767-769.

46. Marigo M, Nouer DF, Genelhu MC, et al. Evaluation of immunologic profile in patients with nickel sensitivity due to use of fixed orthodontic appliances. *Am J Orthod Dentofacial Orthop.* 2003;124:46-52.
47. Summer B, Ständer S, Thomas P. Cytokine patterns in vitro, in particular IL-5/IL-8 ratio, to detect patients with nickel contact allergy. *J Eur Acad Dermatol Venereol.* 2018;32:1542-1548.
48. Bordignon V, Palamara F, Altomonte G, et al. A laboratory test based on determination of cytokine profiles: a promising assay to identify exposition to contact allergens and predict the clinical outcome in occupational allergic contact dermatitis. *BMC Immunol.* 2015;16:4.
49. Moed H, von Blomberg BM, Bruynzeel DP, Scheper RJ, Gibbs S, Rustemeyer T. Regulation of nickel-induced T-cell responsiveness by CD4+CD25+ cells in contact allergic patients and healthy individuals. *Contact Dermatitis.* 2005;53:71-74.
50. Mahnke K, Schönfeld K, Fondel S, et al. Depletion of CD4+CD25+ human regulatory T cells in vivo: kinetics of Treg depletion and alterations in immune functions in vivo and in vitro. *Int J Cancer.* 2007;120:2723-2733.
51. Dittmar D, Uter W, Bauer A, et al. European Surveillance System on Contact Allergies (ESSCA): polysensitization, 2009-2014. *Contact Dermatitis.* 2018;78:373-385.
52. Dyring-Andersen B, Skov L, Løvendorf MB, et al. CD4(+) T cells producing interleukin (IL)-17, IL-22 and interferon- γ are major effector T cells in nickel allergy. *Contact Dermatitis.* 2013;68:339-347.
53. Gibson A, Kim SH, Faulkner L, et al. In vitro priming of naive T-cells with p-phenylenediamine and Bandrowski's base. *Chem Res Toxicol.* 2015;28:2069-2077.
54. Peiser M. Role of Th17 cells in skin inflammation of allergic contact dermatitis. *Clin Dev Immunol.* 2013;2013:261037.
55. Zhao Y, Balato A, Fischelevich R, Chapoval A, Mann DL, Gaspari AA. Th17/Tc17 infiltration and associated cytokine gene expression in elicitation phase of allergic contact dermatitis. *Br J Dermatol.* 2009;161:1301-1306.
56. Abdulmir AS, Hafidh RR, Abubakar F, Abbas KA. Changing survival, memory cell compartment, and T-helper balance of lymphocytes between severe and mild asthma. *BMC Immunol.* 2008;9:73.
57. Dogan E, Erkoc R, Demir C, Sayarlioglu H, Dilek I, Sayarlioglu M. Effect of hormone replacement therapy on CD4+ and CD8+ numbers, CD4+/CD8+ ratio, and immunoglobulin levels in hemodialysis patients. *Ren Fail.* 2005;27:421-424.
58. Valdíglesias V, Sánchez-Flores M, Maseda A, et al. Immune biomarkers in older adults: Role of physical activity. *J Toxicol Environ Health A.* 2017;80:605-620.
59. Lugthart G, van Ostaijen-Ten Dam MM, Jol-van der Zijde CM, et al. Early cytomegalovirus reactivation leaves a specific and dynamic imprint on the reconstituting T cell compartment long-term after hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant.* 2014;20:655-661.
60. McBride JA, Striker R. Imbalance in the game of T cells: What can the CD4/CD8 T-cell ratio tell us about HIV and health? *PLoS Pathog.* 2017;13:e1006624.
61. Clark RA, Chong B, Mirchandani N, et al. The vast majority of CLA+ T cells are resident in normal skin. *J Immunol.* 2006;176:4431-4439.
62. Moed H, Boersma DM, Stoof TJ, et al. Nickel-responding T cells are CD4+ CLA+ CD45RO+ and express chemokine receptors CXCR3, CCR4 and CCR10. *Br J Dermatol.* 2004;151:32-41.
63. Santamaria Babi LF, Picker LJ, Perez Soler MT, et al. Circulating allergen-reactive T cells from patients with atopic dermatitis and allergic contact dermatitis express the skin-selective homing receptor, the cutaneous lymphocyte-associated antigen. *J Exp Med.* 1995;181:1935-1940.
64. Picker LJ, Michie SA, Rott LS, Butcher EC. A unique phenotype of skin-associated lymphocytes in humans. Preferential expression of the HECA-452 epitope by benign and malignant T cells at cutaneous sites. *Am J Pathol.* 1990;136:1053-1068.
65. Turner JE, Wadley AJ, Aldred S, Fisher JP, Bosch JA, Campbell JP. Intensive exercise does not preferentially mobilize skin-homing T cells and NK cells. *Med Sci Sports Exerc.* 2016;48:1285-1293.
66. Avicé MN, Demeure CE, Delespesse G, Rubio M, Armant M, Sarfati M. IL-15 promotes IL-12 production by human monocytes via T cell-dependent contact and may contribute to IL-12-mediated IFN- γ secretion by CD4+ T cells in the absence of TCR ligation. *J Immunol.* 1998;161:3408-3415.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Wicks K, Stretton C, Popple A, et al. T lymphocyte phenotype of contact-allergic patients: experience with nickel and p-phenylenediamine. *Contact Dermatitis.* 2019;81:43-53. <https://doi.org/10.1111/cod.13246>