



UNIVERSITY OF LEEDS

This is a repository copy of *Quality of life in men living with advanced and localised prostate cancer in the UK: a population-based study*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/142866/>

Version: Accepted Version

Article:

Downing, A orcid.org/0000-0002-0335-7801, Wright, P orcid.org/0000-0001-6129-4143, Hounsome, L et al. (23 more authors) (2019) Quality of life in men living with advanced and localised prostate cancer in the UK: a population-based study. *Lancet Oncology*, 20 (3). pp. 436-447. ISSN 1470-2045

[https://doi.org/10.1016/s1470-2045\(18\)30780-0](https://doi.org/10.1016/s1470-2045(18)30780-0)

© 2019 Elsevier Ltd. All rights reserved. Licensed under the Creative Commons Attribution-Non Commercial No Derivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Reuse

See Attached



Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Figure 1: Urinary, bowel and vitality/hormone function in men diagnosed with prostate cancer 18-42 months previously



Figures represent predicted adjusted proportions (adjusted for age, socio-economic deprivation and no. of other long-term conditions)

ADT: Androgen deprivation therapy

Figure 2: Sexual function in men diagnosed with prostate cancer 18-42 months previously




Figure 2a represents predicted adjusted proportions (adjusted for age, socio-economic deprivation and no. of other long-term conditions)










Figure 2b represents predicted adjusted proportions (adjusted for socio-economic deprivation and no. of other long-term conditions)

Figure 3: Use of interventions to improve sexual function by men diagnosed with prostate cancer 18-42 months previously



Figures represent unadjusted proportions

Figure 4: Self-assessed health ratings in men diagnosed with prostate cancer 18-42 months previously



Figures represent predicted adjusted mean scores (adjusted for socio-economic deprivation and no. of other long-term conditions)

ADT: Androgen deprivation therapy

Results should be considered alongside the previously estimated minimally important difference for EQ-5D Self-assessed health (7 points).