



This is a repository copy of *Patient reported outcome measures of quality of life in Duchenne muscular dystrophy (DMD): a systematic review of content and structural validity using COSMIN.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/143299/>

Version: Published Version

Monograph:

Powell, P. orcid.org/0000-0003-1169-3431, Carlton, J., Woods, H. et al. (1 more author) (2019) Patient reported outcome measures of quality of life in Duchenne muscular dystrophy (DMD): a systematic review of content and structural validity using COSMIN. Report. SchARR HEDS Discussion Papers . School of Health and Related Research , University of Sheffield.

© 2019 The Author(s). Article available under the terms of the CC-BY-NC-ND 4.0 licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



The
University
Of
Sheffield.

School of
Health
And
Related
Research

HEALTH ECONOMICS & DECISION SCIENCE

Discussion Paper Series

Title: Patient reported outcome measures of quality of life in Duchenne muscular dystrophy (DMD): a systematic review of content and structural validity using COSMIN

Authors: Philip A Powell, Jill Carlton, Helen Buckley Woods, Paolo Mazzone

Corresponding author: Dr Philip Powell, SchARR, University of Sheffield, Regent Court, 30 Regent Street, Sheffield, S1 4DA, UK. Tel: +44 (0) 114 222 0794, Email: p.a.powell@shef.ac.uk

Disclaimer:

This series is intended to promote discussion and to provide information about work in progress. The views expressed in this series are those of the authors. Comments are welcome, and should be sent to the corresponding author.

This project was funded by project HERCULES. This project is funded by Duchenne UK, Pfizer, PTC Therapeutics, Roche, Summit Therapeutics plc, Sarepta Therapeutics Inc, Wave Lifesciences USA Inc, Solid Biosciences, Catabasis Pharmaceuticals and Santhera Therapeutics. Two of the authors (Philip Powell and Jill Carlton) are employed on a related project that is developing a new preference-based patient reported outcome measure of quality of life for people with DMD.

Date: 25th February 2019

Prepared for: Duchenne UK



Table of contents

| Chapter | Page |
|---|------|
| Abbreviations | 3 |
| Executive summary | 5 |
| 1 Introduction | 7 |
| 2 Objectives | 10 |
| 3 Methodology | 11 |
| 3.1 Protocol registration | 11 |
| 3.2 Search strategy and information resources | 11 |
| 3.2.1 Searches | 11 |
| 3.2.2 Electronic databases | 12 |
| 3.2.3 Search strings | 12 |
| 3.2.4 Additional searches | 14 |
| 3.3 Eligibility criteria | 14 |
| 3.4 Selection process | 15 |
| 3.5 Data extraction and quality appraisal | 16 |
| 3.6 Evidence synthesis | 16 |
| 4 Results | 18 |
| 4.1 Identification of included studies | 18 |
| 4.2 PROMs used to measure quality of life in DMD | 20 |
| 4.3 Content validity | 29 |
| 4.3.1 Appraisal of PROM development studies | 29 |
| 4.3.2 Appraisal of content validity studies | 39 |
| 4.3.3 Evidence synthesis | 41 |
| 4.4 Structural validity | 43 |
| 4.4.1 Appraisal of structural validity studies | 43 |
| 4.4.2 Evidence synthesis | 45 |
| 4.5 Quality assurance of the review | 46 |
| 5 Discussion and conclusion | 48 |
| References | 54 |
| Appendix | 60 |



Abbreviations

| | |
|------------------|--|
| AUQEI | Autoquestionnaire Qualité de vie Enfant Imagé |
| BASC | Behavior Assessment System for Children |
| BDI | Beck Depression Inventory |
| CALI | Child Activity Limitations Interview |
| CAPE | Children's Assessment of Participation and Enjoyment |
| CHQ-PF50 | Child Health Questionnaire - Parent Form 50 |
| CMQM | Comprehensive Model of QoL in Muscular Dystrophy |
| COSMIN | COnsensus- based Standards for the selection of health Measurement INstruments |
| DCGM-37 | DISABKIDS generic module |
| DIKJ | Depressions-Inventar für Kinder und Jugendliche |
| DMD | Duchenne muscular dystrophy |
| DUC-25 | Dutch Children AZL/TNO Questionnaire Quality of Life Short Form |
| EQ-5D(-3L) | Euroqol instrument, Euroqol-5 dimension (3 level) |
| EQ-5D-Y | Euroqol instrument, Euroqol-5 dimension (youth version) |
| FSS | Fatigue Severity Scale |
| GAD-7 | Generalized Anxiety Disorder Scale 7-item |
| GRADE | Grading of Recommendations, Assessment, Development and Evaluations |
| HADS | Hospital Anxiety and Depression Scale |
| HUI | Health Utilities Index |
| INQOL | Individualized Neuromuscular Quality of Life Questionnaire |
| LSIA | Life Satisfaction Index for Adolescents |
| MDCHILD | Muscular Dystrophy Child Health Index of Life with Disabilities |
| Neuro-QoL | Neurological Disorders Quality of Life Questionnaire |
| NICE | National Institute for Health and Care Excellence |
| OSIQ | Offer Self-Image Questionnaire |
| PBM | Preference based measure |
| PedsQL 3.0 DMD | Pediatric Quality of Life Inventory (PedsQL) 3.0 DMD module |
| PedsQL 3.0 MFS | PedsQL 3.0 Multidimensional fatigue scale |
| PedsQL 3.0 NMM | PedsQL 3.0 Neuromuscular module |
| PedsQL 4.0 GCS | Pediatric Quality of Life Inventory (PedsQL) 4.0 Generic Core Scales |
| PedsQL 4.0 SF-15 | PedsQL 4.0 Generic Short-form |
| PHQ-9 | Patient Health Questionnaire Depression Scale |
| PODCI | Pediatric Outcomes Data Collection Instrument |
| PROM | Patient reported outcome measure |
| PROSPERO | International Prospective Register of Systematic Reviews |
| PSQI | Pittsburgh Sleep Quality Index |
| QoL | Quality of life |
| Rasch PCM | Rasch Partial-credit model |
| SDQ | Strengths and Difficulties Questionnaire |
| SF-36 | 36-item Short-Form Health Survey |
| SOLE | Strips of Life with Emoticons Questionnaire |



| | |
|-------------|---|
| SWLS | Satisfaction with Life Scale |
| TAAQoL | TNO-AZL Questionnaire for Adult's Health-Related Quality of Life |
| TACQOL | TNO-AZL Questionnaire for Children's Health-Related Quality of Life |
| WHOQOL-BREF | World Health Organisation Quality of Life Scale-Brief Version |



Executive summary

Duchenne muscular dystrophy (DMD) is an inherited neuromuscular disorder that predominantly affects boys and men. There is no known cure, so current clinical efforts are focused on improving the health-related quality of life (QoL) of people with DMD. Of the validated methods for assessing QoL, patient-reported outcome measures (PROMs) are the most common. In a field of many options, the choice of which PROM to use should be based predominantly on their validity and reliability for the construct, population, and context of use of interest. In the current report we critically reviewed the content and structural validity of PROMs used to assess QoL in people with DMD, using robust, updated CONsensus-based Standards for the selection of health Measurement INSTRUMENTS (COSMIN) guidelines.

In this review, we defined QoL as a multidimensional construct involving physical, psychological, and social components. We conducted a systematic search of the published literature for self-reported, multi-item PROMs assessing at least one aspect of QoL in a sample of at least 75% boys and men with DMD. We further refined these results for studies with evidence of content and structural validity, including development studies. The resulting PROMs and related studies were systematically rated, in terms of methodological quality, and evidence for content and structural validity, as per the latest COSMIN procedures.

From an initial 1,752 records, and 5 additional records identified through citation tracking, 60 published primary research articles were identified that had used a PROM to assess at least one aspect of QoL in people with DMD, of which 5 articles presented evidence on content or structural validity. A further 36 articles were identified through Google Scholar searching and citation tracking presenting content validity information on the development of the PROMs, resulting in a final selection of 41 articles for review. From the articles identified, 40 PROMs were extracted, of which 26 were taken forward for COSMIN assessment.

The results of the assessment for content validity revealed a modal COSMIN quality rating of inadequate, primarily due to the PROM development study not being performed in a sample of patients representing the target population (of people with DMD). The second most common rating was one of doubtful, due to at least some unclear details/suspected problems within the qualitative methods used. Only the KIDSCREEN family of measures received an adequate rating for concept elicitation and PROM design. Only two published articles had independently assessed the content



validity of the QoL PROMs (LSIA, PedsQL 3.0 NMM) in samples of people with DMD, and both received ratings of doubtful due to at least some unclear details/suspected problems with the qualitative methods used. In terms of the evidence on content validity assessed, the KIDSCREEN measures and the LSIA were the only PROMs to receive satisfactory results for all three dimensions of content validity. These reflect relevance (whether items are relevant for the construct, target, population, and context of use of interest); comprehensiveness (the extent to which all key aspects of the construct of interest are covered); and comprehensibility (the understanding of items and response options by the population of interest).

Two studies had investigated the structural validity of the included PROMs (PedsQL 4.0 GCS, PedsQL 3.0 NMM) in people with DMD, of which one received a very good COSMIN quality rating for its methodological content. Nevertheless, an assessment of the evidence for structural validity revealed a rating of indeterminate for the PedsQL 4.0 GCS, as key details of the results from the Rasch model were not reported, and a rating of unsatisfactory for the PedsQL 3.0 NMM, due to psychometric criteria for good measurement properties not being met.

The results of this review suggest that evidence on the content and structural validity of PROMs assessing QoL in people with DMD is sparse, and further research is needed. In the absence of further evidence, we recommend that the KIDSCREEN is used to assess QoL in children and adolescents with DMD. It is difficult to recommend an adult measure, as insufficiencies are evident in content and/or structural validity, but in terms of precedent, the PedsQL is widely used, which has a young adult and adult variant. Furthermore, these two PROMs can be used in cost effectiveness analyses via mapping algorithms. Limitations of this review include the potential harshness of the worst score counts COSMIN system of assessment of methodological quality; the restriction to two, albeit the most important, measurement properties; and the potential restrictive nature of the inclusion criteria, which could be broadened to consider related neuromuscular disorders or mixed samples, if deemed theoretically appropriate.



1 Introduction

Duchenne muscular dystrophy (DMD) is an inherited neuromuscular disorder that predominantly affects boys and men. It has an estimated incidence of 1:3800 to 1:6300 in live births.¹ The disease causes progressive muscle weakness due to an absence of the dystrophin protein, which functions to help keep muscle cells intact. Diagnostic symptoms and functional impairment are evident from as early as two years old and average life expectancy of people with DMD is approximately 25 years,² although increasingly people with DMD are surviving into their fourth and even fifth decades.³ The disease progresses through four recognised clinical stages characterised by increased muscle weakness, impaired ambulation and motor functioning, and cardiovascular and respiratory problems.⁴ There is no known cure for the disease. Current clinical efforts are thus focused on improving the health-related quality of life (QoL) of people with DMD, and health interventions are necessarily evaluated for their cost effectiveness against this objective.

In order to attempt to measure QoL in people with DMD a number of condition-specific and generic questionnaires are used. For NICE, the institution responsible for making decisions on the funding of NHS health interventions, there is a stated preference that health-related QoL data comes from the EQ-5D.⁵ The EQ-5D is a generic, preference-based measure of health-related QoL with 5 dimensions covering mobility, self-care, usual activities, pain/discomfort, and anxiety and depression. In the case of DMD, however, there are a number of concerns that measures like the EQ-5D are insufficient to adequately assess QoL in this population, based on the aspects of QoL that matter to people living with DMD.⁶ For example, DMD is recognised to have an impact on aspects of daily life, such as participation, friendships, independence and dignity which may not be fully captured by generic measures, such as the EQ-5D (or its equivalent for children, the EQ-5D-Y).

In order to satisfy NICE requirements, in cases where the EQ-5D is considered insufficient as a measure of health-related QoL for a particular population, and thus other measures may be more appropriate, evidence must be provided. Such evidence largely centres on the demonstrable validity of a measure, including its content validity and/or psychometric performance in the population of interest. Where it can be evidenced that the EQ-5D is inappropriate as a measure of health-related QoL, alternative preference-based measures, with superior validity, can be used to generate utility values, including condition-specific preference-based measures.



Given that a number of generic and condition-specific questionnaires are available for use in attempting to assess QoL in people with DMD, evidence is desperately needed on the relative validity and psychometric performance of these instruments, when it comes to assessing QoL in DMD. While we are aware of a number of reviews exploring QoL and associated measures in DMD, with some providing very basic information on their psychometric properties,^{6,7} no reviews to date have appropriately evaluated the content validity of available measures when it comes to assessing QoL in DMD. This is a striking omission; content validity has been defined by the COnsensus-based Standards for the selection of health Measurement INstruments (COSMIN) group as the most important property of a patient reported outcome measure (PROM).⁸ Furthermore, prior reviews on QoL in neuromuscular disorders have either not referred to, or used an outdated version of, COSMIN guidance, which is considered a rigorous approach to the systematic assessment(s) of the validity and reliability of PROMs. Updated and expanded COSMIN guidance and documentation for the evaluation of PROMs was published earlier this year, and, as a consequence of its importance in determining measure selection, this included a dedicated manual on assessing the content validity of PROMs.⁹

Content validity refers to the extent that the content of a PROM adequately reflects the target construct that is being measured.¹⁰ It can be meaningfully subdivided into the relevance, comprehensiveness, and comprehensibility of a PROM, for assessing the construct of interest in a target population and context.⁹ Here, relevance of a PROM refers to whether the items are relevant for the construct, target population, and context of use of interest; the response options and recall period of a PROM should also be appropriate and relevant. Comprehensiveness is used to describe the extent to which all key aspects of the construct of interest are covered in the PROM. Finally, comprehensibility pertains to understanding of the items and response options by the population of interest.⁹

A thorough assessment of a PROM's content validity should crucially include studies presenting information on content validity in the population of interest, but also consider the initial PROM development paper(s) and the content of the PROM itself.⁹ Content validity should form the first step of the assessment of the validity of a PROM, as it is integral to that PROM's usefulness in doing the job it was designed to do, and influences all other measurement properties.¹¹ For example, a psychometrically responsive and internally consistent instrument is of little use if it is not measuring what it is intended to measure in the first place.



According to COSMIN guidance, the second most important form of the validity assessment of a PROM is structural validity.¹¹ Structural validity describes the extent that scores derived from a measure adequately reflect the dimensionality of the construct being measured.¹² Quality of life is usually defined, and thus measured, as a multidimensional construct. Therefore, PROMs that feature multiple dimensions of quality of life should be dutifully assessed to check they accurately represent the multidimensional structure of quality of life in the population of interest. If PROMs are designed to target a single dimension of quality of life, assessments should be undertaken to empirically demonstrate their unidimensional nature in the target population. If such assessments are not undertaken, subsequent interpretation of the data (e.g. through generating dimensional scores) may be inaccurate.

This systematic review has been designed to evaluate the content and structural of QoL measures used in people with DMD using updated COSMIN guidance.^{9,13} A similar approach has recently been undertaken and published by members of the COSMIN group when evaluating PROMs for physical functioning used in people with low back pain.¹¹ It forms part of an ongoing project funded by Duchenne UK on producing a preference-based measure (PBM) for people with DMD as part of the Project HERCULES initiative.

For the purposes of this review, we define QoL as a multidimensional construct involving physical (e.g., pain, fatigue), psychological (e.g., mood, self-efficacy), and social (e.g., participation, stigma) components, based on the Comprehensive Model of QoL in Muscular Dystrophy (CMQM).⁶ However, here we operationalise QoL as a subjective construct and do not include purely functional performance or assessment scales that may impact on QoL. In this review, we consider multi-item PROMs that assess *at least* one aspect of QoL in people with DMD.



2 Objectives

- To identify PROMs that are used in people with DMD to measure QoL
- To assess the content validity of PROMs that are used in people with DMD
- To assess the structural validity of PROMs that are used in people with DMD
- To synthesise the evidence and make a recommendation on the use of available PROMs to assess QoL in people with DMD



3 Methodology

3.1 Protocol registration

The protocol was registered with PROSPERO, an international database of prospectively registered systematic reviews in health and social care, welfare, public health, education, crime, justice and international development, where there is a health related outcome. The protocol is accessible at:

https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=93062

3.2 Search strategy and information sources

3.2.1 Searches

A SchARR information specialist was consulted in developing the appropriate search strategy and was responsible for conducting the main database searches.

Search terms in this review included:

- I) Duchenne muscular dystrophy (and derivatives);
- II) A robust search filter developed by the PROM Group at the University of Oxford to identify PROMs;
- III) Patient-reported outcome measures known to be used with people with DMD based on an earlier rapid review of the literature; and
- IV) A robust search filter by the COSMIN Group for identifying studies on measurement properties, as recommended by the COSMIN Group.¹⁴

A two-stage search strategy was used, where in the first stage, the search terms (I) AND ((II) OR (III)) were combined to identify articles using PROMs to measure QoL in DMD. In the second stage, search terms (I) AND (III) AND (IV) were combined with the names of all PROMs identified in Stage one of the search to identify articles reporting on the measurement properties of these instruments in DMD. No restrictions on time or language were applied to the search strategy. The two-stage search strategy has the advantage of allowing us to identify which PROMs have been used in published, peer-reviewed articles on DMD, in the absence of any evidence of content or structural validity for their use in this population.



3.2.2 Electronic databases

The electronic databases searched for the systematic review are outlined in Table 1. All databases were searched from inception.

Table 1. Electronic databases for the primary searches.

| Database | Platform | Span of search | Date searched (Stage 1) | Date searched (Stage 2) |
|------------------|-----------|----------------|-------------------------|-------------------------|
| EMBASE | Ovid SP | From 1974 | April 2018 | September 2018 |
| MEDLINE | Ovid SP | From 1946 | April 2018 | September 2018 |
| CINAHL | EBSCOhost | From 1981 | April 2018 | September 2018 |
| PsycINFO | Ovid SP | From 1967 | April 2018 | September 2018 |
| Cochrane library | Wiley | From 1989 | April 2018 | September 2018 |

3.2.3 Search strings

3.2.3.1 Stage 1 search strategy:

Database: Ovid MEDLINE

Platform: Ovid SP

Table 2. MEDLINE search string.

| | Query | Results |
|-----|---|---------|
| #1 | Muscular Dystrophy, Duchenne/ | 4700 |
| #2 | duchenne*.mp. | 11822 |
| #3 | #1 or #2 | 11822 |
| #4 | (HR-PRO or HRPRO or HtaRQL or HRQoL or QL or QoL).ti,ab. or quality of life.mp. or (health index* or health indices or health profile*).ti,ab. or health status.mp. or ((patient or self or child or parent or carer or proxy) adj (appraisal* or appraised or report or reported or reporting or rated or rating* or based or assessed or assessment*).ti,ab. or ((disability or function or functional or functions or subjective or utility or utilities or wellbeing or well being) adj2 (index or indices or instrument or instruments or measure or measures or questionnaire* or profile or profiles or scale or scales or score or scores or status or survey or surveys)).ti,ab. | 649346 |
| #5 | 'Pediatric Quality of Life Inventory'.mp. (| 884 |
| #6 | PedsQL.mp. | 1094 |
| #7 | (SF-36 or EQ-5D*).mp. | 24322 |
| #8 | "World Health Organization Quality of Life".mp. | 1373 |
| #9 | (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI).mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms] | 2710 |
| #10 | "Childrens Assessment of Participation and Enjoyment".mp. | 96 |
| #11 | (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*).mp.11 | 17397 |



| | | |
|-----|--|--------|
| #12 | (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease").mp. | 10638 |
| #13 | (QoL-NMD DISABKIDS or "Depressionsinventar für kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM).mp. | 17368 |
| #14 | (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).mp. | 4980 |
| #15 | (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).mp. | 23191 |
| #16 | #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 | 80226 |
| #17 | #4 or #16 | 685940 |
| #18 | #3 and #17 | 523 |

3.2.3.2 Stage 2 search strategy:

Database: Ovid MEDLINE

Platform: Ovid SP

Table 3. MEDLINE search string.

| | Query | Results |
|-----|--|---------|
| #1 | Muscular Dystrophy, Duchenne | 4638 |
| #2 | duchenne*.mp. | 11693 |
| #3 | #1 or #2 | 11693 |
| #4 | Autoquestionnaire Qualite de vie Enfant Image.mp. | 18 |
| #5 | (Behavior Assessment System for Children or BASC or Parent Form 50 or PF50 or DUX-25 or EuroQoL 5-domain or Functional Independence Measure* or FIM).mp. | 4877 |
| #6 | (WeeFIM or Life Satisfaction Index or LSI or LSIA or Neurological Disorders Quality of Life Questionnaire or NeuroQOL or pediatric NeuroQOL or Offer Self-Image Questionnaire for Adolescents or OSIQ).mp. | 1602 |
| #7 | (Pediatric Outcomes Data Collection Instrument or PODCI or Neuromuscular module or DMD module or Multidimensional Fatigue Scale or Generic Short-Form or SF15).mp. | 411 |
| #8 | (SDQ or 'Strips of Life with Emoticons Questionnaire' or SOLE or 'World Health Organisation Quality of Life Scale-Brief Version' or WHOQOL-BREF).mp. | 53728 |
| #9 | ('Strength and Difficulties Questionnaire').mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms] | 223 |
| #10 | "Individualized Neuromuscular Quality of Life Questionnaire".mp. | 9 |
| #11 | (INQOL or Child Activity Limitations Interview or CALL or "Satisfaction with Life Scale" or SWLS).mp. | 1461 |
| #12 | 'Pediatric Quality of Life Inventory'.mp. | 871 |
| #13 | ("Pediatric Orthopedic Society of North America Pediatric Musculoskeletal Functional Health Questionnaire" or POSNA or Pittsburgh Sleep Quality Index).mp. | 3411 |
| #14 | PedsQL.mp. | 1078 |



| | | |
|-----|--|--------|
| #15 | (SF-36 or EQ-5D*).mp. | 23965 |
| #16 | "World Health Organization Quality of Life".mp. | 1349 |
| #17 | (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI).mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms] | 2640 |
| #18 | "Childrens Assessment of Participation and Enjoyment".mp. | 96 |
| #19 | (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*).mp. | 17206 |
| #20 | (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease").mp. | 10441 |
| #21 | (QoL-NMD DISABKIDS or "Depressionsinventar fur kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM).mp. | 17136 |
| #22 | (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).mp. | 4886 |
| #23 | (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).mp. | 22934 |
| #24 | #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 | 140059 |
| #25 | #3 and #24 | 82 |

Full copies of the search strategies for Stage 1 and Stage 2 can be found in Appendix A.

3.2.4 Additional searches

Following establishes procedures,¹¹ we searched Google Scholar (last searched 30th January 2019) with the names of the PROMs identified in Stage 1 in order to identify potential development papers for the assessment of content validity. The first 100 hits on Google Scholar were screened for inclusion. Citation tracking, by screening of references and Google Scholar citations of included articles, was conducted on the full text articles meeting eligibility criteria at Stage 2 (last searched 6th February 2019), as a supplementary measure to identify any additional studies not captured by the database searching.¹⁵

3.3 Eligibility criteria

The following selection criteria was applied to the search results at Stage 1 (identifying PROMs):

- Published in English as a full-text original research article (i.e. not including abstracts, editorials, or reviews).



- Used a self-reported, multi-item PROM to assess *at least one aspect of* QoL in boys and/or men diagnosed with DMD (assisted or proxy-reported versions of PROMs were considered for inclusion so long as a self-report version of that PROM exists).
- At least 75% of the sample, on which data from the PROM is reported, was male diagnosed with DMD.

Additional selection criteria were applied at Stage 2 (content and structural validity):

- Described data on the content and/or structural validity of the PROMs identified in Stage 1 in boys and/or men diagnosed with DMD.
- Development studies on the PROMs identified in Stage 1, to assist with the assessment of content validity, were included in any published form (i.e. journal article, book chapter, user manual).

3.4 Selection process

In order to apply the eligibility criteria for the selection of papers from search results, the following steps were performed:

- I) Inclusion criteria (Stage 1) were applied to the titles and abstracts of hits from the Stage 1 searches (and any additional papers identified in the Stage 2 searches or through citation tracking) by two independent reviewers. Records were selected for full-text review if deemed relevant, potentially relevant, or if doubt existed. Any discrepancy was resolved by a third reviewer.
- II) Full-text articles identified in (I) were screened for selection using the Stage 1 inclusion criteria by two independent reviewers. Any disagreements were resolved by a third reviewer through discussion.
- III) The PROMs identified in the articles in (II) were reviewed to ensure they met the requisite inclusion criteria (i.e. assessing an aspect of QoL). If a validated English PROM was not available for review, the corresponding articles were excluded.
- IV) Full-text articles identified in (II) and (III) meeting the Stage 1 inclusion criteria AND identified as containing measurement properties using the COSMIN filter were screened for content and structural evidence using the Stage 2 inclusion criteria by two independent reviewers. Any disagreements were resolved by a third reviewer.
- V) In order to identify key development papers for the PROMs identified in (II) and (III), Google Scholar was searched with the names of the PROMs and the first 100 hits were screened for inclusion. Results of the searches were screened for inclusion by two reviewers.



VI) Finally, citation tracking of all eligible articles at Stage 2 was conducted by reviewing references and citations on Google Scholar for any articles not identified in the initial searches that may meet the eligibility criteria for Stage 1 and/or Stage 2. Results of the searches were screened for inclusion by two reviewers.

3.5 Data extraction and quality appraisal

Data extraction was carried out by two reviewers using a pre-prepared data extraction sheet, with consensus on any ambiguities reached through discussion. The data extraction sheet was informed by the tools developed by COSMIN on reporting guidance:

<https://www.cosmin.nl/tools/guideline-conducting-systematic-review-outcome-measures/>

The methodological quality of the PROM development papers, and studies on content and structural validity were assessed using up-to-date COSMIN standards via the new COSMIN risk of bias checklist.¹⁶ Thirty-five items are used to assess the development studies, comprised of a separate rating of the quality of the concept elicitation process with patients (i.e. item generation for a new PROM), and a rating of the quality of the cognitive interview study (if present) evaluating comprehensiveness and comprehensibility of the PROM with patients.¹⁶ Thirty-eight items are used to assess studies on content validity, made of one set of items assessing studies with patients about relevance, comprehensiveness, and comprehensibility, and the other set assessing studies with professionals (if present) about relevance and comprehensiveness.¹⁶ A total rating for relevance, comprehensiveness, and comprehensibility of a PROM is determined separately. A separate rating is also determined for studies with patients or professionals. Finally, four items are used to assess the methodological quality of a structural validity study.¹⁶

When rating the methodological quality of the studies, each COSMIN standard (or item) is ranked on a 4-point scale: “very good”, “adequate”, “doubtful”, and “inadequate”. Total ratings are determined using the lowest rating for any item for that study (i.e. worst score counts).¹⁷ Studies were initially rated independently by two reviewers, and, in the case of divergence, consensus was reached in a subsequent face-to-face meeting.

3.6 Evidence synthesis

In order to synthesise and assess evidence on content validity, two reviewers independently rated the results of PROM development studies, content validity studies, and the content of the PROM itself on



10 COSMIN criteria, split into 5 on relevance, 1 on comprehensiveness, and 4 on comprehensibility.⁹ Ratings for each source of evidence could either be positive (+), negative (-), or indeterminate (?). Following this an overall judgment on the relevance, comprehensiveness, and comprehensibility of each PROM was made, which could be sufficient (+), insufficient (-), or inconsistent (\pm). Evidence on structural validity was assessed against the updated COSMIN criteria for good measurement properties, using the same rating scale as above.¹³ Finally, the quality of the evidence was graded using a modified GRADE approach,¹⁸ as either “high”, “moderate”, “low”, or “very low”. The GRADE approach takes into account the risk of bias of studies (or study quality); (in)consistency across studies; imprecision (based on sample sizes); and indirectness (of evidence).¹³

For quality assurance purposes, the quality of this systematic review itself was appraised against a recently developed COSMIN checklist to assess the quality of systematic reviews of health-related PROMs.¹⁴



4 Results

4.1 Identification of included studies

From an initial 1,752 records identified through database searching at Stage 1 and 2, 1,536 were excluded at the title/abstract review stage, leaving 216 papers for full-text review. Of these 216 papers, 87 were excluded as they were not full-text published research articles; 26 did not meet the required sample criteria of at least 75% of the sample being boys or men with DMD; 21 were judged not to be assessing QoL; 16 were not published in English; and finally 11 papers did not feature a multi-item PROM. Accordingly, a total of 55 records from the initial searches met the eligibility criteria for Stage 1. The PROMs extracted from these papers are summarised in Section 4.2 below. The observed proportionate agreement between reviewers was 92.4% at title/abstract, with Cohen's $\kappa = 0.51$ or "moderate agreement" and is similar to other published reviews.¹⁹⁻²⁰ At full-text review, the observed proportionate agreement was 93.5% with Cohen's $\kappa = 0.82$ or "almost perfect agreement". A further 5 articles that met the eligibility criteria for Stage 1 were added as a result of citation tracking.

Of the papers that initially met the review criteria at Stage 1, subsequent to a review of the PROM itself, 5 were excluded as not containing content assessing QoL; 4 were not taken forward as no free/review copy was available; 3 were excluded because no validated English copy of the PROM was available; and 1 was excluded as the particular variant of the PROM used (of a potential large item bank) was not clear. Of the resultant 47 records, 20 were identified by the COSMIN measurement properties filter¹⁴ as potentially containing measurement properties. Following review, 11 of these records were excluded for containing information on measurement properties other than content and structural validity; and 4 were excluded as not containing information on measurement properties. The remaining 5 papers featured evidence on content validity ($n = 3$, of which one was classified as a development paper) and structural validity ($n = 2$). Finally, 33 PROM development papers were identified through a review of Google Scholar search results and 3 PROM development papers were identified through citation tracking, resulting in a final selection of 41 papers that met the eligibility for Stage 2 (see Figure 1).






Figure 1. Flow diagram of search strategy and selection of papers.



4.2 PROMs used to measure quality of life in DMD

Table 4 summarises the PROMs used to assess QoL in DMD from the full-texts meeting the initial eligibility criteria at Stage 1 ($n = 60$).



Table 4. PROMs assessing QoL identified in published articles ($n = 60$) including samples of people with DMD.

| PROM | Respondent type | Recall Period | N dimensions (items) | Dimensions of quality of life assessed | Response options | Total score range | Origin | Validated English copy and development papers freely available for review? |
|---|--------------------------------|---------------------|--|---|------------------------------------|---------------------------|--------|--|
| 36-item Short-Form Health Survey (SF-36) v1.0 ²¹⁻²⁸ | Adult self-report | Varies by dimension | 8 + a single item of perceived change in health (36 items) | Physical functioning; bodily pain; role limitations due to physical health problems; role limitations due to personal or emotional problems; emotional well-being; social functioning; energy/fatigue; general health perceptions | Varies by dimension | No total score calculated | USA | Yes |
| SF-36 Health Survey v2.0 ²⁹ | Adult self-report | Varies by dimension | 8 + a single item of perceived change in health (36 items) | Assumed same as SF-36 v1.0 | Varies by dimension | No total score calculated | USA | No |
| Autoquestionnaire Qualité de vie Enfant Imagé (AUQEI) ³⁰ | Child self-report | Unknown/undefined | 4 + a total score (26 items) | Autonomy; leisure; functioning; family; total | 0 - 3 rating scale (with pictures) | 0 – 78 (raw) | France | No |
| Beck Depression Inventory I (BDI) ²⁵ | Adult interview or self-report | Present/today | 1 total score (21 items) | Depression | 0 - 3 rating scale | 0 – 63 (raw) | USA | Yes |



| | | | | | | | | |
|--|--|---|---|---|---|---|---------|-----|
| Behavior Assessment System for Children (BASC) first edition ³¹ | Child self-report Parent report Teacher report | Unknown (withdrawn from use, superseded by BASC-II and BASC-III) | Unknown (withdrawn from use, superseded by BASC-II and BASC-III) | Unknown (withdrawn from use, superseded by BASC-II and BASC-III) | Unknown (withdrawn from use, superseded by BASC-II and BASC-III) | Unknown (withdrawn from use, superseded by BASC-II and BASC-III) | USA | No |
| Child Activity Limitations Interview (CALI) ³¹ | Child interview or self-report | Last 4 weeks | 1 total score (8 items chosen from a set of 21) | Activity limitations | 0 - 4 rating scale | 0 – 32 (raw) | USA | Yes |
| Child Health Questionnaire - Parent Form 50 (CHQ-PF50) ³²⁻³³ | Parent self-report | Last 4 weeks (past year for change in health) | 14 (50 items) | Physical functioning; role/social limitations – physical; role/social limitations – emotional; role/social limitations – behavioral; general health perceptions; bodily pain/discomfort; family activities; parent impact – time; parent impact – emotion; self-esteem; mental health; behaviour; family cohesion; change in health | Varies by dimension | No total score calculated | USA | No |
| Children’s Assessment of Participation and Enjoyment (CAPE) ³⁴⁻³⁶ | Child and young adults self-report or interview | Unknown/undefined | 5 + a total score (55 items) | Diversity of activities; intensity of activities (frequency of participation); enjoyment of activities; with whom; where; total participation | Varies by dimension | 0 – 55 (raw) | USA | No |
| Depressions-Inventar für Kinder und Jugendliche (DIKJ) 2 nd edition ²⁵ | Child self-report | Unknown/undefined | 1 total score (26 items) | Depression | Unknown/undefined | 0 – 46 (raw) | Germany | No |



| | | | | | | | | |
|--|--|-------------------------|--|---|---|-----------------------------|---------------|-----|
| DISABKIDS generic module (DCGM-37) ²⁵ | Child self-report Proxy report | Past 4 weeks | 6 + a total score (37 items) | Independence; emotion; social inclusion; social exclusion; limitation; treatment; total | 1 – 4 rating scale | 37 – 148 (raw) | Multi-country | Yes |
| DISABKIDS – Smileys ²⁵ | Child self-report Proxy report | Assumed same as DCGM-37 | Assumed same as DCGM-37 (12 items) | Assumed same as DCGM-37 | Assumed same as DCGM-37 (with pictures) | 12 – 48 (raw) | Multi-country | No |
| Dutch Children AZL/TNO Questionnaire Quality of Life Short Form (DUC-25) ³⁶ | Child self-report | Unknown/undefined | 4 + a total score (25 items) | Physical; emotional; social; home functioning; total | 0 - 4 rating scale | 0 – 100 (raw) | Netherlands | No |
| EuroQoL 5-domain 3-Level (EQ-5D-3L) ³⁷⁻³⁸ | Adult self-report Proxy report | Present/today | 5 (5 items) + self-rated health VAS | Mobility; self-care; usual activities; pain/discomfort; anxiety/depression | 1 – 3 rating scale | –0.594 – 1 (utility scores) | Multi-country | Yes |
| Fatigue Severity Scale (FSS) ²⁷ | Adult self-report | Within last week | 1 total score (9 items) + global fatigue VAS | Fatigue severity | 1 – 7 rating scale | 1 – 63 (raw) | USA | Yes |
| Generalized Anxiety Disorder Scale 7-item (GAD-7) ³⁹ | Adult self-report | 2 weeks | 1 total score (7 items) | Anxiety | 0 – 3 rating scale | 0 – 21 (raw) | USA | Yes |
| Health Utilities Index Questionnaire mark 2 (HUI-2) 15Q ⁴⁰⁻⁴¹ | Child self-report Adult self-report Proxy report | During past 4 weeks | 7 (15 items) | Sensation; mobility; emotion; cognition; self-care; pain; fertility | Varies by dimension | –0.03 – 1 (utility scores) | Canada | Yes |



| | | | | | | | | |
|--|--|---------------------|-------------------------------|--|---|-------------------------------|---------------|-----|
| Health Utilities Index Questionnaire mark 3 (HUI-3) 15Q ⁴⁰⁻⁴¹ | Child self-report Adult self-report Proxy report | During past 4 weeks | 8 (15 items) | Vision; hearing; speech; ambulation; dexterity; emotion; cognition; pain | Varies by dimension | -0.36 – 1 (utility scores) | Canada | Yes |
| Hospital Anxiety and Depression Scale (HADS) ^{27,38} | Adult self-report | Last week | 2 (14 items) | Anxiety; depression | 0 – 3 rating scale | No total score calculated | UK | Yes |
| Individualized Neuromuscular Quality of Life Questionnaire (INQOL) ²⁹ | Adult self-report | At the moment | 10 + a total score (45 items) | Weakness; locking; pain; fatigue; activities; independence; social relationships; emotions; body image; treatment; total | 7-point rating scale, varies by dimension | Scoring unclear | UK | Yes |
| KIDSCREEN-10 ⁴² | Child self-report Proxy report | Last week | 1 total score (10 items) | Health-related quality of life | 1 – 5 rating scale | 10 – 50 (raw) | Multi-country | Yes |
| KIDSCREEN-27 ⁴³ | Child self-report Proxy report | Last week | 5 (27 items) | Physical well-being; psychological well-being; autonomy and parent relation; social support and peers; school environment | 1 – 5 rating scale | No total score calculated | Multi-country | Yes |
| KIDSCREEN-52 ⁴⁴ | Child self-report Proxy report | Last week | 10 (52 items) | Physical well-being; psychological well-being; moods and emotions; self-perception; autonomy; parent relation and home life; financial resources; social support and peers; school environment; social acceptance (bullying) | 1 – 5 rating scale | No total score calculated | Multi-country | Yes |



| | | | | | | | | |
|---|--|---------------------|--|---|-----------------------------------|---------------------------|--------|-----|
| Life Satisfaction Index for Adolescents (LSIA) ⁴⁵⁻⁴⁸ | Child and young adults self-report | At present | 5 + a total score (45 items) | General well-being; interpersonal relationships; personal development; personal fulfilment; leisure and recreation; total | 1 – 5 rating scale (plus 0 = N/A) | 0 – 225 (raw) | Canada | Yes |
| Muscular Dystrophy Child Health Index of Life with Disabilities (MDCHILD) ⁴⁹ | Child self-report | Past 4 weeks | 7 + a total score (47 items) | Activities of daily living & independence; positioning, transferring, & mobility; comfort & endurance; emotions & behaviour; social interaction & school; health; your overall quality of life; total | Varies by dimension | 0 – 100 (transformed) | Canada | Yes |
| Neurological Disorders Quality of Life Questionnaire (Neuro-QoL) ²⁶ | Adult self-report | Varies by dimension | Up to 16 (up to 564 items in item banks) | Ability to participate in social roles and activities; anxiety; bowel function; cognitive function; communication; depression; emotional and behavioral dyscontrol; fatigue; lower extremity function – mobility; positive affect and well-being; satisfaction with social roles and activities; sleep disturbance; sexual function; stigma; upper extremity function – fine motor, ADL; urinary/bladder function | 1 – 5 rating scale | No total score calculated | USA | Yes |
| Offer Self-Image Questionnaire for Adolescents (OSIQ) ^{45,48} | Child and young adult self-report or interview | Unknown/undefined | 11 + a total score (130 items) | Impulse control; emotional tone; body and self-image; social relationships; morals; vocational and educational goals; family relationships; mastery of the external world; psychopathology; superior adjustment; total | 1 – 6 rating scale | 130 – 780 (raw) | USA | No |



| | | | | | | | | |
|--|--|---|--|---|---------------------|---------------------------|-----|-----|
| Patient Health Questionnaire 9-item (PHQ-9) ³⁹ | Adult self-report | 2 weeks | 1 total score (9 items) | Depression | 0 – 3 rating scale | 0 – 27 (raw) | USA | Yes |
| Pediatric Neurological Disorders Quality of Life Questionnaire (Pediatric Neuro-QoL) ²⁶ | Child self-report | Varies by dimension | Up to 11 (up to 161 items in item banks) | Anger; anxiety; cognitive function; depression; fatigue; lower extremity – mobility; pain; social relations – interaction with adults; social relations – interaction with peers; stigma; upper extremity – fine motor, ADL | 1 – 5 rating scale | No total score calculated | USA | Yes |
| Pediatric Outcomes Data Collection Instrument (PODCI) ⁵⁰⁻⁵³ | Child self-report Proxy report | Varies by dimension | 7 (86 items) | Global function & comfort; upper extremity function; physical function and sport; transfers and mobility; comfort; POSNA happy and satisfied; POSNA expectations | Varies by dimension | No total score calculated | USA | Yes |
| Pediatric Quality of Life Inventory (PedsQL) 3.0 DMD module ⁵⁴⁻⁵⁵ | Child and young adult self-report Proxy report | Past month or past 7 days (acute version) | 4 (18 items) | Daily activities; treatment barriers; worry; communication | 0 – 4 rating scale | No total score calculated | USA | Yes |
| PedsQL 3.0 Multidimensional fatigue scale (MFS) ⁵⁵⁻⁵⁶ | Adult self-report Child and young adult self-report Proxy report | Past month or past 7 days (acute version) | 3 + a total score (18 items) | General fatigue; sleep/rest fatigue; cognitive fatigue; total fatigue | 0 – 4 rating scale | 0 – 100 (transformed) | USA | Yes |
| PedsQL 3.0 Neuromuscular module (NMM) ^{26,41-42,55-61} | Child and young adult self-report Proxy report | Past month or past 7 days (acute version) | 3 + a total score (25 items) | About my/my child's neuromuscular disease; communication; about our family resources; total | 0 – 4 rating scale | 0 – 100 (transformed) | USA | Yes |



| | | | | | | | | |
|--|--|---|---|---|------------------------------------|-----------------------|-------|-----|
| Pediatric Quality of Life Inventory (PedsQL) 4.0 Generic Core Scales (GCS) ^{26,34,42,50,53-57,59,62-69} | Adult self-report Child and young adult self-report Proxy report | Past month or past 7 days (acute version) | 5 + a total score (23 items) | Physical health; psychosocial health; emotional functioning; social functioning; school functioning; total | 0 – 4 rating scale | 0 – 100 (transformed) | USA | Yes |
| PedsQL 4.0 Generic Short-form (SF-15) ⁷⁰ | Adult self-report Child and young adult self-report Proxy report | Past month or past 7 days (acute version) | 5 + a total score (15 items) | Physical health; psychosocial health; emotional functioning; social functioning; school functioning; total | 0 – 4 rating scale | 0 – 100 (transformed) | USA | Yes |
| Pittsburgh Sleep Quality Index (PSQI) ^{26,66} | Adult self-report | Past month | 8 + a total score (10 items) | Subjective sleep quality; sleep latency; sleep duration; sleep efficiency; sleep disturbance; use of sleep medication; daytime dysfunction; total | Varies by dimension | 0 – 21 (raw) | USA | Yes |
| Satisfaction with Life Scale (SWLS) ^{39,45} | Adult self-report | Undefined/present time | 1 total score (5 items) | Life satisfaction | 1 – 7 rating scale | 5 – 35 (raw) | USA | Yes |
| Strength and Difficulties Questionnaire (SDQ) ⁷¹⁻⁷³ | Child self-report Proxy report | Last 6 months | 5 + a total score (25 items) + an impact supplement | Emotional symptoms; conduct problems; hyperactivity/inattention; peer relationship problems; total difficulties; prosocial behaviour | 0 – 2 rating scale | 0 – 40 (raw) | UK | Yes |
| Strips of Life with Emoticons Questionnaire (SOLE) ⁷⁴ | Child self-report | Specific scenarios | 1 total score (33 items) | Quality of life | 0 – 2 rating scale (with pictures) | 0 – 66 (raw) | Italy | No |



| | | | | | | | | |
|---|--|--------------------|--|---|---------------------|---------------------------|---------------|-----|
| TNO-AZL Children's Quality of Life questionnaire (TACQoL) ⁷⁵ | Child self-report Proxy report | The last few weeks | 7 (56 items) | Physical functioning; motor functioning; independent daily functioning; cognitive functioning and school performance; social contacts; positive moods; negative moods | Varies by dimension | No total score calculated | Netherlands | No |
| TNO-AZL Adult Quality of Life questionnaire (TAAQoL) ⁷⁵ | Adult self-report | In the last month | 12 (45 items) | Gross motor functioning; fine motor functioning; cognition; sleep; pain; social contacts; daily activities; sex; vitality; happiness; depressive mood; anger | Varies by dimension | No total score calculated | Netherlands | No |
| World Health Organisation Quality of Life Scale-Brief Version (WHOQOL-BREF) ^{24,26-28} | Adult self-report or interview Proxy report | 2 weeks | 4 (26 items, 24 items make up domain scores) | Physical health; psychological; social relationships; environment | 1 – 5 rating scale | No total score calculated | Multi-country | Yes |

Note. References next to PROM names represent published studies where the PROM has been used in a sample of people with DMD.



A total of 40 PROMs used to assess at least one aspect of QoL in DMD were identified in published research articles through database searching (the two HUI classification systems use the same 15-item PROM). The majority of the PROMs were multidimensional ($n = 32$), designed to assess a range of different facets of QoL. The remaining unidimensional scales were designed to assess: activity limitations (CALI); anxiety (GAD-7); depression (BDI, DIKJ, PHQ-9); fatigue severity (FSS); life satisfaction (SWLS); or quality of life/health-related quality of life unidimensionally (KIDSCREEN, SOLE). Twenty-four of the PROMs had versions designed for completion by adult or young adult respondents, and 26 had versions designed for children. The most popular PROMs used in published research articles assessing QoL in people with DMD were the PedsQL 4.0 GCS (18 articles); PedsQL 3.0 NMM (10 articles); and the SF-36 (8 articles).

In the current review, 26 PROMs were taken forward for COSMIN quality assessment on content and structural validity in DMD. The remaining 14 PROMs were not assessed for the following reasons: a copy of the PROM itself and/or necessary development papers were not freely accessible for review (CAPE, CHQ-PF50, DISABKIDS Smileys, OSIQ, SF-36 v2); no formally validated English copy of the PROM was available or in use (AUQEI, DIKJ, DUC-25, SOLE, TAAQoL, TACQoL); the PROM was no longer available or recommended for use (BASC 1st edition, which has been superseded by the BASC 2); or it was unclear from the study which of a large number of possible variants of a PROM were used (pediatric Neuro-QoL, Neuro-QoL). Table 5 lists the PROMs taken forward for review.

4.3 Content validity

4.3.1 Appraisal of PROM development studies

Table 5 summarises key characteristics and COSMIN quality assessment of the development of the PROMs included in the review, this includes the definition of the construct intended to be measured, target population, and intended context of use of the PROM. Five PROMs were developed to be intended for use specifically within neuromuscular disorders (INQoL, PedsQL 3.0 NMM) or DMD (LSIA, MDCHILD, PedsQL 3.0 DMD module; Table 5). Eleven PROMs either had no patients involved in their development, or it was unclear if patients were involved.

The joint most common COSMIN quality rating assigned to the PROMs for concept elicitation was inadequate ($n = 12$). This was primarily due to: the PROM development study not being performed in a sample of patients representing the target population (BDI, EQ-5D-3L, GAD-7, HADS, HUI 15Q,



PedsQL 3.0 MFS, PHQ-9, SDQ, SF-36, and SWLS); or inadequacies within the details of the qualitative methods used (FSS, INQoL). The concept elicitation study of 11 further PROMs was rated as doubtful due to at least some unclear details/suspected problems within the qualitative methods used (CALI, DCGM-37, LSIA, MDCHILD, PODCI, PedsQL 3.0 NMM, PedsQL 3.0 DMD, PedsQL 4.0 GCS, PedsQL 4.0 SF-15, PSQI, WHOQOL-BREF). Only the KIDSCREEN family of measures ($n = 3$) received an adequate rating for concept elicitation and PROM design. However, the KIDSCREEN measures received a doubtful rating for the overall PROM development study, for failing to provide evidence that comprehensibility and comprehensiveness were assessed in the cognitive interview/pilot study of the PROM.



Table 5. Characteristics and assessment of development papers for PROMs included in the review.

| PROM | Reference(s) | Original language | Construct definition | Target population | Intended context of use | Concept elicitation study | |
|-----------|---|----------------------------------|---|--|---|---------------------------|-------------------------|
| | | | | | | COSMIN quality rating | Were patients involved? |
| BDI | Beck et al. 1961 ⁷⁶ | English (US) | "the items were chosen on the basis of their relationship to the overt behavioral manifestations of depression and do not reflect any theory regarding the etiology or the underlying psychological processes in depression" | Adult patients with suspected symptoms of depression | Quantitative assessment of the intensity of depression in diagnostic and research settings | Inadequate | No |
| CALI | Palermo et al. 2004 ⁷⁷ | English (US) | "functional impairment, defined as difficulty in performing age-appropriate physical, mental, and social activities in daily life due to physical health status (...) functional impairment due to pain (...) specific areas of functioning that are important to children and adolescents with recurrent and chronic pain" | School-age children and adolescents with recurrent and chronic pain | Research and clinical care | Doubtful | Yes |
| DCGM-37 | Petersen et al. 2005 ⁷⁸ Ravens-Sieberer et al. 2007 ⁷⁹ | English (UK) | "a multidimensional construct with social, physical, emotional, and functional domains" | Children aged 4-7 years and 8-16 years with chronic health conditions | Clinical studies or surveys | Doubtful | Yes |
| EQ-5D-3L* | EuroQol Group 1990 ⁸⁰ Brooks et al. 1996 ⁸¹ | Multiple, including English (UK) | "Health-related quality of life" | "Large-scale surveys of the community and (...) for use in postal surveys" | "Complement other quality of life measures, collection of common data set for reference. Generate cross-national comparisons of health state valuations." | Inadequate | No |
| FSS | Krupp et al. 1989 ⁸² | English (US) | "Fatigue" | Patients with "clinical disorders" | Clinical research studies and surveys | Inadequate | No |



| | | | | | | | |
|---------------------|---|--------------|--|--|---|------------|---------|
| GAD-7 | Spitzer et al. 2006 ⁸³ | English (US) | "We first selected potential items for a brief GAD [Generalized Anxiety Disorder] scale (...) that reflected all of the <i>Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV)</i> symptom criteria for GAD and (...) on the basis of review of existing anxiety scales." | General adult population | Clinical practice and research | Inadequate | No |
| HADS | Zigmond & Snaith 1983 ⁸⁴ | English (UK) | "depression subscale were largely based on the anhedonic state (...) psychic manifestations of anxiety neurosis" | Patients under investigation and treatment in medical and surgical departments in non-psychiatric hospital departments | Clinical/screening use within non-psychiatric hospital departments | Inadequate | No |
| HUI-2 / HUI-3 (15Q) | Feeny et al. 1995 ⁸⁵ Torrance et al. 1996 ⁸⁶ | English (US) | "The HUI Mark II and Mark III systems are based on concepts of functional capacity rather than performance (...) generic health profile measures that also permit the computation of a single summary score quantifying health-related quality of life" | Originally survivors of childhood cancer (HUI-2), extended to adults | Clinical evaluative and population health survey studies, in clinical trials, and cost-utility analyses | Inadequate | Unknown |
| INQoL | Vincent et al. 2007 ⁸⁷ | English (UK) | "The structure of INQoL was based on the ICDH-2 model of disease incorporating the concepts of Impairment, Activities, and Participation." | Adults with neuromuscular disorders (16+ years) | Clinical and research use | Inadequate | Yes |



| | | | | | | | |
|--------------|--|----------------------------------|--|---|---|----------|-----|
| KIDSCREEN-52 | Ravens-Sieberer et al. 2001 ⁸⁸ Ravens-Sieberer et al. 2005 ⁸⁹ Detmar et al. 2006 ⁹⁰ | Multiple, including English (UK) | “Health-related quality of life is described as a multidimensional construct covering physical, emotional, mental, social, and behavioral components of well-being and function as perceived by patients and/or individuals (...) agreement was reached that the questionnaire should aim to measure HRQOL as a generic construct in largely healthy children, thus more emphasis was given to the inclusion of psychosocial domains, and less to domains of physical functioning or symptoms such as pain.” | Healthy and chronically-ill children and adolescents between 8 and 18 years | Epidemiological and paediatric studies, clinical settings (healthcare system), and health services research | Adequate | Yes |
| KIDSCREEN-27 | Ravens-Sieberer et al. 2006 ⁹¹ | Assumed the same as KIDSCREEN-52 | Assumed the same as KIDSCREEN-52 | Assumed the same as KIDSCREEN-52 | Assumed the same as KIDSCREEN-52 | Adequate | Yes |
| KIDSCREEN-10 | Ravens-Sieberer et al. 2006 ⁹¹ | Assumed the same as KIDSCREEN-52 | Assumed the same as KIDSCREEN-52 | Assumed the same as KIDSCREEN-52 | Assumed the same as KIDSCREEN-52 | Adequate | Yes |



| | | | | | | | |
|----------------|--|--------------|---|--|--|------------|-----|
| LSIA | Reid & Renwick 1994 ⁴⁵ | English (US) | "quality of life is to conceptualize it as a subjective phenomenon. Specifically, it is viewed in terms of the individual's feelings and evaluations of his or her life circumstances. Many researchers who study quality of life within this perspective emphasize the importance of measuring the individual's degree of life satisfaction. In other words, they are interested in how pleased an individual feels about particular aspects of his or her life" | "Individuals between the ages of 12 and 19 years who have DMD" | Research instrument and potentially useful as a clinical measure | Doubtful | Yes |
| MDCHILD | Propp, 2017 ⁹² Propp et al. 2019 ⁴⁹ | English (UK) | "Health-related priorities for children with DMD (...) defined as concerns, desires, and expectations arising from the lived experience of that condition" | Children with DMD (assumed 5 – 18 years) | Cohort studies, clinical trials, and clinical decision-making | Doubtful | Yes |
| PedsQL 3.0 DMD | Uzark et al. 2012 ⁵⁴ | English (US) | "Health-related quality of life (QoL), a multidimensional construct that includes physical, psychological, and social functioning, has emerged as an important outcome in pediatric populations with chronic health conditions." | Children with DMD from 2 – 18 years | Assumed the same as PedsQL 4.0 GCS | Doubtful | Yes |
| PedsQL 3.0 MFS | Varni et al. 2002 ⁹³ | English (US) | "designed to measure child and parent perceptions of fatigue in pediatric patients" | Assumed the same as PedsQL 4.0 GCS | "may be utilized as outcome measures in pediatric cancer clinical trials, research, and clinical practice for HRQOL" | Inadequate | Yes |



| | | | | | | | |
|------------------|--------------------------------------|--------------|---|---|---|----------|-----|
| PedsQL 3.0 NMM | Iannaccone et al. 2009 ⁹⁴ | English (US) | "HRQOL is a multidimensional construct, consisting at the minimum of physical, psychological (including emotional and cognitive), and social health dimensions delineated by the World Health Organization. HRQOL has emerged as the most appropriate term for quality of life dimensions that represent a patient's perceptions of the impact of an illness and its treatment on their own functioning and well-being and which are within the scope of healthcare services and medical products." | Children and young people with neuromuscular disorders, in particular spinal muscular atrophy | Assumed the same as PedsQL 4.0 GCS | Doubtful | Yes |
| PedsQL 4.0 GCS | Varni et al. 1999 ⁹⁵ | English (US) | "The PedsQL measures the patient's and the parent's perceptions of the patient's HRQOL, as defined in terms of the impact of disease and treatment on an individual's physical, psychological, and social functioning, and by disease/treatment-specific symptoms." | Children aged 8 – 18 across various pediatric chronic health conditions | Epidemiological studies, clinical trials, and performance improvement studies | Doubtful | Yes |
| PedsQL 4.0 SF-15 | Varni et al. 1999 ⁹⁵ | English (US) | Assumed the same as PedsQL 4.0 GCS | Assumed the same as PedsQL 4.0 GCS | Assumed the same as PedsQL 4.0 GCS | Doubtful | Yes |



| | | | | | | | |
|-------|--|--------------|--|---|---|------------|---------------|
| PHQ-9 | Spitzer et al. 1999 ⁹⁶ Kroenke et al. 2001 ⁹⁷ | English (US) | "Depression (...) using diagnostic criteria from the <i>Diagnostic and Statistical Manual of Mental Disorders, Revised Third Edition (DSM-III-R)</i> and <i>Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV)</i> ." | General adult population | Clinical practice and research | Inadequate | No |
| PODCI | Daltroy et al. 1998 ⁹⁸ | English (US) | "The POSNA outcomes instrument scales assess upper extremity function, transfers and mobility, physical function and sports, comfort (painfree), happiness and satisfaction, and expectations for treatment. A POSNA global scale combines the three function subscales and comfort." | Children aged 2-18 years with musculoskeletal disorders | "Patient-based instrument" | Doubtful | Yes (assumed) |
| PSQI | Buysse et al. 1989 ⁹⁹ | English (US) | "sleep quality is a readily accepted clinical construct, it represents a complex phenomenon that is difficult to define and measure objectively. 'Sleep quality' includes quantitative aspects of sleep, such as sleep duration, sleep latency, or number of arousals, as well as more purely subjective aspects, such as "depth" or "restfulness" of sleep" | Clinical/psychiatric populations | Psychiatric clinical practice and research activities | Doubtful | No |
| SDQ | Goodman 1997 ¹⁰⁰ | English (UK) | "young people's behaviours, emotions, and relationships" | Children and young people (aged 4-16 years) | "to meet the needs of researchers, clinicians, and educationalists" | Inadequate | No |



| | | | | | | | |
|-------------|---|--------------|--|-----------------------------------|--|------------|----|
| SF-36 v1.0* | Ware & Sherbourne, 1992 ¹⁰¹ Hays et al. 1993 ¹⁰² Jenkinson et al. 1999 ¹⁰³ Ware 2000 ¹⁰⁴ | English (US) | ““Health”, eight concepts: physical functioning, social and role functioning, mental health, general health perceptions, bodily pain, and vitality.” | “General population and patients” | “Clinical practice and research, healthy policy evaluations, and general population surveys” | Inadequate | No |
| SWLS | Diener et al. 1985 ¹⁰⁵ | English (US) | "Life satisfaction refers to a cognitive, judgmental process. Shin and Johnson (1978) define life satisfaction as "a global assessment of a person's quality of life according to his chosen criteria" (p. 478)" | Unclear | Unclear | Inadequate | No |



| | | | | | | | |
|-------------|---|----------------------------------|---|--|--|----------|-----|
| WHOQOL-BREF | WHOQOL Group 1994 ¹⁰⁶ WHOQOL Group 1995 ¹⁰⁷ Skevington et al. 1997 ¹⁰⁸ WHOQOL Group 1998 ¹⁰⁹ | Multiple, including English (UK) | " It is a broad ranging concept incorporating, in a complex way, the person's physical health, psychological state, level of independence, social relationships, personal beliefs, and relationship to salient features of the environment (...) At minimum, quality of life includes the following dimensions: physical (individuals' perception of their physical state), psychological (individuals' perception of their cognitive and affective state) and social (individuals' perception of the interpersonal relationship relationships and social roles in their life). (...) The WHOQOL includes a spiritual dimension (the person's perception of 'meaning in life', or the overarching personal beliefs that structure and qualify experience)." | "assess the quality of life of chronic disease sufferers, informal caregivers of the sick and disabled, people living in high-stress conditions like refugees, and 'healthy' people" | "in routine clinical work, large scale epidemiological studies and in clinical trials" | Doubtful | Yes |
|-------------|---|----------------------------------|---|--|--|----------|-----|

Note. *PROM development information obtained from prior COSMIN review¹¹⁰, not re-extracted or re-rated in this review, based on COSMIN guidance⁹



4.3.2 Appraisal of content validity studies

Aside from development studies, only 2 published articles had independently assessed the content validity of the QoL PROMs in samples of people with DMD, and the details of these studies and their COSMIN quality assessment are described in Table 6. Neither of these studies were conducted in a UK context, and instead were cross-cultural validation studies. Hu et al. (2013)⁵⁹ assessed the relevance, comprehensiveness, and comprehensibility of the PedsQL 3.0 NMM in Chinese children with DMD. Simon et al. (2017)⁴⁷ assessed comprehensibility of the LSIA in Brazilian children with DMD, and comprehensiveness in professionals. However, both of these studies received ratings of doubtful due to at least some unclear details/suspected problems within the qualitative methods used.



Table 6. Characteristics and assessment of content validity papers in DMD samples for PROMs included in the review.

| PROM | Reference | Language (Country) | DMD sample characteristics | | | COSMIN rating | | | Results (synthesis) |
|----------------|---------------------------------|-------------------------------|----------------------------|-----------------------|-------------------------------------|---------------|-------------------|-------------------|--|
| | | | N | Age (years, \pm SD) | % ambulatory | Relevance | Comprehensiveness | Comprehensibility | |
| LSIA | Simon et al. 2017 ⁴⁷ | Brazilian Portuguese (Brazil) | 43 | 11.4 \pm 3.38 | Not stated | / | Doubtful | Doubtful | The level of comprehension reached via the final Probe technique was 97% for the parent version and 95% for the patient version, which is above the minimum of 85% required. |
| PedsQL 3.0 NMM | Hu et al. 2013 ⁵⁹ | Chinese (China) | 56 | 7.54 \pm 4.06 | 37 children "could climb stairs" | Doubtful | Doubtful | Doubtful | Cognitive debriefing was conducted with six children with DMD and their parents to confirm that the final Chinese version was understandable and acceptable. |

Note. / = content validity aspect not evaluated.



4.3.3 Evidence synthesis

The evidence from the PROM development papers and content validity studies was combined with reviewer ratings of the PROMs to produce a synthesis of the available evidence using the 10 COSMIN criteria for good content validity.⁹ Most of the quality of the evidence was downgraded from High to Low or Very Low due to the assessment being based on development studies of doubtful or inadequate quality, respectively.⁹ Only the LSIA and the PedsQL 3.0 NMM had moderate supporting evidence, featuring independent content validity studies as well as development papers. The KIDSCREEN measures and the LSIA were the only PROMs to receive satisfactory results for all three dimensions of content validity: relevance; comprehensiveness; and comprehensibility, based on the evidence available. Full synthesised results are presented in Table 7.



Table 7. Evidence synthesis on the content and structural validity of PROMs used to assess QoL in people with DMD.

| PROM | Content validity | | | Structural Validity | | |
|---------------------|------------------|-------------------|-------------------|---------------------|-------------------|---------------------|
| | Relevance | Comprehensiveness | Comprehensibility | Quality of evidence | Rating of results | Quality of evidence |
| BDI | ± | - | ± | Very low | ? | ? |
| CALI | + | - | ± | Low | ? | ? |
| DCGM-37 | ± | + | + | Low | ? | ? |
| EQ-5D-3L | + | - | + | Very low | ? | ? |
| FSS | ± | - | ± | Very low | ? | ? |
| GAD-7 | + | - | + | Very low | ? | ? |
| HADS | - | - | - | Very low | ? | ? |
| HUI-2 / HUI-3 (15Q) | - | - | - | Very low | ? | ? |
| INQoL | ± | ± | + | Very low | ? | ? |
| KIDSCREEN-52 | + | + | + | Low | ? | ? |
| KIDSCREEN-27 | + | + | + | Low | ? | ? |
| KIDSCREEN-10 | + | + | + | Low | ? | ? |
| LSIA | + | + | + | Moderate | ? | ? |
| MDCHILD | ± | + | + | Low | ? | ? |
| PedsQL 3.0 DMD | ± | ? | ± | Very low | ? | ? |
| PedsQL 3.0 MFS | ± | - | ± | Very low | ? | ? |
| PedsQL 3.0 NMM | ± | ? | ± | Moderate | - | High |
| PedsQL 4.0 GCS | ± | + | ± | Low | ? | Very low |
| PedsQL 4.0 SF-15 | ± | + | ± | Low | ? | ? |
| PHQ-9 | + | - | ± | Very low | ? | ? |
| PODCI | ± | + | ± | Very low | ? | ? |
| PSQI | ± | - | ± | Very low | ? | ? |
| SDQ | - | - | + | Very low | ? | ? |
| SF-36 v1.0 | + | + | ± | Very low | ? | ? |
| SWLS | - | - | ± | Very low | ? | ? |
| WHOQOL-BREF | + | + | ± | Very low | ? | ? |

Note. + = satisfactory results; - = unsatisfactory results; ± = inconsistent results; ? = indeterminate.



4.4 Structural validity

4.4.1 Appraisal of structural validity studies

Two studies had assessed the structural validity of the PROMs included in this review in samples of people with DMD (Table 8). Both of these were conducted using English versions of the PROMs and either in the UK or USA. Lim et al. (2014)⁶⁴ assessed the structural validity of the PedsQL 4.0 GCS using an unspecified Rasch model in 63 boys with DMD. This study received a COSMIN quality rating of doubtful because it was doubtful that the sample size included in the analysis was adequate. Landfeldt et al. (2018)⁵⁸ assessed the structural validity of the PedsQL 3.0 NMM using a Rasch partial-credit model (PCM) in 278 people with DMD. This study received a very good COSMIN quality rating for its methodological content.



Table 8. Characteristics, assessment, and results of structural validity papers in DMD samples for PROMs included in the review.

| PROM | Reference | Country (language) | Patient characteristics | | | PROM score | COSMIN Quality Rating | Analysis – model | Results (synthesis) |
|----------------|-------------------------------------|--------------------|--|-----------------|--|---|-----------------------|-----------------------------|---|
| | | | N | Age (yr, M ±SD) | % ambulatory | | | | |
| PedsQL 4.0 GCS | Lim et al. 2014 ⁶⁴ | USA (English) | 63 boys with DMD (and up to 50 parents, not necessarily matched) | 10.2 ± 2.5 | 95.24 | Child: M = 64.5, SD = 15.3. Parent: M=56.2, SD=12.9. | Doubtful | Rasch (model not specified) | Model misfit for items determined with infit > 1.4 and outfit > 2.0 MnSq values and standardized scores > 2.0. All items fit in parent proxy-reports of physical health scale and child self-reports of psychosocial health scale. 2 out of 8 items showed high infit statistics in child self-reports of the physical health scale (taking a bath or shower; doing chores around the house). In addition 2 out of 15 items showed high infit for the parent proxy-reports of the psychosocial health scale (trouble sleeping; keep up with school work). |
| PedsQL 3.0 NMM | Landfeldt et al. 2018 ⁵⁸ | UK / USA (English) | 278 (95 UK) | 16 ± 7 | 40% not "full-time wheelchair dependent" | Not reported | Very good | Rasch PCM | Eight items displayed inadequate fit (χ^2 : $p > 0.01$). Six items had fit residuals ≤ -2.5 or ≥ 2.5 (4 significant at $p < .05$). Inadequate overall fit (χ^2 item-trait interaction: $p = < .001$). Disordered thresholds for 22 of 25 items. Suboptimal targeting. |



4.4.2 Evidence synthesis

Of the 2 studies that assessed the structural validity of the PedsQL 4.0 GCS and PedsQL 3.0 NMM in people with DMD, neither provided satisfactory results (Table 7). First, the structural validity of the PedsQL 4.0 GCS in people with DMD received an indeterminate rating, as key details of the results from the Rasch model denoting good measurement properties were not reported. Due to the risk of bias assessment of Lim et al. (2014)⁶⁴ the quality of the evidence supporting this indeterminate conclusion was rated as very low. Second, the structural validity of the PedsQL 3.0 NMM in people with DMD received an unsatisfactory rating, as the psychometric criteria for good measurement properties were not met. The favourable risk of bias assessment for Landfeldt et al. (2018)⁵⁸ meant that the quality of evidence supporting this conclusion was graded as high.



4.5 Quality assurance of the review

The quality of this review was self-assessed against a newly derived COSMIN checklist¹⁴, designed to evaluate the quality of systematic reviews of health-related PROMs. The results are displayed in Table 9. In general, the review meets numerous quality indicators as defined by the COSMIN team, including the elements included in the research aim, search strategies, article selection, and assessment of measurement properties and quality. In a couple of instances, criteria have been partly met. For example, in this review all instruments were included where a *validated English copy was freely available for review*. It is possible that additional instruments could have been included if licenses were paid for to access the relevant PROMs and development materials. Second, citation tracking (i.e. reference checking) was conducted on the final set of articles eligible at Stage 2 of the searches ($n = 41$), but not on results eligible for inclusion at Stage 1.



Table 9. Quality assessment of this systematic review against COSMIN guidance.

| COSMIN criteria | Review meets criteria |
|--|-----------------------|
| Elements included in the research aim: | |
| Construct of interest | + |
| Population of interest | + |
| Type of measurement instrument of interest | + |
| Measurement properties of interest | + |
| All available instruments included | ± |
| Only instruments included that have at least some evidence of measurement properties | + |
| Search strategy described | + |
| No search terms or validated search filter used for: | |
| Measurement properties | - |
| Type of instrument | - |
| Number of databases searched: | 5 |
| Search in at least 2 databases | + |
| MEDLINE/PubMed | + |
| EMBASE | + |
| Additional databases | + |
| Reference checking used | ± |
| No time limits used or good arguments for a time limit | + |
| No language restrictions used | + |
| Inclusion and exclusion criteria clearly described | + |
| Reasons for excluding articles reported | + |
| Abstract selection by at least 2 reviewers? | + |
| Full-text article selection by at least 2 reviewers? | + |
| Abstract and full-text article selection by at least 2 reviewers? | + |
| Methodological quality of studies assessed | + |
| Quality assessment of studies done by at least 2 reviewers | + |
| Data on measurement properties extracted by at least 2 reviewers | + |
| Quality of the instrument (measurement properties) assessed | + |
| Quality assessment of the instrument by at least 2 reviewers | + |
| Results from multiple studies on the same instrument somehow combined (e.g., best evidence synthesis or pooling) | + |
| Data synthesis was performed: | |
| Per measurement property | + |
| Only for domains (reliability, validity, responsiveness) | |
| Only for the whole instrument | |
| Recommendation provided for the best instrument: | |
| One instrument is recommended per construct | + |
| More instruments are recommended per construct | |
| No recommendation for the best instrument | |
| Results for the measurement properties reported as raw data | + |
| Number of measurement properties reported | 2 |
| Conflict of interest or funding source declared | + |
| One of the authors of the review is also the developer of one of the instruments evaluated in the review | - |

Note. + = criterion met; - = criterion not met; ± = criterion partly met



5 Discussion and conclusion

In this systematic review, the published scientific evidence on the content and structural validity of PROMs used to measure at least one aspect of QoL in people with DMD was thoroughly evaluated. The overriding theme was one of sparse evidence. Many PROMs that are being used to assess aspects of QoL in people with DMD are being utilised without the accompanying good quality evidence that supports their validity for this task. Only five of the PROMs uncovered in this review were specifically designed for use in people with neuromuscular problems (three for DMD), and only two of these have had their content and/or structural validity independently assessed in this population (with the content validity studies involving translated versions). When the evidence is available, most of it is either of a low quality, featuring insufficient detail in the published articles to make thorough and comprehensive assessments of content and structural validity as demanded by COSMIN,¹³ leading to doubtful ratings. Indeed, one of the highest quality pieces of evidence reviewed in terms of reported methodology, Landfeldt et al. (2018)⁵⁸, reported insufficient structural validity of the PedsQL 3.0 Neuromuscular module (NMM) in DMD.

The results from the review should not be viewed as surprising. Many of the PROMs identified are what could be described as “legacy” measures. They were developed at a time when the science of construct and item generation was largely overlooked. The content of instruments was largely defined by clinical or expert opinion, with little explanation of what that entailed. The reporting of such stages in publications or questionnaire manuals was not commonplace. The transparency of reporting on the early stages of PROM development has only gained traction in the last decade or so. Whilst this is a positive step for researchers, clinicians and users alike, progress can be limited by journal restrictions on word count and remit. It is however possible for such legacy measures to be appropriately validated (or have their validity assessed) in properly designed studies assessing content or structural validity in modern samples of people with DMD. The problem observed in this review is that researchers are likely using such measures as a consequence of precedent or tradition, rather than a supportive evidence base. Thus the first recommendation from this review is for more research into the content and structural validity of QoL PROMs used in DMD, and, if the PROMs are found to be insufficient on these criteria, for additional PROM development in DMD samples.

The limitations of sparse evidence notwithstanding, some PROMs performed better than others under COSMIN assessment. First, the KIDSCREEN instrument (all versions) does show some evidence of applicability given that it covers many aspects of QoL. The PROM development study for the



KIDSCREEN instrument was the only one rated as adequate, it was designed to assess QoL in children and adolescents with chronic illnesses, and the ratings for the content validity of the measure were positive (based on the available evidence in the measure’s development). However, it must also be borne in mind that there is little or no direct evidence to support the content or structural validity in DMD, specifically. The original KIDSCREEN instrument (52-item version) was designed to assess multiple aspects of QoL, namely: physical well-being; psychological well-being; moods and emotions; self-perception; autonomy; parent relation and home life; financial resources; social support and peers; school environment; and social acceptance (bullying), covering much of the CMQM framework.⁶ The conceptual framework of the instrument is thus intuitively applicable to the Duchenne community; however the measurement of impact may be limited due to the target age range of the PROM itself (8 - 18 years). While this is not uncommon (i.e. differences in measuring QoL from child to adulthood), there is some question of the applicability for the broader DMD population given the lower age target. While we would recommend this PROM, above all others tested in the review, for measuring QoL in children and adolescents with Duchenne, more research is needed to definitively support the use of KIDSCREEN (and its derivatives) within DMD.

The second-best performing PROM in this review was the LSIA, which received a satisfactory score for relevance, comprehensiveness, and comprehensibility in terms of content validity, based on the information available and reviewers’ ratings of the PROM itself. However, the development study for this paper lacked key details necessary in good PROM development, and thus was rated as doubtful. Furthermore, while the LSIA was one of few measures to feature a content validity study, it was a cross-cultural adaptation study of a Brazilian version of the measure, and the results of the formal assessment of this study were doubtful. Thus, the LSIA is possibly a good candidate for measuring QoL in DMD, but more evidence is needed. While the measure is comprehensive, it only comes in a 45-item version, which is potentially quite burdensome. Furthermore, the measure is designed for use in children and young adults only, and may not generalise to adults with DMD.

The most recent PROM developed specifically for use in children and adolescents with DMD was the MDCHILD. Although the PROM is designed to measure “health-related priorities”,⁴⁹ much of the content maps onto the CMQM framework⁶ and thus covers QoL. While the MDCHILD had many commendable strengths in PROM design, the overall rating of the PROM development, based on the COSMIN worst score counts system,¹⁷ was rated as doubtful due to lack of details reported in the development papers. For example, it was unclear if skilled interviewer(s) were used; to what degree data was coded independently; and to what degree, if at all, at least two researchers were involved in



the data analysis. This led to a low quality of evidence. Further, because the target population of interest was not clearly defined (i.e. age ranges were not specified), despite performing well in other areas, the PROM received an inconsistent rating for relevance. These results speak to the potential harshness of a worst score counts system advocated by COSMIN, which we discuss further below. Further, because the PROM is new, there is a lack of published content validity studies that may improve the quality of evidence for the MDCHILD going forward, such as that contained in Chapter 7 of a non-peer-reviewed thesis,⁹² not eligible for inclusion in the current review.

The PedsQL and associated modules were the most commonly used out of all the PROMs identified within the review. It should be noted that the development studies of the PedsQL were rated as doubtful. There was little evidence to support the content validity of the neuromuscular module of the PedsQL 3.0 (NMM). Furthermore, the psychometric properties of the NMM were not well supported by Landfeldt et al. (2018)⁵⁸. The inclusion of PedsQL within clinical practice, cohort studies or pragmatic trials in DMD thus appears to be based upon precedent and common use, rather than published empirical evidence of suitability, based on content and structural validity. A notable advantage of the PedsQL (and its derivatives) is the young child (via proxy report), child (self and proxy report), young adult forms (self-report), and adult forms, which have now been developed. A further consideration is that the PedsQL scales are designed to be used in parallel (e.g. the generic core scales with the NMM or DMD modules), but were assessed individually under COSMIN guidance. Thus comprehensiveness may be improved by using these scales together. Nevertheless, in the absence of further evidence, it is difficult to recommend the routine use of the PedsQL to assess QoL in people with DMD on content and structural grounds. Instead, the findings of this review support the need for further PROM development, which is able to accurately assess the impact of DMD upon people's QoL.

The search identified some PROM instruments that we were unable to obtain. Access to the PROM and/or associated development papers was limited due to licensing requirements, and therefore it was not possible to include these instruments within the review. It is unlikely that these instruments are commonly used within research and/or clinical practice due to the difficulties around access. Their suitability for the DMD population cannot formally be determined; however, their use is likely to be limited by a lack of accessibility derived from license restrictions, reflected in the few citations in which they appeared.

Another consideration relevant to the selection of QoL PROMs for use in the DMD population is whether the PROM has accompanying utility weights to make it preference-based and amenable to



use in economic valuation. Most of the PROMs assessed in this review, except the EQ-5D-3L, HUI, and SF-36 (via the SF-6D)¹¹¹ were not preference-based measures. Further, these preference-based measures did not perform well on assessments of content validity. However, alternative strategies are possible in order to obtain utility weights for some of the alternative PROMs discussed in this review, including our recommended PROM (KIDSCREEN) and the most widely-used PROM (PedsQL). These include mapping algorithms to a preference-based measure,¹¹² enabling their use in cost-effective analysis. Nevertheless, these kinds of solutions are to a degree imperfect, and there is clearly room for the development of a new preference-based measure of QoL in people with DMD.

This review adopted guidance developed by the COSMIN initiative, and has adhered to their recommended methods in identification of evidence, data extraction, data assessment and data synthesis. Whilst the appropriateness of these robust methods cannot be questioned, this has resulted in relatively low ratings of the PROMs included within the review. It is important to recognise that this does not suggest categorically that the instruments used within published and/or current studies are not appropriate or fit for purpose; content and structural validity only form one component of PROM suitability within a population. Furthermore, as stated, many of the instruments were developed at a time when instrument development methods and procedures were not reported – that is not to say the development of the instruments is flawed, just that an assessment of them cannot be made. The COSMIN appraisal tools assume a worst score counts system for the rating of the methodological quality of studies.¹⁷ This means that, in theory, a study could be rated as very good or adequate on all but one criteria, on which it is rated as doubtful or inadequate, and the overall score is thus reduced to the latter lower-quality rating. Sometimes this can be because key details, such as whether skilled interviewers were used, are not reported. We thus think that there is room for constructive debate on the usefulness of retaining a worst score counts system, as opposed to an alternative procedure that better reflects the variance or range of ratings across the COSMIN criteria. An example of this could be to use numerical scores to represent ratings for each of the COSMIN criteria and to calculate a (weighted) total, within which ranges or bands of scores reflect sequential improvements in methodological quality, from inadequate to very good.

Given that DMD is a rare condition, the development and validation of PROMs that measure the impact of the condition on QoL is challenging. The number of participants included within various phases of PROM development and validation will be lower than that of a condition such as diabetes, asthma or eczema. Some of the studies identified within the review included participants with other similar conditions. As part of the methods employed within the review, the inclusion criteria stated at



least 75% of the sample were to include men/boys with DMD. It is possible to expand the scope of the search to include a more “relaxed” criterion; however, it is not known how appropriate this would be. It can be postulated that other neuromuscular disorders could imply similar impacts upon QoL, however this has not been explored within the context of this review.

The rarity of DMD also limits assessment of cross-cultural applicability of items, concepts and indeed the PROMs. Very little (or no) data was identified that explored this issue. Few studies reported on participant ethnicity, nor discussed any potential issues that may differ between different ethnic groups. Some instruments were identified within the search that were not included within the review as no English version of the questionnaire was available or had been developed. Such instruments may be applicable to the DMD population, however this has yet to be determined and can only be assessed following a robust translational study.

The focus of this review was to report on the content and structural validity of PROM instruments that have been used to quantify the impact of DMD on individual’s QoL. However, content and structural validity only address some aspects of PROM suitability, and further work could be undertaken to formally appraise the instruments described. Other measurement properties, such as psychometric performance, could be considered. Furthermore, the inclusion of subsidiary samples such as other neuromuscular disorders, may be of interest. One of the inclusion criteria of this review was a 75% DMD sample (as recommended within the COSMIN guidance), however given the rarity of DMD it may be appropriate to relax this approach to include a wider sample of neuromuscular disorders.

This review is not without its limitations. While the methodological approach of the review is robust and follows the recommendations of COSMIN and that of other published reviews, it must be acknowledged that the rating criteria of the PROMs identified can be viewed as harsh. The COSMIN approach encourages researchers and reviewers to critically appraise evidence of PROM development – however the presence of evidence within published literature is sparse. That is not to say that the development phases did not occur, merely that they are not reported and/or not reported in sufficient detail as required by COSMIN assessment. To critique a PROM’s applicability using this criterion could be perceived as being unduly critical; more recent PROMs tend to report the early stages of instrument development, and we are assessing all PROMs by modern standards. Similarly, the descriptions of PROMs themselves are often lacking. Basic information such as number of items, recall period, domain structure and scoring procedure were noted to be sporadically reported, although better in recent literature. The COSMIN-recommended reviewer rating of the identified PROMs for suitability for DMD (as reported in Table 7) has a large subjective component. Whilst this was completed as per the



COSMIN guidelines (with two reviewers and discrepancies reconciled following discussion), some of the ratings are at risk of bias based on the team of raters. For example, it is not known whether similar ratings of suitability would be achieved if reviewed by an individual with DMD, a family member or carer of a person with DMD, or a clinician. This is further exacerbated when we consider what QoL is – for the purpose of this review it was a multidimensional construct, PROMs that measure a subset of interest (such as depression) may be appropriate to include within studies as part of a host/suite of measures.



References

1. Mendell JR, Schilling C, Leslie ND, et al. Evidence-based path to newborn screening for Duchenne muscular dystrophy. *Ann Neurol*. 2012;71:304-313.
2. Strehle EM, Straub V. Recent advances in the management of Duchenne muscular dystrophy. *Arch Dis Child* 2015;100:1173-1177.
3. Kieny PI, Chollet S, Delalande P, et al. Evolution of life expectancy of patients with Duchenne Muscular Dystrophy at AFM Yolaine de Kepper centre between 1981 and 2011. *Ann Phys Rehabil Med* 2013;56:443-454.
4. Landfeldt E, Lindgren P, Bell CF, et al. Compliance to care guidelines for Duchenne muscular dystrophy. *J Neuromuscul Dis* 2015;2:63-72.
5. NICE (National Institute for Health and Care Excellence). Guide to the methods of technology appraisal. London, UK: NICE, 2013.
6. Bann CM, Abresch RT, Biesecker B, et al. Measuring quality of life in muscular dystrophy. *Neurology* 2015;84:1034-1042.
7. Wei Y, Speechley K, Campbell C. Health-related quality of life in children with Duchenne muscular dystrophy: A review. *J Neuromuscul Dis* 2015;2:313-324.
8. Prinsen CAC, Mokkink LB, Bouter LM, et al. COSMIN guideline for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018;27:1147-1157.
9. Terwee CB, Prinsen CA, Chiarotto A, et al. COSMIN methodology for assessing the content validity of PROMs - user manual. 2018; <http://www.cosmin.nl>. Accessed 12 July 2018.
10. Mokkink LB, Terwee CB, Patrick DL, et al. The COSMIN study reached international consensus on taxonomy, terminology, and definitions of measurement properties for health-related patient-reported outcomes. *J Clin Epidemiol* 2010;63:737-745.
11. Chiarotto A, Ostelo RW, Boers M, et al. A systematic review highlights the need to investigate the content validity of patient-reported outcome measures for physical functioning in patients with low back pain. *J Clin Epidemiol* 2018;95:73-93.
12. Prinsen CA, Vohra S, Rose MR, et al. How to select outcome measurement instruments for outcomes included in a "Core Outcome Set" – a practical guideline. *Trials* 2016;17:449.
13. Mokkink LB, Prinsen CA, Patrick DL, et al. COSMIN methodology for systematic reviews of Patient-Reported Outcome Measures (PROMs) - user manual. 2018; <http://www.cosmin.nl>. Accessed 12 July 2018.
14. Terwee CB, Prinsen CAC, Ricci Garotti MG, et al. The quality of systematic reviews of health-related outcome measurement instruments. *Qual Life Res* 2016;25:767-779.
15. Papaioannou D, Sutton A, Carroll C, et al. Literature searching for social science systematic reviews: consideration of a range of search techniques. *Health Info Libr J* 2010;27:114-122.
16. Mokkink LB, de Vet HCW, Prinsen CAC, et al. COSMIN Risk of Bias checklist for systematic reviews of Patient-Reported Outcome Measures. *Qual Life Res* 2018;27:1171-1179.
17. Terwee CB, Mokkink LB, Knol DL, et al. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Qual Life Res* 2012;21:651-657.
18. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924.
19. Manser RL, Irving LB, Byrnes G, et al. Screening for lung cancer: a systematic review and meta-analysis of controlled trials *Thorax* 2003;58:784-789.
20. Mateen FJ, Oh J, Tergas AI, et al. Titles versus titles and abstracts for initial screening of articles for systematic reviews. *Clin Epidemiol* 2013;5:89-95.
21. Kohler M, Clarenbach CF, Böni L, et al. Quality of life, physical disability, and respiratory impairment in Duchenne muscular dystrophy. *Am J Respir Crit Care Med* 2005;172:1032-1060.
22. Simonds AK, Muntoni F, Heather S, et al. Impact of nasal ventilation on survival in hypercapnic Duchenne muscular dystrophy. *Thorax* 1998;53:949-952.



23. Catalano A, Vita GL, Russo M, et al. Effects of teriparatide on bone mineral density and quality of life in Duchenne muscular dystrophy related osteoporosis: a case report. *Osteoporos Int* 2016;27:3655-3659.
24. Lue YJ, Chen SS, Lu YM. Quality of life of patients with Duchenne muscular dystrophy: from adolescence to young men. *Disabil Rehabil* 2017;39:1408-1413.
25. Elsenbruch S, Schmid J, Lutz S, et al. Self-reported quality of life and depressive symptoms in children, adolescents, and adults with Duchenne muscular dystrophy: a cross-sectional survey study. *Neuropediatrics* 2013;44:257-264.
26. McDonald CM, Henricson EK, Abresch RT, et al. The Cooperative International Neuromuscular Research Group Duchenne Natural History Study-A Longitudinal Investigation In The Era Of Glucocorticoid Therapy: Design Of Protocol And The Methods Used. *Muscle and Nerve* 2013;48:32-54.
27. Pangalila RF, van den Bos GA, Bartels B, et al. Prevalence of fatigue, pain, and affective disorders in adults with duchenne muscular dystrophy and their associations with quality of life. *Arch Phys Med Rehabil* 2015;96:1242-1247.
28. Pangalila RF, van den Bos GA, Bartels B, et al. Quality of life of adult men with Duchenne muscular dystrophy in the Netherlands: implications for care. *J Rehabil Med* 2015;47:161-166.
29. Leung DG, Herzka DA, Reid Thompson W, et al. Sildenafil does not improve cardiomyopathy in Duchenne/Becker muscular dystrophy. *Ann Neurol* 2014;76:541-549.
30. de Moura MCDS, Wutzki HC, Voos MC, et al. Is functional dependence of Duchenne muscular dystrophy patients determinant of the quality of life and burden of their caregivers? *Arq Neuropsiquiatr* 2015;73:52-57.
31. Zebracki K, Drotar D. Pain and activity limitations in children with Duchenne or Becker muscular dystrophy. *Dev Med Child Neurol* 2008;50:546-552.
32. Bray P, Bundy AC, Ryan MM, et al. Health status of boys with Duchenne muscular dystrophy: a parent's perspective. *J Paediatr Child Health* 2011;47:557-562.
33. Baiardini I, Minetti C, Bonifacino S, et al. Quality of life in Duchenne muscular dystrophy: the subjective impact on children and parents. *J Child Neurol* 2011;26:707-713.
34. Bendixen RM, Senesac C, Lott DJ, et al. Participation and quality of life in children with Duchenne muscular dystrophy using the International Classification of Functioning, Disability, and Health. *Health Qual Life Outcomes* 2012;10:43.
35. Bendixen RM, Lott DJ, Senesac C, et al. Participation in daily life activities and its relationship to strength and functional measures in boys with Duchenne muscular dystrophy. *Disabil Rehabil* 2014;36:1918-1923.
36. Beenakker EA, Fock JM, Van Tol MJ, et al. Intermittent prednisone therapy in Duchenne muscular dystrophy: a randomized controlled trial. *Arch Neurol* 2005;62:128-132.
37. Cavazza M, Kodra Y, Armeni P, et al. Social/economic costs and health-related quality of life in patients with Duchenne muscular dystrophy in Europe. *Eur J Health Econ* 2016;17 Suppl 1:19-29.
38. Pangalila RF, van den Bos GA, Stam HJ, et al. Subjective caregiver burden of parents of adults with Duchenne muscular dystrophy. *Disabil Rehabil* 2012;34:988-996.
39. Graham CD, Rose MR. What explains high life satisfaction in men living with Duchenne muscular dystrophy? A preliminary study to inform psychological intervention. *Muscle Nerve* 2017;56:163-166.
40. Landfeldt E, Lindgren P, Bell CF, et al. The burden of Duchenne muscular dystrophy: an international, cross-sectional study. *Neurology* 2014;83:529-536.
41. Landfeldt E, Lindgren P, Bell CF, et al. Health-related quality of life in patients with Duchenne muscular dystrophy: a multinational, cross-sectional study. *Dev Med Child Neurol* 2016;58:508-515.
42. Otto C, Steffensen BF, Hojberg AL, et al. Predictors of Health-Related Quality of Life in boys with Duchenne muscular dystrophy from six European countries. *J Neurol* 2017;264:709-723.



43. Zamani G, Heidari M, Azizi Malamiri R, et al. The quality of life in boys with Duchenne muscular dystrophy. *Neuromuscul Disord* 2016;26:423-427.
44. Houwen-van Opstal S, Jansen M, van Alfen N, et al. Health-related quality of life and its relation to disease severity in boys with Duchenne muscular dystrophy: Satisfied boys, worrying parents- A case-control study. *J Child Neurol* 2014;29:1486-1495.
45. Reid DT, Renwick RM. Preliminary validation of a new instrument to measure life satisfaction in adolescents with neuromuscular disorders. *Int J Rehabil Res* 1994;17:184-188.
46. Simon VA, Dutra Resende MB, Simon MAVP, et al. Duchenne muscular dystrophy: Quality of life among 95 patients evaluated using the life satisfaction index for adolescents. *Arq Neuropsiquiatr* 2011;69:19-22.
47. Simon VA, Zanoteli E, Simon MAVP, et al. Translation and validation of the life satisfaction index for adolescents scale with neuromuscular disorders: LSI-A Brazil. *Arq Neuropsiquiatr* 2017;75:553-562.
48. Reid DT, Renwick RM. Relating familial stress to the psychosocial adjustment of adolescents with Duchenne muscular dystrophy. *Int J Rehabil Res* 2001;24:83-93.
49. Propp R, et al. Development and content validation of the Muscular Dystrophy Child Health Index of Life with Disabilities questionnaire for children with Duchenne muscular dystrophy. *Dev Med Child Neurol* 2019;61:75-81.
50. McDonald CM, McDonald DA, Bagley A, et al. Relationship between clinical outcome measures and parent proxy reports of health-related quality of life in ambulatory children with Duchenne muscular dystrophy. *J Child Neurol* 2010;25:1130-1144.
51. McDonald CM, Henricson EK, Abresch RT, et al. Long-term effects of glucocorticoids on function, quality of life, and survival in patients with Duchenne muscular dystrophy: a prospective cohort study. *Lancet* 2018;391:451-461.
52. Ruiz-Cortes X, Ortiz-Corredor F, Mendoza-Pulido C. Reliability of home-based, motor function measure in hereditary neuromuscular diseases. *J Int Med Res* 2017;45:261-271.
53. Henricson E, Abresch R, Han JJ, et al. The 6-Minute Walk Test and Person-Reported Outcomes in Boys with Duchenne Muscular Dystrophy and Typically Developing Controls: Longitudinal Comparisons and Clinically-Meaningful Changes Over One Year. *PLoS Curr* 2013;5:08.
54. Uzark K, King E, Cripe L, et al. Health-related quality of life in children and adolescents with Duchenne muscular dystrophy. *Pediatrics* 2012;130:e1559-1566.
55. Wei Y, Speechley KN, Zou G, et al. Factors Associated With Health-Related Quality of Life in Children With Duchenne Muscular Dystrophy. *J Child Neurol* 2016;31:879-886.
56. Messina S, Vita GL, Sframeli M, et al. Health-related quality of life and functional changes in DMD: A 12-month longitudinal cohort study. *Neuromuscul Disord* 2016;26:189-196.
57. Davis SE, Hynan LS, Limbers CA, et al. The PedsQL™ in pediatric patients with Duchenne muscular dystrophy: Feasibility, reliability, and validity of the pediatric quality of life inventory neuromuscular module and generic core scales. *J Clin Neuromuscul Dis* 2010;11:97-109.
58. Landfeldt E, Mayhew A, Straub V, et al. Psychometric analysis of the pediatric quality of life inventory 3.0 neuromuscular module administered to patients with Duchenne muscular dystrophy: A rasch analysis. *Muscle Nerve* 2018;21:21.
59. Hu J, Jiang L, Hong S, et al. Reliability and validity of the Chinese version of the pediatric quality of life inventory™ (PedsQL™) 3.0 neuromuscular module in children with Duchenne muscular dystrophy. *Health and Quality of Life Outcomes* 2013;11:no pagination.
60. Hu J, Kong M, Ye Y, et al. Serum miR-206 and other muscle-specific microRNAs as non-invasive biomarkers for Duchenne muscular dystrophy. *J Neurochem* 2014;129:877-883.
61. Hu J, Ye Y, Kong M, et al. Daily prednisone treatment in Duchenne muscular dystrophy in southwest China. *Muscle Nerve* 2015;52:1001-1007.
62. Bray P, Bundy AC, Ryan MM, et al. Health-related quality of life in boys with Duchenne muscular dystrophy: agreement between parents and their sons. *J Child Neurol* 2010;25:1188-1194.



63. Bray P, Bundy AC, Ryan MM, et al. Can in-the-moment diary methods measure health-related quality of life in Duchenne muscular dystrophy? *Qual Life Res* 2017;26:1145-1152.
64. Lim Y, Velozo C, Bendixen RM. The level of agreement between child self-reports and parent proxy-reports of health-related quality of life in boys with Duchenne muscular dystrophy. *Qual Life Res* 2014;23:1945-1952.
65. McDonald CM, Henricson EK, Abresch RT, et al. The 6-minute walk test and other clinical endpoints in duchenne muscular dystrophy: Reliability, concurrent validity, and minimal clinically important differences from a multicenter study. *Muscle and Nerve* 2013;48:357-368.
66. Ozyurt G, Bayram E, Karaoglu P, et al. Quality of life and sleep in children diagnosed with Duchenne Muscular Dystrophy and their mothers' level of anxiety: A case-control study. *Dusunen Adam: Journal of Psychiatry and Neurological Sciences* 2015;28:362-368.
67. Schreiber-Katz O, Klug C, Thiele S, et al. Comparative cost of illness analysis and assessment of health care burden of Duchenne and Becker muscular dystrophies in Germany. *Orphanet J Rare Dis* 2014;9:210.
68. Wei Y, Speechley KN, Zou G, et al. The relationship between quality of life and health-related quality of life in young males with Duchenne muscular dystrophy. *Dev Med Child Neurol* 2017;59:1152-1157.
69. Teoh LJ, et al. Health care utilization and costs for children and adults with Duchenne muscular dystrophy. *Muscle Nerve* 2016;53:877-884.
70. Sienko S, Buckon C, Fowler E, et al. Prednisone and Deflazacort in Duchenne Muscular Dystrophy: Do They Play a Different Role in Child Behavior and Perceived Quality of Life? *PLoS Curr* 2016;8:17.
71. Colombo P, Nobile M, Tesei A, et al. Assessing mental health in boys with Duchenne muscular dystrophy: Emotional, behavioural and neurodevelopmental profile in an Italian clinical sample. *Europ J Paediatr Neurol* 2017;21:639-647.
72. Garralda ME, McConachie H, Le Couteur A, et al. Emotional impact of genetic trials in progressive paediatric disorders: a dose-ranging exon-skipping trial in Duchenne muscular dystrophy. *Child: Care, Health & Development* 2013;39:449-455.
73. Darke J, et al. Survey of behaviour problems in children with neuromuscular disorders. *Eur J Paediatr Neurol* 2006;10:129-134.
74. Orcesi S, Ariaudo G, Mercuri E, et al. A new self-report quality of life questionnaire for children with neuromuscular disorders: presentation of the instrument, rationale for its development, and some preliminary results. *J Child Neurol* 2014;29:167-181.
75. Grootenhuis MA, et al. Living with muscular dystrophy: health related quality of life consequences for children and adults. *Health Qual Life Outcomes* 2007;5:31.
76. Beck AT, Ward CH, Mendelson M, et al. An inventory for measuring depression. *Archives of general psychiatry*. 1961;4:561-571.
77. Palermo TM, Witherspoon D, Valenzuela D, et al. Development and validation of the Child Activity Limitations Interview: a measure of pain-related functional impairment in school-age children and adolescents. *Pain* 2004;109:461-470.
78. Petersen C, Schmidt S, Power M, et al. (2005). Developing and pilot testing a chronic generic quality of life questionnaire for children with chronic conditions. *Qual Life Res* 73:1065-1075.
79. Ravens-Sieberer U, Schmidt S, Gosch A, et al. Measuring subjective health in children and adolescents: results of the European KIDSCREEN/DISABKIDS Project. *Psychosoc Med* 2007;4:Doc08.
80. EuroQol Group. EuroQol--a new facility for the measurement of health-related quality of life. *Health policy (Amsterdam, Netherlands)* 1990;16:199.
81. Brooks R. EuroQol: the current state of play. *Health Policy* 1996;37:53-72.
82. Krupp LB, LaRocca NG, Muir-Nash J, et al. The fatigue severity scale. Application to patients with multiple sclerosis and systemic lupus erythematosus. *Arch Neurol*. 1989;46:1121-1123.



83. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing Generalized Anxiety Disorder: The GAD-7. *Arch Intern Med.* 2006;166:1092–1097.
84. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-370.
85. Feeny D, Furlong W, Boyle M, et al. Multi-attribute health status classification systems. *Health Utilities Index. Pharmacoeconomics* 1995;7:490-502.
86. Torrance GW, Feeny DH, Furlong WJ, et al. Multiattribute utility function for a comprehensive health status classification system. *Health Utilities Index Mark 2. Med Care* 1996;34:702-722.
87. Vincent KA, Carr AJ, Walburn J, et al. Construction and validation of a quality of life questionnaire for neuromuscular disease (INQoL). *Neurology* 2007;68:1051-1057.
88. Ravens-Sieberer U, Gosch A, Abel T, et al. Quality of life in children and adolescents: a European public health perspective. *Soz Praventivmed* 2001;46:294-302.
89. Ravens-Sieberer U, Gosch A, Rajmil L, et al. KIDSCREEN-52 quality-of-life measure for children and adolescents. *Expert Rev Pharmacoecon Outcomes Res* 2005;5:353-364.
90. Detmar SB, Bruil J, Ravens-Sieberer U, et al. The use of focus groups in the development of the KIDSCREEN HRQL questionnaire. *Qual Life Res* 2006;15:1345-1353.
91. Ravens-Sieberer U, Europe KG. The Kidscreen questionnaires: quality of life questionnaires for children and adolescents; handbook: Pabst Science Publ; 2006.
92. Propp, R. Development and Psychometric Evaluation of the Muscular Dystrophy Child Health Index of Life with Disabilities (MDCHILD) Questionnaire in Children with Duchenne Muscular Dystrophy. 2017. <http://hdl.handle.net/1807/92986>
93. Varni JW, Burwinkle TM, Katz ER, et al. The PedsQL™ in pediatric cancer: reliability and validity of the pediatric quality of life inventory™ generic core scales, multidimensional fatigue scale, and cancer module. *Cancer* 2002;94:2090-2106.
94. Iannaccone ST, Hynan LS, Morton A, et al. The PedsQL™ in pediatric patients with spinal muscular atrophy: Feasibility, reliability, and validity of the pediatric quality of life inventory™ generic core scales and neuromuscular module. *Neuromuscular Disorders* 2009;19:805-812.
95. Varni JW, Seid M, Rode CA. The PedsQL™: measurement model for the pediatric quality of life inventory. *Medical care* 1999:126-139.
96. Spitzer RL, Kroenke K, Williams JBW, et al. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *JAMA.* 1999;282:1737-1744.
97. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16:606-613.
98. Daltroy LH, Liang MH, Fossel AH, et al. The POSNA pediatric musculoskeletal functional health questionnaire: report on reliability, validity, and sensitivity to change. Pediatric Outcomes Instrument Development Group. *Pediatric Orthopaedic Society of North America. J Pediatr Orthop* 1998;18:561-571.
99. Buysse DJ, Reynolds CF, 3rd, Monk TH, et al. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193-213.
100. Goodman R. The Strengths and Difficulties Questionnaire: a research note. *J Child Psychol Psychiatry* 1997;38:581-586.
101. Ware Jr J, Sherbourne C. The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Medical Care* 1992;30:473-483.
102. Hays RD, Sherbourne CD, Mazel RM. The RAND 36-Item Health Survey 1.0. *Health Econ* 1993;2:217-227.
103. Jenkinson C, Stewart-Brown S, Petersen S, et al. Assessment of the SF-36 version 2 in the United Kingdom. *J Epidemiol Community Health* 1999;53:46-50.
104. Ware Jr JE. SF-36 health survey update. *Spine* 2000;25:3130-3139.
105. Diener E, Emmons RA, Larsen RJ, et al. The Satisfaction With Life Scale. *J Pers Assess* 1985;49:71-75.



106. WHOQoL Group. Development of the WHOQOL: Rationale and current status. *International Journal of Mental Health* 1994;23:24-56.
107. WHOQoL Group. The World Health Organization quality of life assessment (WHOQOL): position paper from the World Health Organization. *Social Science & Medicine*. 1995;41:1403-1409.
108. Skevington SM, MacArthur P, Somerset M. Developing items for the WHOQOL: An investigation of contemporary beliefs about quality of life related to health in Britain. *Brit J Health Psych*. 1997;2:55-72.
109. WHOQoL Group. Development of the World Health Organization WHOQOL-BREF quality of life assessment. *Psychological medicine* 1998;28:551-558.
110. Chiarotto A, Terwee CB, Kamper SJ, et al. Evidence on the measurement properties of health-related quality of life instruments is largely missing in patients with low back pain: A systematic review. *J Clin Epidemiol* 2018;102:23-37.
111. Brazier J, Roberts J, Deverill M. The estimation of a preference-based measure of health from the SF-36. *J Health Econ* 2002;21:271-292.
112. Wailoo AJ, Hernandez-Alava M, Manca A, et al. Mapping to estimate health-state utility from non-preference-based outcome measures: An ISPOR good practices for outcomes research task force report. *Value Health* 2017;20:18-27.



Appendix A: Full search strategies

Embase Stage One

1. Muscular Dystrophy/
2. duchenne*.mp.
3. 1 or 2
4. (HR-PRO or HRPRO or HRQL or HRQoL or QL or QoL).ti,ab. or quality of life.mp. or (health index* or health indices or health profile*).ti,ab. or health status.mp. or ((patient or self or child or parent or carer or proxy) adj (appraisal* or appraised or report or reported or reporting or rated or rating* or based or assessed or assessment*)).ti,ab. or ((disability or function or functional or functions or subjective or utility or utilities or wellbeing or well being) adj2 (index or indices or instrument or instruments or measure or measures or questionnaire* or profile or profiles or scale or scales or score or scores or status or survey or surveys)).ti,ab.
5. 'Pediatric Quality of Life Inventory'.mp.
6. PedsQL.mp.
7. (SF-36 or EQ-5D*).mp.
8. "World Health Organization Quality of Life".mp.
9. (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
10. "Childrens Assessment of Participation and Enjoyment".mp.
11. (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*).mp.
12. (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease").mp.
13. (QoL-NMD DISABKIDS or "Depressionsinventar fur kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM).mp.
14. (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).mp.
15. (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).mp.
16. 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15
17. 4 or 16
18. 3 and 17

Embase Stage Two

1. Duchenne muscular dystrophy/
2. duchenne*.mp.
3. 1 or 2
4. Autoquestionnaire Qualite de vie Enfant Image.mp.



5. (Behavior Assessment System for Children or BASC or Parent Form 50 or PF50 or DUX-25 or EuroQoL 5-domain or Functional Independence Measure* or FIM).mp.
6. (WeeFIM or Life Satisfaction Index or LSI or LSIA or Neurological Disorders Quality of Life Questionnaire or NeuroQOL or pediatric NeuroQOL or Offer Self-Image Questionnaire for Adolescents or OSIQ).mp.
7. (Pediatric Outcomes Data Collection Instrument or PODCI or Neuromuscular module or DMD module or Multidimensional Fatigue Scale or Generic Short-Form or SF15).mp.
8. (SDQ or 'Strips of Life with Emoticons Questionnaire' or SOLE or 'World Health Organisation Quality of Life Scale-Brief Version' or WHOQOL-BREF).mp.
9. ('Strength and Difficulties Questionnaire').mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word, candidate term word]
10. "Individualized Neuromuscular Quality of Life Questionnaire".mp.
11. (INQOL or Child Activity Limitations Interview or CALI or "Satisfaction with Life Scale" or SWLS).mp.
12. 'Pediatric Quality of Life Inventory'.mp.
13. ("Pediatric Orthopedic Society of North America Pediatric Musculoskeletal Functional Health Questionnaire" or POSNA or Pittsburgh Sleep Quality Index).mp.
14. PedsQL.mpp.
15. (SF-36 or EQ-5D*).mp.
16. "World Health Organization Quality of Life".mp.
17. (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word, candidate term word]
18. "Childrens Assessment of Participation and Enjoyment".mp.
19. (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*).mp.
20. (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease").mp.
21. (QoL-NMD DISABKIDS or "Depressionsinventar fur kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM).mp.
22. (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).mp.
23. (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).mp.
24. 12 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23

PsychINFO Stage One

1. Muscular Dystrophy/
2. duchenne*.mp.
3. 1 or 2



4. (HR-PRO or HRPRO or HRQL or HRQoL or QL or QoL).ti,ab. or quality of life.mp. or (health index* or health indices or health profile*).ti,ab. or health status.mp. or ((patient or self or child or parent or carer or proxy) adj (appraisal* or appraised or report or reported or reporting or rated or rating* or based or assessed or assessment*)).ti,ab. or ((disability or function or functional or functions or subjective or utility or utilities or wellbeing or well being) adj2 (index or indices or instrument or instruments or measure or measures or questionnaire* or profile or profiles or scale or scales or score or scores or status or survey or surveys)).ti,ab.
5. 'Pediatric Quality of Life Inventory'.mp.
6. PedsQL.mp.
7. (SF-36 or EQ-5D*).mp.
8. "World Health Organization Quality of Life".mp.
9. (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
10. "Childrens Assessment of Participation and Enjoyment".mp.
11. (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*).mp.
12. (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease").mp.
13. (QoL-NMD DISABKIDS or "Depressionsinventar fur kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM).mp.
14. (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).mp.
15. (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).mp.
16. 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15
17. 4 or 16
18. 3 and 17

PsycINFO Stage Two

1. Muscular Dystrophy/
2. duchenne*.mp.
3. 1 or 2
4. Autoquestionnaire Qualite de vie Enfant Image.mp.
5. (Behavior Assessment System for Children or BASC or Parent Form 50 or PF50 or DUX-25 or EuroQoL 5-domain or Functional Independence Measure* or FIM).mp.
6. (WeeFIM or Life Satisfaction Index or LSI or LSIA or Neurological Disorders Quality of Life Questionnaire or NeuroQOL or pediatric NeuroQOL or Offer Self-Image Questionnaire for Adolescents or OSIQ).mp.
7. (Pediatric Outcomes Data Collection Instrument or PODCI or Neuromuscular module or DMD module or Multidimensional Fatigue Scale or Generic Short-Form or SF15).mp.
8. (SDQ or 'Strips of Life with Emoticons Questionnaire' or SOLE or 'World Health Organisation Quality of Life Scale-Brief Version' or WHOQOL-BREF).mp.
9. ('Strength and Difficulties Questionnaire').mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]



10. "Individualized Neuromuscular Quality of Life Questionnaire".mp.
11. (INQOL or Child Activity Limitations Interview or CALL or "Satisfaction with Life Scale" or SWLS).mp.
12. 'Pediatric Quality of Life Inventory'.mp.
13. ("Pediatric Orthopedic Society of North America Pediatric Musculoskeletal Functional Health Questionnaire" or POSNA or Pittsburgh Sleep Quality Index).mp.
14. PedsQL.mp.
15. (SF-36 or EQ-5D*).mp.
16. "World Health Organization Quality of Life".mp.
17. (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
18. "Childrens Assessment of Participation and Enjoyment".mp.
19. (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*).mp.
20. (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease").mp.
21. (QoL-NMD DISABKIDS or "Depressionsinventar fur kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM).mp.
22. (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).mp.
23. (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).mp.

Cochrane Library Stage One

Search Name: Duchenne for PP

Last Saved: 11/04/2018 09:19:13

Comment:

- | ID | Search |
|-----|---|
| #1 | MeSH descriptor: [Muscular Dystrophy, Duchenne] explode all trees |
| #2 | duchenne*:ti,ab,kw |
| #3 | #1 or #2 |
| #4 | (HR-PRO or HRPRO or HRQL or HRQoL or QL or QoL):ti,ab |
| #5 | (quality of life):ti,ab,kw |
| #6 | (health index* or health indices or health profile*):ti,ab,kw |
| #7 | (health status):ti,ab,kw |
| #8 | ((patient or self or child or parent or carer or proxy) near (appraisal* or appraised or report or reported or reporting or rated or rating* or based or assessed or assessment*)):ti,ab |
| #9 | ((disability or function or functional or functions or subjective or utility or utilities or wellbeing or well being) near/2 (index or indices or instrument or instruments or measure or measures or questionnaire* or profile or profiles or scale or scales or score or scores or status or survey or surveys)) .ti,ab |
| #10 | #4 or #5 or #6 or #7 or #8 or #9 |
| #11 | ("Pediatric Quality of Life Inventory" or PedsQL or SF-36 or EQ-5D* or "World Health Organization Quality of Life" or WHOQoL or KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI or "Children's Assessment of Participation and Enjoyment" or CAPE or "Child Health |



Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI* or Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease" or QoL-NMD DISABKIDS or "Depressionsinventar fur kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM):ti,ab,kw

#12 (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve):ti,ab,kw

#13 (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six):ti,ab,kw

#14 #11 or #12 or #13

#15 #10 or #14

#16 #3 and #15

Cochrane Library Stage Two

Search Name: Duchenne Stage Two for PP September 2018

Last Saved: 18/09/2018 15:23:27

Comment:

ID Search

#1 MeSH descriptor: [Muscular Dystrophy, Duchenne] this term only

#2 (duchenne*):ti,ab,kw

#3 #1 or #2

#4 (Autoquestionnaire Qualite de vie Enfant Image):ti,ab,kw

#5 ("Behavior Assessment System for Children"):ti,ab,kw

#6 (BASC):ti,ab,kw

#7 (Parent Form 50):ti,ab,kw

#8 (PF50 or DUX-25):ti,ab,kw

#9 ("EuroQoL 5-domain"):ti,ab,kw

#10 (Functional Independence Measure* or FIM):ti,kw,ab

#11 (WeeFIM or Life Satisfaction Index or LSI or LSIA or Neurological Disorders Quality of Life Questionnaire or NeuroQOL or pediatric NeuroQOL or Offer Self-Image Questionnaire for Adolescents or OSIQ):ti,ab,kw

#12 (Pediatric Outcomes Data Collection Instrument or PODCI or Neuromuscular module or DMD module or Multidimensional Fatigue Scale or Generic Short-Form or SF15):ti,ab,kw

#13 (SDQ or 'Strips of Life with Emoticons Questionnaire' or SOLE or 'World Health Organisation Quality of Life Scale-Brief Version' or WHOQOL-BREF):ti,ab,kw

#14 ('Strength and Difficulties Questionnaire'):ti,ab,kw

#15 ("Individualized Neuromuscular Quality of Life Questionnaire"):ti,ab,kw

#16 (INQOL or Child Activity Limitations Interview or CALL or "Satisfaction with Life Scale" or SWLS):ti,ab,kw

#17 ('Pediatric Quality of Life Inventory'):ti,ab,kw

#18 ("Pediatric Orthopedic Society of North America Pediatric Musculoskeletal Functional Health Questionnaire" or POSNA or Pittsburgh Sleep Quality Index):ti,ab,kw

#19 (PedsQL):ti,ab,kw

#20 (SF-36 or EQ-5D*):ti,ab,kw

#21 ("World Health Organization Quality of Life"):ti,ab,kw

#22 (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI):ti,ab,kw

#23 ("Childrens Assessment of Participation and Enjoyment"):ti,ab,kw

#24 (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*):ti,ab,kw



- #25 (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease"):ti,ab,kw
- #26 (QoL-NMD DISABKIDS or "Depressionsinventar fur kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM):ti,ab,kw
- #27 (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve):ti,ab,kw
- #28 (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six):ti,ab,kw
- #29 #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28
- #30 #3 and #29
- #31 "construct validity" or "content validity" or "criterion validity" or "inter rater reliability" or "interrater reliability"
- #32 #30 and #31

CINAHL Stage One

| Search ID# | Search Terms | Search Options | Last Run Via | Results |
|------------|-------------------------------------|-------------------------------|--|---------|
| S1 | (MH "Muscular Dystrophy, Duchenne") | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S2 | duchenne* | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S3 | S1 OR S2 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S4 | HR-PRO | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - | Display |



| | | | | |
|----|--|-------------------------------|---|---------|
| | | | Advanced Search Database - CINAHL with Full Text | |
| S5 | ((disability or function or functional or functions or subjective or utility or utilities or wellbeing or well being) N2 (index or indices or instrument or instruments or measure or measures or questionnaire* or profile or profiles or scale or scales or score or scores or status or survey or surveys)) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S6 | HRPRO or HRQL or HRQoL | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S7 | quality of life or health index* or health indices or health profile* or health status | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S8 | TI QL | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S9 | TI QoL | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |



| | | | | |
|-----|---|-------------------------------|---|---------|
| S10 | ((patient or self or child or parent or carer or proxy) and (appraisal* or appraised or report or reported or reporting or rated or rating* or based or assessed or assessment*)) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S11 | S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S12 | (Pediatric Quality of Life Inventory or PedsQL or SF-36 or EQ-5D* or World Health Organization Quality of Life or WHOQoL or KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI or Childrens Assessment of Participation and Enjoyment or CAPE or Child Health Questionnaire or CHQ* or Health Utilities Index Questionnaire or HUI* or Fatigue Severity Scale or FSS or Hospital Anxiety and Depression Scale or HADS or COPE Inventory or Quality of Life in Neuromuscular Disease or QoL-NMD DISABKIDS or Depressionsinventar fur kinder und Jugendliche or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or Life Satisfaction Index for Adolescents or LSI-A or Quality of Life | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |



| | | | | |
|-----|---|-------------------------------|---|---------|
| | Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM) | | | |
| S13 | (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S14 | (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S15 | S12 OR S13 OR S14 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S16 | S11 OR S15 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | Display |
| S17 | S3 AND S16 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 754 |

CINAHL Stage Two



| Search ID# | Search Terms | Search Options | Last Run Via | Results |
|------------|--|-------------------------------|--|---------|
| S1 | (MH "Muscular Dystrophy, Duchenne") | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 1,330 |
| S2 | duchenne* | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 1,800 |
| S3 | S1 OR S2 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 1,800 |
| S4 | Autoquestionnaire Qualite de vie Enfant Image | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 3 |
| S5 | (Behavior Assessment System for Children or BASC or Parent Form 50 or PF50 or DUX-25 or EuroQoL 5-domain or Functional Independence Measure* or FIM) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 3,288 |
| S6 | (WeeFIM or Life Satisfaction Index or LSI or LSIA or Neurological Disorders Quality of Life Questionnaire or NeuroQOL or pediatric | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search | 559 |



| | | | | |
|-----|--|-------------------------------|---|-------|
| | NeuroQOL or Offer Self-Image Questionnaire for Adolescents or OSIQ) | | Database - CINAHL with Full Text | |
| S7 | (Pediatric Outcomes Data Collection Instrument or PODCI or Neuromuscular module or DMD module or Multidimensional Fatigue Scale or Generic Short-Form or SF15) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 197 |
| S8 | (SDQ or 'Strips of Life with Emoticons Questionnaire' or SOLE or 'World Health Organisation Quality of Life Scale-Brief Version' or WHOQOL-BREF) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 6,403 |
| S9 | ('Strength and Difficulties Questionnaire') | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 948 |
| S10 | "Individualized Neuromuscular Quality of Life Questionnaire" | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 3 |
| S11 | (INQOL or Child Activity Limitations Interview or CALI or "Satisfaction with Life Scale" or SWLS) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 837 |
| S12 | 'Pediatric Quality of Life Inventory' | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - | 521 |



| | | | | |
|-----|--|-------------------------------|---|--------|
| | | | Advanced Search Database - CINAHL with Full Text | |
| S13 | ("Pediatric Orthopedic Society of North America Pediatric Musculoskeletal Functional Health Questionnaire" or POSNA or Pittsburgh Sleep Quality Index) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 1,548 |
| S14 | PedsQL | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 651 |
| S15 | (SF-36 or EQ-5D*) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 16,749 |
| S16 | "World Health Organization Quality of Life" | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 623 |
| S17 | (KIDSCREEN or Pittsburgh Sleep Quality questionnaire or PSQI) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 1,040 |
| S18 | "Childrens Assessment of Participation and Enjoyment" | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - | 0 |



| | | | | |
|-----|--|-------------------------------|---|---------|
| | | | Advanced Search Database - CINAHL with Full Text | |
| S19 | (CAPE or "Child Health Questionnaire" or CHQ* or Health Utilities Index Questionnaire or HUI*) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 3,751 |
| S20 | (Fatigue Severity Scale or FSS or "Hospital Anxiety and Depression Scale" or HADS or COPE Inventory or "Quality of Life in Neuromuscular Disease") | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 530,689 |
| S21 | (QoL-NMD DISABKIDS or "Depressionsinventar für kinder und Jugendliche" or DIKJ or Beck Depression Inventory or BDI or CARE-NMD or State-Trait Anxiety Inventory or STAI or "Life Satisfaction Index for Adolescents" or LSI-A or Quality of Life Evaluation Scale or AUQUEI or Activity Limitations Questionnaire or ACTIVLIM) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 9,541 |
| S22 | (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve) | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 524 |
| S23 | (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - | 2,253 |



| | | | | |
|-----|--|-------------------------------|---|---------|
| | thirty six or short form thirtysix or short form thirty six) | | Advanced Search Database - CINAHL with Full Text | |
| S24 | S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14 OR S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S21 OR S22 OR S23 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 561,269 |
| S25 | S3 AND S24 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 250 |
| S26 | TX "construct validity" or "content validity" or "criterion validity" or "inter rater reliability" or "interrater reliability" | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 55,290 |
| S27 | S25 AND S26 | Search modes - Boolean/Phrase | Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL with Full Text | 5 |

