UNIVERSITY OF LEEDS

This is a repository copy of *Mutation of the Ferric Uptake Regulator (FUR) Severely Impairs Toxin Production in a Human in vitro Gut Model of Clostridium Difficile Infection.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/143379/>

Version: Accepted Version

Proceedings Paper:

Monaghan, TM, Chilton, CH orcid.org/0000-0002-8076-1699, Fit, M et al. (4 more authors) (2016) Mutation of the Ferric Uptake Regulator (FUR) Severely Impairs Toxin Production in a Human in vitro Gut Model of Clostridium Difficile Infection. In: Gut. British Society of Gastroenterology Annual General Meeting, 20-23 Jun 2016, Liverpool. BMJ Publishing Group , A25-A26.

<https://doi.org/10.1136/gutjnl-2016-312388.43>

© 2015 BMJ Publishing Group Ltd and British Society of Gastroenterology. This is an author accepted version of a conference abstract published in Gut. Uploaded in accordance with the publisher's archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

OC-043 Mutation of the Ferric Uptake Regulator (FUR) Severely Impairs Toxin Production in a Human in vitro Gut Model of Clostridium Difficile Infection

TM Monaghan¹, CH Chilton², M Fit³, AM Buckley², SA Kuehne³, MH Wilcox², A Cockayne³

¹NIHR Biomedical Research Unit in Gastrointestinal and Liver Diseases, University of Nottingham, Nottingham

²Leeds Institute for Molecular Medicine, University of Leeds, Leeds

³Clostridia Research Group, Centre for Biomolecular Sciences, University of Nottingham, Nottingham, UK

Introduction Regulation of iron uptake and utilisation is critical for bacterial growth and for prevention of iron toxicity. To date, little research has been carried out on iron uptake mechanisms and their regulation in *Clostridium difficile*. However, analysis of available *C. difficile* genome sequences reveals the presence of multiple iron-uptake systems and regulators underlining the importance of iron acquisition for clostridial growth. In this study, we investigated the phenotypic effects of the ferrous iron uptake system FeoB1 and the ferric uptake regulator and iron-dependent global gene regulator Fur in *C. difficile*.

Methods ClosTron mutagenesis was used to generate knockout mutants in a single *feoB1* and *fur* homologue in *C. difficile* 630Δerm, which were then inoculated into an in vitro human gut model to investigate relative propensity to induce *C. difficile* infection (CDI). Three parallel triple-stage chemostat gut models were primed with human faecal emulsions and spiked with *C. difficile* spores (~10⁷ spores) from each mutant in addition to the 630Δerm wild-type parental strain. Bacterial populations were allowed to equilibrate before simulated CDI was induced by instillation of clindamycin (33.9 mg/L, four times daily for 7 days). Serial samples were collected for enumeration of microflora populations, *C. difficile* vegetative cells, spores and measurement of cytotoxin titres.

Results Cytotoxicity assays revealed that the *fur* mutant strain produced considerably lower toxin levels (~1000 fold lower) than the *feoB1* and wild type strain. Following clindamycin exposure, all three *C. difficile* strains germinated and exhibited sustained vegetative proliferation (~5.5/6 log₁₀ cfu/mL). The *feoB1* mutant strain germinated slightly earlier than the other strains, which may have been influenced by the slightly lower clindamycin levels in the *feoB1* model. Indeed, compared to wild type, higher minimum inhibitory concentrations were observed for both mutant strains, indicating reduced susceptibility to clindamycin. In all three models, the introduction of clindamycin caused a decline in Bifidobacteria (3.5 log₁₀ cfu/mL), Clostridia (~3 log₁₀ cfu/mL) and Lactobacilli (~2 log₁₀ cfu/mL) with increases in Enterococci and Enterobacteriaceae (2–4 log₁₀ cfu/mL). However, no specific microflora changes correlated with the strain of *C. difficile* used in each of the models.

Conclusion These findings reveal the important role of the Fur system in regulating the expression of *C. difficile* toxins. Modulation of iron homeostasis may represent a potential novel therapeutic or preventative strategy against CDI.

Disclosure of Interest None Declared