

1 STAT3 Differential Scanning Fluorimetry and Differential Scanning Light Scattering Assays: addressing  
2 a missing link in the characterization of STAT3 Inhibitor interactions

3 Matthieu Desroses<sup>1</sup>, Sander Busker<sup>2</sup>, Juan Astorga-Wells,<sup>3</sup> Sanaz Attarha,<sup>1</sup> Iryna kolosenko<sup>2</sup>, Roman  
4 Zubarev,<sup>3</sup> Thomas Helleday<sup>1</sup>, Dan Grandér<sup>2</sup>, Brent D. G. Page<sup>1\*</sup>

5 <sup>1</sup>Karolinska Institute, Science for Life Laboratory, Department of Oncology and Pathology, 171 21,  
6 Stockholm, Sweden

7 <sup>2</sup>Karolinska Institute, Department of Oncology-Pathology, Cancer Center Karolinska, R8:03, 171 76,  
8 Stockholm, Sweden

9 <sup>3</sup>Karolinska Institute, Medical Biochemistry and Biophysics, 171 65, Stockholm, Sweden

10  
11 To whom correspondence should be addressed: [brent.page@scilifelab.se](mailto:brent.page@scilifelab.se)

12  
13 *Abstract*

14 STAT3 protein is an established target for the development of new cancer therapeutic agents.  
15 Despite lacking a traditional binding site for small molecule inhibitors, many STAT3 inhibitors have  
16 been identified and explored for their anti-cancer activity. Because STAT3 signaling is mediated by  
17 protein-protein interactions, indirect methods are often employed to determine if proposed STAT3  
18 inhibitors bind to STAT3 protein. While established STAT3 inhibition assays (such as the fluorescence  
19 polarization assay, electrophoretic mobility shift assay and ELISAs) have been used to identify novel  
20 inhibitors of STAT3 signaling, methods that directly assess STAT3 protein-inhibitor interactions could  
21 facilitate the development of novel inhibitors. In this context, we herein report new STAT3 binding  
22 assays, based on differential scanning fluorimetry (DSF) and differential scanning light scattering  
23 (DSLS) to characterize interactions between STAT3 protein and inhibitors. Several peptide and small  
24 molecule STAT3 inhibitors have been evaluated, and new insight into how these compounds may  
25 interact with STAT3 is provided.

26 *Introduction*

27 Signal transducer and activator of transcription 3 (STAT3) protein is a widely explored target for anti-  
28 cancer drug development.<sup>1</sup> This protein possesses several biological characteristics that make it an

attractive target for therapeutic intervention in cancer. Overactive STAT3 signaling drives proliferation, survival and immune system evasion in cancer cells, but healthy cells have transient STAT3 activation and can survive in the absence of STAT3 function.<sup>2-4</sup> While the biology of STAT3 suggests it is a good anti-cancer target, the protein itself is notoriously difficult to target with small molecule inhibitors. STAT3 does not possess a typical enzyme active site, and its activity is mediated by protein-protein and protein-DNA interactions that involve large, relatively flat areas of the protein surface. Selectively disrupting these interactions with small, drug-like molecules remains an evasive scientific challenge. In spite of this, many STAT3 inhibitors have been identified and new inhibitors continually flow into the scientific literature.<sup>2</sup> While early inhibitors, such as STATTIC,<sup>5</sup> S3I-201<sup>6</sup> and peptide inhibitors,<sup>7</sup> failed to progress into clinical testing, more recently identified STAT3 inhibitors have reached this prestigious goal, including STA-21,<sup>8</sup> STAT3 decoy-oligonucleotides,<sup>9</sup> and OPB-51602<sup>10, 11</sup>.

While the development of STAT3 inhibitors has continued at a rapid pace, the implementation of new techniques to evaluate these inhibitors has fallen behind. There are several established assays that are commonly used to measure STAT3 inhibition *in vitro* and in tumor models.<sup>2</sup> While these assays are often used to characterize and optimize the activity of STAT3 inhibitors, there are still gaps in technology that limit the understanding of how proposed STAT3 inhibitors interact with STAT3 protein.

Consequently, after two decades of research, an interesting trend has formed. Many reported STAT3 inhibitors have the propensity to act as electrophilic alkylating agents. This has recently been highlighted using mass spectrometry<sup>13, 14</sup> and fluorescence tagging<sup>15</sup> techniques with some of the most widely used STAT3 inhibitors. The majority of published STAT3 inhibitors are reported as selective Src Homology 2 (SH2) domain antagonists, however, the assays that are used to support SH2 domain binding may be also sensitive to compounds that can alkylate STAT3. As outlined in **Figure 1A**, inhibition of STAT3 may produce a protein-inhibitor complex that is energetically

1 favorable, where the protein-inhibitor complex has a lower free energy than the inhibitor and  
2 protein apart. In this case, STAT3 is inhibited (exemplified by a dimmed color compared to the  
3 brightly-colored native STAT3), and the complex is thermodynamically stable. Alternatively, a reactive  
4 inhibitor may covalently modify residues on the surface of STAT3 and induce conformational changes  
5 that alter or destabilize STAT3's tertiary structure. In these cases, STAT3 would also be inhibited  
6 (represented by the dimmed color in **Figure 1A**), however the modified tertiary structure may not  
7 bind to traditional STAT3 binding partners, or the induced instability may cause STAT3 to denature  
8 and precipitate (as depicted by the mesh surface representation). This may be especially important  
9 for *in vitro* STAT3 assays because recombinant STAT3 protein is known to be poorly soluble, unstable  
10 and difficult to work with.<sup>16</sup> In commonly used STAT3 inhibition assays, (including the fluorescence  
11 polarization (FP) assay, electrophoretic mobility shift assay (EMSA) and ELISAs), reactive compounds  
12 that chemically modify STAT3 to impair its stability or binding interactions would induce the same  
13 response as potent but non-reactive STAT3 inhibitors (described in **Figure 1B**).<sup>12</sup>

14 When used in cell-based assays, these reactive compounds may non-specifically alkylate cellular  
15 components to induce toxicity. A particularly hazardous manifestation of this would be in cancer cell  
16 proliferation assays where reactive compounds (that show inhibitory activity in traditional *in vitro*  
17 STAT3 assays) would inhibit cancer cell proliferation and could modify cell signaling networks  
18 because of their inherent toxicity, and not necessarily because they bind selectively to STAT3 or  
19 another protein of interest. Thus, differentiating between selective STAT3 binders and compounds  
20 that may non-specifically react with STAT3 in biochemical assays is an important challenge within the  
21 current STAT3 inhibitor development landscape.

22 In this context, thermal stability assays, such as differential scanning fluorimetry (DSF) or differential  
23 scanning light scattering (DSLs), can differentiate between stabilizing and destabilizing interactions  
24 between a protein of interest and potential inhibitors.<sup>16</sup> The traditional DSF assay relies on a polarity  
25 sensitive fluorescent dye (such as Sypro Orange™), which increases its fluorescence when exposed to

hydrophobic environments. Thus, thermal denaturing of a recombinant protein can be monitored by measuring Sypro Orange™ fluorescence while increasing the temperature. Capillary DSF or so called nanoDSF experiments typically use a tryptophan (Trp) fluorescence ratio to track protein unfolding. When Trp residues are buried in the hydrophobic core of the protein, their fluorescence maxima occurs around 330 nm, however when they are exposed at the protein surface, this is shifted to 350 nm. Thus, by tracking the ratio of  $F_{350}/F_{330}$  one can also monitor protein unfolding. Major advantages of the nanoDSF technique include low sample volume and no exogenous dye is added to the samples. Finally protein aggregation can also be measured by turbidometric scattering of the sample by differential scanning light scattering (DSL). Using the same set up as for nanoDSF (nanoDSL), a capillary tube is loaded with an inhibitor and the protein of interest, and thermal denaturing can be simply monitored by measuring absorbance as a representation of the turbidity of the solution.

In thermal stability assays, interactions between the recombinant protein and inhibitory small molecule may alter the melting temperature ( $T_m$ ) of the protein (the temperature at which half-maximal fluorescence or light scattering is reached). Generally, non-covalent binding of an inhibitor to a recombinant protein will increase the  $T_m$  in thermal stability assays, while covalent modifications to the protein tend to alter the tertiary structure and destabilize the protein, resulting in a lower  $T_m$ .

Recent efforts have demonstrated that commonly used STAT3 inhibitors BP1-102<sup>14</sup>, STATIC<sup>13</sup> and S3i-201<sup>15</sup> can alkylate STAT3 protein *in vitro*. It is proposed that these covalent modifications may induce conformational changes which impair interactions between STAT3 and its binding partners in biochemical assays. Thus, unlike the typically employed STAT3 assays, a STAT3 DSF assay could distinguish between the stabilizing or destabilizing interactions of proposed STAT3 inhibitors. Therefore, we aimed to generate STAT3 thermostability assays and focused on testing established STAT3 SH2 domain binders, some of which have also been implicated as possible alkylating agents (**Figure 1C**). Peptide-based STAT3 inhibitors, Ac-pYLPQTV (gp130), Ac-pYKPQMH (LIFR) and pYLKTK

(STAT3 consensus sequence, or STAT3c), were also evaluated as these represent validated SH2 domain binders that do not possess reactive groups.<sup>7</sup>

### *Materials and Methods*

#### Protein Production and Purification

BL21 DE3(T1R) pRARE2 cells were transformed with the STAT3<sup>127-688</sup> and STAT3<sup>127-465</sup> constructs and 1.5l TB (supplemented with 8 g/L Glycerol, 50 µg/mL Kanamycin, 34 µg/mL Chloramphenicol) cultures were started from overnight cultures (grown in the same medium at 30°C). The cultures were grown using a LEX bioreactor (Epiphyte3) at 37°C until an OD (600nm) of approximately 2 was reached. The temperature was then reduced to 18°C and after a further hour the cultures were induced with IPTG (0.5 mM). After incubation overnight the cultures were harvested by centrifugation at 4500g for 10min. The cell pellets were resuspended in Lysis buffer (100 mM HEPES, 500 mM NaCl, 10% glycerol, 10 mM imidazole, 0.5 mM TCEP, pH 8.0, 5 µL Benzonase Nuclease (Sigma), Complete EDTA free protease inhibitor cocktail tablet (Roche)) and then frozen at -80°C. The thawed suspended cells were lysed by sonication on ice in ~40 mL samples (Sonics, Vibra Cell 1:45min, 4 sec on, 12 sec off) and the cell debris removed by centrifugation at 49000g x 20 min before filtration of the supernatant through 0.45 µm filters. Chromatography was carried out using an Äkta Xpress (GE HEALTHCARE) at 8°C. 5 mL HisTrap (GE HEALTHCARE) columns equilibrated with Wash Buffer 1 (20 mM HEPES, 500 mM NaCl, 10% glycerol, 10 mM imidazole, 0.5 mM TCEP, pH 7.5) were loaded with the filtered supernatant and then washed with Wash buffer 1 and Wash Buffer 2 (20 mM HEPES, 500 mM NaCl, 10% glycerol, 50 mM imidazole, 0.5 mM TCEP, pH 7.5) before being eluted with Elution Buffer (20 mM HEPES, 500 mM NaCl, 10% glycerol, 500 mM imidazole, 0.5 mM TCEP, pH 7.5). The eluate was loaded on to a HiLoad 16/60 Superdex 75 (GE HEALTHCARE) column which had been equilibrated with gel Filtration buffer (20 mM HEPES, 500 mM NaCl, 10% glycerol, 0.5 mM TCEP, pH 7.5, 2 mM TECP) and eluted using the same buffer. After SDS-PAGE analysis,

fractions containing the desired protein were pooled and concentrated using Vivaspin concentrators (Sartorius). The protein was frozen in aliquots using liquid nitrogen and stored at -80°C.

### STAT3 Circular Dichroism (CD) Spectroscopy

Far ultraviolet (UV) CD spectra of STAT3<sup>127-688</sup> & STAT3<sup>127-465</sup> were recorded on a Jasco J-810 spectrometer (Jasco Spectroscopic Company, Japan) at 20 °C in a buffer containing (2 mM HEPES, 50 mM NaCl, 1 % glycerol, 0.05 mM TCEP, pH 7.5, 0.2 mM TECP). The CD Spectra were recorded over a wavelength range of 260-190 nm, with a step size of 1.0 nm, a bandwidth of 1 nm and an averaging time of 2.0 sec. Measurements were performed in a 2 mm path length quartz glass cell using a 0.1 mg/ml concentration of proteins. Five scans were applied continuously and the data were averaged. The CD spectra were smoothed and processed after baseline subtraction using Pro-Data Viewer software (Applied Photophysics, UK).

### STAT3 Fluorescence Polarization (FP) Assay

The STAT3 FP assay was carried out as previously described with subtle modifications.<sup>17, 18</sup> A corning 384 well black flat bottom plate was loaded with serial dilutions of STAT3 protein or truncated variants and 10 nM of FP assay probe (5-FAM-GpYLPQTV) in buffer containing 50 mM NaCl, 10 mM HEPES, 1 mM DTT at pH 7.5. After 10 minutes of incubation, polarized fluorescence was measured using a HidexSense reader ( $\lambda_{\text{ex}}$  = 492 nm,  $\lambda_{\text{em}}$  = 535 nm, medium lamp intensity, 10 flashes). Experiments with phosphopeptide inhibitors were carried out with 150 nM STAT3 protein. IC<sub>50</sub> values were determined by plotting concentration of probe *versus* polarized fluorescence and fitting the data to a one site binding curve.

For competition experiments, data points were plotted using GraphPad Prism and curves were fit using non-linear regression analysis for competitive binding according to the formula below. Experiments were performed in duplicate or triplicate and repeated in at least two independent experiments.

$$y = Bottom + \frac{(Top - Bottom)}{1 + 10^{(x - Log(IC_{50}))}}$$

Standard error for IC<sub>50</sub> values were transformed from the standard error in the Log(IC<sub>50</sub>) values, by adding or subtracting the standard log(IC<sub>50</sub>) errors from the log(IC<sub>50</sub>) value and then transforming those into upper and lower limits for the IC<sub>50</sub> error values. The larger of these errors was reported for the error in IC<sub>50</sub> value.

#### Hydrogen-Deuterium Exchange (HDX) Mass Spectrometry

Binding-interface characterization of the STAT3-gp130 complex was performed by differential HDX MS. First, the complex was prepared in  $\approx$  1:4 protein/ligand molar ratio by mixing 25  $\mu$ L of 5.8 mg/mL STAT3 protein in 50 mM HEPES, 1 mM EDTA, 50 mM NaCl at pH 7.5 with 0.9  $\mu$ L of 10 mM peptide stock solution in 100 % DMSO. Each individual HDX labeling reaction was initiated by mixing 2  $\mu$ L of complex solution with 18  $\mu$ L of deuterated buffer at room temperature (22°C). All the experiments were done in triplicates using a 10 min HDX labeling reaction time. For labeling, the deuterated buffer contained the same ionic composition than the protein solution, but water was replaced by heavy water (D<sub>2</sub>O). The control experiment was prepared by mixing 25  $\mu$ L of 5.8 mg/mL STAT3 protein with 0.9  $\mu$ L of DMSO, and labeled in the same manner than the STAT3-gp130 complex. After 10 min HDX labeling, each reaction was stopped by adding 30  $\mu$ L of an ice-cold 100 mM phosphate buffer pH 2.3 containing 100 mM TCEP, 3.3 M Urea, and snap-frozen in liquid nitrogen.

#### Sample Analysis – LC MS

Each labeled and quenched sample was analyzed in a semi-automated HDX-MS system (Biomotif AB, Stockholm, Sweden) in which manually injected samples were automatically digested, cleaned and separated at 2°C. Deuterated samples were digested using an in-house packed immobilized pepsin column (2.1×30 mm from ACE HPLC Columns packed with pepsin-agarose from porcine gastric mucosa obtained from Sigma-Aldrich) by a 75 seconds at 70  $\mu$ L/min flow protocol, followed by an on-line desalting step using a 2 mm I.D x 10 mm length C-18 pre-column (ACE HPLC Columns, Aberdeen, UK) using 0.05% TFA at 350  $\mu$ L/min for 3 min. Peptic peptides were then separated by a LC MS gradient method using an aqueous Solution A containing 5% of acetonitrile (ACN) in 0.3% formic acid

and an organic Solution B containing 95% of ACN and 0.3% of formic acid. The LC MS gradient profile consisted in 2-20 % ACN in 5 min, 20-30 % ACN from 5 to 20 min, 30-95% ACN from 20 to 23 min, followed by 95% ACN for 1.5 min, and 95 to 2 % ACN B in 1.5 min. The analytical column was a 2 mm I.D x 50 mm length HALO C18/1.8  $\mu$ m operated at 100  $\mu$ L/min flow. An Orbitrap XL mass spectrometer (Thermo Fisher Scientific) operated at 60,000 resolution at  $m/z$  400 was used for analysis. Peptic peptide identification was performed by 3 independent LC MS/MS analysis of an undeuterated protein sample using the same methodology than for the deuterated samples. The HDExaminer software (Sierra Analytics, USA) was used to process all HDX-MS data.

#### STAT3 Differential Scanning Fluorimetry Assay

The DSF assay was optimized according to procedures outlined previously.<sup>19</sup> Optimal conditions were achieved using buffer containing 100 mM Tris-Cl, 40 mM NaCl, 10 mM MgCl<sub>2</sub> at pH 7.4. STAT3 proteins were used at a final concentration of 1  $\mu$ M and Sypro Orange™ at “5x” (from a stock concentration of “5000x”). Inhibitor solutions were added to protein prior to the addition of Sypro Orange™. Experiments were conducted on a Biorad C1000 Thermal Cycler with CFX96 Real Time System. Heating was conducted from using gradients from 25-95 °C or 30 to 80°C (increasing 1°C per minute). Collected data was normalized to maximum and minimum values then fit using GraphPad Prism non-linear regression to a Boltzman sigmoidal curve with the formula:

$$y = Bottom + \frac{(Top - Bottom)}{1 + e^{\left(\frac{T_m - x}{Slope}\right)}}$$

Data sets were trimmed for curve fitting to include 3 additional points from minimal and maximal fluorescence values of the melt curve (which were set to 0 and 100 % relative fluorescence for STAT3 truncations, the 30 °C fluorescence value was set as the baseline for full-length STAT3 experiments). Full melt curves are shown in **Supplementary Figures 1 and 2**.  $K_d$  values were calculated from dose-response curves of the  $T_m$  values as previously described.<sup>20</sup> Briefly  $T_m$  values were plotted against ligand concentration and fitted to the following equation to find the  $K_d$  value:



$$y = Bottom + (Top - Bottom) \times \left( 1 - \frac{P - K_d - x + \sqrt{(P + x + K_d)^2 - 4Px}}{2P} \right)$$

where P represents the protein concentration, x represents the ligand concentration and y is the experimentally determined  $T_m$ .<sup>20</sup>

#### STAT3 nanoDSF and nanoDSLs Assays

NanoDSF standard glass capillaries were filled with the truncated STAT3 variants (with or without compound) in 100 mM Tris-HCl buffer, pH 7.5, containing 40 mM NaCl and 10 mM MgCl<sub>2</sub>. Using a Prometheus NT.48 (Nanotemper) instrument, the mixtures were subjected to a thermal gradient from 20 to 95°C at a heating rate of 1° C per minute. For nanoDSF experiments, Trp fluorescence at 350 and 330 nm was recorded.  $T_m$  values were determined by plotting normalized fluorescence ratio ( $F_{350}/F_{330}$ ) versus temperature and fitting to a Boltzman Sigmoidal curve as above. Data sets were trimmed to include points from 35-75 °C prior to fitting. Simultaneously, the intensity of the back-reflected light was analyzed to assess protein aggregation by nanoDSLs which served as a secondary output for this method. Again  $T_m$  values were determined from a plot of relative scattering versus temperature (from 35-75 °C) and fitting to a Boltzman Sigmoidal curve.

#### Statistical Analysis

Statistical analysis was performed using GraphPad Prism 7.0 using. 1way ANOVA was used to generate p values which are displayed as ns = p > 0.05, \* = p ≤ 0.05, \*\* = p ≤ 0.01, \*\*\* = p ≤ 0.001 and \*\*\*\* = p ≤ 0.0001.

#### Chemical Reagents

All reagents and inhibitors were purchased from commercial suppliers and used without further modifications. Fluorescent and non-labeled phosphopeptide sequences were purchased from Innovagen AB (Lund, Sweden) or Biomatik (Cambridge, Canada) and diluted in DMSO or H<sub>2</sub>O prior to use. Full-length STAT3 protein (amino acids 1-770) was purchased from NordicBioSite (Täby,

Sweden). Cloning and production of truncated STAT3 proteins was conducted at the Karolinska Institute Protein Science Facility (Solna, Sweden).

### Results

Full-length STAT3 protein (STAT3<sup>Full</sup>) is known to be difficult to produce, store and handle.<sup>16</sup> Thus, it was suspected that STAT3<sup>Full</sup> might have problematic instability at the elevated temperatures needed for thermal stability assays. More stable STAT3 variants had previously been reported for protein crystallography experiments, where N- and C-terminal truncations afforded a STAT3 variant that could be crystalized.<sup>16</sup>

Because many reported STAT3 inhibitors claim to bind to the SH2 domain, truncated STAT3 variants were designed so that SH2 domain binding could be assessed. Several truncated proteins were developed in matched pairs to include or exclude the SH2 domain. Of the possibilities analyzed in test expressions, STAT3<sup>127-465</sup> (containing the coiled-coil domain (CCD) and DNA binding domain (DBD)) and STAT3<sup>127-688</sup> (CCD to SH2 domain) gave high expression of soluble protein (schematics are shown in **Figure 2A**). The corresponding variant from CCD to Linker domain (STAT3<sup>127-578</sup>) was not soluble and could only be detected at low levels in test expressions. STAT3<sup>Full</sup> could not be isolated under these conditions and instead was purchased from a commercial supplier. STAT3<sup>127-465</sup> and STAT3<sup>127-688</sup> were analyzed by circular dichroism (CD) to determine if they possessed appropriate secondary structures (**Supplementary Figure 3**). Indeed, both truncations had well-defined CD spectra with mostly alpha helical character indicating that these truncated STAT3 variants still formed folded structures.

STAT3<sup>127-688</sup> was further analyzed for SH2 domain integrity using the STAT3 FP assay<sup>17</sup> and hydrogen-deuterium exchange (HDX) experiments.<sup>21</sup> The STAT3 FP assay utilizes fluorescently tagged peptide probe (5-aminofluorescein-GpYLPQTV, referred to herein as 5-FAM-gp130) to assess SH2 domain binding.<sup>17</sup> Typically, the FP assay is used to assess the ability of proposed SH2 domain binders to displace 5-FAM-gp130 from the SH2 domain of STAT3. In this case, STAT3<sup>127-688</sup>, STAT3<sup>127-465</sup> and

STAT3<sup>Full</sup> were titrated against 5-FAM-gp130 and polarized fluorescence was measured to ensure that the SH2 domain of STAT3<sup>127-688</sup> was intact. As shown in **Figure 2B**, STAT3<sup>127-688</sup> (which contains the SH2 domain) retained its ability to bind 5-FAM-gp130, as increasing concentrations of STAT3<sup>127-688</sup> resulted in greater polarized fluorescence output. As well, STAT3<sup>127-688</sup> gave greater FP signal compared to full-length STAT3 and had a lower  $K_d$  value ( $61 \pm 6$  nM *versus*  $550 \pm 230$  nM for STAT3<sup>Full</sup>). As anticipated, STAT3<sup>127-465</sup>, which lacks the SH2 domain, showed no binding to 5-FAM-gp130.

To further assess the integrity of the SH2 domain of STAT3<sup>127-688</sup>, known SH2 domain-binding peptide sequences from LIFR, gp130 and STAT3c were also assessed by FP assay (**Figure 2C**).<sup>7</sup> As expected, 5-FAM-gp130 binding was inhibited by these peptides with  $IC_{50}$  values of  $0.7 \pm 0.15$   $\mu$ M,  $0.66 \pm 0.09$   $\mu$ M and  $130 \pm 30$   $\mu$ M for gp130, LIFR and STAT3c, respectively.

As another confirmation that the SH2 domain of STAT3<sup>127-688</sup> was intact, HDX mass spectrometry was used to identify the interaction site between STAT3<sup>127-688</sup> and gp130 (**Figure 2D**). Indeed, exposure of STAT3<sup>127-688</sup> to gp130 decreased the incorporation deuterium within the STAT3 SH2 domain, indicating binding. Thus, the truncations employed to generate STAT3<sup>127-688</sup> did not affect its ability to interact with known STAT3 SH2 domain binders in biochemical assay settings.

STAT3<sup>Full</sup> and the truncated variants were then assessed for their suitability for use in thermal stability assays (**Figure 3A-E**). Under the described DSF conditions, STAT3<sup>Full</sup> was quite unstable towards elevated temperatures and gave a  $T_m$  of just 37.2 °C (**Figure 3A**). Moreover, gradual increased fluorescence emission at sub-physiological temperatures (30-37 °C) was detected which may reflect the inherent instability of the recombinant STAT3<sup>Full</sup> protein. Both STAT3 truncations were much more stable towards thermal degradation (STAT3<sup>127-465</sup>  $T_m$  = 53.8 °C and STAT3<sup>127-688</sup>  $T_m$  = 53.5 °C, representative curves shown in **Figure 3B** and **C**, respectively and multiple experiments are summarized in **Figure 3D** and **E**).

1 To determine if thermal stability could be affected by SH2 domain binding, the truncated and full-  
2 length proteins were analyzed by DSF in the presence of 1 mM gp130. Unexpectedly, the  $T_m$  of full-  
3 length STAT3 was not shifted in the presence of gp130 (**Figure 3A**). This may reflect the presence of  
4 unstable regions within N- and C-terminal domains that may instigate the thermal denaturing  
5 process which cannot be stabilized by SH2 domain binding. The gp130 sequence also did not stabilize  
6 STAT3<sup>127-465</sup> which lacks the SH2 domain (**Figure 3B**), however gp130 induced a positive  $T_m$  shift ( $\Delta T_m$ )  
7 of 8.2 °C with STAT3<sup>127-688</sup>, indicating a direct stabilizing interaction (**Figure 3C**).

8 To expand on these initial results, LIFR and STAT3c were also analyzed by DSF with STAT3<sup>127-688</sup>. Like  
9 gp130, these inhibitors also stabilized STAT3<sup>127-688</sup>. The  $T_m$  shifts for these inhibitors were in  
10 accordance with reported inhibitory values<sup>7</sup> (representative curves are shown in **Figure 3F** and  
11 multiple experiments are summarized in **Figure 3G**). Dose response experiments with the inhibitory  
12 peptides (from 2.4  $\mu$ M to 5 mM at 2-fold dilution steps) were also performed to generate  $K_d$  values  
13 from the DSF assay. The observed  $K_d$  values also corresponded to known inhibitory constants for  
14 STAT3 with these peptide inhibitors (**Figure 3H**).<sup>7</sup> No interactions were detected between these  
15 peptides and Sypro Orange<sup>TM</sup> when run without STAT3 proteins in control experiments  
16 (**Supplementary Figure 4**).

17 Having demonstrated that our DSF assay could be used for assessing direct interactions between the  
18 STAT3 SH2 domain and known peptide inhibitors, established small molecule STAT3 inhibitors were  
19 also tested. Of the plethora of small molecule STAT3 inhibitors present in the scientific literature,  
20 STATTIC,<sup>5</sup> S3I-201,<sup>6</sup> BP1-102<sup>22, 23</sup> and STA-21<sup>24</sup> were selected for analysis. All four of these agents are  
21 proposed STAT3 SH2 domain binders.<sup>5, 6, 22-24</sup> Furthermore, STATTIC, S3I-201 and BP1-102 have been  
22 identified as probable covalent modifiers of STAT3 protein,<sup>13-15</sup> adding more interest to their  
23 evaluation in the DSF assay. Binding of BP1-102 to STAT3<sup>127-688</sup> was also confirmed by FP assay  
24 (**Supplementary Figure 5**).

1 Unexpectedly, no overt stabilizing (or destabilizing) interactions were detected between STAT3<sup>127-688</sup>  
2 S3I-201 or STA-21 (**Figure 4A**). BP1-102 and STATTIC caused dose-dependent decrease in the  $T_m$  of  
3 STAT3<sup>127-688</sup>. This fits with theories suggesting that these agents may function *via* covalent  
4 modification of STAT3 *in vitro*, which could result in decreased overall stability.

5 To assess if these destabilizing effects were mediated by specific binding to the STAT3 SH2 domain,  
6 these same inhibitors were assessed for binding to STAT3<sup>127-465</sup> which lacks the SH2 domain. Like  
7 STAT3<sup>127-688</sup>, dose-dependent destabilization of STAT3<sup>127-465</sup> was observed with BP1-102 or STATTIC  
8 (**Figure 3B**). This indicates that these compounds may interact with STAT3 at other locations than  
9 only its SH2 domain. Only very small  $T_m$  shifts (< 1 °C) were observed upon treatment of STAT3<sup>127-465</sup>  
10 with S3i-201 or STA-21, and although some variation proved to be statistically significant (likely do to  
11 the high level of reproducibility between experiments), it is believed that the fraction of a degree  
12 difference in STAT3<sup>127-465</sup>  $T_m$  values between S3i-201 and STA-21 versus the DMSO control would not  
13 have major biological implications.

14 Surprisingly, in control experiments where the small molecule inhibitors were incubated with Sypro  
15 Orange<sup>TM</sup> alone, BP1-102 was found to interact with the dye and alter its fluorescence properties in a  
16 temperature dependent manner (**Supplementary Figure 6**). This may help to explain the high degree  
17 of variability with the BP1-102 samples in these experiments which contributed to a lack of statistical  
18 significance upon analysis with STAT3<sup>127-688</sup>. Thus, to further clarify if the observed  $T_m$  shifts were due  
19 to interactions with Sypro Orange<sup>TM</sup>, thermal denaturing curves were generated using the intrinsic  
20 fluorescence of Trp residues and by turbidometric scattering to monitor protein aggregation (so-  
21 called nanoDSF and nanoDSLS experiments, respectively). STAT3<sup>127-465</sup> has only 3 Trp residues, which  
22 prevented accurate analysis of thermal denaturing based on Trp fluorescence ratios (**Supplementary**  
23 **Figure 7a**). Therefore melt curves for STAT3<sup>127-465</sup> could only be generated by measuring scattering  
24 from the capillary solution with increasing temperature (**Supplementary Figure 7b**). STAT3<sup>127-688</sup>

melting curves were measured using Trp fluorescence ratios or scattering as shown in **Supplementary Figure 7a-b**.

Using these additional methods, interactions between truncated STAT3 proteins and inhibitors were assessed (representative melting curves are shown in **Supplementary Figure 8** and **Figure 4C-F** summarizes multiple experiments). NanoDSF and nanoDLS methods confirmed the results from the DSF assay with Sypro Orange<sup>TM</sup>, showing that peptide inhibitors (gp130, LIFR and STAT3c) stabilized STAT3<sup>127-688</sup> but not STAT3<sup>127-465</sup> towards thermal degradation. As well, nanoDSF and nanoDLS experiments confirmed that BP1-102 and STATTIC destabilized both STAT3<sup>127-688</sup> and STAT3<sup>127-465</sup> suggesting that these agents might function differently than the peptide inhibitors and may interact with STAT3 at sites beyond its SH2 domain. The nanoDSF experiments could also detect a very subtle destabilizing effect of DMSO on STAT3<sup>127-688</sup>, again it is suspected that this likely would not have biological significance.

### Discussion

Using the reported STAT3 thermal stability assays, stabilizing interactions were detected between peptide STAT3 inhibitors and STAT3<sup>127-688</sup>. In contrast, small molecule STAT3 inhibitors failed to induce positive  $T_m$  shifts. Instead, BP1-102 and STATTIC decreased the  $T_m$  of STAT3<sup>127-688</sup>, consistent with reports indicating that these agents may act *via* covalent modification of STAT3 protein.<sup>13, 14</sup> Surprisingly, S3i-201, which has also been implicated as a potential covalent modifier of STAT3,<sup>15</sup> did not decrease the  $T_m$  of STAT3<sup>127-688</sup>. While STA-21 and S3i-201 induced minute shifts to the  $T_m$  of STAT3<sup>127-688</sup> (and STAT3<sup>127-465</sup>) in the DSF assay, one cannot rule out that these agents may still bind directly to STAT3 protein, perhaps at regions beyond residues 127-688 or in a modality that does not affect the  $T_m$  of the protein.

Ideally, one would like to use STAT3<sup>Full</sup> in the DSF assay, however STAT3<sup>Full</sup> had a very low  $T_m$  which could not be appreciably shifted by high concentrations of gp130 (one of the most potent STAT3 inhibitors discovered) or by small molecule inhibitors BP1-102 and STATTIC (all  $\Delta T_m < 1$  °C)

(Supplementary Figure 9). By truncating the N- and C-termini of the protein, more stable STAT3 variants were produced, as indicated by higher  $T_m$  values in the DSF assay. The  $T_m$  of STAT3<sup>127-688</sup> was shifted by peptide STAT3 inhibitors (gp130, LIFR and STAT3c), however they did not shift the  $T_m$  of STAT3<sup>127-465</sup>, supporting their use as selective SH2 domain antagonists. Beyond the DSF assay, STAT3<sup>127-688</sup> was shown to be active in the STAT3 FP assay and could also be used in HDX experiments.  $\Delta T_m$  values for STAT3<sup>127-688</sup> in the DSF assay were confirmed using nanoDSF and nanoDSLS to rule out possible interactions between Sypro Orange<sup>TM</sup> and the inhibitors. While BP1-102 demonstrated some interaction with the Sypro Orange<sup>TM</sup>, the destabilizing nature of BP1-102 was confirmed using nanoDSF and nanoDSLS using intrinsic Trp fluorescence and protein aggregation in place of the exogenously added Sypro Orange<sup>TM</sup> dye.

Similar to these thermal stability assays, another thermofluorescence assay was recently reported for STAT proteins.<sup>25</sup> This assay used displacement of 5-FAM-gp130 from the STAT3 SH2 domain to track protein degradation by DSF. The authors demonstrated that upon heating, STAT proteins were denatured which resulted in the probe being unable to bind to the protein. When the probe was displaced, its fluorescence decreased due to solvent quenching effects. The authors contended that STAT3 inhibitors, including BP1-102, displaced the probe, leading to decreased fluorescence intensity as measured by area under the first derivative of fluorescence curve. While further exploration into this topic is recommended, our findings suggest an alternative explanation for the observed results. Instead of specifically displacing the probe, we suggest that BP1-102 destabilizes STAT3 and enhances its thermal degradation. This which prevent the probe from binding and could explain the observed decreases in fluorescence intensity.

## Conclusions

Unlike many commonly used STAT3 inhibition assays, the described DSF assay can differentiate between inhibitors that form more stable complexes with STAT3 *versus* potentially reactive compounds that destabilize the protein or alter its tertiary structure. Advantageously, this assay can

1 identify compounds that directly interact with STAT3, providing important information about the  
2 binding region as well as the mechanism of action of such compounds. While developed for STAT3  
3 inhibitors, the same platform can be applied to identify binders of other STAT proteins, becoming a  
4 valuable tool for the discovery of novel STAT inhibitors with a broad spectrum of applications.

#### 5 *Acknowledgements*

6 The study was supported by the grants from the Swedish Society for Medical Research (SSMF) and  
7 the David and Astrid Hegelén Foundation (BDGP), with additional support from the Swedish Cancer  
8 Society (TH, DG), the Swedish Childhood Cancer Society (TH, DG), Radiumhemmet Research  
9 Foundation (DG) and Swedish Research Council (TH, DG). SB is supported by KID funding from the  
10 Karolinska Institute. Part of this work was facilitated by the Protein Science Facility at Karolinska  
11 Institute and Scilifelab (<http://psf.ki.se>). We would like to thank Dr. Nicholas Valerie for assistance  
12 with statistical analysis.

#### 13 *Figure Captions*

14 **Figure 1. A)** Description of possible consequences of STAT3 inhibitor binding. Native STAT3 can be  
15 recruited to activated receptors and interact with binding partners to fulfill its cellular functions. A  
16 STAT3 inhibitor (STAT3i) may block STAT3 function through a variety of mechanisms. **i)** Selective non-  
17 covalent binding of an inhibitor preserves the tertiary structure and stability of STAT3 but inhibits  
18 STAT3 function (represented by the dimmed surface color). **ii)** Covalent modification to STAT3 may  
19 decrease the protein's stability (indicated by a mesh surface representation) or **iii)** could alter the  
20 tertiary structure of the protein and prevent interactions with binding partners (which would likely  
21 impact the protein's stability as well). **B)** Descriptions of commonly used STAT3 inhibition assays that  
22 show confounding results with compounds that destabilize or modify the tertiary structure of the  
23 protein. Non-specific binding or reactivity with STAT3 protein in these biochemical assays may  
24 masquerade as selective STAT3 inhibition. **C)** Established small molecule STAT3 inhibitors STATTC,  
25 STA-21, S3i-201 and BP-1-102.



1

2 **Figure 2. A)** Representation of STAT3 proteins produced for thermal stability studies. Full length  
3 STAT3 (STAT3<sup>Full</sup>) includes the N-terminal domain (ND), coiled-coil domain (CCD), DNA binding  
4 domain (DBD), linker domain, Src homology 2 domain (SH2) and the transactivation domain (TAD).  
5 STAT3<sup>127-688</sup> (CCD to SH2 domains) and STAT3<sup>127-465</sup> (CCD to DBD) were easily produced using  
6 traditional bacterial expression and purification techniques. **B)** Fluorescence polarization (FP) assay  
7 showing STAT3<sup>Full</sup>, and STAT3<sup>127-688</sup> but not STAT3<sup>127-465</sup> can bind to 5-FAM-gp130 resulting in  
8 increased FP signal at higher concentrations of protein. **C)** STAT3<sup>127-688</sup> FP assay using peptide STAT3  
9 inhibitors. STAT3<sup>127-688</sup> maintained strong binding affinity for known phosphopeptide sequences  
10 from gp130, LIFR and the STAT3 consensus sequence. **D)** Hydrogen-deuterium exchange assay with  
11 STAT3<sup>127-688</sup> and gp130. Relative change in deuterium incorporation per retrieved peptide after gp130  
12 peptide binding to STAT3<sup>127-688</sup> is plotted against the amino acid sequence with functional domains  
13 indicated. Deuterium incorporation was significantly decreased in peptides extending below the red  
14 line, confirming that gp130 peptide binds specifically to the SH2 domain of STAT3<sup>127-688</sup>. Data is an  
15 average of three independent replicates.

16

17 **Figure 3. STAT3 Differential Scanning Fluorimetry Assays. A)** A representative DSF assay showing  
18 STAT3<sup>Full</sup> has poor thermal stability ( $T_m = 37.2\text{ }^{\circ}\text{C}$ ) and it was not stabilized by the addition of 1 mM  
19 gp130. **B)** A representative experiment showing STAT3<sup>127-465</sup> is thermostable,  $T_m = 53.8\text{ }^{\circ}\text{C}$ , and no  
20 stabilization of STAT3<sup>127-465</sup> was observed upon treatment with 1 mM gp130 as expected because  
21 STAT3<sup>127-465</sup> lacks the SH2 domain. **C)** A representative plot of melting curves, STAT3<sup>127-688</sup> was  
22 stabilized by treatment with the gp130 peptide sequence resulting in a large  $T_m$  shift ( $\Delta T_m = 8.2\text{ }^{\circ}\text{C}$ ).  
23 **D)** Graphical summary of  $T_m$  shifts induced by gp130 from two independent experiments. Only the  $T_m$   
24 for STAT3<sup>127-688</sup> was significantly shifted by the addition of gp130 compared to the DMSO control. **E)**  
25 Table of  $T_m$  values from **D**. **F)** Analysis of known peptide inhibitors of STAT3 protein: gp130, LIFR and

the STAT3 consensus sequence induced positive  $T_m$  shifts in STAT3<sup>127-688</sup>, a representative experiment is shown. **G)** Graph and table summarizing three independent experiments investigating  $T_m$  shifts with peptide inhibitors. **H)** Dose-response analysis of  $T_m$  values with peptide STAT3 inhibitors (2.4  $\mu$ M to 5 mM, 2 fold dilutions). Larger shifts in  $T_m$  were observed with increasing concentrations of inhibitors until saturation at high concentrations.  $K_d$  values derived from dose-response experiments are also shown.

**Figure 4. A)** Dose-response analysis of known STAT3 inhibitors in the DSF assay using STAT3<sup>127-688</sup>. BP1-102 and STATTIC caused a dose-dependent decrease in the  $T_m$  of STAT3<sup>127-688</sup>. S3I-201 and STA-21 did not shift the  $T_m$  of STAT3<sup>127-688</sup>. The gp130 sequence caused a dose-dependent positive  $T_m$  shift with STAT3<sup>127-688</sup>. **B)** STATTIC and BP1-102 also decreased the  $T_m$  STAT3<sup>127-465</sup> which does not contain the SH2 domain. STA-21, S3I-201 and the gp130 sequence did not affect the  $T_m$  of STAT3<sup>127-465</sup>. **C-F)** NanoDSF and nanoDSLS experiments with STAT3<sup>127-688</sup> and STAT3<sup>127-465</sup>. Inhibitory peptides (STAT3c, LIFR and gp130, 1 mM) stabilized STAT3<sup>127-688</sup> towards thermal degradation when analyzed by Trp fluorescence ratio ( $F_{350}/F_{330}$ ) **C)** or by light scattering **D)**. Small molecule STAT3 inhibitors BP1-102 (80  $\mu$ M) and STATTIC (80  $\mu$ M) destabilized STAT3<sup>127-688</sup> towards thermal degradation. **E)**  $T_m$  values for STAT3<sup>127-465</sup> in the nanoDSLS assay monitoring protein denaturing by scattering. Trp fluorescence ratio could not be used to monitor STAT3<sup>127-465</sup> degradation due to the low number of Trp residues in STAT3<sup>127-465</sup>. **F)** Table summarizing the results from the nanoDSF experiments depicted in **C-E)**.

## References

1. Lai, P. S.; Rosa, D. A.; Magdy Ali, A.; Gómez-Biagi, R. F.; Ball, D. P.; Shouksmith, A. E.; Gunning, P. T., A STAT inhibitor patent review: progress since 2011. *Expert Opin Ther Pat* **2015**, 25 (12), 1397-421.
2. Furtek, S. L.; Backos, D. S.; Matheson, C. J.; Reigan, P., Strategies and Approaches of Targeting STAT3 for Cancer Treatment. *ACS Chem Biol* **2016**, 11 (2), 308-18.
3. Gao, J.; McConnell, M. J.; Yu, B.; Li, J.; Balko, J. M.; Black, E. P.; Johnson, J. O.; Lloyd, M. C.; Altioek, S.; Haura, E. B., MUC1 is a downstream target of STAT3 and regulates lung cancer cell survival and invasion. *Int J Oncol* **2009**, 35 (2), 337-45.
4. Bromberg, J. F.; Wrzeszczynska, M. H.; Devgan, G.; Zhao, Y.; Pestell, R. G.; Albanese, C.; Darnell, J. E., Stat3 as an oncogene. *Cell* **1999**, 98 (3), 295-303.

5. Schust, J.; Sperl, B.; Hollis, A.; Mayer, T. U.; Berg, T., Stattic: a small-molecule inhibitor of STAT3 activation and dimerization. *Chem Biol* **2006**, *13* (11), 1235-42.
6. Siddiquee, K.; Zhang, S.; Guida, W. C.; Blaskovich, M. A.; Greedy, B.; Lawrence, H. R.; Yip, M. L.; Jove, R.; McLaughlin, M. M.; Lawrence, N. J.; Sebt, S. M.; Turkson, J., Selective chemical probe inhibitor of Stat3, identified through structure-based virtual screening, induces antitumor activity. *Proc Natl Acad Sci U S A* **2007**, *104* (18), 7391-6.
7. Ren, Z.; Cabell, L. A.; Schaefer, T. S.; McMurray, J. S., Identification of a high-affinity phosphopeptide inhibitor of Stat3. *Bioorg Med Chem Lett* **2003**, *13* (4), 633-6.
8. Miyoshi, K.; Takaishi, M.; Nakajima, K.; Ikeda, M.; Kanda, T.; Tarutani, M.; Iiyama, T.; Asao, N.; DiGiovanni, J.; Sano, S., Stat3 as a therapeutic target for the treatment of psoriasis: a clinical feasibility study with STA-21, a Stat3 inhibitor. *J Invest Dermatol* **2011**, *131* (1), 108-17.
9. Sen, M.; Thomas, S. M.; Kim, S.; Yeh, J. I.; Ferris, R. L.; Johnson, J. T.; Duvvuri, U.; Lee, J.; Sahu, N.; Joyce, S.; Freilino, M. L.; Shi, H.; Li, C.; Ly, D.; Rapireddy, S.; Etter, J. P.; Li, P. K.; Wang, L.; Chiosea, S.; Seethala, R. R.; Gooding, W. E.; Chen, X.; Kaminski, N.; Pandit, K.; Johnson, D. E.; Grandis, J. R., First-in-human trial of a STAT3 decoy oligonucleotide in head and neck tumors: implications for cancer therapy. *Cancer Discov* **2012**, *2* (8), 694-705.
10. Ogura, M.; Uchida, T.; Terui, Y.; Hayakawa, F.; Kobayashi, Y.; Taniwaki, M.; Takamatsu, Y.; Naoe, T.; Tobinai, K.; Munakata, W.; Yamauchi, T.; Kageyama, A.; Yuasa, M.; Motoyama, M.; Tsunoda, T.; Hatake, K., Phase I study of OPB-51602, an oral inhibitor of signal transducer and activator of transcription 3, in patients with relapsed/refractory hematological malignancies. *Cancer Sci* **2015**, *106* (7), 896-901.
11. Wong, A. L.; Soo, R. A.; Tan, D. S.; Lee, S. C.; Lim, J. S.; Marban, P. C.; Kong, L. R.; Lee, Y. J.; Wang, L. Z.; Thuya, W. L.; Soong, R.; Yee, M. Q.; Chin, T. M.; Cordero, M. T.; Asuncion, B. R.; Pang, B.; Pervaiz, S.; Hirpara, J. L.; Sinha, A.; Xu, W. W.; Yuasa, M.; Tsunoda, T.; Motoyama, M.; Yamauchi, T.; Goh, B. C., Phase I and biomarker study of OPB-51602, a novel signal transducer and activator of transcription (STAT) 3 inhibitor, in patients with refractory solid malignancies. *Ann Oncol* **2015**, *26* (5), 998-1005.
12. Furtek, S. L.; Matheson, C. J.; Backos, D. S.; Reigan, P., Evaluation of quantitative assays for the identification of direct signal transducer and activator of transcription 3 (STAT3) inhibitors. *Oncotarget* **2016**, *7* (47), 77998-78008.
13. Heidelberger, S.; Zinzalla, G.; Antonow, D.; Essex, S.; Basu, B. P.; Palmer, J.; Husby, J.; Jackson, P. J.; Rahman, K. M.; Wilderspin, A. F.; Zloh, M.; Thurston, D. E., Investigation of the protein alkylation sites of the STAT3:STAT3 inhibitor Stattic by mass spectrometry. *Bioorg Med Chem Lett* **2013**, *23* (16), 4719-22.
14. Ali, A. M.; Gómez-Biagi, R. F.; Rosa, D. A.; Lai, P. S.; Heaton, W. L.; Park, J. S.; Eiring, A. M.; Vellore, N. A.; de Araujo, E. D.; Ball, D. P.; Shouksmith, A. E.; Patel, A. B.; Deininger, M. W.; O'Hare, T.; Gunning, P. T., Disarming an Electrophilic Warhead: Retaining Potency in Tyrosine Kinase Inhibitor (TKI)-Resistant CML Lines While Circumventing Pharmacokinetic Liabilities. *ChemMedChem* **2016**, *11* (8), 850-61.
15. Ball, D. P.; Lewis, A. M.; Williams, D.; Resette, D.; Wilson, D. J.; Gunning, P. T., Signal transducer and activator of transcription 3 (STAT3) inhibitor, S3I-201, acts as a potent and non-selective alkylating agent. *Oncotarget* **2016**.
16. Becker, S.; Groner, B.; Müller, C. W., Three-dimensional structure of the Stat3beta homodimer bound to DNA. *Nature* **1998**, *394* (6689), 145-51.
17. Schust, J.; Berg, T., A high-throughput fluorescence polarization assay for signal transducer and activator of transcription 3. *Anal Biochem* **2004**, *330* (1), 114-8.
18. Page, B. D.; Croucher, D. C.; Li, Z. H.; Haftchenary, S.; Jimenez-Zepeda, V. H.; Atkinson, J.; Spagnuolo, P. A.; Wong, Y. L.; Colaguri, R.; Lewis, A. M.; Schimmer, A. D.; Trudel, S.; Gunning, P. T., Inhibiting aberrant signal transducer and activator of transcription protein activation with tetrapodal, small molecule Src homology 2 domain binders: promising agents against multiple myeloma. *J Med Chem* **2013**, *56* (18), 7190-200.

19. Mashalidis, E. H.; Śledź, P.; Lang, S.; Abell, C., A three-stage biophysical screening cascade for fragment-based drug discovery. *Nat Protoc* **2013**, *8* (11), 2309-24.
20. Vivoli, M.; Novak, H. R.; Littlechild, J. A.; Harmer, N. J., Determination of protein-ligand interactions using differential scanning fluorimetry. *J Vis Exp* **2014**, (91), 51809.
21. Eiring, A. M.; Page, B. D.; Kraft, I. L.; Mason, C. C.; Vellore, N. A.; Resetca, D.; Zabriskie, M. S.; Zhang, T. Y.; Khorashad, J. S.; Engar, A. J.; Reynolds, K. R.; Anderson, D. J.; Senina, A.; Pomicter, A. D.; Arpin, C. C.; Ahmad, S.; Heaton, W. L.; Tantravahi, S. K.; Todic, A.; Colaguori, R.; Moriggl, R.; Wilson, D. J.; Baron, R.; O'Hare, T.; Gunning, P. T.; Deininger, M. W., Combined STAT3 and BCR-ABL1 inhibition induces synthetic lethality in therapy-resistant chronic myeloid leukemia. *Leukemia* **2015**, *29* (3), 586-97.
22. Zhang, X.; Yue, P.; Page, B. D.; Li, T.; Zhao, W.; Namanja, A. T.; Paladino, D.; Zhao, J.; Chen, Y.; Gunning, P. T.; Turkson, J., Orally bioavailable small-molecule inhibitor of transcription factor Stat3 regresses human breast and lung cancer xenografts. *Proc Natl Acad Sci U S A* **2012**, *109* (24), 9623-8.
23. Page, B. D.; Fletcher, S.; Yue, P.; Li, Z.; Zhang, X.; Sharmeen, S.; Datti, A.; Wrana, J. L.; Trudel, S.; Schimmer, A. D.; Turkson, J.; Gunning, P. T., Identification of a non-phosphorylated, cell permeable, small molecule ligand for the Stat3 SH2 domain. *Bioorg Med Chem Lett* **2011**, *21* (18), 5605-9.
24. Song, H.; Wang, R.; Wang, S.; Lin, J., A low-molecular-weight compound discovered through virtual database screening inhibits Stat3 function in breast cancer cells. *Proc Natl Acad Sci U S A* **2005**, *102* (13), 4700-5.
25. de Araujo, E. D.; Manaswiyoungkul, P.; Israelian, J.; Park, J.; Yuen, K.; Farhangi, S.; Berger, A.; Abu-Jazar, L.; Gunning, P. T., High-throughput thermofluor-based assays for inhibitor screening of STAT SH2 domains. *J Pharm Biomed Anal* **2017**.