



UNIVERSITY OF LEEDS

This is a repository copy of *Incomplete and Systemic Lupus Erythematosus Reveal A Different Pattern of Interferon-Stimulated Genes Up-Regulation*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/144372/>

Version: Accepted Version

Proceedings Paper:

Md Yusof, MY orcid.org/0000-0003-3131-9121, El-Sherbiny, Y orcid.org/0000-0003-4791-3475, Psarras, A et al. (4 more authors) (2016) Incomplete and Systemic Lupus Erythematosus Reveal A Different Pattern of Interferon-Stimulated Genes Up-Regulation. In: *Annals of the Rheumatic Diseases*. Annual European Congress of Rheumatology (EULAR 2016), 08-11 Jun 2016, London, UK. BMJ Publishing Group , pp. 285-286.

<https://doi.org/10.1136/annrheumdis-2016-eular.6040>

(c) 2016, Published by the BMJ Publishing Group Limited. This is an author produced version of an abstract published in *Annals of the Rheumatic Diseases*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Medical or Research Professionals/Clinicians

Topic area: Basic and translational research

Topic: 8. SLE, Sjögren's and APS - etiology, pathogenesis and animal models

EULAR16-6040

INCOMPLETE AND SYSTEMIC LUPUS ERYTHEMATOSUS REVEAL A DIFFERENT PATTERN OF INTERFERON-STIMULATED GENES UP-REGULATION

M. Y. Md Yusof^{1,2}, Y. El-Sherbiny¹, A. Psarras¹, E. M. Hensor^{1,2}, M. Wittmann¹, P. Emery^{1,2}, E. M. Vital^{1,2}

¹Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, ²NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom

My abstract has been or will be presented at a scientific meeting during a 12 months period prior to EULAR 2016:

Yes

Abstract presented or will be presented at (meeting): EWRR 2016

Is the first author applying for a travel bursary and/or an award for undergraduate medical students?: No

Background: Type I interferons (IFN-I) play a central role in the pathogenesis systemic lupus erythematosus (SLE). IFN-I activity has been used as a biomarker by measuring interferon-stimulated gene (ISG) transcripts in the peripheral blood of SLE patients, but its relationship with clinical features still remains unclear. Incomplete lupus erythematosus (ILE) describes individuals with new onset of features suggestive of SLE, but do not fulfil diagnostic criteria. Up to 20% of these patients eventually progress to SLE.

Objectives: To investigate ISG expression in patients with ILE and explore possible qualitative differences between established and early stages of SLE.

Methods: A meta-analysis of known ISGs indicated 33 genes of importance[1]. The expression of 33 ISGs was measured using qPCR in PBMCs from individuals with SLE (n=54), ILE (n=27), and healthy controls (HC; n=14). SLE was defined using 2012 ACR/SLICC criteria. ILE was defined as ANA +ve, 1-2 clinical ACR/SLICC criteria, and symptom duration <12 months. Factor analysis (FA) was used to reduce expression data to a limited set of factors, which were compared between patient groups using ANCOVA test.

Results: FA on SLE patients indicated two factor scores explaining 80% of the data variance. The majority of variability was explained by Factor F1; however, Factor F2 appeared more relevant to the presence of fully established SLE. F1 and F2 were significantly different between patient groups; p=0.005 and p=0.044 respectively. F1 was similarly high in both SLE [SLE:HC=4.22 (1.80, 9.88), p=0.001] and ILE [ILE:HC=2.96 (1.20, 7.32), p=0.019]. In contrast, F2 was increased only in SLE [SLE:HC=1.38 (0.95, 2.00), p=0.086] but not in ILE [ILE:HC=1.02 (0.69, 1.51), p=0.917]; a significant difference was observed between SLE and ILE patients [SLE:ILE=1.35 (1.04, 1.77), p=0.026]. In total, 16 and 14 genes loaded onto F1 and F2 respectively. Most of the genes were involved in IFN signalling pathways. Genes related to apoptosis and ubiquitination were present in both F1 and F2. Additionally, F1 loaded genes were mainly associated with antiviral immunity, complement regulation, and Th2 responses; genes loaded onto F2 were associated with regulation of IFN-I, dsDNA binding, chemotaxis, and Th1 responses.

Conclusions: IFN-I activity is present in ILE. However, the majority of measured ISG expression (F1) cannot distinguish ILE and SLE. F1 represents genes that distinguish healthy individuals, but show little variation at different stages of disease development. We define a subset of ISGs (F2), the expression of which is only increased in patients with confirmed clinical SLE. ISG expression is not unidimensional: qualitative differences in expression of distinct ISGs can contribute to clinical progression after disease initiation.

References: [1] Chiche L et al, A&R 2014

Acknowledgement: Dr Alaa Mohamed, Dr Ahmed Zayat and Dr Adewonuola Alase

Disclosure of Interest: None declared