



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/144435/>

Version: Accepted Version

Proceedings Paper:

De Vecchis, D, Reithmeier, RA and Kalli, A (2019) Molecular Simulations Reveal the Dynamics of the Band 3 Anion Transporter in a Model Native Red Blood Cell Membrane. In: Biophysical Journal. 63rd Annual Meeting of the Biophysical-Society, 02-06 Mar 2019, Baltimore, MD. Elsevier. Article no: 2253-Plat, 455A. ISSN: 0006-3495. EISSN: 1542-0086.

<https://doi.org/10.1016/j.bpj.2018.11.2456>

This article is protected by copyright. Licensed under the Creative Commons Attribution-Non Commercial No Derivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Molecular simulations reveal the dynamics of the band 3 anion transporter in a model native red blood cell membrane

Dario De Vecchis^{1,2}, Reinhart Reithmeier³ and Antreas Kalli^{1,2}

¹ Leeds School of Medicine, University of Leeds, Leeds, United Kingdom.

² Astbury Centre for Structural Molecular Biology, University of Leeds, Leeds, United Kingdom.

³ Department of Biochemistry, University of Toronto, Toronto, Canada.

d.devecchis@leeds.ac.uk

Band 3, the red blood cell anion exchanger (AE1/SLC4A1), is responsible for the rapid transport of bicarbonate and chloride across the red blood cell plasma membrane, a process necessary for efficient respiration. Human Band 3 is comprised of a cytosolic domain and a membrane domain that contains 14 transmembrane helices. Although structural data are available for both isolated domains, the structure of the complete Band 3 remains elusive. This is a limiting factor in the study of Band 3 anion transport and of its interactions with cytosolic proteins. By integrating molecular modelling and molecular dynamics simulations at the coarse-grained and all-atom resolutions, we have constructed a model of Band 3 that consists of both the transmembrane and the cytosolic domains. This model enabled us to identify the orientation of the cytosolic domain relative to the transmembrane domain and the role of the linker regions that connect these two domains in their interactions. Our model was validated using functional data and molecular simulations. Simulations were performed in complex bilayers that resemble the native red cell plasma membrane, containing a full complement of phospholipids, sphingomyelin and cholesterol. Our results provide novel molecular insights into the interactions of the lipid environment with Band 3. Specific lipids, e.g. cholesterol, have been found to localize in the Band 3 dimer interface possibly stabilizing the dimer. Specific lipid head groups were also found to interact with the cytosolic domain regulating its orientation relative to the transmembrane domain. Moreover, our modeling approach identified key residues involved in anion binding. Understanding the dynamics and interactions of Band 3 in a model native red cell membrane provides new insights into the function of this important human transporter.