

This is a repository copy of *Protecting women and children from second hand tobacco smoke: a public health priority*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/145726/>

Version: Published Version

---

**Article:**

Siddiqi, Kamran orcid.org/0000-0003-1529-7778 (Accepted: 2019) Protecting women and children from second hand tobacco smoke: a public health priority. *Pakistan Journal of Chest Medicine*. pp. 1-2. ISSN 2309-9844 (In Press)

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## Protecting women and children from second hand tobacco smoke; A public health priority

**Kamran Siddiqi**

Address for correspondence  
Department of Health Sciences,  
University of York, York, United Kingdom  
E-mail: kamran.siddiqi@york.ac.uk

This editorial may be cited as: Siddiqi K. Protecting women and children from second hand tobacco smoke; A public health priority. Pak J Chest Med 2019; 25(01): 1-2

**S**econd hand tobacco smoke (SHS) is a mixture of exhaled and side stream smoke released from a burning cigarette or other forms of tobacco such as shisha, cigar, pipe, bidi, etc. SHS contains 4,000 chemicals, 70 of which can cause cancer. Children and non-smoking adults exposed to SHS end up inhaling carcinogens and other toxic components that are present in it. SHS exposure is a serious health hazard to non-smokers, leading to an estimated 890,000 deaths and a loss of 10.9 million disability-adjusted life years (DALYs) globally every year.<sup>1</sup>

Women and children are worst affected by SHS exposure: 28% of deaths from SHS exposure occur in children.<sup>2</sup> SHS exposure impairs children's lung development and causes immune dysregulation; therefore, increasing their risk of acquiring lower respiratory tract infections,<sup>3</sup> tuberculosis, and incident cases, recurrent episodes, and exacerbations of asthma.<sup>4</sup> Parental smoking is also associated with an increased risk of their children's admissions to hospital.<sup>3</sup> Moreover, SHS exposure in children and adolescents leads to poor cognitive functions and academic achievements.<sup>5</sup> Children living in smoking households are at high risk of becoming adult smokers later.<sup>6</sup> SHS exposure during pregnancy is a key and avoidable risk to foetal development and contributes to adverse perinatal and postnatal outcomes, often with a lasting and negative impact during infancy and beyond.

Unfortunately, 40% of children are exposed to SHS worldwide amounting to a major public health threat.<sup>2</sup> Situation is worse in low- and middle-income countries where almost 56% children could be exposed to SHS.<sup>8</sup> This is in stark contrast to figures from the UK and Canada where only a minority of children are now exposed to such risks. Smoking is now less visible in public places, and most children and non-smoking adults are living in smoke-free environments. Consequently, hospital admissions due to chest infections and heart diseases have come

down too.

The south and south-east Asia region has the highest burden of disease attributable to SHS in the world. According to the Global Tobacco Surveillance System data and Demographic Health Surveys, the majority of women and children living in Bangladesh and Pakistan are exposed to SHS.<sup>9</sup> In 2014 in Pakistan, approximately 86% of people using restaurant and 76% using public transportation might have been exposed to SHS.<sup>10</sup> Nearly half of all adolescents (13-15 years old) in Pakistan could be exposed to SHS.<sup>8</sup> In a recent survey in 12 schools in Dhaka, Bangladesh, we found that 95% of 9-11 year old children had salivary cotinine levels consistent with recent exposure to SHS.<sup>11</sup> In addition to public places, children are also exposed to SHS in their private homes and cars. The prevalence of daily SHS exposure during pregnancy was found greater than active tobacco use in pregnancy across many low- and middle-income countries ranging from 6% (95% CI 5% to 7%) (Nigeria) to 73% (95% CI 62% to 81%) (Armenia).<sup>9</sup> In Pakistan, nearly 40% of pregnant women get exposed to SHS on a daily basis.<sup>9</sup>

If the above survey findings are a true reflection of the picture for most low- and middle-income countries including Pakistan, quite clearly, current measures are failing to protect the vast majority of women and children from SHS and the risks it poses. This level of exposure may well be contributing towards poor health and development of children in these countries

Smoking in indoor public spaces and workplaces is now banned in many countries, including Bangladesh and Pakistan. Where comprehensive and enforced, these bans have resulted in a significant reduction in SHS exposure and associated morbidity and mortality.<sup>12</sup> However, compliance to the smoke-free legislation is problematic in countries like Pakistan.<sup>1</sup> Smoking on public transport and in the workplace is still commonplace, and there is no restriction on smoking in the home. There is also little evidence on

the effectiveness of non-legislative interventions to protect women and children from SHS exposure.

Implementing smoke-free laws should remain a priority to curb exposure for non-smokers in low- and middle-income countries, including policies to promote smoke-free environments in multi-unit housing, in vehicles and in open public spaces such as children's play areas. Public awareness campaigns are also needed to raise awareness about the harms of SHS exposure in women and children. In low- and middle-income countries, nongovernmental organisations should support a grassroots movement to change smoking norms in communities.

An important first step would be to recognise that protecting women and children from exposure to SHS is key to improving maternal and child health. Healthcare professionals working with pregnant women and children should be aware of these issues and be proactive in offering advice and support to women and children who are exposed to SHS.

1. World Health Organization. WHO report on the global tobacco epidemic, 2017: Monitoring tobacco use and prevention policies. World Health Organization, 2017.
2. Oberg M, Jaakkola MS, Woodward A, Peruga A, Pruss-Ustun A. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet* (London, England). 2011; 377(9760): 139-46.
3. Cook DG, Strachan DP. Health effects of passive smoking: Summary of effects of parental smoking on the respiratory health of children and implications for research. *Thorax*. 1999; 54(4): 357-66.
4. Burr ML, Anderson HR, Austin JB, Harkins LS, Kaur B, Strachan DP, et al. Respiratory symptoms and home environment in children: a national survey. *Thorax*. 1999; 54(1): 27-32.
5. Chen R, Clifford A, Lang L, Anstey KJ. Is exposure to secondhand smoke associated with cognitive parameters of children and adolescents?--a systematic literature review. *Ann Epidemiol*. 2013; 23(10): 652-61.
6. Consortium PHR. A review of young people and smoking in England. 2009.
7. Rogers JM. Tobacco and pregnancy. *Reprod Toxicol*. 2009; 28(2): 152-60.
8. Xi B, Liang Y, Liu Y, Yan Y, Zhao M, Ma C, et al. Tobacco use and second-hand smoke exposure in young adolescents aged 12-15 years: data from 68 low-income and middle-income countries. *Lancet Glob Health*. 2016; 4(11): e795-e805.
9. Reece S, Morgan C, Parascandola M, Siddiqi K. Secondhand smoke exposure during pregnancy: a cross-sectional analysis of data from Demographic and Health Survey from 30 low-income and middle-income countries. *Tob Control*. 2018.
10. Saqib MAN, Rafique I, Qureshi H, Munir MA, Bashir R, Arif BW, et al. Burden of Tobacco in Pakistan: Findings From Global Adult Tobacco Survey 2014. *Nicotine Tob Res*. 2019; 21(1): 136.
11. Shah S, Kanaan M, Huque R, Sheikh A, Dogar O, Thomson H, et al. Secondhand Smoke Exposure in Primary School Children: A Survey in Dhaka, Bangladesh. *Nicotine & Tobacco Research*. 2017: ntx248-ntx.
12. Been JV, Nurmatov UB, Cox B, Nawrot TS, van Schayck CP, Sheikh A. Effect of smoke-free legislation on perinatal and child health: a systematic review and meta-analysis. *Lancet*. 2014; 383(9928): 1549-60.
13. Siddiqi K, Sarmad R, Usmani RA, Kanwal A, Thomson H, Cameron I. Smoke-free homes: an intervention to reduce second-hand smoke exposure in households. *Int J Tuberc Lung Dis*. 2010; 14(10): 1336-41.
14. Siddiqi K, Khan A, Ahmad M, Dogar O, Kanaan M, Newell JN, et al. Action to stop smoking in suspected tuberculosis (ASSIST) in Pakistan: a cluster randomized, controlled trial. *Ann Intern Med*. 2013; 158(9): 667-75.
15. Baxi R, Sharma M, Roseby R, Polnay A, Priest N, Waters E, et al. Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke. In: *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd, 2014.
16. Daly JB, Mackenzie LJ, Freund M, Wolfenden L, Roseby R, Wiggers JH. Interventions by Health Care Professionals Who Provide Routine Child Health Care to Reduce Tobacco Smoke Exposure in Children: A Review and Meta-analysis. *JAMA Pediatr*. 2016; 170(2): 138-47.
17. Rosen LJ, Myers V, Winickoff JP, Kott J. Effectiveness of Interventions to Reduce Tobacco Smoke Pollution in Homes: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health*. 2015; 12(12): 16043-59.