UNIVERSITY OF LEEDS

This is a repository copy of *Exposure to bioaerosols at open dumpsites: A case study of bioaerosols exposure from activities at Olusosun open dumpsite, Lagos Nigeria.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/147584/>

Version: Supplemental Material

Article:

Akpeimeh, GF, Fletcher, LA and Evans, BE orcid.org/0000-0001-9815-3141 (2019)
Exposure to bioaerosols at open dumpsites: A case study of bioaerosols exposure from activities at Olusosun open dumpsite, Lagos Nigeria. *Waste Management*, 89. pp. 37-47. ISSN 0956-053X

<https://doi.org/10.1016/j.wasman.2019.03.058>

© 2019. Published by Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

**Supplementary Information for: Exposure to Bioaerosols at Open Dumpsites:
A Case study of bioaerosols exposure from activities at Olusosun Open
Dumpsite, Lagos Nigeria.**

Table S.1. Media preparations and incubating conditions regarding sampled microorganisms

Table S.2. Meteorological conditions during ambient sampling

**Table S.3. Summary of the bioaerosol exposure assessment from activities at Olusosun
Dumpsite**

**Figure S.1. Waste sorting by one of the volunteers during Activity-based sampling with SKC
button sampler attached**

**Figure S.2. Comparison of exposure of (A) Total Bacteria, (B) Gram-negative (C) *Aspergillus
fumigatus* from activities at Olusosun dumpsite to environmental sampling.**

Table S1 Media preparations and incubating conditions regarding sampled microorganisms

Bioaerosols	Agar	Specific growth inhibitor	Incubation Temperature	Duration of incubation
Total Mesophilic bacteria	14 g L ⁻¹ nutrient agar and 10 g L ⁻¹ bacteriological agar	Cycloheximide, 100 mg L ⁻¹	37 °C	48 h
Gram negative bacteria	52 g L ⁻¹ MacConkey Agar	Cycloheximide, 200 mg L ⁻¹	37 °C in the dark	3 -7 days
<i>Aspergillus fumigatus</i>	20 g L ⁻¹ each of malt extract agar and bacteriological agar.	Streptomycin, 50 mg L ⁻¹ and Novobiocin, 10 mg L ⁻¹	40 °C	48 h
Total fungi	20 g L ⁻¹ each of malt extract agar and bacteriological agar.	Streptomycin, 50 mg L ⁻¹ and Novobiocin, 10 mg L ⁻¹	40 °C	48 h

Table S2 Meteorological conditions during ambient sampling

Visit days	Mean air Temperature (°C)	Mean Wind speed (m/s)	Relative humidity %
1	34.5 (32-37)	2.9	72.5
2	35.7 (32-38)	1.5	64.3
3	33.9 (32-36)	2.5	70.0
4	34.5 (32-37)	1.8	70.3
5	34.5 (32-36)	1.7	69.5
6	30.6 (29-32)	1.7	68.3
7	31.3 (30-32)	3.5	75.5
8	32.0 (29-34)	2.3	85.8
9	27.0 (25-29)	2.8	88.0
10	29.5 (28-30)	4.2	73.5
11	31.5 (31-33)	1.2	73.0
12	29.1 (28-30)	2.4	69.5
13	27.8 (26-30)	2.7	72.5

Table S3. Summary of the bioaerosol exposure assessment from activities at Olusosun Dumpsite

Variable	Scavengers	Waste sorters	Waste workers
Tidal volume (ml)	500		
Respiration rate	12 breaths per minute		
Working hours	11 hours		
Total volume of air inhaled (m ³)	3.96		
Total number of total bacteria inhaled (CFU)	1.9 x 10 ⁶	3.8 x 10 ⁵	9.5 x 10 ⁵
Total number of gram negative bacteria inhaled (CFU)	5.54 x 10 ⁶	3.17 x 10 ⁶	3.84 x 10 ⁵
Total number of <i>Aspergillus fumigatus</i> inhaled (CFU)	2.06 x 10 ⁵	2.74 x 10 ⁵	9.11 x 10 ⁴

Figure S2. Comparison of exposure of (A) Total Bacteria, (B) Gram-negative (C) Aspergillus fumigatus from activities at Olusosun dumpsite to environmental sampling.