

This is a repository copy of *Exploring the added value of mental health social work in multidisciplinary community teams: An analysis of a national service user survey*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/148119/>

Version: Accepted Version

---

**Article:**

Boland, Jennifer, Abendstern, Michele, Wilberforce, Mark Robson  
[orcid.org/0000-0001-6977-4483](https://orcid.org/0000-0001-6977-4483) et al. (3 more authors) (Accepted: 2019) Exploring the added value of mental health social work in multidisciplinary community teams: An analysis of a national service user survey. *Journal of Social Work*. ISSN 1468-0173 (In Press)

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Journal of Social Work

## Mental health social work in multidisciplinary community teams: An analysis of a national service user survey

Journal:	<i>Journal of Social Work</i>
Manuscript ID	JSW-18-0116.R2
Manuscript Type:	Original Article
Keywords:	Mental health, Community services, Social work values, Social work skills, Social work practice
Abstract:	<p>ABSTRACT</p> <p>Summary: The article addresses the continued lack of clarity about the role of the mental health social worker within CMHTs for working age adults and particularly the limited evidence regarding this from the perspective of service users. It compares findings from the literature, found to originate from a predominantly professional viewpoint, with secondary analysis of a national survey of service users to assess their views.</p> <p>Findings: Three particular aspects of mental health social workers' role identified in the literature were, to some extent, also located within the national survey and can be summarised as: approaches to practice, nature of involvement, and scope of support. The presence of these features was largely not substantiated by the survey results, with few differences evident between service users' experiences of mental health social workers compared with other mental health staff. When nurses and social workers were compared, results were either the same for both professions or favoured nurses. The findings point both to the difficulty of articulating the social work contribution and to the limitations of the secondary data.</p> <p>Application: The findings are a useful benchmark, highlighting the limited evidence base and the need for further research to improve both the understanding of the mental health social work role and how it is experienced by service users. The profession is keen to emphasise its specific contribution. Research evidence is required to underscore this and to ensure that the role is not subsumed within generic practice.</p>

SCHOLARONE™  
Manuscripts

**Table 1:** Service user characteristics by professional group

		Social worker n (%)	CMHN n (%)	$\chi^2$	p
<b>Gender</b>	Female	388 (57)	1430 (55)	.403	.544
	Male	294 (43)	1145 (44)		
<b>Respondent</b>	Service User	355 (54)	1685 (68)	63.63	<.001
	A Friend/Relative	174 (26)	419 (17)		
	Service User & Friend/Relative	58 (9)	251 (10)		
	Service User & Professional	66 (10)	124 (5)		
<b>Age</b>	18-35	98 (14)	339 (13)	3.43	.329
	36-50	168 (24)	615 (24)		
	51-65	194 (28)	687 (27)		
	66+	222(33)	934 (37)		
<b>Time in services</b>	< 1 Year	99 (15)	392 (15)	1.48	.686
	1-5 Years	239 (36)	869 (34)		
	6-10 Years	92 (14)	319 (13)		
	10+ Years	237 (35)	943 (37)		
<b>Last seen</b>	< Month	413 (61)	1867 (72)	39.45	<.001
	1-3 Months	156 (23)	443 (17)		
	4-6 Months	82 (12)	202 (8)		
	7-12 Months	31 (4)	63 (2)		
<b>Total</b>		695	2589		

**Table 2: Communication**

	Social workers n (%)	CMHN n (%)	$\chi^2$	p
<b>Did the person listen carefully to you?</b>				
	n=674	n=2542		
Yes definitely	485 (72)	1983 (78)	10.94	.004
Yes, to some extent	153 (23)	451 (18)		
No	36 (5)	108 (4)		
<b>Were you given enough time to discuss your needs and treatment?</b>				
	n=661	n=2490		
Yes definitely	437 (66)	1809 (73)	11.06	.004
Yes, to some extent	175 (25)	524 (21)		
No	49 (7)	157 (6)		
<b>Did the person or people you saw understand how your mental health needs affect other areas of your life?</b>				
	n=657	n=2478		
Yes definitely	410 (62)	1653 (67)	8.73	.013
Yes, to some extent	177 (27)	643 (26)		
No	70 (11)	182 (7)		

**Table 3: Co-production**

	Social workers n (%)	CMHN n (%)	$\chi^2$	p
<b>Have you agreed with someone from the NHS mental health service what care you will receive? (N=663/2516)</b>				
Yes definitely	363 (55)	1550 (62)		
Yes, to some extent	227 (34)	788 (3)	15.79	.000
No	73 (11)	178 (7)		
<b>Were you involved as much as you wanted to be in agreeing what care you will receive? (n=577/2315)</b>				
Yes definitely	332 (57)	1459 (63)		
Yes, to some extent	207 (36)	709 (31)	11.98	.007
No, but I wanted to be	26 (4)	125 (5)		
No, but I did not want to be	12 (2)	22 (1)		
<b>Does this agreement on what care you will receive take your personal circumstances into account? (N=581/2294)</b>				
Yes definitely	373 (64)	1540 (67)		
Yes, to some extent	178 (31)	655 (27)	2.04	.361
No	30 (5)	99 (4)		
<b>In the last 12 months have you had a formal meeting with someone from the NHS mental health services to discuss how your care is working? (N=553, N=2066)</b>				
Yes	474 (86)	1752 (85)		
No	79 (14)	314 (15)	0.29	.593
<b>Were you involved as much as you wanted to be in discussing how your care is working? (N=462/1729)</b>				
Yes definitely	289 (63)	1178 (68)		
Yes, to some extent	143 (31)	451 (26)	5.70	.127
No, but I wanted to be	26 (6)	91 (5)		
No, but I did not want to be	4 (1)	9 (1)		
<b>Did you feel the decisions were made together by you and the person you saw during this discussion? (N=456/1716)</b>				
Yes definitely	278 (61)	1138 (66)		
Yes, to some extent	144 (32)	465 (27)	6.53	.088
No, but I wanted to be	27 (6)	100 (6)		
No, but I did not want to be	7 (2)	13 (1)		
<b>Have the NHS mental health services involved a member of your family or someone else close to you as much as you would like? (N=602/2246)</b>				
Yes Definitely	308 (51)	1187 (53)		
Yes, to some extent	126 (21)	400 (18)		
No, not as much as I would like	60 (10)	206 (9)	8.03	.154
No, they have involved them too much	6 (1)	43 (2)		
They did not want to be involved	24 (4)	70 (3)		
I didn't want them involved	78 (13)	340 (15)		

**Table 4:** Finance and employment advice

	Social workers n (%)	CMHN n (%)	$\chi^2$	p
<b>In the last 12 months, did the NHS mental health services give you any help or advice finding support for financial advice or benefits?</b>				
	N=648	N=2467		
Yes definitely	208 (46)	682 (48)	2.7 4	.254
Yes, to some extent	137 (30)	371 (26)		
No, but I would have liked help	107 (24)	356 (25)		
<b>In the last 12 months, did the NHS mental health services give you any help or advice with finding support for finding or keeping work?</b>				
	N=361	N=1332		
Yes definitely	83 (40)	254 (41)	1.5 1	.470
Yes, to some extent	74 (36)	197 (31)		
No, but I would have liked help	51 (24)	175 (28)		

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Mental health social work in multidisciplinary community teams: An analysis of a national service user survey**

For Peer Review

## Introduction

Social workers have played a major role in the development of community mental health services for adults in England. Deinstitutionalisation policies from the mid-20<sup>th</sup> Century demanded enhanced community support services (Department of Health [DH], 1962; Mental Health Act, 1959), which in turn drew on key social work values linking successful support with the need for understanding of the dynamics between patients, their families, communities and wider social forces. Additionally, social work's long history of individual casework orientated to people in their own homes and communities naturally lent itself to the needs of new services. More specifically, the 1959 Mental Health Act gave additional responsibilities to social workers including post hospital discharge support (Burns, 2014). A decade later, the Medical-Psychological Association (1969) recommended that a new body of mental health social workers with additional training was now required to support the growing numbers of people with profound and complex mental health needs living in the community (Godin, 1996). The introduction of the Approved Social Worker in 1983 (MHA, 1983) conferred on this group of staff further duties and responsibilities to conduct assessments where formal detention was considered. Within this role their duty included investigating the feasibility of community alternatives to avoid hospital admission (Rapaport, 2005). These factors reinforced the shift away from genericism in the social work role (Challis & Ferlie, 1987, 1988). Furthermore, government policies which sought to reorient mental health services towards care in the community often included an enhanced role for social workers within integrated services (e.g. DH, 1975, 1989, 1990, 1995, 1998; Health Act, 1999).

New approaches to support involved multidisciplinary services and focused on early intervention and the maintenance of independence (Anthony, 1993; Hibbard & Gilbert, 2014). Today such services are the norm, with social work joining psychiatry, nursing, psychology and occupational therapy in a spectrum of specialist community teams (Malone, Marriott, Newton-Howes, Simmonds & Tyrer, 2007). Such teams are increasingly prevalent across Europe, North America, and Australasia (Draper & Anderson, 2010; Evans et al., 2012; Ng, Herrman & Chiu, 2009). These multi-disciplinary teams encompass the early assessment and diagnosis of psychiatric conditions and the coordination of long-term support and care to meet specific needs.

However, as social workers have been included within the wider mental health system, boundaries between professionals have blurred with a 'creeping genericism' gradually eroding traditional roles (Brown, Crawford & Darongkamas, 2000, p426). Role blurring and the



1  
2  
3 erosion of traditional professional practices have become a salient issue for many practitioners  
4 (Jones, 2014). This is evidenced, for example, by clinical psychologists helping to organise  
5 accommodation for service users, and social workers implementing psychotherapeutic  
6 interventions (Abendstern et al., 2014; Brown et al., 2000; Wall, 1998). The increasing  
7 pressure on social workers and other members of community mental health teams (CMHTs)  
8 for adults to move towards more generic roles has furthered a lack of clarity regarding what  
9 social workers should do compared to other professionals, whilst roles that were specific to  
10 them historically, such as the approved social work role, have been opened up to others  
11 (Bailey & Liyanage, 2012). Social work has long been recognised as difficult to define (Allen,  
12 2014; Howe, 1979; Rode, 2017) and more recently role blurring has added to the challenge  
13 of articulating its unique contribution to mental health.  
14  
15  
16  
17  
18  
19  
20  
21

22 This poses distinct challenges for the practice and organisation of mental health social work.  
23 First, service users may be unclear about the roles and remit of social work in their mental  
24 health care, which may undermine confidence and impede the contribution that social workers  
25 can then make (ComRes, 2017). Second, repeated studies have found that mental health  
26 social work staff in multidisciplinary environments have relatively poor job satisfaction and face  
27 significant risk of stress (Evans et al., 2005; Onyett 2011). This is empirically linked to social  
28 worker perceptions that their core skills and knowledge are not well matched with those  
29 demanded in their role (Wilberforce et al., 2013). Third, there appears to be a growing trend  
30 towards the removal of mental health social workers from multidisciplinary environments, at  
31 least based on anecdotal reports (ADASS 2018; Lilo 2016; McNicoll, 2016), for fears that  
32 social workers are not being utilised appropriately. Such decisions are (inevitably) being taken  
33 without appropriate evidence asserting their unique role.  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 This article aims to articulate the unique contribution of the social work role in mental health  
44 through a synthesis of two processes. First a focussed review of the literature was undertaken  
45 to identify the features of the social worker role in mental health care. Second, a new analysis  
46 of nationally collected data from the Care Quality Commission (CQC) was employed to identify  
47 service user perspectives of social work in mental health. These data enabled a comparison  
48 to be made between the experiences of service users supported by social workers and those  
49 supported by other professionals.  
50  
51  
52  
53  
54  
55  
56

## 57 **Methods**

### 58 ***Review of literature***

1  
2  
3  
4  
5 This review sought to identify and synthesise the literature on the distinct contribution of the  
6 social work role in mental health. It was designed to serve the needs of a wider research  
7 study investigating the contribution of social work in community mental health teams  
8 (underway at the time of writing, and which required the collation of a list of attributes to  
9 incorporate into data collection tools). 'Contribution' was defined by the research team as  
10 including both 'what' they do and 'how' they do it, acknowledging that social workers may do  
11 similar tasks to other professions, but in a different way. The pragmatic aims meant there was  
12 no requirement for the review to be exhaustive. Nevertheless good practice in literature  
13 reviewing was followed drawing upon rapid review methods as a means of expediting the  
14 identification and synthesis of existing literature. Whilst no formal definition for a rapid review  
15 exists, the process adopted used Tricco and colleagues (2015) working definition which states  
16 that they are "a type of knowledge synthesis in which components of the systematic review  
17 process are simplified or omitted to produce information in a short period of time" (Tricco et  
18 al., 2015, p225). In line with this this, they tend to be characterised by a restriction of searches  
19 to one or two databases, limiting search terms, and a presentation of results within a narrative  
20 summary with no quality appraisal.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 Three different sources of literature were reviewed. The first identified six authoritative  
32 textbooks which described mental health practice and generic social work (Briar & Miller, 1971;  
33 Davies, 2012; Goldstein, 1973; Golightley & Geomans, 2014; Moxley, 1989; Raiff & Shore,  
34 1993). These texts were selected by two of the authors (DE, JH), with professional social work  
35 qualifications and the most extensive experience from within the research team, making a  
36 pragmatic choice from existing textbooks known to them. The second consisted of existing  
37 research and practice documents known to the authors to be closely aligned to the intent of  
38 the present review, that is, the nature of social work practice today (e.g. All Party Parliamentary  
39 Group 2016; Moriarty & Manthorpe, 2016). Finally a bespoke literature search of two  
40 databases was undertaken (Web of Science and PsychInfo) using the search terms "social  
41 work\*" AND (role OR function) AND "community mental health", restricted to the period 1999  
42 to 2017. Broad inclusion criteria enabled the capture of evidence of social work attributes  
43 within both generic and specialist mental health social work. All included texts were required  
44 to include descriptions of one or more social work attribute, in line with the aim of the review  
45 which sought to collate a list of such features. This yielded 85 references after duplicates were  
46 removed. Titles and abstracts were reviewed by all the research team and any abstract  
47 identified as relevant by any member (n=44) were obtained, bar five items that could not be  
48 sourced. Those with exclusive focus on older adults services were also excluded (n=2).  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 For each included source, a short summary narrative of social workers' contributions was  
4 produced and entered into MS Excel (RP). These were then discussed by the wider authorship  
5 group who, through an iterative process, identified three broad areas under which the findings  
6 were collated: 'approaches to practice'; 'nature of involvement'; and 'scope of support'. Within  
7 these headings, RP and MW jointly devised sub-themes by reviewing all Excel entries, revising  
8 and updating these as the analysis progressed. The final step involved providing a narrative  
9 commentary of each theme and sub-theme, with an example of social work practice illustrating  
10 each in practice. The aim was to draw out distinctions where possible. Decisions were  
11 pragmatic rather than definitive with overlap acknowledged, indicative of the characteristics of  
12 social work practice.  
13  
14  
15  
16  
17  
18  
19  
20

### 21 **Secondary data: Community Mental Health Service User Survey 2016**

22  
23  
24 The CQC (2016) annual Community Mental Health Survey data were selected for analysis, as  
25 a standardised survey with national coverage. Its primary aim is to find out what service users  
26 think about the NHS healthcare services they use, to highlight good care and to identify the  
27 potential risks to the quality of services. The survey consists of 47 questions with 41 of these  
28 asking about the service user's specific experience of the care they receive. Twelve of these  
29 questions were analysed, being those most relevant to the core aim of the article: to articulate  
30 the social work role within these services (see Tables 2-4 for details). These were organised  
31 according to the literature review themes: approaches to practice (n=3); nature of involvement  
32 (n=7); and scope of support (n=2). The responses are collected using Likert-type categories.  
33  
34 The survey also asks who is the main person organising the service user's care, providing an  
35 opportunity for comparison between social workers and community mental health nurses  
36 (CMHNs) as care coordinators.  
37  
38  
39  
40  
41  
42  
43  
44

45 **Settings:** All Trusts providing community mental health services in England were eligible to  
46 take part in the survey. Fifty eight providers of NHS mental health services in England,  
47 including combined mental health and social care trusts, foundation trusts and community  
48 healthcare social enterprises commissioned by Trusts, provided mental health services.  
49  
50 Fieldwork for the survey took place between February and June 2016.  
51  
52  
53

54 **Respondents:** Each NHS Trust providing or commissioning mental health services drew a  
55 random sample from their records of 850 people receiving services. Service users were  
56 eligible to complete the Community Mental Health User Survey if they were over 18 years of  
57 age and had received specialist care or treatment for a mental health condition from a  
58  
59  
60

1  
2  
3 community-based treatment or care service, delivered through a Mental Health Trust during  
4 the sampling period. This also included those who received care under the Care Programme  
5 Approach (CPA). Several exclusions were applied by the authors, including specialist services  
6 for people with learning disabilities, drug and alcohol problems, and forensic psychiatry.  
7  
8 Further, those only seen for assessment; those who were inpatients at the time of the survey;  
9  
10 and people seen exclusively by Improving Access to Psychological Therapies services were  
11  
12 excluded. Full details are provided in CQC (2016).  
13  
14  
15

16  
17 The CQC dataset comprised 12,522 people who had seen a mental health practitioner in the  
18 previous 12 months; 2,739 of whom were care coordinated by a CMHN and 802 by a social  
19 worker. Some respondents perceived that more than one professional acted as care  
20 coordinator and were removed from the analysis. The final sample comprised 2,575 and 682  
21 people whose care was coordinated by a CMHN and social worker respectively.  
22  
23  
24  
25

26  
27 **Analysis:** Data analysis on the selected CQC questions, which were grouped into three  
28 themes, was conducted using SPSS statistical software. Effect sizes were also calculated,  
29 however, only as a guide since the data were categorical not continuous. This permitted  
30 quantification of the difference between two groups, and is helpful in large samples where  
31 small differences may be statistically significant. The effect size is the standardised mean  
32 difference between the two groups. By convention, an effect size of 0.2 or less is 'small'  
33 (Cohen, 1977).  
34  
35  
36  
37  
38

## 39 **Results**

40  
41  
42 Following a description of the characteristics of the literature and survey sample, the results  
43 are described under three sub-headings, each of which contains findings from the literature  
44 followed by the results of the secondary data analysis of the related survey questions. A final  
45 section about overall satisfaction contains survey data only.  
46  
47  
48  
49

### 50 **Literature characteristics**

51  
52 Together the three sources produced 59 articles and books published between 1971 and  
53 2016. The majority (n=32) came from the UK with a substantial number (n=17) being published  
54 in the US. A smaller amount came from a range of European countries (n=10). The  
55 descriptions of social worker contributions were organised under the three interlinked themes  
56  
57  
58  
59  
60

1  
2  
3 already noted. The CQC data was then also arranged under those headings. Each theme  
4 contained a number of sub-themes which are outlined below. The included literature was  
5 dominated by non-empirical texts in the form of published books and articles (n=28) and grey  
6 literature (n=11). Twenty empirical research articles were included of which 13 reported data  
7 collected from social workers. Only three reported the perspectives of service users. Unless  
8 otherwise stated, the data reflects a mix of empirical research and opinion from social work  
9 experts within academia and/or policy environments.  
10  
11  
12  
13  
14  
15

### 16 ***Survey sample characteristics***

17  
18  
19 Service user characteristics are displayed in Table 1. There were no significant differences  
20 between the two groups (those supported by a social worker and those supported by a CMHN)  
21 with regard to gender, age or time spent in services. However, on average service users on  
22 CMHN caseloads had been seen more recently compared to service users on social worker's  
23 caseloads. Also significantly more service users on CMHN caseloads had completed the  
24 questionnaire themselves, compared to service users on social workers caseloads.  
25  
26  
27  
28  
29

30 <Insert Table 1 about here >  
31  
32

### 33 ***Approaches to practice***

34  
35  
36 ***The literature:*** Three areas were identified within the literature on this theme: that social work  
37 theory and practice is situated within an understanding of society as socially biased against  
38 the vulnerable; starts from a holistic perspective; and prioritises good working relationships  
39 with individuals to support positive change.  
40  
41  
42  
43

44  
45 Social work practice in mental health and other services was reported to be intentionally non-  
46 neutral; framed by an assumption that people with mental health problems are vulnerable to  
47 abuses of their human rights (Ife, 2012) and face greater difficulties in accessing health and  
48 welfare services, education and training, employment, housing, and participation in civic  
49 society (Ahmedani, 2011). Further, discrimination due to mental health was recognised in  
50 social work texts as inseparable from other forms of injustice, for example with regards to  
51 ethnicity and culture or sexual and gender identities (Allen, 2014; Faust, 2008; Golightley &  
52 Geomans, 2014; Ramon, 2010).  
53  
54  
55  
56  
57

58  
59 Social workers were also said to be guided by an awareness and understanding of how  
60 individual wellbeing is inextricably linked to their social environment (Goldstein, 1973; Raiff &

1  
2  
3 Shore, 1993). As a consequence social work has been identified as being cautious about the  
4 medical model of psychiatry, as insufficient to explain causes and consequences of mental  
5 health problems (Carpenter, Schneider, Brandon & Wooff, 2003), described as impeding  
6 mental health recovery due to an overriding focus on deficits' alone (Davies, 2012; Ramon,  
7 2010; Stanley et al., 2003; Stromwall & Hurdle, 2003). A holistic approach that takes into  
8 account a persons' wider needs and social context was said to be valued by service users  
9 (Beresford, 2007).

10  
11  
12  
13  
14  
15  
16 More generally, the social worker's approach was described as prioritising a positive working  
17 relationship with clients and their families built on compassion (Ramon, 2010), trust and clear,  
18 uncomplicated, communication (Allen, 2014; Golightley & Geomans, 2014; Hardiker & Barker,  
19 1999; Herman, 2014; Peck & Norman, 1999). Mental health social work training has long-  
20 included relationship work as one component of their duties (Perlman, 1979). More recently,  
21 this has received attention as part of recovery principles, allied with concepts of 'hope',  
22 'strengths' and 'control' to improve social functioning and promote engagement in the wider  
23 community (Allen, 2014; Pahwa, Smith, McCullagh, Hoe & Brekke, 2016). An emphasis  
24 on self-awareness including limited self-disclosure (Golightley & Geomans, 2014) and the  
25 ability to actively listen and empathise with service users (Faust, 2008; Penhale & Young,  
26 2015) were also skills acknowledged to be required to build positive working relationships.  
27 Two publications which focussed on users' views stressed that the social worker's approach  
28 to practice, including kindness, sensitivity, reliability and a non-judgemental attitude, was  
29 paramount to service users' satisfaction with social work (Beresford, 2007; Penhale & Young,  
30 2015).

31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41 **The secondary data:** The three questions within this theme spanned careful listening,  
42 whether service users were given enough time, and how well they thought they were  
43 understood by their key worker. Descriptive statistics for these questions are displayed in  
44 Table 2. There were significant differences for all three questions, with respondents in the  
45 CMHN group answering more positively compared to those in the social worker group,  
46 although the effect sizes were small (all  $d \leq 0.13$ ).

47  
48  
49  
50  
51  
52 <insert table 2 about here>

### 53 54 55 56 **Nature of involvement**

57  
58  
59  
60

1  
2  
3 **The literature:** This theme includes the sub-themes of advocacy; anti-oppressive practice;  
4 and the exercise of care coordination. Advocacy has been described as a routine element of  
5 social work ensuring that service users' rights are upheld and respected (Cummings & Cassie,  
6 2015; Davis & Jung, 2012; Manktelow et al., 2002). Social workers were characterised in the  
7 literature as promoting social justice, giving the powerless a voice (Faust, 2008) and  
8 supporting people to express themselves so that they could be "recognised on equal terms  
9 with others" (Parrott, 2014: p105). Anti-oppressive practice in social work, linked to  
10 challenging discrimination along all lines of difference (Beresford, 2007) has emerged from  
11 social work training that articulated theories and practices related to resolving differential  
12 power relationships within families, social networks, public services, and communities.  
13  
14  
15  
16  
17  
18  
19

20  
21 One of the key vehicles through which social workers engage service users and their social  
22 networks in promoting self-determination is through their involvement in care coordination,  
23 which in social work is also geared towards changing the power balance between the  
24 supported and those supporting them (Herman, 2014; Penhale & Young, 2015; Ring, 2001).  
25 In terms of assessment activity, social workers were found to make key links between  
26 psychiatric, psychological and social functioning together with reviewing risk and physical  
27 health needs (Aschbrenner et al., 2015; DH, 1999). In relation to care planning and  
28 coordination, social workers play a key role in creating comprehensive and personalised care  
29 plans which reflect an individual's needs, preferences and strengths and enable individuals to  
30 live more independently (Allen, 2014; Raiff & Shore, 1993). This role is reported to involve  
31 arranging, purchasing and monitoring social care packages and referrals (Marshall, Lockwood  
32 & Gath, 1995; Moxley, 1989) and therefore involves liaising, mediating and negotiating with  
33 other professionals and agencies to ensure continuation of care (Cummings & Cassie, 2015;  
34 Janlov et al., 2015). Social workers were also reported to liaise closely with nurses, care  
35 agencies and voluntary organisations, GPs and hospitals, and specialist psychological support  
36 services (Golightley & Geomans, 2014) to coordinate care and ensure service users' and  
37 carers' needs are appropriately met (Hardiker & Barker, 1999).  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 Mulhall (2000) found that service users wanted to be treated with respect, to be involved in  
50 planning their own care and to be listened to, all of which have been identified as core skills  
51 of social workers. Additionally, Beresford (2007) reported that service users valued social  
52 workers for their focus on supporting independence and participation rather than dependence.  
53  
54  
55

56  
57 **The secondary data:** Seven questions were linked to the theme of coproduction covering the  
58 level of involvement of the service user and their wider network in planning and reviewing their  
59  
60

1  
2  
3 care. The questions focused on the extent of involvement in discussions about care needs,  
4 formulating plans, and agreeing decisions. Descriptive statistics are displayed in Table 3.  
5  
6  
7

8 Significant differences emerged in relation to the level of involvement in agreeing the care the  
9 service user would receive, with service users on the CMHN caseload seeming more satisfied,  
10 although effect sizes were small ( $d_s \leq 0.12$ ). No other significant differences were found  
11 within this theme, suggesting that both CMHNs and social workers involved the service users  
12 and their wider networks in decisions surrounding their care to the same extent.  
13  
14  
15  
16  
17

18 <Insert table 3 about here>  
19  
20

### 21 ***The scope of social work support***

22  
23  
24  
25 ***The literature:*** The literature described three elements of support provided by social workers:  
26 knowledge of and ability to access a broad range of resources; direct interventions; and  
27 statutory roles requiring specific knowledge and skills.  
28  
29  
30

31 A central contribution of social work to mental health care, as noted above, is its wide frame  
32 of reference compared to a more medicalised model: if a person is to be viewed holistically,  
33 then the range of support must not be constrained to clinical resources. Social workers were  
34 found to be knowledgeable about services available through the local authority, including  
35 social care and housing (Mitchell & Patience, 2002). King and colleagues' (2002) survey  
36 comparing different professionals in community mental health services found social workers  
37 to have significantly greater knowledge of employment support than other staff. Social workers  
38 were also found to routinely liaise with wider groups including the police and offenders'  
39 services, immigration, jobcentres, benefits support and local community support groups  
40 spanning a range of potential needs (Allen, 2014; Stromwall & Hurdle, 2003). Outreach work  
41 took social workers into hospitals, jails, and communities where long-term goals were created  
42 based on the individual's stage of readiness (Dumaine, 2003). To this end, social workers  
43 were reported to be adept at multidisciplinary working, and to offer unified and integrated  
44 services that enable individuals with mental health needs to improve their social and  
45 community functioning (All Party Parliamentary Group, 2013; Stromwall & Hurdle, 2003).  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

56 Social workers were reported to implement a spectrum of interventions (All Party  
57 Parliamentary Group, 2016). Practical interventions dominated accounts in the literature, such  
58 as Priebe and colleagues' (2005) who found that 82 per cent of social workers in London  
59  
60



1  
2  
3 reported that support in, and training of, daily living skills was one of their main roles. Other  
4 interventions targeted social functioning with the aim of improving engagement in the  
5 community and enabling individuals to enter meaningful vocations (Pahwa et al., 2016;  
6 Ramon, 2010; Stromwall & Hurdle, 2003). Social workers used psychological interventions  
7 (Davis & Jung, 2012), including counselling (Beresford, 2007; Lang et al., 2011; Peck &  
8 Norman, 1999), psycho-education around medication effectiveness and side effects (Davies,  
9 2012; Pahwa et al., 2016), and emotional support to individuals in crisis (Marshall et al., 1995;  
10 Raiif & Shore, 1993).

11  
12  
13  
14  
15  
16  
17 The literature also described the statutory roles of social workers within mental health  
18 (Golightley & Geomans, 2014; Ramon, 2010). It illustrated how social workers exercised  
19 professional judgement over ethical dilemmas and risk when supporting individuals and  
20 families with the most serious needs, spanning safeguarding, domestic abuse, child  
21 protection, criminality, homelessness and substance use issues (Goldstein, 1973; Gould,  
22 2016; Rubin & Parrish, 2012). Social workers were reported to require assessment and  
23 decision making skills under circumstances where full information was either not available,  
24 was uncertain, and/or within fast-moving and volatile situations (Davies, 2012).

25  
26  
27  
28  
29  
30  
31  
32 One study by Cree and Davis (2007) highlighted service users' views on the scope of mental  
33 health support from social workers. The authors conducted four service user and two carer  
34 interviews about the social worker input into their care. They found that both the service users  
35 and carers identified that social workers liaised with other services on their behalf, introduced  
36 other treatment options, for example, Cognitive Behavioural Therapy, and involved family  
37 members where appropriate

38  
39  
40  
41  
42 **The secondary data:** This theme was limited to two questions regarding the scope of support  
43 service users received in relation to financial and employment advice. Descriptive statistics  
44 can be found in Table 4. For both questions, no significant differences emerged, ( $ds \leq 0.02$ ),  
45 suggesting both social workers and CMHNs provided the same level of support.

46  
47  
48  
49  
50 **Overall Satisfaction:** The CQC questionnaire also included a question about service users'  
51 overall satisfaction with their experience of NHS mental health services, using a 10 point Likert  
52 scale ranging from 0 (I had a very poor experience) to 10 (I had a very good experience). A t  
53 test revealed a significant difference between CMHNs and social workers  $t(3048) = 3.75, p <$   
54  $.001$ . Service users on CMHN caseloads rated their experience as significantly higher than  
55 service users on social workers' caseloads.

1  
2  
3 <Insert table 4 about here>  
4  
5

## 6 **Discussion**

7

8  
9  
10 Despite decades of debate about the importance of social work in mental health, the  
11 profession appears to have made little progress in establishing a clear evidence-base for its  
12 role. This article reviewed literature on the mental health social work role and provided a new  
13 analysis of secondary data on service user perceptions thereof. The discussion considers the  
14 implications of the findings for mental health social work going forward, focusing on the  
15 importance of developing a clearer role definition which can be understood by all, including  
16 service users, alongside a fuller comprehension of the service user perspective of this. The  
17 strengths and limitations of the data used for this study are also explored.  
18  
19  
20  
21  
22

### 23 ***Social work today and the service user voice***

24

25  
26  
27 The review of the literature undertaken for this study indicated that social workers operated  
28 within a value-based approach recognising societal influences on the individual; perceived the  
29 promotion of self-determination of vulnerable individuals as central to their work; and  
30 undertook a broad range of support including advocacy; direct interventions and the ability to  
31 access others, alongside statutory responsibilities. Social workers were found to recognise  
32 the importance of the individual participating as fully as possible in decision-making (Golightley  
33 & Geomans, 2014; Herman, 2014; Penhale & Young, 2015; Ring, 2001); to play a key role in  
34 creating comprehensive and personalised care plans, reflecting individuals' needs and  
35 preferences (Allen, 2014; Raiff & Shore, 1993); and to understand the need to develop trusting  
36 relationships to support these ends (e.g. Allen, 2014; Beresford, 2007).  
37  
38  
39  
40  
41  
42  
43

44 These principles are the foundations of current social work training in England, with the  
45 ThinkAhead initiative being one example of the drive to promote graduate entry to mental  
46 health specialist training (Clifton & Thorley, 2014). Its publicity highlights key features of social  
47 work including building relationships with people, providing guidance and therapy, arranging  
48 support and care, ensuring people's safety, standing up for people's rights, and improving  
49 community services. They describe the role of a mental health social worker as someone who  
50 empowers individuals through therapy, support and advocacy, building resilience in  
51 individuals, their networks, and their communities, thus transforming people's wellbeing and  
52 improving our society and economy (ThinkAhead.org, 2018).  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The evidence presented in this article identifies that within the literature whilst social workers,  
4 educators and other professionals are (broadly) able to discern the unique contribution social  
5 work provides as part of community teams (see also ADASS 2018), there is a surprising lack  
6 of testimony or articulation of the service user perspective in relation to their role in mental  
7 health care. Interestingly, even the Barclay Report of the 1980s on Social Workers' Roles and  
8 Tasks (Barclay, 1982) makes no mention of mental health care in their chapter on views of  
9 social work. Given the profession's position as advocating for, and empowering, the service  
10 user, these findings are puzzling. This is not unique to mental health. Penhale and Young's  
11 (2015) review of research spanning service user views of social work in general found a  
12 paucity of such evidence but it was notable that evidence specific to mental health services  
13 was even scarcer. Two reports that did focus on users' views, although not on mental health  
14 services specifically, recorded a range of attributes that service users valued and which they  
15 identified with social workers (Beresford, 2007; Penhale & Young, 2015). These studies, which  
16 are by no means definitive, mirrored the professional perspective of the social worker's unique  
17 contribution and included recognising and respecting diversity; seeing the client as a unique  
18 individual with unique needs; being non-judgemental; and being trustworthy and honest.

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Despite the paucity of service user-based research, there is also an argument that existing  
data is not used to its full capability. In this article, secondary analysis of the CQC Community  
Mental Health Survey data was undertaken, in part to redress the imbalance found in the  
literature. It enabled a comparison of those supported by a social worker with those supported  
by a mental health nurse. Perhaps surprisingly, minimal evidence was found to support the  
views identified in the literature and noted above as being particular to social workers. Indeed,  
where differences between mental health nurses approaches and those of social workers did  
emerge, they largely favoured nurses. This merits some reflection: why does the unique  
contribution of social work articulated in the literature not percolate through to evidence of  
service user experiences? What is obscuring its visibility?

First, there continues to be widespread misunderstanding of the social work role which may  
influence service user expectations. For example, ThinkAhead recently commissioned the  
polling company ComRes to find out what the public thought social workers did. They  
interviewed 2,033 adults online across Great Britain, in March 2017. Key findings included that  
only 41 per cent thought that social workers were an important provider of mental health  
support, that the most likely type of support provided by social workers to people with mental  
health conditions was to assess practical needs (65%); and that only 33 per cent thought that  
social workers were involved in the detention of individuals under the Mental Health Act  
(ComRes, 2017). This is not a new debate: that social work struggles to make clear its

1  
2  
3 purpose to the general public has been discussed at length, both at home and abroad  
4 (Barclay, 1982; LeCroy & Stinson, 2004). However the implications may be profound. Service  
5 user misunderstanding can undermine confidence that social workers can help them; can  
6 affect the social worker's own belief in their capacity to make a difference; and together form  
7 a self-fulfilling prophecy (Legood et al., 2016).  
8  
9  
10

11  
12 A second reason may be that social work in community mental health is culturally and  
13 numerically subsumed within a health-dominated framework. Mental health social workers  
14 remain a minority within a medically dominated workforce (Evans et al., 2012) meaning that  
15 when teams and services are faced with managing crises in an increasingly austere  
16 environment it might become more difficult for social workers to "argue the importance of ... a  
17 person's right to accommodation, building social networks and buffers, or the use of social  
18 interventions" (Woodbridge-Dodd 2017, p3). This view is also supported by evidence from  
19 social workers themselves who have described themselves as being isolated within NHS  
20 Trusts (Morriss, 2016). The same study found that those who did not have social worker  
21 managers were described as being unable to make their contribution visible through  
22 supervision. This was corroborated by earlier research, albeit with CMHTs for older people,  
23 where social workers supervised and managed by non-social workers were reported to feel  
24 less well understood and their contribution less valued and supported when compared with  
25 those managed and supervised by social workers (Abendstern et al., 2014).  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 Evidence also suggests that within multidisciplinary teams social workers are increasingly  
37 seen as generic mental health professionals whose roles overlap with other professionals  
38 more than in the past (Wilberforce et al., 2013). These trends are exemplified in legislative  
39 changes, whereby approved mental health professionals (AMHPs) have replaced the  
40 approved social worker role, although in practice 95 per cent remain social workers (ADASS,  
41 2018). Interestingly, Beresford (2007) reported an expression of concern from service users  
42 regarding a possible reduction in the helpful practice provided by social workers in this field  
43 with the ending of the approved social worker role. This provides some evidence or suggestion  
44 of service user recognition of difference between the approaches taken and roles of different  
45 professional groups and a preference for those of social workers.  
46  
47  
48  
49  
50  
51  
52  
53

### 54 ***Study limitations***

55  
56  
57 A number of study limitations must be acknowledged. The literature review was not fully  
58 systematic and its findings must therefore be treated with some caution. In particular, the six  
59  
60

1  
2  
3 authoritative textbooks were personally selected and others might have been chosen (e.g.  
4 Karban, 2011; Tew, 2011). Data extraction was also limited to the collection of social work  
5 attributes rather than delineating the voices from whence these data came. The narrative  
6 approach to reporting these data, nonetheless, helped to identify particular dimensions of  
7 practice for further analysis via the secondary data. These data, however, also had various  
8 shortcomings. Firstly, it is possible that service user questionnaires are not sensitive enough  
9 to detect experiential differences between being supported by a social worker or other  
10 professional. Additionally, the researchers had no control over what questions were asked in  
11 the survey and consequentially their mapping to the literature was approximate. The true  
12 distinctiveness of social work is perhaps more nuanced than the survey questions would allow.  
13 If there is a distinction to be detected, other research methods may be necessary to draw this  
14 out. Secondly and more specifically, the CQC questionnaire was not part of a controlled  
15 experiment and therefore differences in case mix supported by different professional groups  
16 that might impact on the findings could not be measured. For example, other research has  
17 demonstrated that social workers often work with different groups to CMHNs, including people  
18 with the most complex needs and circumstances (Allen, 2014; Huxley and Kerfoot, 1993;  
19 Penhale & Young, 2015). One study found that social workers in CMHTs tend to carry  
20 caseloads of those with higher levels of severity of mental illness and impairment than CMHNs  
21 (Huxley et al., 1998). This is not inconsistent with the fact that fewer of the social worker  
22 supported respondents had self-completed the survey. Such a difference in case mix could  
23 affect the perceived satisfaction of service users thereby confounding attempts to compare  
24 experiences between respondent groups.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39  
40 Thirdly, the CQC data also incorporated a range of different services, including crisis teams,  
41 recovery teams, and outpatient services. Satisfaction is perhaps more attainable in long-term  
42 care, possibly because service users who are involved with a service over a longer period of  
43 time develop relationships with their staff enabling them to respond more meaningfully to  
44 questions about satisfaction with their input. In the current survey, however, the particular  
45 service used by the respondent was not identified. Finally, a significant difference was found  
46 between when service users were last seen by a social work compared to a mental health  
47 nurse, with those supported by a mental health nurse having been seen more recently.  
48 Although it is not clear from the data why this was the case, its occurrence could also have  
49 detrimentally affected their satisfaction, whether because they were dissatisfied with the level  
50 of contact received or simply because they could not remember the nature of the contact due  
51 to the time elapsed since it had occurred.  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Conclusion

1  
2  
3  
4  
5 The mixed methods approach used for this study had both strengths and weaknesses. The  
6 former lies in its ability to access and analyse existing large-scale data that would not  
7 otherwise be available. Its limitation, however, is whether the data source was sufficient to  
8 illuminate the social work role, which has been shown to be notoriously difficult to articulate.  
9 Future research will need to pay heed to these limitations to shed more light on whether there  
10 are any true distinctions between the experiences of service users supported by social workers  
11 and other professionals. New research funded by the NIHR School for Social Care Research  
12 is currently underway which aims to address this through an investigation of service user and  
13 staff perspectives of the value of the social work role within Community Mental Health Teams  
14 (CMHTs) using a variety of tools and methods. It will be important to identify any distinctions  
15 found between service users supported by social workers and other CMHT practitioners and  
16 explanations for this. In addition, empirical research to understand the exercise of mental  
17 health social work in practice is required to compare with the literature expounding its optimal  
18 attributes. Such research might usefully consider the voices of social workers themselves as  
19 well as service users and carers. Whilst the focus of this article has been on social work in  
20 mental health it is worth commenting that the contribution of social workers for other service  
21 users (e.g. older people) and in other settings (e.g. intermediate care) is also difficult to  
22 articulate. Thus social work in general as well as mental health social work in particular is in  
23 need of research which helps to articulate its role and value.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 **Research ethics** Ethical approval was granted by the North of Scotland (2) Research Ethics  
38 Committee: grant number: 17/NS/0127.  
39

40  
41 **Funding** This article presents independent research funded by the National Institute for Health  
42 Research (NIHR) School for Social Care Research (SSCR). The views expressed in this  
43 article are those of the authors and not necessarily those of the NIHR SSCR or the Department  
44 of Health, NIHR, or NHS.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

Abendstern, M., Tucker, S., Wilberforce, M., Jasper, R., Brand, C., & Challis, D. (2014). Social workers as members of community mental health teams for older people: what is the added value? *The British Journal of Social Work*, 46, 63-80. doi:10.1093/bjsw/bcu122

ADASS (2018). *AMHPs, Mental Health Act assessments & the mental health social care workforce: National findings – England*. ADASS, England. <https://www.adass.org.uk/media/6428/nhsbn-and-adass-social-care-national-report.pdf>.

Ahmedani, B. K. (2011). Mental health stigma: society, individuals, and the profession. *Journal of Social Work Values and Ethics*, 8(2), 4-16. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3248273/>

All Party Parliamentary Group. (2013). *Inquiry into the state of social work report*. <https://www.basw.co.uk/appg/>

All Party Parliamentary Group. (2016). *Report of the inquiry into adult mental health services in England*. [http://cdn.basw.co.uk/upload/basw\\_75200-9.pdf](http://cdn.basw.co.uk/upload/basw_75200-9.pdf)

Allen, R. (2014). *The role of the social worker in adult mental health services*. London: The College of Social Work. <https://www.bl.uk/britishlibrary/~media/bl/global/social-welfare/pdfs/non-secure/r/o//role-of-the-social-worker-in-adult-mental-health-services.pdf>

Anthony, W. A. (1993). Recovery from mental illness: The guiding vision of the mental health service system in the 1990s. *Psychosocial Rehabilitation Journal*, 16(4), 11-23. doi:10.1037/h0095655

Aschbrenner, K., Mueser, K., Bartels, S., Carpenter-Song, E., Pratt, S., Barre, L. & Kinney, A. (2015). The other 23 hours: A qualitative study of fitness provider perspectives on social support for health promotion for adults with mental illness. *Health & Social Work*, 40, 91-99. doi: 10.1093/hsw/hlv006

Bailey, D. and Liyanage, L., 2012. The role of the mental health social worker: Political pawns in the reconfiguration of adult health and social care. *British Journal of Social Work*, 42, 1113-1131. doi:10.1093/bjsw/bcs069

1  
2  
3 Barclay, P. (1982) *Social workers: Their roles and tasks*, London: Bedford Square Press.  
4  
5

6 Beresford, P. (2007). *The changing roles and tasks of social work from service users'*  
7 *perspectives: A literature informed discussion paper*. London: Shaping Our Lives National  
8 User Network.  
9  
10

11  
12 Briar, S., Miller, H. (1971). *Problems and issues in social casework*. New York: Columbia  
13 University Press.  
14  
15

16  
17 Brown, B., Crawford, P., & Darongkamas, J. (2000). Blurred roles and permeable boundaries:  
18 the experience of multidisciplinary working in community mental health. *Health & Social Care*  
19 *in the Community*, 8, 425-435. doi: 10.1046/j.1365-2524.2000.00268.x  
20  
21

22  
23 Burns, T. (2014) *Our necessary shadow: The nature and meaning of psychiatry*. London:  
24 Penguin.  
25  
26

27  
28 Care Quality Commission, Picker Institute Europe. (2016). *Community mental health service*  
29 *user survey, 2016*. UK Data Service. SN: 8109.  
30  
31

32  
33 Carpenter, J., Schneider, J., Brandon, T. & Wooff, D. (2003). Working in multidisciplinary  
34 community mental health teams: The impact on social workers and health professionals of  
35 integrated mental health care. *British Journal of Social Work*, 33, 1081-1103. doi:  
36 10.1093/bjsw/33.8.1081  
37  
38

39  
40 Challis, D. & Ferlie, E. (1987). Changing Patterns of Fieldwork: Organization: II: The Team  
41 Leaders' View. *The British Journal of Social Work*, 17, 147-167. doi:  
42 10.1093/oxfordjournals.bjsw.a055321  
43  
44

45  
46 Challis, D. & Ferlie, E. (1988). The Myth of Generic Practice: Specialisation in Social Work.  
47 *Journal of Social Policy*, 17, 1-22. doi: 10.1017/S0047279400016421  
48  
49

50  
51 Clifton, J. & Thorley, C. (2014). *Think Ahead: Meeting the workforce challenges in mental*  
52 *health social work*. Institute for Public Policy Research.  
53 [https://www.ippr.org/publications/think-ahead-meeting-the-workforce-challenges-in-mental-](https://www.ippr.org/publications/think-ahead-meeting-the-workforce-challenges-in-mental-health-social-work)  
54 [health-social-work](https://www.ippr.org/publications/think-ahead-meeting-the-workforce-challenges-in-mental-health-social-work)  
55  
56  
57

58  
59 Cohen, J. (1977). *Statistical power analysis for the behavioural sciences*. UK: Routledge.  
60



1  
2  
3  
4  
5 ComRes (2017). *Think Ahead social work poll*. March 2017 [online] Available from  
6 [https://thinkahead.org/wp-content/uploads/2017/03/Think-Ahead\\_headline-report-social-](https://thinkahead.org/wp-content/uploads/2017/03/Think-Ahead_headline-report-social-)  
7 [work-poll-170317.pdf](https://thinkahead.org/wp-content/uploads/2017/03/Think-Ahead_headline-report-social-).  
8  
9

10  
11 Cree, V. E. and Davis, A. (2007). *Social work: Voices from the inside*. London: Routledge.  
12

13  
14 Cummings, S. and Cassie, K. (2015). Community mental health centres. In: D. Kaplan and B.  
15 Berkman, ed., *The Oxford handbook of social work in health and aging* (2nd ed). Oxford, UK:  
16 Oxford University Press.  
17  
18

19  
20 Davies, M. B. (2012). *Social work with adults*. UK: Palgrave Macmillan.  
21

22  
23 Davis, K. & Jung, H. (2012). The mental health field of practice. In: Dulmus, C.N. and Sowers.  
24 *Social work fields of practice: Historical trends, professional issues, and future opportunities*.  
25 USA: John Wiley & Sons.  
26  
27

28  
29 Department of Health. (1962). *A hospital plan for England and Wales* (Cmnd 1604). London:  
30 HMSO.  
31  
32

33  
34 Department of Health. (1975). *Better services for the mentally ill* (Cmnd 6233). London:  
35 HMSO.  
36  
37

38  
39 Department of Health. (1989). *Caring for people* (Cm 849). London: HMSO.  
40  
41

42  
43 Department of Health. (1990). *The Care Programme Approach for people with a mental health*  
44 *illness referred to specialist psychiatric services*, HC(90)23/LASSL(90)11. London:  
45 Department of Health.  
46  
47

48  
49 Department of Health. (1995). *Building bridges: A guide to inter-agency working for the care*  
50 *and protection of severely mentally ill people*. London: Department of Health.  
51  
52

53  
54 Department of Health. (1998). *Modernising social services* (Cm 4169). London: The Stationery  
55 Office.  
56  
57

58  
59 Department of Health. (1999). *A national service framework for mental health*. London:  
60 Department of Health.

1  
2  
3  
4  
5 Draper, B., & Anderson, D. (2010). The baby boomers are nearly here – but do we have  
6 sufficient workforce in old age psychiatry? *International Psychogeriatrics*, 22, 947-949.  
7 doi:10.1017/S1041610210000566  
8  
9

10  
11 Dumaine, M.L. (2003). Meta-analysis of interventions with co-occurring disorders of severe  
12 mental illness and substance abuse: implications for social work practice. *Research on Social*  
13 *Work Practice*, 13,142-165. doi: 10.1177/1049731502250403  
14  
15

16  
17 Evans, S., Huxley, P., Webber, M., Katona, C., Gately, C., Mears, A. & Kendall, T. (2005). The  
18 impact of 'statutory duties' on mental health social workers in the UK. *Health & Social Care in*  
19 *the Community*, 13, 145-154. doi: 10.1111/j.13652524.2005.00540.x  
20  
21  
22

23  
24 Evans, S., Huxley, P., Baker, C., White, J., Madge, S., Onyett, S. and Gould, N. (2012). The  
25 social care component of multidisciplinary mental health teams: A review and national survey.  
26 *Journal of Health Services Research and Policy*, 17, 23-29. doi:10.1258/jhsrp.2012.011117 D  
27  
28

29  
30 Faust, J. R. (2008). Clinical social worker as patient advocate in a community mental health  
31 center. *Clinical Social Work Journal*, 36, 293-300. doi: 10.1007/s10615-007-0118-0  
32  
33

34  
35 Godin, P. (1996). The development of community psychiatric nursing: a professional project?  
36 *Journal of Advanced Nursing*, 23, 925-934. doi:10.1046/j.1365-2648.1996.10110.x  
37  
38

39  
40 Goldstein, H. (1973). *Social work practice: A unitary approach*. South Carolina: University of  
41 South Carolina Press.  
42  
43

44  
45 Golightley, M. & Geomans, R. (2014). *Social work and mental health*. London: Learning  
46 Matters.  
47  
48

49  
50 Gould, N. (2016). *Mental health social work in context*. Routledge.  
51

52  
53 Hardiker, P. & Barker, M. (1999). Early steps in implementing the new community care: the  
54 role of social work practice. *Health & Social Care in the Community*, 7, 417-426. doi:  
55 10.1046/j.1365-2524.1999.00205.x  
56  
57

58  
59 Health Act (1999). England and Wales, C8 S31 [http://www.legislation.gov.uk/ukpga/](http://www.legislation.gov.uk/ukpga/1999/8/pdfs/ukpga_19990008_en.pdf)  
60 [1999/8/pdfs /ukpga\\_19990008\\_en.pdf](http://www.legislation.gov.uk/ukpga/1999/8/pdfs/ukpga_19990008_en.pdf).

1  
2  
3  
4  
5 Herman, D. B. (2014). Transitional support for adults with severe mental illness: Critical time  
6 intervention and its roots in assertive community treatment. *Research on Social Work*  
7 *Practice, 24*, 556-563. doi: 10.1177/1049731513510976  
8  
9

10  
11 Hibbard, J., & Gilbert, H. (2014). *Supporting people to manage their health: an introduction to*  
12 *patient activation*. London: The King's Fund.  
13 [https://www.kingsfund.org.uk/sites/default/files/field/field\\_publication\\_file/supporting-people-](https://www.kingsfund.org.uk/sites/default/files/field/field_publication_file/supporting-people-manage-health-patient-activation-may14.pdf)  
14 [manage-health-patient-activation-may14.pdf](https://www.kingsfund.org.uk/sites/default/files/field/field_publication_file/supporting-people-manage-health-patient-activation-may14.pdf).  
15  
16  
17

18  
19 Howe, D. (1979) Agency Function and Social Work Principles. *British Journal of Social Work*,  
20 9(1), 29-47. doi: 10.1093/oxfordjournals.bjsw.a057057  
21  
22

23  
24 Huxley, P., Evans, S., Gately, C., Webber, M., Mears, A., Pajak, S., Kendal, T., Medina, J., &  
25 Katona, C. (2005). Stress and pressures in mental health social work: The worker  
26 speaks. *British Journal of Social Work, 35*, 1063-1079. doi: 10.1093/bjsw/bch218  
27  
28

29  
30 Huxley, P. and Kerfoot, M. (1993) The mental health workforce in the community: a  
31 comparison of national Community Psychiatric Nursing and Approved Social Work survey  
32 results. *Health and Social Care in the Community, 1*, 169-174. doi: 10.1111/j.1365-  
33 2524.1993.tb00212.x  
34  
35

36  
37  
38 Huxley, P., Reilly, S., Mohamad, H. and Harrison, J. (1998) Severe mental illness: comparing  
39 CPNs' and social workers' input. *Mental Health Nursing 18*, 14-17.  
40  
41

42  
43 Ife, J. (2012). *Human rights and social work: Towards rights-based practice*. Cambridge:  
44 Cambridge University Press.  
45  
46

47  
48 Janlöv, A. C., Granskär, M. & Berg, A. (2015). Challenges Highlighted During Peer  
49 Supervision by Mental Health Nurses and Social Workers Recently Trained as Case  
50 Managers in Sweden. *Issues in Mental Health Nursing, 36*, 809-816. doi:  
51 10.3109/01612840.2015.1047073  
52  
53

54  
55 Jones, R. (2014). The best of times, the worst of times: Social work and its moment. *British*  
56 *Journal of Social Work, 44*, 485-502. doi: 10.1093/bjsw/bcs157  
57  
58

59  
60 Karban, K. (2011). *Social work and mental health*. Cambridge: Polity.

1  
2  
3  
4  
5 King, R., Yellowlees, P., Nurcombe, B., Spooner, D., Sturk, H., Spence, S., & Le Bas, J.  
6 (2002). Psychologists as mental health case managers. *Australian Psychologist*, 37, 118-122.  
7 doi: 10.1080/00050060210001706756  
8  
9

10  
11 Lang, L., Orton, S., Sallah, D., Hewitt-Moran, T., Zhang, D., Cullen, S., Dixon, S., Bell, B., Bell,  
12 D., Meeson, L. & Chen, R. (2011). A survey of engagement and competence levels in  
13 interventions and activities in a community mental health workforce in England. *BMC Health*  
14 *Services Research*, 11, 352-362. doi: 10.1186/1472-6963-11-352  
15  
16  
17

18  
19 LeCroy, C.W. and Stinson, E.L. (2004). The Public's Perception of Social Work: Is It What We  
20 Think It Is? *Social Work*, 49, 164–174. doi: 10.1093/sw/49.2.164  
21  
22

23  
24 Legood, A., McGrath, M., Searle, R. and Lee, A. (2016). Exploring How Social Workers  
25 Experience and Cope with Public Perception of Their Profession. *The British Journal of Social*  
26 *Work*, 46, 1872–1889. doi: 10.1093/bjsw/bcv139  
27  
28  
29

30  
31 Lilo, E. (2016). *Mental Health Integration Past, Present and Future: A Report of National*  
32 *Survey into Mental Health Integration in England*. Mersey Care NHS Trust Integrated Care  
33 Demonstrator Site Project NHS North West Health Education Workforce Transformation  
34 Initiative. NHS Health Education North West and Mersey Care NHS Trust.  
35  
36  
37

38  
39 Malone, D., Marriott, S. V., Newton-Howes, G., Simmonds, S., & Tyrer, P. (2007). Community  
40 mental health teams (CMHTs) for people with severe mental illnesses and disordered  
41 personality. *Cochrane Database of Systematic Reviews*, (3). doi:  
42 10.1002/14651858.CD000270.pub2  
43  
44  
45

46  
47 Manktelow, R., Hughes, P., Britton, F., Campbell, J., Hamilton, B. and Wilson, G. (2002). The  
48 experience and practice of approved social workers in Northern Ireland. *British Journal of*  
49 *Social Work*, 32, 443-461. doi: 10.1093/bjsw/32.4.443  
50  
51

52  
53 Marshall, M., Lockwood, A. & Gath, D. (1995). Social services case-management for long-  
54 term mental disorders: a randomised controlled trial. *The Lancet*, 345(8947), 409-412. doi:  
55 10.1016/S0140-6736(95)90399-2  
56  
57  
58  
59  
60

1  
2  
3 McNicoll, A. (2016). *Withdrawing social workers from NHS will 'boost Care Act compliance',*  
4 *claims council.* [http://www.communitycare.co.uk/2016/01/08/withdrawing-social-workers-](http://www.communitycare.co.uk/2016/01/08/withdrawing-social-workers-nhs-will-boost-care-act-compliance-claims-council/)  
5 [nhs-will-boost-care-act-compliance-claims-council/](http://www.communitycare.co.uk/2016/01/08/withdrawing-social-workers-nhs-will-boost-care-act-compliance-claims-council/)  
6  
7

8  
9 Medical-Psychological Association (1969) *Report of the Committee on Local Authority and*  
10 *Allied Personal Social Services.* Royal Medical-Psychological Association, London.  
11  
12

13  
14 Mental Health Act (1959). UK. London: HMSO.  
15  
16

17 Mitchell, F. & Patience, D. A. (2002). Conjoint multi-disciplinary assessment in a community  
18 mental health team: The impact on the social work role. *Social Work in Health Care*, 35, 605-  
19 613. doi: 10.1300/J010v35n01\_14  
20  
21

22  
23 Moriarty, J., Baginsky, M., & Manthorpe, J. (2015). *Literature review of roles and issues within*  
24 *the social work profession in England.* Social Care Workforce Research Unit, King's College,  
25 London. Available at: [http://socialwelfare.bl.uk/subject-areas/services-activity/social-work-](http://socialwelfare.bl.uk/subject-areas/services-activity/social-work-care-services/socialcareworkforceresearchunit/174392moriarty-et-al.-2015-PSA)  
26 [care-services/socialcareworkforceresearchunit/174392moriarty-et-al.-2015-PSA.](http://socialwelfare.bl.uk/subject-areas/services-activity/social-work-care-services/socialcareworkforceresearchunit/174392moriarty-et-al.-2015-PSA)  
27  
28  
29

30  
31 Moriarty, J. and Manthorpe, J. (2016). *The effectiveness of social work with adults. A*  
32 *systematic scoping review.* London: Social Care Workforce Research Unit, King's College  
33 London.  
34  
35

36  
37 Morriss, L. (2016). Being seconded to a mental health trust: The (in) visibility of mental health  
38 social work. *British Journal of Social Work*, 47, 1344-1360. doi.org/10.1093/bjsw/bcw022  
39  
40

41  
42 Moxley, D. P. (1989). *Practice of Case Management* (Vol. 58). Thousand Oaks, CA: Sage.  
43  
44

45  
46 Mulhall, J. (2000). Using community mental health services: Asking users. *Practice*, 12(4), 27-  
47 36. doi: 10.1080/09503150008415196  
48  
49

50  
51 Ng, C., Herrman, H., & Chiu, E. (2009). Community mental health care in the Asia-Pacific  
52 region: Using current best-practice models to inform future policy. *World Psychiatry*, 8, 49-55.  
53 doi: 10.1002/j.2051-5545.2009.tb00211.x  
54  
55

56  
57 Onyett, S. (2011). Revisiting job satisfaction and burnout in community mental health  
58 teams. *Journal of Mental Health*, 20, 198-209. doi: 10.3109/09638237.2011.556170  
59  
60

1  
2  
3 Pahwa, R., Smith, M. E., McCullagh, C. A., Hoe, M. & Brekke, J. S. (2016). Social support-  
4 centered versus symptom-centered models in predicting functional outcomes for individuals  
5 with schizophrenia. *Journal of the Society for Social Work and Research*, 7, 247-268. doi:  
6 10.1086/686770  
7  
8  
9

10  
11 Parrott, L. (2014). *Values and ethics in social work practice*. Exeter, UK: Learning Matters.  
12  
13

14 Peck, E. & Norman, I. J. (1999). Working together in adult community mental health services:  
15 exploring inter-professional role relations. *Journal of Mental Health*, 8, 231-242. doi:  
16 10.1080/09638239917391  
17  
18  
19

20 Penhale, B. & Young, J. (2015). *A review of the literature; public and service users' opinions*  
21 *about the conduct and competence of social workers*. The Professional Standards Authority.  
22  
23  
24

25 Perlman, H.H. (1979). *Relationship: The heart of helping people*. Chicago: University of  
26 Chicago Press.  
27  
28  
29

30 Priebe, S., Fakhoury, W.K., Hoffmann, K. and Powell, R.A. (2005). Morale and job perception  
31 of community mental health professionals in Berlin and London. *Social Psychiatry and*  
32 *Psychiatric Epidemiology*, 40, 223-232. doi: 10.1007/s00127-005-0880-7  
33  
34  
35

36 Raiff, N.R. & Shore, B. K. (1993). *Advanced case management: New strategies for the*  
37 *nineties* (No. 66). Thousand Oaks, CA: Sage.  
38  
39  
40

41 Ramon, S. (2010). Adult mental health in a changing international context: The relevance to  
42 social work. *British Journal of Social Work*, 39, 1615-1622. doi: 10.1093/bjsw/bcp066  
43  
44  
45

46 Rapaport, J., 2005. Policy swings over thirty-five years of mental health social work in England  
47 and Wales 1969–2004. *Practice*, 17, 43-56. doi: 10.1080/09503150500058041  
48  
49  
50

51 Ring, C. (2001). Quality assurance in mental healthcare: a case study from social work. *Health*  
52 *& Social Care in the Community*, 9, 383-390. doi: 10.1046/j.0966-0410.2001.00324.x  
53  
54  
55

56 Rode, N. (2017) Defining social work is a never-ending story. *European Journal of Social*  
57 *Work*, 20, 64-75. doi: 10.1080/13691457.2016.1185704  
58  
59  
60

1  
2  
3 Rubin, A. and Parrish, D.E. (2012). Comparing social worker and non-social worker  
4 outcomes: A research review. *Social Work*, 57, 309-320. doi: 10.1093/sw/sws015  
5  
6

7  
8 Stanley, N. (2003). Working on the interface: identifying professional responses to families  
9 with mental health and child-care needs. *Health & Social Care in the Community*, 11, 208-  
10 218. doi: 10.1046/j.1365-2524.2003.00428.x  
11  
12

13  
14 Stromwall, L. K. & Hurdle, D. (2003). Psychiatric rehabilitation: An empowerment-based  
15 approach to mental health services. *Health & Social Work*, 28, 206-213. doi:  
16 10.1093/hsw/28.3.206  
17  
18

19  
20 Tew, J. (2011.) *Social approaches to mental distress*. Basingstoke: Palgrave Macmillan.  
21  
22

23  
24 Think Ahead (2018). [https://thinkahead.org/about-the-programme/about-mental-health-  
25 social-work/#what-mental-health-social-workers-do](https://thinkahead.org/about-the-programme/about-mental-health-social-work/#what-mental-health-social-workers-do).  
26  
27

28  
29 Tricco, A. C., Antony, J., Zarin, W., Striffler, L., Ghassemi, M., Ivory, J. & Straus, S. E. (2015).  
30 A scoping review of rapid review methods. *BMC medicine*, 13, 224-239. doi: 10.1186/s12916-  
31 015-0465-6  
32  
33

34  
35 Wall, A. (1998). Organisations. Knowing who's boss. *Health Service Journal*, 108 (5630), 22-  
36 24. <https://www.ncbi.nlm.nih.gov/pubmed/10186195>  
37  
38

39  
40 Wilberforce, M., Tucker, S., Abendstern, M., Brand, C., Giebel, C.M. and Challis, D. 2013).  
41 Membership and management: structures of inter-professional working in community mental  
42 health teams for older people in England. *International Psychogeriatrics*, 25, 1485-1492. doi:  
43 10.1017/S104161021300077X  
44  
45

46  
47 Woodbridge-Dodd, K. (2017) *Integrating mental health and social care*. Centre for Mental  
48 Health, London. <http://www.infocoponline.es/pdf/integrating-social-care.pdf>  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60