

This is a repository copy of *Activated carbons of varying pore structure eliminate the bioavailability of 2,3,7,8-tetrachlorodibenzo-p-dioxin to a mammalian (mouse) model*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/150942/>

Version: Accepted Version

Article:

Sallach, J. Brett orcid.org/0000-0003-4588-3364, Crawford, Robert, Li, Hui et al. (4 more authors) (2019) Activated carbons of varying pore structure eliminate the bioavailability of 2,3,7,8-tetrachlorodibenzo-p-dioxin to a mammalian (mouse) model. *Science of the Total Environment*. pp. 2231-2238. ISSN 1879-1026

<https://doi.org/10.1016/j.scitotenv.2018.09.270>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Activated carbons of varying pore structure eliminate the bioavailability of 2,3,7,8-**
2 **tetrachlorodibenzo-p-dioxin to a mammalian (mouse) model**

3 J. Brett Sallach^{a,1}, Robert Crawford^b, Hui Li^a, Cliff T. Johnston^c, Brian J. Teppen^a, Norbert E.
4 Kaminski^{b,d}, Stephen A. Boyd^{a*},

5
6 ^aDepartment of Plant, Soil, and Microbial Sciences, Michigan State University, East Lansing,
7 Michigan, USA 48824

8 ^bInstitute for Integrative Toxicology, Michigan State University, East Lansing, Michigan, USA
9 48824

10 ^cCrop, Soil, and Environmental Science, Purdue University, West Lafayette, Indiana, USA
11 47907

12 ^dDepartment of Pharmacology and Toxicology, Michigan State University, East Lansing,
13 Michigan, USA 48824

14 ¹Environment Department, University of York, Heslington, York, United Kingdom, YO10 5NG

15

16 J. Brett Sallach (brett.sallach@york.ac.uk), Robert Crawford (crawfo28@msu.edu), Hui Li
17 (lihui@msu.edu), Cliff T. Johnston (cliffjohnston@purdue.edu), Brian J. Teppen
18 (teppen@msu.edu), Norbert E. Kaminski (kamins11@msu.edu), Stephen A. Boyd
19 (boyds@msu.edu),

20

21

22

23

24

25

26

27

28 *Corresponding author. Tel: 517 355-0271. E-mail: boyds@msu.edu (Stephen Boyd)

29 **Abstract**

30 The use of activated carbon (AC) as an in situ sorbent amendment to sequester
31 polychlorinated-dibenzo-p-dioxins and furans (PCDD/Fs) present in contaminated soils and
32 sediments has recently gained attention as a novel remedial approach. This remedy could be
33 implemented at much lower cost while minimizing habitat destruction as compared to traditional
34 remediation technologies that rely on dredging/excavation and landfilling. Several prior studies
35 have demonstrated the ability of AC amendments to reduce pore water concentrations and hence
36 bioaccumulation of PCDD/Fs in invertebrate species. However, our recent study was the first to
37 show that AC had the ability to sequester 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in a form
38 that eliminated bioavailability to a mammalian (mouse) model. Here we show that three
39 commercially available ACs, representing a wide range of pore size distributions, were equally
40 effective in eliminating the bioavailability of TCDD based upon two sensitive bioassays, hepatic
41 induction of cyp1A1 mRNA and immunoglobulin M antibody-forming cell response. These
42 results provide direct evidence that a wide range of structurally diverse commercially available
43 ACs may be suitable for use as in situ sorbent amendments to provide a cost-effective remedy for
44 PCDD/F contaminated soils and sediments. Potentially, adaption of this technology would
45 minimize habitat destruction and be protective of ecosystem and human health.

46 **Key Words**

47 TCDD, immune response, remediation, sorbent amendments

48

49

50

51

52

53

54 **1. Introduction**

55 The ubiquitous occurrence of polychlorinated dibenzo-p-dioxins and polychlorinated
56 dibenzofurans (PCDD/Fs) in the environment results from their formation as unintentional
57 byproducts of chemical manufacturing, including pesticide production and the historic chlor-
58 alkali process, and from both anthropogenic (incineration) and natural (forest fires and volcanic
59 activities) combustion [1,2]. The natural in situ formation of predioxins and octachlorodibenzo-
60 p-dioxin may also occur on the surfaces of ball clays [3]. In recent decades, significant
61 technological and regulatory improvements have limited the anthropogenic release of these
62 compounds to the environment. However, their widespread distribution and recalcitrance in soils
63 and sediments, coupled with their high toxicity at low levels of exposure, contributes to their
64 high priority for remediation throughout the world [4,5].

65 Human exposure of PCDD/Fs is potentially associated with many adverse health effects
66 including cardiovascular disease, diabetes, cancer, porphyria, endometriosis, altered hormone
67 levels and reproductive health, skin, tooth, and nail abnormalities amongst others [5, 6]. Perhaps
68 most alarming, exposure to PCDDs at levels only a single order of magnitude greater than current
69 mean background levels for the general population (viz. 15 ppt serum lipid basis) manifests
70 negative health outcomes [7]. Exposure to PCDD/Fs has been linked to prenatal mortality in a
71 number of mammalian species including mice, rabbits and mink [8]. Interestingly, the
72 proliferation of antibiotic resistance genes in the gut microbiota of mice has been associated with
73 the immune response induced by TCDD exposure [9].

74 Remediation of PCDD/F contaminated soils and sediments often involves removal by
75 excavation or dredging and disposal in hazardous waste landfills, with varied degrees of

76 effectiveness [10,11]. This traditional remedy is associated with high cost and substantial habitat
77 destruction, for example detrimental effects on benthic ecosystems, and can result in re-
78 distribution of contaminated sediments [12]. Therefore, efforts have been made to develop new
79 remediation technologies that are less expensive and destructive while being protective of
80 ecosystem and human health. The use of activated carbon (AC) sorbent amendments has
81 emerged as a particularly promising treatment alternative [13].

82 A select number of studies showing reductions in pore water concentrations of PCDDs
83 and subsequent reductions in bioaccumulation amongst benthic organisms and soil invertebrates
84 has provided the impetus for further scientific investigation of this technology [13–17].
85 However, from a public policy standpoint, mammalian exposure and bioavailability has been
86 considered in order to make decisions protective of human health. In 2012, based on evidence
87 that 16-28% of measured PCDD/Fs in Midland bulk soils were orally bioavailable to mammals,
88 Dow Chemical (Midland, Michigan, USA) was granted a site-specific variance in soil
89 remediation targets (from 90 to 250 ppt TEQ) by the Michigan Department of Environmental
90 Quality (MDEQ) [18,19]. The significance of the direct oral exposure pathway has been
91 established through studies on the advertent and inadvertent ingestion of soil documented in
92 humans, especially children, and wild animals [20-22].

93 We recently showed that a commercial AC has the ability to sequester PCDD/Fs in a
94 form that eliminates mammalian bioavailability. When TCDD was sequestered by AC it failed to
95 elicit a hallmark of TCDD exposure, i.e. suppression of immune system response; other sorbent
96 materials including silica and smectite (e.g. montmorillonite) clays did not reduce TCDD
97 bioavailability to the mammalian (mouse) model [23–25]. In addition, TCDD sequestered by AC
98 was shown to eliminate characteristic TCDD influences on the gut microbiome [26]. The

99 beneficial effect of AC in reducing mammalian toxicant exposure in the gut has motivated its
100 recommended use for livestock and humans following acute exposures [27-29].

101 The efficacy of AC as a sorbent for organic contaminants is well established accounting
102 for its use in many treatment processes including both water and gas flu treatment [30,31]. This
103 is especially true in the case of planar hydrophobic compounds which are intrinsically suitable
104 for hydrophobic pore-filling processes coupled with van der Waals attraction characteristic of
105 contaminant sorption by AC [32]. The sorption capacity of ACs have been shown to be orders of
106 magnitude greater than the primary native soil/sediment sorptive component for hydrophobic
107 contaminants, namely amorphous organic matter [33]. Pore structure is known to play an
108 important role in the sorption potential of all porous media across a variety of scales [34]. The
109 pore structure of ACs is dependent on the source material as well as physical and chemical
110 processes utilized in their formation. Because of this, the pore structure of ACs varies
111 significantly among different commercially available products, likely affecting their suitability
112 for specific applications [35].

113 Pore characteristics play a significant role in determining the irreversibility of
114 contaminant sorption, or the kinetic release, by ACs. For example, a common assumption is that
115 a pore size of 1.3-1.7 times the molecular (kinetic) diameter of a particular compound manifests
116 the greatest sorption energy and preferential sorption [36]. This concept has been utilized in the
117 pharmaceutical industry to help modulate drug delivery via a porous silica media [37]. Dynamic
118 molecular simulations show that the energetics of sorption are most favorable with pores slightly
119 larger than TCDD molecules [38]. As the pore size increases, the mean potential energy of
120 sorption for the TCDD molecule decreases. Molecular simulations also suggest that the water
121 density within individual pores decrease as pore size decreases. The resulting sub-aqueous

122 environment would plausibly be energetically favorable for hydrophobic compounds such as
123 TCDD [29].

124 The goal of the current study was to investigate the effectiveness of AC materials
125 representing a wide range of pore structure distributions, as well as specific surface areas, in
126 reducing the bioavailability of TCDD using a mammalian (mouse) model. The mouse has been
127 extensively characterized with respect to its biological and toxicologic responses to PCDD and
128 dioxin-like compounds with induction of hepatic enzyme, cytochrome P-4501A1 (cyp1A1), and
129 suppression of the primary IgM antibody response being among the most sensitive to PCDD/F
130 exposure. For this reason the mouse and these specific responses were assayed when
131 determining bioavailability of PCDDs. In addition to the WPC AC used in our previous study,
132 two additional AC materials were selected for study (Table 1; FM1 and G60). The three ACs
133 were loaded with TCDD via the incipient wetness method, and delivered to mice via oral gavage.
134 Bioavailability in mice was determined through enumeration of the anti-sheep erythrocyte
135 (sRBC) IgM antibody forming cells (AFC) and induction of cyp1A1 mRNA, two hallmark
136 responses of TCDD exposure in mammals.

137 **2. Materials and Methods**

138 2.1 Selection of ACs

139 In a previous study, five activated carbons were characterized using nitrogen absorption to
140 determine specific surface area and pore size distribution [23]. Of the five ACs, three were
141 selected for use in the current study (Table 1). WPC, used in the previous feeding study,
142 represented a microporous dominant AC while Darco FM-1 (Cabot Corp) represented a
143 mesoporous dominant AC. Specific surface area was also considered for selection of AC
144 materials. The specific surface area of FM-1 was smaller than WPC while Darco G60 (Cabot

145 Corp) had increased specific surface area compared to WPC and with a more even distribution of
146 mesopores and micropores.

147 < Table 1. Structural properties of activated carbon >

148

149

150 2.2 Preparation of ACs

151 Loading the three ACs with the required amount of 2,3,7,8-TCDD via aqueous sorption
152 was impractical due to TCDD's extremely low solubility in water (0.2-0.3 $\mu\text{g/L}$) [39]. Therefore,
153 the incipient wetness method was employed to load TCDD into the pore structures of the three
154 AC sorbents as conducted and validated in previous studies [23-25].

155 Two 500 mg portions of each of the three ACs were measured in Corex glass centrifuge
156 tubes (30 mL). TCDD spike solutions were prepared in DMSO at concentrations (47.7, 63.37, &
157 100 $\mu\text{g/mL}$) necessary to deliver equivalent masses of TCDD in DMSO volumes (336, 253, 160
158 μL) equivalent to the pore volume of the respective ACs (G60, FM-1, WPC) listed in Table 1.
159 Removal of DMSO followed the method described in our previous study [23]. The procedure
160 resulted in TCDD-AC of either 0 or 32 $\mu\text{g/g}$ for each of the three ACs. Thermogravimetric
161 analysis (TGA) was used to quantify mass loss during heating. Approximately 30 mg of AC was
162 placed in a ceramic crucible and placed in the TGA (model TGA/SDTA851e, Mettler Toledo,
163 OH). The samples were heated from 25 to 1000 $^{\circ}\text{C}$ at an increment rate of 20 $^{\circ}\text{C}$ per minute in a
164 N_2 atmosphere. The TGA data are reported as percent mass loss of the initial mass and the 1st
165 derivative of the mass loss curves which shows the temperature ranges where mass-loss events
166 occur.

167 Aqueous suspensions, necessary to administer the samples to the mice via oral gavage,
168 were prepared by combining 156.25 mg of each TCDD-AC material with 5 mL of deionized
169 water in 20 mL glass scintillation vials.

170 2.3 TCDD Analysis

171 Samples of each of the three AC materials, loaded with TCDD following the incipient
172 wetness method described above, were analyzed by Pacific Rim Laboratories INC (Surrey, BC,
173 Canada) following the EPA 1613b standard reference method [40]. Briefly, 0.5 g samples of
174 each AC underwent 64 hours of Soxhlet extraction using toluene. Resulting extracts were
175 brought up to 100 mL with toluene. A 10 μ L aliquote of sample extract was combined with 0.5
176 ng of $^{13}\text{C}_{12}$ -2,3,7,8-TCDD and 1.0 ng of $^{13}\text{C}_{12}$ -1,2,3,4-TCDD and made up to a final volume of
177 50 μ L prior to HRGC/HRMS analysis. Results from the analyses of all three AC materials are
178 provided in Table 2.

179 < Table 2. Analytical detection and recoveries of TCDD from activated carbon materials >

180 2.4 Animals

181 Eight to twelve week old pathogen-free B6C3F1 female mice, purchased from Charles
182 River Breeding Laboratories (Portage, MI, USA), were randomly divided into 9 experimental
183 groups (5 mice per group). Each group was placed into its own plastic cage containing sawdust
184 bedding. Prior to the start of the experiment, the mice were acclimated for two weeks to allow
185 their body weights to reach approximately 20 g. Animal holding rooms were operated with 12-
186 hour light/dark cycles at temperatures of 21-24°C and 40-60 % relative humidity. Water and
187 food (Purina Certified Laboratory Chow) were provided without restraint and all procedures

188 involving mice were in accordance with the Michigan State University Institutional Animal Care
189 and Use Committee.

190 2.5 Seven Day Feeding Trials

191 Following the feeding protocol previously established [17–19], a 7-day feeding study
192 comprising 4 TCDD treatments was performed. Treatment groups included those receiving 1.0
193 $\mu\text{g}/\text{mL}$ TCDD in either corn oil (TCDD-CO) or in the three different AC solutions (TCDD-AC).
194 In addition, control groups consisting of the vehicles (corn oil or AC) only were prepared. The
195 final group was kept naïve, receiving no treatment regimen. Details of the treatment matrix are
196 provided in Table 3.

197 < Table 3. Treatment groups and experiment timeline >

198 On days one through four, mice received 200 μL aliquots of their respective treatment via
199 oral gavage, with mice in groups 5-8 receiving a mass of TCDD (10 $\mu\text{g}/\text{kg}$ bw/day) in their
200 respective vehicles daily. On day 3, Groups 1-8 mice were sensitized with 1×10^9 sheep red
201 blood cells (sRBC) by intraperitoneal injection to initiate a T cell dependent humoral immune
202 response. Mice were euthanized by cervical dislocation 4 days post sensitization (day 7). Body
203 weight was immediately determined prior to resection of the liver (for cyp1A1 induction) and
204 spleen (for AFC response). Each liver and spleen were then weighed individually. Mouse
205 feeding trials were repeated approximately 6 months apart to confirm reproducibility. Results are
206 representative of the 2 separate experiments.

207 Antibody forming cell response

208 Enumeration of anti-sRBC IgM antibody forming cells (AFCs) was performed using the
209 Jerne plaque assay [41] following the method described previously [23]. Duplicate assays were

210 prepared for each mouse sample (5 mice per treatment) resulting in 10 assays per treatment
211 group. AFC counts were normalized with total cell counts enumerated with a ZI Coulter particle
212 counter (Beckman Coulter, Pasadena, CA, USA) and figures are presented as anti-sRBC IgM
213 AFC/ 1×10^6 splenocytes.

214 2.6 Cyp1A1 gene expression

215 Induction of cyp1A1 mRNA was quantified by real time polymerase chain reaction
216 (PCR). Sacrificed mouse livers, stored at -70C in TRI Reagent (Sigma-Aldrich, St. Louis, MO,
217 USA), were homogenized then phase separated using bromochlorophenol. RNA was precipitated
218 from the aqueous phase using isopropanol. Extraction, purification, and DNase treatment
219 followed using a Promega SV total RNA isolation system. A high capacity cDNA reverse
220 transcription kit (Applied Biosystems, Foster City, CA, USA) was employed for reverse
221 transcription of total RNA using random primers. Amplification of the of the cDNA using a
222 Taqman primer/probe set for mouse cyp1A1 (Applied Biosystems) preceded analysis with a 7900
223 HT fast real-time polymerase chain reaction (PCR) system (Applied Biosystems). Fold change
224 values were calculated using the $\Delta\Delta C_T$ method [42].

225 2.7 Statistical Analysis

226 Real-time PCR statistical analysis was performed on ΔC_T values using Prism version 4.0a
227 (Graphpad, La Jolla, CA, USA). Statistically significant differences between treatment groups
228 and controls were determined by Dunnett's two tailed t test.

229 3. Results and Discussion

230 3.1 Analytical assessment of TCDD-AC material

231 Analytical determination of the concentration of PCDD/Fs and similar compounds in
232 carbonaceous materials is complicated by the lack of an established reference method that
233 provides adequate recoveries. Currently, EPA Method 1613 serves as the standard method for
234 the extraction and quantification of tetra- through octa- chlorinated dioxins and furans from
235 numerous matrices including soils and sediments [40]. Following the preparation of ACs by the
236 incipient wetness method, AC samples from each of the three study materials were analyzed by
237 Pacific Rim Laboratories using EPA Method 1613. Total calculated concentrations ranged from
238 7.5 – 8.7 $\mu\text{g/g}$ with corresponding percent recoveries of 23.5-27.1 % (Table 2). Interestingly,
239 extraction efficiency seemed to increase with decreasing percentage of micropore volume. These
240 low results are consistent with extraction efficiencies for PCDD/Fs and similar compounds and
241 from similar carbonaceous materials reported elsewhere, and highlights the irreversibility of
242 TCDD binding and the ineffectiveness of current standard methods for the extraction and
243 analysis of PCDD/Fs from graphitic porous materials [43–45]. Furthermore, the inefficiency of
244 TCDD extraction directly reflects the sequestering ability of ACs and corresponding reduction in
245 bioavailability.

246 As pyrogenic carbonaceous materials are natural constituents of all soils and sediments,
247 prior environmental assessments using standard methodology may underestimate the actual
248 environmental abundance of these compounds. Our prior published studies on the bioavailability
249 of TCDD sorbed by silica and smectite clay [24,25] followed the incipient wetness method.
250 Results from these studies showed no loss of TCDD associated with the incipient wetness
251 method. Specifically, dose dependent responses in mice were identical when the equivalent doses
252 of TCDD were administered directly in corn-oil or as TCDD–clay and –silica complexes. In the
253 experiments with AC, the extracted concentrations of 7.5 – 8.7 $\mu\text{g/g}$ determined using the

254 standard method and quantified via HR-GC/MS, would be sufficient to elicit a significant
255 bioresponse by both bioassays (i.e. AFC response and *cyp1A1* mRNA induction) in the current
256 and previous studies if that mass of TCDD was bioavailable. In fact, we have shown repeatedly
257 that exposure to TCDD at levels as low as 0.01 $\mu\text{g}/\text{mL}$, which would correspond to TCDD-AC
258 concentrations of 0.32 $\mu\text{g}/\text{g}$, would result in a significant bioresponse in both bioassays assuming
259 the TCDD was bioavailable [23-25]. Therefore, thermal gravimetric analysis was used to verify
260 loading via incipient wetness, and the two sensitive bioassays utilized in our prior published
261 work [23-25] were used to measure bioavailability following oral exposure of the mammalian
262 (mouse) model.

263 3.2 Confirmation of pore filling

264 Based on our working hypothesis that smaller micropores sorb TCDD more strongly than
265 larger mesopores, we hypothesized that bioavailability would increase with increasing proportion
266 of mesoporosity. A benefit inherent in the incipient wetness method is that sorption of TCDD
267 dissolved in DMSO is directly related to pore filling, since the volume of DMSO solvent
268 (containing dissolved TCDD) added corresponds to the pore volume of each AC. When added to
269 the AC, the material is mixed rigorously until all the solvent has been internalized within the AC
270 pore structure. Thus, the TCDD-DMSO fills both mesopores and micropores.

271 To quantify the pore filling process, thermogravimetric analysis (TGA) was performed on
272 AC samples loaded with DMSO via the incipient wetness method, both before the 2-hour 200°C
273 solvent removal and after. Thermograms of the mass removal curves (TG) and their derivatives
274 (DTG) are shown in Figure 1. The amount of DMSO added was equal to the pore volume of the
275 mass of AC used based on the following measured pore volumes: WPC (32 %) < FM1 (50.5 %)

276 < G60 (67.1 %). The percent removals (Figure 1) confirm that the masses of DMSO removed
277 (WPC<FM1<G60) correlates with the masses of DMSO added and the AC pore volumes. The
278 derivatives of the mass loss thermograms (DTG) (Figure 1) provide evidence of pore filling for
279 both micro and meso pores in the three ACs. The large negative peak in the DTG curves at 150
280 °C for both G60 and FM1 corresponds to rapid removal of DMSO from larger mesopores (2-50
281 nm). In addition, the two ACs with significant micropore volumes, WPC and G60, both showed
282 significant tailing in the DTG curve at higher temperatures extending from 150 to 300 °C. This
283 tailing is characteristic of the removal of DMSO from micropores (<2 nm), which requires more
284 time and energy.

285 < Figure 1. Thermogravimetric analysis (TGA) of DMSO infused activated carbons via the
286 incipient wetness method. Thermograms (TG) of mass loss loss (top) and their derivatives
287 (DTG) (bottom) following a heating ramp to 500°C. >

288
289 When TGA analysis was performed on the AC materials after DMSO removal at 200°C
290 for 2 hours, an insignificant mass loss was observed demonstrating the effectiveness of the
291 heating protocol for DMSO removal. The early peaks in the derivative curves for all three ACs is
292 likely associated with a small amount of moisture that condensed on the sample during the
293 cooling process after the material has been heated. Taken together, data from TGA analysis
294 provides good evidence that the incipient wetness method was effective in pore-filling of both
295 meso- and micro- pores. In addition, these data also confirmed that the heating protocol, 200°C
296 for 2 hours, was effective in removing DMSO from AC materials.

297 3.3 Activated carbon effect on reducing bioavailability via cyp1A1 induction and the Jerne
298 plaque Assay

299 The induction of *cyp1A1* is a hallmark response of AhR agonists including PCDDs and
300 was measured using liver tissue of mice. When exposed to TCDD via the corn oil vehicle,
301 *cyp1A1* mRNA expression increased by more than 4000 fold compared to that of mice exposed
302 to the corn oil vehicle with no TCDD (Figure 2). This response is in excellent agreement with the
303 *cyp1A1* expression in mice exposed to the same levels of TCDD in corn oil from previous work
304 [23]. However, when the same mass of TCDD was delivered to mice in each of the three AC
305 vehicles, no significant response was detected (Figure 2). Likewise, compared to the corn oil
306 control, no significant difference in *cyp1A1* expression resulted from administering any of the
307 three AC vehicles without TCDD. Importantly, sequestration of TCDD by any of the three ACs
308 eliminated TCDD bioavailability to bind the AhR, whereas TCDD freely available in corn oil
309 was clearly bioavailable and resulted in increased expression of the *cyp1A1* gene.

310 < Figure 2. *Cyp1a1* mRNA fold expression in mouse liver after treatment with corn oil or
311 activated carbon of various pore structures with and without 2,3,7,8 - tetrachlorodibenzo-p-
312 dioxin. * indicates a significant difference at $p < 0.05$ level compared to the respective vehicle
313 control group. >

314
315 Suppression of immune function is another hallmark of TCDD toxicity in mammals.
316 Hence, in addition to *cyp1A1* induction in the liver, the Jerne Plaque Assay was employed to
317 evaluate TCDD-induced suppression of humoral immune function in mice, and its elimination
318 via reduction in bioavailability, through quantification of antigen-specific T cell dependent IgM
319 AFC response. Mice were sensitized to the antigen, sRBC, on day three of the seven day feeding
320 trial (Table 3). Our work has demonstrated that AC materials, alone, do not interfere with the
321 IgM responses in mice [23]. Therefore, in experiments where the mammalian (mouse) model
322 was exposed to corn oil, TCDD and AC, alone and in various combinations, suppression of the
323 anti-sRBC IgM AFC response results from exposure to TCDD, and establishes the

324 bioavailability of TCDD. As expected, the AFC response was significantly suppressed in mice
325 following exposure to TCDD in the corn oil vehicle (Figure 3). However, the AFC response in
326 mice exposed to TCDD-AC showed no evidence of suppression compared to the corresponding
327 groups exposed to each AC with no TCDD. This was true for all three TCDD-AC materials
328 despite substantial differences in pore size distribution, i.e. the relative percentages of micro- and
329 meso-pores. These results confirm and expand our prior findings [23].

330 < Figure 3. Suppression of humoral immunity observed in response to 2,3,7,8 -
331 tetrachlorodibenzo-p-dioxin administered by oral gavage in either corn oil or sorbed on to
332 activated carbon. * indicates a significant difference at $p < 0.05$ level compared to the respective
333 vehicle control group.>

334
335 Previous studies have shown that TCDD exposure in mice can impact organ mass relative
336 to body weight [23-25]. In agreement with these prior observations, TCDD exposure in corn oil
337 resulted in an increased liver weight ratio (to body mass) and decreased spleen weight ratio
338 compared to the corn oil vehicle (Figure 4). Mice fed TCDD sequestered by the AC materials did
339 not manifest this characteristic response, again indicating the elimination of TCDD
340 bioavailability. In fact, TCDD-G60 resulted in a significant reduction in liver weight ratio
341 compared to G60 alone.

342 < Figure 4. Organ to total body weight ratios for the liver (top) and spleen (bottom) of mice after
343 treatment with corn oil or activated carbon with and without 2,3,7,8 - tetrachlorodibenzo-p-
344 dioxin. >

345 3.4 Environmental implications

346 These results demonstrate that TCDD sequestration by structurally diverse ACs eliminate
347 its oral bioavailability to a mammalian (mouse) model. This result was not evident a priori since

348 molecular simulations of TCDD interactions with pores suggested more favorable energetics
349 with smaller micropores [38]. Variations in the pore structure of the ACs tested showed no
350 impact on the observed elimination of TCDD bioavailability. However, in actual practice at
351 remediation sites, other confounding interactions must be considered. For instance, soil/sediment
352 constituents have been attributed to the clogging of micropores and reduced contaminant
353 sorption capacity of certain ACs [46]. Likewise, pore clogging by natural organic matter (NOM)
354 in sediments over time has also been shown to attenuate contaminant sorption by AC [47],
355 although other studies including our own have shown that NOM additions enhanced uptake of
356 dioxin by ACs [48]. To be clear, contaminant sorption/sequestration alone is insufficient to
357 ensure concomitant reduction in mammalian (mouse) bioavailability. For example, TCDD
358 intercalated in the smectite clay saponite was equally bioavailable to the mammalian (mouse)
359 model as TCDD dissolved in corn oil [24], i.e. sorption by clay manifested no reduction in the
360 oral bioavailability of TCDD. Also, the solubilization of certain biochar components was
361 implicated as being responsible for increased bioaccessibility of sorbed polychlorinated
362 biphenyls in a simulated gastric fluid [49]. Having confirmed that structurally diverse ACs are
363 equally effective in eliminating the bioavailability of TCDD to a mammalian (mouse) model,
364 selection of AC materials for soil and sediment remediation should be further evaluated based on
365 other environmental processes relevant to in-situ application of AC sorbent amendments,
366 including the fouling of ACs by NOM and other materials as well as optimizing the mass transfer
367 of contaminants from environmental geosorbents and media to ACs used in this new remediation
368 technology.

369 **4. Conclusions**

370 The rapid acceptance of remediation strategies that employ sorbent amendments to
371 sequester contaminants in forms that reduce or eliminate bioavailability, specifically involving
372 AC, continues despite a paucity of studies that have evaluated their effectiveness with
373 appropriate mammalian models. Such studies are needed to establish that this remedy effectively
374 reduces PCDD bioavailability to mammals and hence mammalian exposure, and by inference is
375 protective of human health.

376 Recently, we demonstrated one commercial AC material, WPC AC, selected to maximize
377 (viz. higher proportion of micro vs meso pores) the irreversible binding of TCDD, could
378 sequester TCDD in a form that eliminated its bioavailability to an appropriate mammalian
379 (mouse) model. Not only were the results of the prior study using only WPC AC replicated, the
380 ability to eliminate TCDD bioavailability is apparently characteristic of AC materials
381 irrespective of their specific pore structures; three ACs with micropore volume ranging from
382 43.3 to 90.5 percent each eliminated TCDD bioavailability. By measuring cyp1A1 mRNA
383 induction in the liver, anti-sRBC IgM AFC response in the spleen, and organ to body weight
384 ratios, our results showed that ACs comprised of widely differing pore structures were equally
385 effective in the elimination of TCDD bioavailability. This suggests that chemisorption,
386 interactions with the material's specific surface, may be a driving factor, rather than simply pore
387 isolation.

388 This study also highlights the fact that existing standard methods for the extraction and
389 quantification of TCDD in soils and sediments are seemingly ineffective for use with porous
390 high surface area carbonaceous materials, or samples that contain such materials. One
391 implication is that previous surveys of PCDD/F contamination may underestimate their
392 abundance, especially in soils and sediments enriched with chars or black carbon. Development

393 of an extraction method for the efficient removal of PCDD/Fs from carbonaceous materials is
394 urgently needed.

395 In terms of the development of a new remediation technology, this study indicates that
396 AC materials, of various structures, have strong potential for use as in situ sorbent amendments
397 for soils and sediments impacted by PCDDs and similar poorly water-soluble organic
398 contaminants. With this understanding, identification of ideal AC materials for use in
399 remediation should focus on other factors including reducing environmental interactions such as
400 biofouling and maximizing sorption kinetics in environmental matrices. Understanding the mass
401 transfer kinetics of PCDD/Fs from contaminated soils and sediments into AC amendments is an
402 essential step in the further development and acceptance of this emerging remediation
403 technology.

404 **Acknowledgements**

405 The authors would like to acknowledge Ashwini Phadnis-Moghe, Jingeng Li, Natalia Kovalova,
406 Mike Rizzo, Joseph Henriquez, and Jiajun Zhou for their assistance with mouse handling as well
407 as Premachandra Gnanasiri for assistance with activated carbon preparations. Research reported
408 in this publication was supported by AgBio Research at Michigan State University and the
409 National Institute of Environmental Health Sciences of the National Institutes of Health under
410 Award Number P42ES004911. The content is solely the responsibility of the authors and does
411 not necessarily represent the official views of the National Institutes of Health.

412

413

414

415

416

417 **References**

- 418 [1] P.S. Kulkarni, J.G. Crespo, C.A.M. Afonso, Dioxins sources and current remediation
419 technologies - A review, *Environ. Int.* 34 (2008) 139–153.
420 doi:10.1016/j.envint.2007.07.009.
- 421 [2] R.E. Alcock, K.C. Jones, Dioxins in the environment: A review of trend data, *Environ.*
422 *Sci. Technol.* 30 (1996) 3133–3143. doi:10.1021/es960306z.
- 423 [3] C. Gu, H. Li, B.J. Teppen, S.A. Boyd, Octachlorodibenzodioxin formation on Fe(III)-
424 montmorillonite clay, *Environ. Sci. Technol.* 42 (2008) 4758–4763.
425 doi:10.1021/es7029834.
- 426 [4] M. Van Den Berg, L. Birnbaum, A.T.C. Bosveld, B. Brunström, P. Cook, M. Feeley, et
427 al., Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife,
428 *Environ. Health Perspect.* 106 (1998) 775–792. doi:10.1289/ehp.98106775.
- 429 [5] USEPA, An Inventory of Sources and Environmental Releases of Dioxin-Like
430 Compounds in the United States for the Years 1987, 1995, and 2000, 2006.
- 431 [6] P. Mocarelli, P.M. Gerthoux, D.G. Patterson, S. Milani, G. Limonta, M. Bertona, et al.,
432 Dioxin exposure, from infancy through puberty, produces endocrine disruption and affects
433 human semen quality, *Environ. Health Perspect.* 116 (2008) 70–77.
434 doi:10.1289/ehp.10399.
- 435 [7] S.S. White, L.S. Birnbaum, An overview of the effects of dioxins and dioxin-like
436 compounds on vertebrates, as documented in human and ecological epidemiology, *J.*
437 *Environ. Sci. Heal. - Part C Environ. Carcinog. Ecotoxicol. Rev.* 27 (2009) 197–211.
438 doi:10.1080/10590500903310047.
- 439 [8] S.J. Bursian, J.L. Newsted, M.J. Zwiernik, Polychlorinated Biphenyls, Polychlorinated
440 Dibenzo-P-Dioxins and Polychlorinated Dibenzofurans, in: *Reprod. Dev. Toxicol.*,
441 Academic Press, 2011: pp. 543–567. doi:10.1016/B978-0-12-382032-7.10041-4.
- 442 [9] R.D. Stedtfeld, T.M. Stedtfeld, K.A. Fader, M.R. Williams, P. Bhaduri, J. Quensen, et al.,
443 TCDD influences reservoir of antibiotic resistance genes in murine gut microbiome,
444 *FEMS Microbiol. Ecol.* 93 (2017). doi:10.1093/femsec/fix058.
- 445 [10] T.S. Bridges, K.E. Gustavson, Paul Schroeder, S.J. Ells, D. Hayes, S.C. Nadeau, et al.,
446 Dredging processes and remedy effectiveness: Relationship to the 4 Rs of environmental
447 dredging, *Integr. Environ. Assess. Manag.* 6 (2010) 619–630. doi:10.1002/ieam.71.
- 448 [11] N.R. Council, Bioavailability of Contaminants in Soils and Sediments, National
449 Academies Press, Washington, D.C., 2003. doi:10.17226/10523.

- 450 [12] A. Akcil, C. Erust, S. Ozdemiroglu, V. Fonti, F. Beolchini, A review of approaches and
451 techniques used in aquatic contaminated sediments: Metal removal and stabilization by
452 chemical and biotechnological processes, *J. Clean. Prod.* 86 (2015) 24–26.
453 doi:10.1016/j.jclepro.2014.08.009.
- 454 [13] U. Ghosh, R.G. Luthy, G. Cornelissen, D. Werner, C.A. Menzie, In-situ sorbent
455 amendments: A new direction in contaminated sediment management, *Environ. Sci.*
456 *Technol.* 45 (2011) 1163–1168. doi:10.1021/es102694h.
- 457 [14] S. Josefsson, M. Schaanning, G.S. Samuelsson, J.S. Gunnarsson, I. Olofsson, E. Eek, K.
458 Wiberg, Capping efficiency of various carbonaceous and mineral materials for in situ
459 remediation of polychlorinated dibenzo-p-dioxin and dibenzofuran contaminated marine
460 sediments: Sediment-to-water fluxes and bioaccumulation in boxcosm tests, *Environ. Sci.*
461 *Technol.* 46 (2012) 3343–3351. doi:10.1021/es203528v.
- 462 [15] G. Cornelissen, K. Amstaetter, A. Hauge, M. Schaanning, B. Beylich, J.S. Gunnarsson, et
463 al., Large-scale field study on thin-layer capping of marine PCDD/F-contaminated
464 sediments in grenlandfjords, Norway: Physicochemical effects, *Environ. Sci. Technol.* 46
465 (2012) 12030–12037. doi:10.1021/es302431u.
- 466 [16] S.K. Fagervold, Y. Chai, J.W. Davis, M. Wilken, G. Cornelissen, U. Ghosh,
467 Bioaccumulation of polychlorinated dibenzo-p-dioxins/dibenzofurans in *E. fetida* from
468 floodplain soils and the effect of activated carbon amendment, *Environ. Sci. Technol.* 44
469 (2010) 5546–5552. doi:10.1021/es9027138.
- 470 [17] Y. Chai, R.J. Currie, J.W. Davis, M. Wilken, G.D. Martin, V.N. Fishman, U. Ghosh,
471 Effectiveness of activated carbon and biochar in reducing the availability of
472 polychlorinated dibenzo-p-dioxins/dibenzofurans in soils, *Environ. Sci. Technol.* 46
473 (2012) 1035–1043. doi:10.1021/es2029697.
- 474 [18] R.A. Budinsky, J.C. Rowlands, S. Casteel, G. Fent, C.A. Cushing, J. Newsted, J.P. Giesy,
475 M. V. Ruby, L.L. Aylward, A pilot study of oral bioavailability of dioxins and furans from
476 contaminated soils: Impact of differential hepatic enzyme activity and species differences,
477 *Chemosphere.* 70 (2008) 1774–1786. doi:10.1016/j.chemosphere.2007.08.035.
- 478 [19] MDEQ, DEQ Basis of Decision and Response for a Site-Specific Residential Direct
479 Contact Cleanup Criterion (SSRDCC) for Dioxins/Furans (D/F) Toxic Equivalents (TEQ)
480 for Midland Area Soils, 2012. [https://www.michigan.gov/documents/deq/2012-3-
481 9_Final_Midland_SSRDCC_Recommendation_378921_7.pdf](https://www.michigan.gov/documents/deq/2012-3-9_Final_Midland_SSRDCC_Recommendation_378921_7.pdf) (accessed September 13,
482 2018).
- 483 [20] W.N. Beyer, E.E. Connor, S. Gerould, Estimates of Soil Ingestion by Wildlife, *J. Wildl.*
484 *Manage.* 58 (1994) 375. doi:10.2307/3809405.
- 485 [21] D.A. Kreulen, Lick use by large herbivores: a review of benefits and banes of soil
486 consumption, *Mamm. Rev.* 15 (1985) 107–123. doi:10.1111/j.1365-2907.1985.tb00391.x.
- 487 [22] E.J. Stanek, E.J. Calabrese, B. Xu, Meta-Analysis of Mass-Balance Studies of Soil
488 Ingestion in Children, *Risk Anal.* 32 (2012) 433–447. doi:10.1111/j.1539-
489 6924.2011.01673.x.

- 490 [23] S.A. Boyd, J.B. Sallach, Y. Zhang, R. Crawford, H. Li, C.T. Johnston, et al., Sequestration
491 of 2,3,7,8-tetrachlorodibenzo-p-dioxin by activated carbon eliminates bioavailability and
492 the suppression of immune function in mice, *Environ. Toxicol. Chem.* 9999 (2017) 1–8.
493 doi:10.1002/etc.3815.
- 494 [24] S.A. Boyd, C.T. Johnston, T.J. Pinnavaia, N.E. Kaminski, B.J. Teppen, H. Li, et al.,
495 Suppression of humoral immune responses by 2,3,7,8-tetrachlorodibenzo-p-dioxin
496 intercalated in smectite clay, *Environ. Toxicol. Chem.* 30 (2011) 2748–2755.
497 doi:10.1002/etc.701.
- 498 [25] B.L.F. Kaplan, R.B. Crawford, N. Kovalova, A. Arencibia, S.S. Kim, T.J. Pinnavaia, et
499 al., TCDD adsorbed on silica as a model for TCDD contaminated soils: Evidence for
500 suppression of humoral immunity in mice, *Toxicology.* 282 (2011) 82–87.
501 doi:10.1016/j.tox.2011.01.013.
- 502 [26] R. Stedtfeld, J. Sallach, R. Crawford, T. Stedtfeld, M. Williams, H. Waseem, et al., TCDD
503 administered on activated carbon eliminates bioavailability and subsequent shifts to a key
504 murine gut commensal, *Appl Microbiol Biotech.* In press (2017). doi:10.1007/s00253-
505 017-8460-9.
- 506 [27] G.R. Bond, The role of activated charcoal and gastric emptying in gastrointestinal
507 decontamination: A state-of-the-art review, *Ann. Emerg. Med.* 39 (2002) 273–286.
508 doi:10.1067/mem.2002.122058.
- 509 [28] R.M. Cook, K.A. Wilson, Removal of Pesticide Residues from Dairy Cattle, *J. Dairy Sci.*
510 54 (1971) 712–718. doi:10.3168/jds.S0022-0302(71)85913-1.
- 511 [29] K.A. Engebretson, K.L. Davison, Dieldrin accumulation and excretion by rats fed
512 phenobarbital and carbon, *Bull. Environ. Contam. Toxicol.* 6 (1971) 391–400.
513 doi:10.1007/BF01684358.
- 514 [30] D. Mohan, C.U. Pittman, Arsenic removal from water/wastewater using adsorbents-A
515 critical review, *J. Hazard. Mater.* 142 (2007) 1–53. doi:10.1016/j.jhazmat.2007.01.006.
- 516 [31] J.L. Figueiredo, M.F.R. Pereira, M.M.A. Freitas, J.J.M. Órfão, Modification of the surface
517 chemistry of activated carbons, *Carbon N. Y.* 37 (1999) 1379–1389. doi:10.1016/S0008-
518 6223(98)00333-9.
- 519 [32] G. Lesage, M. Sperandio, L. Tiruta-Barna, Analysis and modelling of non-equilibrium
520 sorption of aromatic micro-pollutants on GAC with a multi-compartment dynamic model,
521 *Chem. Eng. J.* 160 (2010) 457–465. doi:10.1016/j.cej.2010.03.045.
- 522 [33] C.T. Johnston, B. Khan, E.F. Barth, S. Chattopadhyay, S.A. Boyd, Nature of the interlayer
523 environment in an organoclay optimized for the sequestration of Dibenzo-p -dioxin,
524 *Environ. Sci. Technol.* 46 (2012) 9584–9591. doi:10.1021/es300699y.
- 525 [34] J. Zachara, S. Brantley, J. Chorover, R. Ewing, S. Kerisit, C. Liu, et al., Internal Domains
526 of Natural Porous Media Revealed: Critical Locations for Transport, Storage, and
527 Chemical Reaction, *Environ. Sci. Technol.* 50 (2016) 2811–2829.
528 doi:10.1021/acs.est.5b05015.
- 529 [35] B. Panella, M. Hirscher, S. Roth, Hydrogen adsorption in different carbon nanostructures,

- 530 Carbon N. Y. 43 (2005) 2209–2214. doi:10.1016/j.carbon.2005.03.037.
- 531 [36] A. Bembnowska, R. Pelech, E. Milchert, Adsorption from aqueous solutions of
532 chlorinated organic compounds onto activated carbons, *J. Colloid Interface Sci.* 265
533 (2003) 276–282. doi:10.1016/S0021-9797(03)00532-0.
- 534 [37] B. Coasne, A. Galarneau, R.J.M. Pellenq, F. Di Renzo, Adsorption, intrusion and freezing
535 in porous silica: the view from the nanoscale, *Chem. Soc. Rev.* 42 (2013) 4141.
536 doi:10.1039/c2cs35384a.
- 537 [38] F. Gao, S. Qu, C. Liu, H. Li, C.T. Johnston, B.J. Teppen, S.A. Boyd, Molecular dynamics
538 study of the influence of activated carbon pore structures on dioxin adsorption, in: 253rd
539 ACS Natl. Meet. San Fr. CA, 2017.
- 540 [39] W. Ying Shlu, F.A.P.C. Gobas, D. Mackay, W. Doucette, A. Andren, Physical-Chemical
541 Properties of Chlorinated Dibenzo-p-dioxins, *Environ. Sci. Technol.* 22 (1988) 651–658.
542 doi:10.1021/es00171a006.
- 543 [40] US Environmental Protection Agency, Method 1613: Tetra- through Octa-Chlorinated
544 Dioxins and Furans by Isotope Dilution HRGC/HRMS, 1994.
- 545 [41] N.K. Jerne, A.A. Nordin, Plaque Formation in Agar by Single Antibody-Producing Cells.,
546 *Science.* 140 (1963) 405. doi:10.1126/science.140.3565.405.
- 547 [42] K.J. Livak, T.D. Schmittgen, Analysis of Relative Gene Expression Data Using Real-
548 Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method, *Methods.* 25 (2001) 402–408.
549 doi:10.1006/meth.2001.1262.
- 550 [43] I. Winald, D.J. Miller, E. De Pauw, S.B. Hawthorne, Supercritical fluid extraction and
551 accelerated solvent extraction of dioxins from high- and low-carbon fly ash, *Anal. Chem.*
552 72 (2000) 3916–3921. doi:10.1021/ac9914972.
- 553 [44] B. Beckingham, U. Ghosh, Field-scale reduction of PCB bioavailability with activated
554 carbon amendment to river sediments, *Environ. Sci. Technol.* 45 (2011) 10567–10574.
555 doi:10.1021/es202218p.
- 556 [45] N. Alexandrou, M.J. Lawrence, J. Pawliszyn, Cleanup of Complex Organic Mixtures
557 Using Supercritical Fluids and Selective Adsorbents, *Anal. Chem.* 64 (1992) 301–311.
558 doi:10.1021/ac00027a011.
- 559 [46] K. Amstaetter, E. Eek, G. Cornelissen, Sorption of PAHs and PCBs to activated carbon:
560 Coal versus biomass-based quality, *Chemosphere.* 87 (2012) 573–578.
561 doi:10.1016/j.chemosphere.2012.01.007.
- 562 [47] A.M.P. Oen, B. Beckingham, U. Ghosh, M.E. Kruså, R.G. Luthy, T. Hartnik, T.
563 Henriksen, G. Cornelissen, Sorption of organic compounds to fresh and field-aged
564 activated carbons in soils and sediments, *Environ. Sci. Technol.* 46 (2012) 810–817.
565 doi:10.1021/es202814e.
- 566 [48] S. Qu, C.T. Johnston, B.J. Teppen, S.A. Boyd, Sorption of dibenzo-p-dioxin/dibenzofuran
567 by high surface area carbonaceous geosorbents, in: 248th ACS Natl. Meet. San Fr. CA,
568 2014.

569 [49] B. Beckingham, U. Ghosh, Differential bioavailability of polychlorinated biphenyls
570 associated with environmental particles: Microplastic in comparison to wood, coal and
571 biochar, *Environ. Pollut.* 220 (2017) 150–158. doi:10.1016/j.envpol.2016.09.033.