

This is a repository copy of *Potential metabolism of pharmaceuticals in radish : Comparison of in vivo and in vitro exposure*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/151017/>

Version: Accepted Version

Article:

Li, Yuanbo, Chuang, Ya Hui, Sallach, J. Brett orcid.org/0000-0003-4588-3364 et al. (3 more authors) (2018) Potential metabolism of pharmaceuticals in radish : Comparison of in vivo and in vitro exposure. *Environmental Pollution*. pp. 962-969. ISSN 1873-6424

<https://doi.org/10.1016/j.envpol.2018.07.060>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Potential metabolism of pharmaceuticals in radish: Comparison of in vivo and**
2 **in vitro exposure**

3 Yuanbo Li, Ya-Hui Chuang, J. Brett Sallach, Wei Zhang, Stephen A. Boyd, and Hui Li*
4 Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing,
5 Michigan 48824, USA

6 * Corresponding Author: Phone: 517-353-0151; fax: 517-355-0270; e-mail: lihui@msu.edu
7

8 **ABSTRACT**

9 Metabolism of pharmaceuticals in plants is important to evaluate their fate and accumulation in
10 vegetables, and subsequently the risks to human health. However, limited knowledge is available
11 to evaluate the metabolism of pharmaceuticals in plants due to the lack of appropriate research
12 approaches. In this study, radish was selected as a model plant to investigate the metabolism of
13 pharmaceuticals in intact plants (in vivo) growing in hydroponic solution and in plant tissue
14 enzyme extracts (in vitro). For caffeine, six phase-I demethylation metabolites identified in the
15 intact radish plant were also found in the plant enzyme extracts. After 7 days of in vivo exposure,
16 the amount of the identified metabolites was about 5.4 times greater than the parent caffeine in
17 radish roots. Furthermore, the metabolism potential of fifteen pharmaceuticals in radish was
18 evaluated on the basis of mass balance. After 7 days of hydroponic exposure, oxytetracycline,
19 trimethoprim, carbamazepine, lincomycin, monensin and tylosin manifested relatively less extent
20 of metabolism with the mass recoveries ranging from 52.3 to 78.2%. In contrast, 17 β -estradiol,
21 sulfamethoxazole, sulfadiazine, estrone, triclosan, acetaminophen, caffeine, carbadox and
22 lamotrigine underwent extensive metabolism with only 3.0 to 32.1% of the parent compound
23 recovered. In the in vitro system, 17 β -estradiol, estrone, triclosan, oxytetracycline,
24 acetaminophen, sulfadiazine and sulfamethoxazole were readily metabolized in radish root

25 enzyme extracts with 1.8 to 34.0% remaining after 96-h exposure. While in the leaf enzyme
26 extracts, only triclosan was rapidly metabolized with 49.2% remaining, and others
27 pharmaceuticals were $\geq 60\%$, indicating that the varying extents of metabolism occurred in
28 different plant parts. This study highlights the importance of pharmaceutical metabolism in
29 plants, and suggests that plant tissue enzyme extracts could serve as an alternative tool to assess
30 pharmaceutical metabolism in plants.

31

32 Capsule: Similar metabolism patterns were observed for rapidly metabolized pharmaceuticals in
33 both in vivo (radish tissue enzyme extracts) and in vivo (the intact plant) exposure.

34

35 Keywords: Plant metabolism; Plant uptake; Pharmaceuticals; Plant tissue enzyme extracts.

36

37 **1. Introduction**

38 Reclaimed water reuse in agricultural irrigation has been increasingly practiced to
39 alleviate the burden of water scarcity, especially in arid and semiarid regions (Bischel et al., 2011;
40 Sato et al., 2013). Biosolids and animal manures are also commonly applied to agricultural lands
41 for their fertility values, and as a convenient disposal approach as well (Clarke and Smith, 2011;
42 Cogger et al., 2013; Kumar et al., 2005). These practices also release pharmaceuticals to the
43 environment, resulting in their ubiquitous presence in soils with concentrations at ng kg^{-1} to μg
44 kg^{-1} levels (Chen et al., 2014; Durán-Alvarez et al., 2009; Kinney et al., 2006; Vazquez-Roig et
45 al., 2010). These pharmaceuticals could enter vegetables from soils via root uptake, which serves
46 as the starting point in the food chain of human and animal dietary consumption. It has been well
47 documented that vegetables could accumulate a range of pharmaceuticals from soils (Malchi et

48 al., 2014; Prosser and Sibley, 2015; Tanoue et al., 2012; Wu et al., 2013). However, only a few
49 attempts have been made to elucidate the metabolism of pharmaceuticals in plants. In addition to
50 uptake and translocation, metabolic transformation of pharmaceuticals is also an important
51 process influencing their residue levels in plants and their potential risk of food safety.

52 Pharmaceuticals could be metabolized in plants by multiple pathways and form a variety
53 of transformation products (Macherius et al., 2012; Miller et al., 2016; Riemenschneider et al.,
54 2017). In general, the metabolic processes involve three phases of reactions. Pharmaceuticals
55 could be mediated by plant enzymes and form more polar and water-soluble products via phase I
56 reactions (oxidation, reduction and hydrolysis). In phase II reactions, pharmaceuticals and their
57 phase-I metabolites could be conjugated to endogenous plant biomolecules such as amino acids,
58 sugars and glutathione, and form relatively larger-sized molecules. Compared to their parent
59 compounds these conjugates generally demonstrate an enhanced water solubility and higher
60 mobility in plants. These conjugates formed by phase II reactions could be sequestered in plant
61 vacuoles or cell walls by phase III reactions (He et al., 2017; Macherius et al., 2012; Sandermann,
62 1992). The metabolic transformations of some pharmaceuticals in plants have been studied such
63 as triclosan (Macherius et al., 2012), benzotriazole (LeFevre et al., 2015), ibuprofen (He et al.,
64 2017), diclofenac (Huber et al., 2012), iopromide (Cui et al., 2017) and carbamazepine
65 (Riemenschneider et al., 2017). Carbamazepine was moderately metabolized in tomato after 35
66 days of exposure, 21 phase I and II transformation products were identified, and their total
67 amount was equivalent to ~ 45% of carbamazepine accumulated in plants (Riemenschneider et
68 al., 2017). Among these metabolites, the 10,11-epoxycarbamazepine manifested even greater
69 toxic potency than carbamazepine (Tomson et al., 1990). In many cases, the amount of formed
70 metabolites exceeded that of the parent compounds remaining in plants. For instance, the sum of

71 eight phase II triclosan conjugates was 5 times greater than triclosan in carrots during two-month
72 growth in greenhouse (Macherius et al., 2012). In another study, three major metabolites of
73 benzotriazole were found to be >1.5 times more than benzotriazole after 8-day exposure in
74 hydroponic Arabidopsis (LeFevre et al., 2015). Besides the intact plants, the metabolism of
75 pharmaceuticals was also studied using plant cell cultures (Fu et al., 2017; Huber et al., 2009;
76 Macherius et al., 2012; Marsik et al., 2017; Sauvêtre et al., 2018; Wu et al., 2016). Triclosan,
77 naproxen, diclofenac, ibuprofen, gemfibrozil, sulfamethoxazole and atorvastatin were found to
78 rapidly disappeared in carrot cell cultures with 0.4 to 47.3% remaining after 90 h of exposure
79 (Wu et al., 2016). The lack of assessment of metabolism could underestimate the total
80 accumulation of pharmaceuticals and metabolites in plants and the potential risk related to the
81 consumption of contaminated agricultural products. Paltiel et al. (2016) recently reported that
82 human consumption of fresh produce irrigated with reclaimed water could lead to the appearance
83 of bioactive metabolites of carbamazepine in their urine.

84 In vivo exposure (intact plants) is an effective way to examine the fate and metabolism of
85 pharmaceuticals in plant-water (or -soil) systems. However, the intact plant experiments
86 conducted in hydroponic solution or soil are usually labor-intensive and time-consuming. The
87 presence of microorganisms and root exudates also influence pharmaceutical metabolism, which
88 cannot be excluded from the in vivo studies (Yu et al., 2013; Zhang et al., 2016). The uptake and
89 transport processes could further complicate the metabolism of pharmaceuticals in plants by
90 governing the input and output of parent compounds/metabolites in a specific plant part. In
91 plants, enzyme-mediated transformation has been considered as a major process to the
92 metabolism of xenobiotics in plants (Huang et al., 2013; Van Eerd et al., 2003). Plant enzyme
93 extracts have been used in the in vitro system to investigate the enzyme-mediated metabolism of

94 polybrominated diphenyl ethers (Huang et al., 2013), estrogens (Card et al., 2013) and metformin
95 (Cui and Schröder, 2016). These studies showed that the metabolites from the reactions with
96 plant enzyme extracts were similar to the products formed in the intact plants. Therefore, plant
97 enzyme extracts may serve as an alternative system to provide a simple and quick approach to
98 evaluate the metabolism of pharmaceuticals in plants, which is minimally affected by plant
99 microbial, uptake and transport processes.

100 The objective of this study was to compare the pharmaceutical metabolism in radish
101 using both in vitro and in vivo experiments. Radish was selected because this root vegetable is
102 usually consumed in raw, and might represent the worst case of exposure scenarios. Caffeine was
103 selected as one representative pharmaceutical for further evaluation of the formed metabolites in
104 both intact radish and radish tissue enzyme extracts because it is one of the most frequently
105 detected pharmaceuticals in vegetables (Goldstein et al., 2014; Malchi et al., 2014; Wu et al.,
106 2014; Wu et al., 2013). In addition, we expanded the experiments to fifteen commonly-used
107 pharmaceuticals with a wide range of physicochemical properties to assess their metabolism
108 potential in radish. This study provides new knowledge on the metabolic transformation of
109 pharmaceuticals in plants. The in vitro exposure method could serve as an alternative to quickly
110 screen the metabolism potential of pharmaceuticals in plants.

111

112 **2. Materials and methods**

113

114 2.1. Chemicals and reagents

115 Fifteen pharmaceuticals were purchased from Sigma-Aldrich (St. Louis, MO, USA)
116 including acetaminophen, caffeine, carbamazepine, sulfadiazine, sulfamethoxazole, lamotrigine,

117 carbadox, estrone, 17 β -estradiol, triclosan, trimethoprim, lincomycin, oxytetracycline, monensin
118 and tylosin. These pharmaceuticals were selected because they are frequently detected in the
119 environment, and demonstrate a wide range of physicochemical properties (Table S1). Caffeine
120 metabolites xanthine, 3-methylxanthine, 7-methylxanthine, theophylline, paraxanthine and
121 theobromine were purchased from Sigma-Aldrich (St. Louis, MO, USA). Ceramic homogenizers,
122 octadecylsilane (C18), primary secondary amine (PSA), and graphitized carbon black (GCB)
123 were purchased from Agilent Technologies (Santa Clara, CA, USA). Disodium
124 ethylenediaminetetraacetate (Na_2EDTA), formic acid, glacial acetic acid, and sodium chloride
125 (NaCl) were purchased from J.T. Baker (Phillipsburg, NJ, USA). All organic solvents were of
126 HPLC grade from Fisher (Fair Lawn, NJ, USA). Water used in this study was produced from a
127 Milli-Q water purification system (Billerica, MA, USA).

128

129 2.2. In vivo hydroponic experiment

130 Radish seeds (*Raphanus sativus*, Burpee & Co., Warminster, PA) were germinated on
131 moistened filter paper for 4 days. The seedlings were then transferred to a plastic container with
132 10 L of nutrient solution (pH 6.5) (Hydrodynamics International, Lansing, MI, USA). After 14
133 days of growth, the radish plants (~10 cm in height) were used in the exposure experiments.
134 Each radish plant was carefully transferred to a 250-mL glass jar containing 230 mL of nutrient
135 solution with spiked pharmaceuticals. This exposure experiment was conducted with two
136 treatments (1) the radish was exposed to 200 ng mL^{-1} of caffeine in nutrient solution; (2) the
137 radish was exposed to a mixture of fifteen pharmaceuticals with the initial concentration of 100
138 ng mL^{-1} for each compound. The solution was continuously aerated via Teflon tubing connected
139 to a fusion air pump. The jars were wrapped with aluminum foil to prevent the potential

140 photolysis of pharmaceuticals and to minimize algae growth. Pharmaceutical-free controls
141 (radish only) and pharmaceutical solution controls (without radish) were also conducted in the
142 experiments. All controls and exposure experiments were performed in triplicate. During the
143 experiment, nutrient solution (without pharmaceuticals) was added daily to each jar to
144 compensate the water loss from plant transpiration. The glass jars were placed in a controlled
145 growth chamber at 20 ± 2 °C. The light cycle was set up as 10 h of fluorescent light every day
146 with the intensity of 150 mmol/m²/s followed by 14 h of darkness. For the treatment with only
147 caffeine, radish roots, leaves and solution were sampled at day 1, 3 and 7. For the exposure of the
148 mixture of pharmaceuticals, aqueous solution, radish roots and leaves were collected after 7 days
149 of growth. The radish samples were thoroughly rinsed with deionized water, wiped with tissue
150 paper, weighed, chopped into small pieces, freeze-dried and ground to fine powders. All samples
151 were stored in freezer at -20 °C prior to the extraction of pharmaceuticals.

152

153 2.3. In vitro reactions with radish enzyme extracts

154 Pharmaceutical metabolism was also examined in vitro using the crude enzyme extracts
155 from radish tissues. Radish enzyme extracts were prepared using the method reported by Card et
156 al. (Card et al., 2013). In brief, fresh radish tissues were separated into roots and leaves, chopped
157 into small pieces, quickly frozen, ground into powders in liquid nitrogen, and then extracted with
158 50 mM potassium phosphate buffer (pH 7.0) at the mass to solution ratio of 1: 2 (g: mL) in ice
159 for 30 min. The homogenate was centrifuged at 8000 g for 40 min at 4 °C. The supernatant of
160 tissue enzyme extracts was collected, passed through a 0.22- μ m cellulose ester membrane
161 (Millipore, Cork, Ireland), and immediately stored in an ice bath prior to use.

162 The in vitro exposure experiment was performed by mixing 200 ng mL⁻¹ of caffeine or a
163 mixture of fifteen pharmaceuticals at 100 ng mL⁻¹ for each chemical with the radish enzyme
164 extracts. The reactant mixtures were incubated in 1.5-mL centrifuge tubes in a water bath (25 °C),
165 and the reaction was quenched by adding 50 µL of glacial acetic acid. The experiments were
166 conducted in triplicate, along with enzyme extracts-free controls. The resultant solution was
167 sampled at the time intervals of 0, 4, 10, 24, 48, 72 and 96 h. An aliquot of the solution (0.5 mL)
168 was diluted to 1.0 mL with methanol and subject to pharmaceutical analysis. The proteins
169 concentration in radish enzyme extracts was used to estimate the activity of the enzyme mixture,
170 which was measured during the experiment using Bradford assay (Bradford, 1976). The proteins
171 concentration in the radish enzyme extracts demonstrated relatively stable during 96 h of
172 reaction (Figure S1).

173

174 2.4. Analysis of pharmaceuticals

175 Pharmaceuticals in radish root and leaf samples were extracted using a modified
176 QuEChERS method (Chuang et al., 2015). The detailed extraction procedure is given in the
177 supporting information (SI). Hydroponic solution was filtered through a 0.22-µm cellulose ester
178 membrane, and an aliquot of the filtrate (0.5 mL) was diluted to 1.0 mL with methanol and
179 injected directly into a LC-MS/MS system. The LC-MS/MS system consisted of a Shimadzu
180 prominence high-performance liquid chromatography (Columbia, MD, USA) coupled to a Sciex
181 4500 triple quadrupole mass spectrometer (Foster City, CA, USA), and a 50 mm × 2.1 mm
182 Agilent C18 column (Torrance, CA, USA). The optimized conditions and quality assurance of
183 the LC-MS/MS is detailed in SI. Multiple reaction monitoring (MRM) parameters are listed in

184 Tables S2 and S3. The corresponding matrix recoveries and method detection limits (MDLs) for
185 the studied pharmaceuticals are summarized in Tables S4-S7.

186 The metabolism of pharmaceuticals in radish enzyme extracts was fit to the first-order
187 kinetic model $C_i = C_0 e^{-kt}$, where C_i and C_0 is pharmaceutical concentration at sampling time t (h)
188 and the beginning time, respectively, and k is the rate constant (h^{-1}). The dissipation half-life ($T_{1/2}$)
189 was calculated as $T_{1/2} = \ln 2/k$.

190

191 **3. Results and discussion**

192 3.1. Biotransformation of caffeine in hydroponic system

193 The mass distribution of caffeine in radish leaves, roots and hydroponic solution is shown
194 in Figure S2. On the basis of the initial amount of caffeine added to the solution, 11.8, 48.9 and
195 82.6% disappeared in the system after 1, 3 and 7 days of exposure, respectively. The quick
196 depletion of caffeine in the solution, compared to the unplanted controls ($< 5\%$ of loss after 7
197 days), suggests that caffeine could be taken up by radish and undergoes relatively rapid
198 transformation in the hydroponic system. Six demethylation metabolites from phase I reactions
199 were identified and quantified using their authentic standards including xanthine, 3-
200 methylxanthine, 7-methylxanthine, theophylline, paraxanthine and theobromine. All six
201 metabolites were detected in both radish roots and leaves. None of the metabolites was found in
202 the radish-free controls indicating that the metabolism is plant-associated. Only a small amount
203 of xanthine was found in the hydroponic solution with radishes at day 3 and 7, and no other
204 metabolites were found in the solution (Table S8). The presence of xanthine in the solution could
205 be attributed to the excretion from plants and/or caffeine transformation by root exudates.

206 The molar fractions of the six metabolites increased with exposure time in both roots and
207 leaves (Figure 1). In the radish roots, the parent compound caffeine gradually decreased from
208 67.1% at day 1 to 49.8% at day 3, and to 15.6% at day 7, indicating that caffeine was
209 substantially metabolized. The product xanthine (loss of three $-CH_3$ groups from caffeine) was
210 the most abundant metabolite with the molar fraction increasing from 16.7% at day 1 to 58.5% at
211 day 7. Theobromine (loss of one $-CH_3$ group) was the major intermediate with the fraction
212 increasing from 9.0% at day 1 to 13.2% at day 7. This time-dependent transformation pattern
213 was also observed for 3- and 7-methylxanthine with their molar fractions up to 5.3% at the end
214 of experiment. Theophylline and paraxanthine (loss of one $-CH_3$ group) were the least abundant
215 metabolites with the molar fractions $< 3.8\%$ (day 3); their fractions increased between day 1 and
216 3 and then decreased from day 3 to 7. Compared with the metabolism occurring in radish roots,
217 caffeine experienced less transformation in the leaves (Figure 1). The parent compound caffeine
218 was predominant with the molar fraction of 94.6% at day 1, 77.5% at day 3, and 68.1% at day 7.
219 Xanthine was the most abundant metabolite at day 1 (2.2%) and day 3 (9.6%), and theobromine
220 was most abundant at day 7 (9.9%). Given that these phase I reaction products all contained the
221 moiety of xanthine that maintains the bioactivity of the parent compound (Goth and Cleaver,
222 1976), the extensive biotransformation of caffeine in radish, particularly in the edible root
223 fraction, could lead to an underestimate of the risk associated with food safety if only
224 considering the presence of the parent compound.

225

226 3.2. Transformation of caffeine in radish enzyme extracts

227 The results of the *in vivo* hydroponic experiment provided the information on the
228 metabolism of caffeine in the intact radish. However, the uptake and translocation of caffeine in

229 radish, as well as radish growth, could add more confounding impacts to elucidating the
230 metabolism process. The in vitro exposure experiments with radish enzyme extracts offer an
231 alternative to investigate the metabolism of pharmaceuticals in plants. The amount of caffeine
232 and its metabolites formed in the radish tissue enzyme extracts as a function of time is shown in
233 Figure 2. During 96-h exposure, the loss of caffeine in root enzyme extracts was 29.9%, which
234 was approximately 2 times that in leaf enzyme extracts (15.4%) (Figure 2). Meanwhile, the total
235 amount of metabolites formed by demethylation reactions in root enzyme extracts was
236 approximately 2 times that in leaf enzyme extracts, suggesting that a larger extent of
237 biotransformation occurred in radish roots. This is consistent with the results of the in vivo
238 hydroponic experiment in which the larger fraction of metabolites was formed in radish roots
239 than in leaves. The amount of the six demethylation metabolites continued to increase over time
240 in both root and leaf enzyme extracts, except for xanthine in the leaf enzyme extracts which
241 decreased slightly from 72 h to 96 h. The amount of metabolites formed in both tissue enzyme
242 extracts at 96 h of exposure ranked in the order of xanthine > theobromine > 3-methylxanthine
243 \approx 7-methylxanthine > theophylline \approx paraxanthine (Figure 2), which is also consistent with the
244 ranking of demethylation metabolites found in the in vivo intact radish. In the in vivo hydroponic
245 experiment, caffeine accumulation in roots and leaves varied with time due to the continuous
246 water movement into radish; therefore, it is not feasible to compare the absolute amount of the
247 formed metabolites between the in vitro and in vivo experiments. However, the consistence in the
248 types of metabolites and their rankings in magnitude in both radish roots and leaves suggests that
249 the in vitro exposure to plant tissue enzyme extracts could provide a convenient alternative to
250 evaluate the metabolism of pharmaceuticals in a specific plant compartment (Card et al., 2013;
251 Cui and Schröder, 2016).

252 In both in vivo and in vitro studies, phase I demethylation reaction is identified as a
253 metabolism pathway for caffeine in radish (Figure 3). The possible transformation route follows
254 stepwise N-demethylation reactions via the loss of one $-CH_3$ group to form paraxanthine,
255 theobromine and theophylline, loss of the second $-CH_3$ group to form 7- and 3-methylxanthine,
256 and loss of all three $-CH_3$ groups to form the end product xanthine. The initial N-demethylation
257 reaction (loss of one $-CH_3$ group) is common in mammals and insects, which is believed to be
258 mediated by cytochrome P450 enzyme systems (i.e. CYP1A2 isoenzyme) (Berthou et al., 1991;
259 Berthou et al., 1992; Coelho et al., 2015; Kot and Daniel, 2008). The N-demethylation reaction
260 of antidiabetic agent metformin was also observed in *Typha latifolia* (Cui and Schröder, 2016).
261 P450 enzymes are also commonly present in plants, and catalyze many metabolic reactions of
262 xenobiotics including some pharmaceuticals and herbicides (Fonne-Pfister et al., 1988; Huber et
263 al., 2009; Siminszky, 2006; Thies et al., 1996). Caffeine is believed to undergo similar enzyme-
264 mediated transformation in radish. In mammals paraxanthine represents the large fraction of
265 the primary metabolites (70-80%) during the first step of demethylation (Berthou et al., 1991;
266 Kalow and Tang, 1993; Kot and Daniel, 2008). However, in this study theobromine was the most
267 abundant intermediate in radish.

268

269 3.3. Mass distribution of pharmaceuticals in hydroponic system

270 Pharmaceuticals have been frequently detected in vegetables grown in fields amended
271 with biosolids or irrigated with reclaimed water (Christou et al., 2017; Malchi et al., 2014; Wu et
272 al., 2014); however, their metabolism in plants still remains largely unknown. To obtain more
273 basic information on the metabolism of pharmaceuticals in vegetables, we further assessed
274 metabolism potential of fifteen pharmaceuticals in radish on the basis of mass balance using the

275 hydroponic experiment. The mass recoveries of pharmaceuticals in the radish-free controls
276 ranged from 92.0 to 108.2% indicating that pharmaceuticals in nutrient solution were relatively
277 stable, with the only exception of oxytetracycline (14.5%) (Table S9). The substantial loss of
278 oxytetracycline could be due to the rapid hydrolysis in the nutrient solution at pH 6.5 (near
279 neutral), which is the most favorable condition for the hydrolysis of oxytetracycline (Xuan et al.,
280 2009).

281 The mass distribution of the fifteen pharmaceuticals in solution, roots and leaves after 7-d
282 exposure is presented in Figure 4. The mass recoveries varied between 3.0 and 78.2% of the
283 initial pharmaceutical input, with the relatively high recoveries for tylosin (78.2%), monensin
284 (76.4%), lincomycin (74.3%), carbamazepine (68.4%), trimethoprim (58.6%) and
285 oxytetracycline (52.3%). Among these six pharmaceuticals, the majority of trimethoprim
286 (36.7%), lincomycin (69.4%), monensin (54.5%), and tylosin (43.8%) remained in the nutrient
287 solution, whereas large fractions of oxytetracycline (27.1%) and carbamazepine (40.8%) were
288 accumulated in radish plants. The metabolism of carbamazepine in vegetables such as sweet
289 potato, carrot, tomato, and cucumber could produce 10,11-epoxycarbamazepine and 10,11-
290 dihydroxycarbamazepine (Goldstein et al., 2014; Malchi et al., 2014). Riemenschneider et al.
291 (2017) found 21 phase-I and II transformation products derived from carbamazepine in tomato
292 after 35 days of exposure, and accounted for ~45% of carbamazepine uptake into plant. The
293 recoveries were relatively low for other nine pharmaceuticals ranging from 3.0 to 32.1%, only
294 3.0-4.0% for 17 β -estradiol, sulfamethoxazole, sulfadiazine and estrone, 11.7-18.4% for triclosan,
295 acetaminophen and caffeine, 20.7 and 32.1% for carbadox and lamotrigine. Similar metabolism
296 potential of triclosan was also observed in carrot by Macherius et al. (2012), who identified 8
297 triclosan conjugation products and their total amounts was about 5 times more than the parent

308 compound triclosan. The total recovered mass was < 80% for all tested pharmaceuticals after 7
309 days of exposure, and the metabolism in radish is expected to be responsible for such
310 discrepancies between the initial input and the remaining amount. The potential risk originating
311 from the metabolites are still unclear, particularly for those quickly and intensively metabolized
312 pharmaceuticals such as estrogens and sulfonamide antibiotics.

303

304 3.4. Metabolism of pharmaceuticals in radish enzyme extracts

305 The metabolism potential of the fifteen pharmaceuticals in radish root and leaf enzyme
306 extracts was examined by measuring their dissipation during 96 h of exposure (Figure 5). In root
307 enzyme extracts the tested pharmaceuticals demonstrated a wide range of variation in mass
308 recoveries from 1.8 to 98.1% of the initial dosage (Figure 5). Lincomycin, monensin, tylosin,
309 carbadox, carbamazepine, lamotrigine, trimethoprim and caffeine showed relatively low to
310 minimal transformation with the mass recoveries ranging between 72.3 and 98.1%.
311 Carbamazepine and trimethoprim demonstrated high stability in an in vitro study, where their
312 concentration remained nearly unchanged in carrot cell cultures during the 90 h of exposure, and
313 < 5% of carbamazepine was transformed after 22 days of incubation (Wu et al., 2016). In another
314 lately in vitro study, about 5% of carbamazepine was transformed in horseradish hairy root cell
315 cultures after 6 days of exposure (Sauvêtre et al., 2018). Sulfamethoxazole, sulfadiazine,
316 acetaminophen, oxytetracycline, triclosan, estrone and 17 β -estradiol manifested a rapid
317 metabolism with the recovered amount of < 34.0% of the initially spiked dosage. The fast
318 metabolism of estrogens, acetaminophen, sulfamethoxazole and triclosan is consistent with
319 previous in vitro studies (Card et al., 2013; Huber et al., 2009; Macherius et al., 2012; Wu et al.,
320 2016). Three conjugates (acetaminophen–glucoside, acetaminophen–glutathione, and cysteine

321 conjugate) of acetaminophen in the cell cultures of *A. rusticana* were identified after 6 h of
322 exposure, and their total amounts was about 4.6 times that of acetaminophen (Huber et al., 2009).
323 Wu et al. (2016) found that about 55.5 and 91.7% of triclosan and sulfamethoxazole disappeared
324 in carrot cell cultures within 90 h. Macherius et al. (2012) reported that ~95% of triclosan was
325 quickly metabolized within 24 h in carrot cell cultures. In this study, the rapidly depleted
326 pharmaceuticals were found for sulfamethoxazole, sulfadiazine, acetaminophen, oxytetracycline,
327 triclosan, estrone and 17 β -estradiol in both intact radish and root enzyme extracts. However, the
328 highly metabolized carbadox and caffeine in intact radish showed less depletion in the enzyme
329 extracts.

330 The extent of metabolism of pharmaceuticals in leaf enzyme extracts was less than that in
331 root extracts with the exception of carbadox and lamotrigine (Figure 5). Among the
332 pharmaceuticals studied, triclosan demonstrated the highest transformation in leaf enzyme
333 extracts (50.8%); however, the remaining fraction was still ~5 times greater than that in root
334 enzyme extracts (9.5%). Sulfadiazine, carbadox and sulfamethoxazole were moderately
335 metabolized in leaf enzyme extracts with the remaining fractions of 59.4%, 61.4% and 72.1%,
336 respectively. For the remaining ten pharmaceuticals, the majority of the initially spiked
337 pharmaceuticals still existed in the leaf enzyme extracts (79.2-98.8%) (Figure 5). The different
338 metabolism rates of pharmaceuticals between root vs leaf enzyme extracts could be related to the
339 types and amount of extracted enzymes, and their metabolic activities in different radish tissues.
340 The changes in metabolism magnitude could alter the distribution and accumulation of
341 pharmaceuticals in different plant parts, especially in the scenario that pharmaceuticals enter
342 plants by root uptake. For instance, only 1.8% of 17 β -estradiol remained in root enzyme extracts
343 after 96-h exposure, while > 90% of 17 β -estradiol was still present in leaf enzyme extracts

344 (Figure 5). In the in vivo experiment, 17 β -estradiol concentration in the leaves was 16 ng g⁻¹
345 which was >10 times greater than that in roots (1.4 ng g⁻¹). The results from the in vitro
346 experiments clearly support the rapid metabolism of 17 β -estradiol in radish roots, which is
347 believed to lead to the corresponding lower concentration in roots in the in vivo experiments.
348 Pharmaceuticals in different plant parts could have varying metabolism rates such as
349 carbamazepine in sweet potato, carrot, and tomato (Malchi et al., 2014; Riemenschneider et al.,
350 2017) and iopromide in cattail (Cui et al., 2017).

351 The rapid metabolism of sulfamethoxazole, sulfadiazine, acetaminophen, oxytetracycline,
352 triclosan, estrone and 17 β -estradiol in radish root enzyme extracts was fit to the first-order
353 kinetic model, and the fittings are shown as curves in Figure 6 with R² = 0.94-0.97 (Table S10).
354 The corresponding half-life was estimated as 17 β -estradiol (15.4 h) < estrone (17.3 h) <
355 triclosan (30.1 h) < acetaminophen (33.0 h) < oxytetracycline (36.5 h) < sulfadiazine (49.5 h) <
356 sulfamethoxazole (63.0 h). The rapid dissipation of triclosan and sulfamethoxazole were also
357 found in the carrot cell cultures with half-life of ~0.2 h and ~7.7 h, respectively (Wu et al., 2016).
358 Compared with the results of carrot cell cultures, triclosan and sulfamethoxazole both exhibited a
359 much slower dissipation rate in the radish enzyme extracts. Triclosan demonstrated a faster
360 dissipation rate than sulfamethoxazole in both in vitro systems. The high removal of these
361 pharmaceuticals in root enzyme extracts is consistent with their low mass recoveries in the in
362 vivo study (Figure 4), suggesting that these biotransformation reactions in root enzyme extracts
363 could also take place in radish roots in the in vivo experiments. Application of plant tissue
364 enzyme extracts to react with pharmaceuticals could serve as an appropriate approach for rapid
365 evaluation of their metabolism potential in plants. This particularly simplifies the evaluation of

366 plant compartment-dependent metabolism, and could also eliminate the impact of uptake and
367 translocation to metabolism process in the intact plant experiments.

368

369 **4. Conclusion**

370 The deficient knowledge on metabolism of pharmaceuticals in plants limits the accurate
371 risk assessment of pharmaceuticals accumulated in vegetables because of the lack of appropriate
372 experimental or technique protocols for identification and quantification of metabolites. In this
373 study, the comparison in caffeine metabolism in radish between in vivo and in vitro exposure
374 experiments demonstrates that the demethylation metabolites identified in radish tissue enzyme
375 extracts (in vitro study) were similar to those formed in the intact radish roots and leaves in the in
376 vivo experiment. Similar metabolism patterns (dissipation of parent compounds) were observed
377 for rapidly metabolized pharmaceuticals including 17 β -estradiol, estrone, acetaminophen,
378 triclosan, oxytetracycline and sulfonamides in both radish tissue enzyme extracts and in the
379 intact plant. These results suggest that the metabolic reactions with plant tissue enzyme extracts
380 could be used as an appropriate approach for rapid examination of metabolism potential of
381 pharmaceuticals in a specific plant tissue, with the minimal impact from uptake and translocation.
382 For those extensively metabolized pharmaceuticals such as 17 β -estradiol, acetaminophen,
383 triclosan and sulfonamides, the measurement of the parent compounds in plants is not sufficient
384 to assess their accumulation and potential risks since the formed metabolites might still carry the
385 bioactive moieties and pose similar adverse impacts to plant and human health. Further
386 investigation of the metabolism of these pharmaceuticals in plants and their related
387 bioactivity/toxicity are needed to warrant the safe use of reclaimed water for irrigation and land
388 application of biosolids and animal manures.

389

390 **Acknowledgements**

391 This study was supported in part by Agriculture and Food Research Initiative Competitive Grant
392 2016-67017-24514 from USDA National Institute of Food and Agriculture, and Michigan
393 AgBioResearch Project GREEN.

394

395 **Appendix A. Supporting Information**

396 Additional details are provide in the Supporting Information including physicochemical
397 properties of pharmaceuticals, analytical methods, mass recovery of pharmaceuticals in the
398 hydroponic controls, protein concentration in enzyme extracts, mass distribution of caffeine in
399 hydroponic-radish system, dissipation curves of pharmaceuticals in radish enzyme extracts, and
400 representative chromatograms of pharmaceuticals.

401

402 **References**

403

- 404 Berthou, F., Flinois, J.-P., Ratanasavanh, D., Beaune, P., Riche, C., Guillouzo, A., 1991.
405 Evidence for the involvement of several cytochromes P-450 in the first steps of caffeine
406 metabolism by human liver microsomes. *Drug Metab. Disposition* 19, 561-567.
- 407 Berthou, F., Guillois, B., Riche, C., Dreano, Y., Jacqz-Aigrain, E., Beaune, P., 1992. Interspecies
408 variations in caffeine metabolism related to cytochrome P4501A enzymes. *Xenobiotica*
409 22, 671-680.
- 410 Bischel, H.N., Simon, G.L., Frisby, T.M., Luthy, R.G., 2011. Management experiences and
411 trends for water reuse implementation in Northern California. *Environ. Sci. Technol.* 46,
412 180-188.
- 413 Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities
414 of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248-254.
- 415 Card, M.L., Schnoor, J.L., Chin, Y.-P., 2013. Transformation of natural and synthetic estrogens
416 by maize seedlings. *Environ. Sci. Technol.* 47, 5101-5108.
- 417 Chen, C., Li, J., Chen, P., Ding, R., Zhang, P., Li, X., 2014. Occurrence of antibiotics and
418 antibiotic resistances in soils from wastewater irrigation areas in Beijing and Tianjin,
419 China. *Environ. Pollut.* 193, 94-101.
- 420 Christou, A., Karaolia, P., Hapeshi, E., Michael, C., Fatta-Kassinos, D., 2017. Long-term
421 wastewater irrigation of vegetables in real agricultural systems: Concentration of

422 pharmaceuticals in soil, uptake and bioaccumulation in tomato fruits and human health
423 risk assessment. *Water Res.* 109, 24-34.

424 Chuang, Y.-H., Zhang, Y., Zhang, W., Boyd, S.A., Li, H., 2015. Comparison of accelerated
425 solvent extraction and quick, easy, cheap, effective, rugged and safe method for
426 extraction and determination of pharmaceuticals in vegetables. *J. Chromatogr. A* 1404, 1-
427 9.

428 Clarke, B.O., Smith, S.R., 2011. Review of ‘emerging’ organic contaminants in biosolids and
429 assessment of international research priorities for the agricultural use of biosolids.
430 *Environ. Int.* 37, 226-247.

431 Coelho, A., Fraichard, S., Le Goff, G., Faure, P., Artur, Y., Ferveur, J.-F., Heydel, J.-M., 2015.
432 Cytochrome P450-dependent metabolism of caffeine in *Drosophila melanogaster*. *PloS*
433 *one* 10, e0117328.

434 Cogger, C.G., Bary, A.I., Kennedy, A.C., Fortuna, A.-M., 2013. Long-term crop and soil
435 response to biosolids applications in dryland wheat. *J. Environ. Qual.* 42, 1872-1880.

436 Cui, H., de Angelis, M.H., Schröder, P., 2017. Iopromide exposure in *Typha latifolia* L.:
437 Evaluation of uptake, translocation and different transformation mechanisms in planta.
438 *Water Res.* 122, 290-298

439 Cui, H., Schröder, P., 2016. Uptake, translocation and possible biodegradation of the antidiabetic
440 agent metformin by hydroponically grown *Typha latifolia*. *J. Hazard. Mater.* 308, 355-
441 361.

442 Durán-Alvarez, J.C., Becerril-Bravo, E., Castro, V.S., Jiménez, B., Gibson, R., 2009. The
443 analysis of a group of acidic pharmaceuticals, carbamazepine, and potential endocrine
444 disrupting compounds in wastewater irrigated soils by gas chromatography–mass
445 spectrometry. *Talanta* 78, 1159-1166.

446 Fonne-Pfister, R., Simon, A., Salaun, J.-P., Durst, F., 1988. Xenobiotic metabolism in higher
447 plants involvement of microsomal cytochrome P-450 in aminopyrine N-demethylation.
448 *Plant Sci.* 55, 9-20.

449 Fu, Q., Zhang, J., Borchardt, D., Schlenk, D., Gan, J., 2017. Direct Conjugation of Emerging
450 Contaminants in *Arabidopsis*: Indication for an Overlooked Risk in Plants? *Environ. Sci.*
451 *Technol.* 51, 6071-6081.

452 Goldstein, M., Shenker, M., Chefetz, B., 2014. Insights into the uptake processes of wastewater-
453 borne pharmaceuticals by vegetables. *Environ. Sci. Technol.* 48, 5593-5600.

454 Goth, R., Cleaver, J., 1976. Metabolism of caffeine to nucleic acid precursors in mammalian
455 cells. *Mutat Res.* 36, 105-113.

456 He, Y., Langenhoff, A.A., Sutton, N.B., Rijnaarts, H.H., Blokland, M.H., Chen, F., Huber, C.,
457 Schröder, P., 2017. Metabolism of ibuprofen by *Phragmites australis*: uptake and
458 phytodegradation. *Environ. Sci. Technol.* 51, 4576-4584.

459 Huang, H., Zhang, S., Wang, S., Lv, J., 2013. In vitro biotransformation of PBDEs by root crude
460 enzyme extracts: Potential role of nitrate reductase (NaR) and glutathione S-transferase
461 (GST) in their debromination. *Chemosphere* 90, 1885-1892.

462 Huber, C., Bartha, B., Harpaintner, R., Schröder, P., 2009. Metabolism of acetaminophen
463 (paracetamol) in plants—two independent pathways result in the formation of a
464 glutathione and a glucose conjugate. *Environ. Sci. Pollut. Res.* 16, 206-213.

465 Huber, C., Bartha, B., Schröder, P., 2012. Metabolism of diclofenac in plants—Hydroxylation is
466 followed by glucose conjugation. *J. Hazard. Mater.* 243, 250-256.

467 Kalow, W., Tang, B.K., 1993. The use of caffeine for enzyme assays: a critical appraisal. *Clin.*
468 *Pharmacol. Ther.* 53, 503-514.

469 Kinney, C.A., Furlong, E.T., Werner, S.L., Cahill, J.D., 2006. Presence and distribution of
470 wastewater-derived pharmaceuticals in soil irrigated with reclaimed water. *Environ.*
471 *Toxicol. Chem.* 25, 317-326.

472 Kot, M., Daniel, W.A., 2008. The relative contribution of human cytochrome P450 isoforms to
473 the four caffeine oxidation pathways: an in vitro comparative study with cDNA-
474 expressed P450s including CYP2C isoforms. *Biochem. Pharmacol.* 76, 543-551.

475 Kumar, K., Gupta, S., Baidoo, S., Chander, Y., Rosen, C., 2005. Antibiotic uptake by plants
476 from soil fertilized with animal manure. *J. Environ. Qual.* 34, 2082-2085.

477 LeFevre, G.H., Müller, C.E., Li, R.J., Luthy, R.G., Sattely, E.S., 2015. Rapid
478 phytotransformation of benzotriazole generates synthetic tryptophan and auxin analogs in
479 *Arabidopsis*. *Environ. Sci. Technol.* 49, 10959-10968.

480 Macherius, A., Eggen, T., Lorenz, W., Moeder, M., Ondruschka, J., Reemtsma, T., 2012.
481 Metabolization of the bacteriostatic agent triclosan in edible plants and its consequences
482 for plant uptake assessment. *Environ. Sci. Technol.* 46, 10797-10804.

483 Malchi, T., Maor, Y., Tadmor, G., Shenker, M., Chefetz, B., 2014. Irrigation of root vegetables
484 with treated wastewater: evaluating uptake of pharmaceuticals and the associated human
485 health risks. *Environ. Sci. Technol.* 48, 9325-9333.

486 Marsik, P., Sisa, M., Lacina, O., Motkova, K., Langhansova, L., Rezek, J., Vanek, T., 2017.
487 Metabolism of ibuprofen in higher plants: A model *Arabidopsis thaliana* cell suspension
488 culture system. *Environ. Pollut.* 220, 383-392.

489 Miller, E.L., Nason, S.L., Karthikeyan, K., Pedersen, J.A., 2016. Root Uptake of
490 Pharmaceuticals and Personal Care Product Ingredients. *Environ. Sci. Technol.* 50, 525-
491 541.

492 Prosser, R., Sibley, P., 2015. Human health risk assessment of pharmaceuticals and personal care
493 products in plant tissue due to biosolids and manure amendments, and wastewater
494 irrigation. *Environ. Int.* 75, 223-233.

495 Riemenschneider, C., Seiwert, B., Moeder, M., Schwarz, D., Reemtsma, T., 2017. Extensive
496 transformation of the pharmaceutical carbamazepine following uptake into intact tomato
497 plants. *Environ. Sci. Technol.* 51, 6100-6109.

498 Sandermann, H., 1992. Plant metabolism of xenobiotics. *Trends Biochem. Sci* 17, 82-84.

499 Sato, T., Qadir, M., Yamamoto, S., Endo, T., Zahoor, A., 2013. Global, regional, and country
500 level need for data on wastewater generation, treatment, and use. *Agric. Water Manage.*
501 130, 1-13.

502 Sauvêtre, A., May, R., Harpaintner, R., Poschenrieder, C., Schröder, P., 2018. Metabolism of
503 carbamazepine in plant roots and endophytic rhizobacteria isolated from *Phragmites*
504 *australis*. *J. Hazard. Mater.* 342, 85-95.

505 Siminszky, B., 2006. Plant cytochrome P450-mediated herbicide metabolism. *Phytochem. Rev.* 5,
506 445-458.

507 Tanoue, R., Sato, Y., Motoyama, M., Nakagawa, S., Shinohara, R., Nomiyama, K., 2012. Plant
508 uptake of pharmaceutical chemicals detected in recycled organic manure and reclaimed
509 wastewater. *J. Agric. Food. Chem.* 60, 10203-10211.

510 Thies, F., Backhaus, T., Bossmann, B., Grimme, L.H., 1996. Xenobiotic biotransformation in
511 unicellular green algae (involvement of cytochrome P450 in the activation and selectivity
512 of the pyridazinone pro-herbicide metflurazon). *Plant Physiol.* 112, 361-370.

513 Tomson, T., Almkvist, O., Nilsson, B.Y., Svensson, J.-O., Bertilsson, L., 1990. Carbamazepine-
514 10, 11-epoxide in epilepsy a pilot study. *Arch. Neurol.* 47, 888-892.

515 Van Eerd, L.L., Hoagland, R.E., Zablotowicz, R.M., Hall, J.C., 2003. Pesticide metabolism in
516 plants and microorganisms. *Weed Sci.* 51, 472-495.

517 Vazquez-Roig, P., Segarra, R., Blasco, C., Andreu, V., Picó, Y., 2010. Determination of
518 pharmaceuticals in soils and sediments by pressurized liquid extraction and liquid
519 chromatography tandem mass spectrometry. *J. Chromatogr. A* 1217, 2471-2483.

520 Wu, X., Conkle, J.L., Ernst, F., Gan, J., 2014. Treated wastewater irrigation: uptake of
521 pharmaceutical and personal care products by common vegetables under field conditions.
522 *Environ. Sci. Technol.* 48, 11286-11293.

523 Wu, X., Ernst, F., Conkle, J.L., Gan, J., 2013. Comparative uptake and translocation of
524 pharmaceutical and personal care products (PPCPs) by common vegetables. *Environ. Int.*
525 60, 15-22.

526 Wu, X., Fu, Q., Gan, J., 2016. Metabolism of pharmaceutical and personal care products by
527 carrot cell cultures. *Environ. Pollut.* 211, 141-147.

528 Xuan, R., Arisi, L., Wang, Q., Yates, S.R., Biswas, K.C., 2009. Hydrolysis and photolysis of
529 oxytetracycline in aqueous solution. *J. Environ. Sci. Heal B.* 45, 73-81.

530 Yu, M., Liu, J., Wang, T., Sun, J., Liu, R., Jiang, G., 2013. Metabolites of 2, 4, 4'-tribrominated
531 diphenyl ether (BDE-28) in pumpkin after in vivo and in vitro exposure. *Environ. Sci.*
532 *Technol.* 47, 13494-13501.

533 Zhang, H., Wen, B., Hu, X., Wu, Y., Pan, Y., Huang, H., Liu, L., Zhang, S., 2016. Uptake,
534 Translocation, and Metabolism of 8: 2 Fluorotelomer Alcohol in Soybean (*Glycine max*
535 *L. Merrill*). *Environ. Sci. Technol.* 50, 13309-13317.

536

537