



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/151515/>

Version: Accepted Version

Proceedings Paper:

Clifford, J., Wostenholme, D., Grindell, C. et al. (2017) Improving services for weight management after stroke : a co-production approach. In: International Journal of Stroke. UK Stroke Forum 2017, 28-30 Nov 2017, Liverpool, UK. SAGE Publications, p. 26. ISSN: 1747-4930. EISSN: 1747-4949.

<https://doi.org/10.1177/1747493017732216>

UK Stroke Forum Abstracts, 2017. (2017). International Journal of Stroke, 12(5_suppl), 7–59. © The Authors. <https://doi.org/10.1177/1747493017732216>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

**Improving services for weight management after stroke: a co-production
approach
UK Stroke Forum 2017**

J Clifford¹, D Wostenholme², C Grindell², A Tod³, T Ryan³ and C Homer⁴

¹*Sheffield Health and Social Care Trust, Sheffield, UK*

²*CLAHRC-Y&H, Sheffield Teaching Hospitals Foundation Trust, Sheffield, UK*

³*The School of Nursing and Midwifery, University of Sheffield, Sheffield, UK*

⁴*Centre for Health and Social Care Research, Sheffield Hallam University, Sheffield UK*

Introduction: Clinicians have a responsibility to promote healthy lifestyle after stroke. Recent research indicates that post-stroke weight-gain can be a problem for some people longer term. A CLAHRC Y&H funded project explored the implications of this for stroke survivors in Sheffield. Stroke survivors, carers, and service providers met for co-production. The aim was to generate ideas for service development.

Methods: 2 workshops based on the principles of co-production took place. Stroke survivors, their carers, and service providers jointly considered their experiences of post-stroke weight management services and their wishes for an ideal service.

Results: Ideas for service development included: staff training, individualised stroke-specific services, improved access to dietetic support, and weight-gain management resource development. Stroke pathway and other weight management services would benefit from being more 'joined up'. Staff could be trained in post-stroke lifestyle promotion. A single accessible information resource for stroke survivors and carers would help. A collaborative document to monitor and assist with weight management throughout the pathway was suggested.

Conclusion: The workshops generated ideas for service development that could improve patient experience of weight management after stroke. Further funding will be sought to develop, trial and evaluate these ideas on the stroke pathway.