



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/153874/>

Version: Published Version

Proceedings Paper:

Biggs, K., Hind, D., Bradburn, M. et al. (2019) Lessons learnt from a multi-centre type 3 surgical trial. In: Trials. 5th International Clinical Trials Methodology Conference (ICTMC 2019), 06-09 Oct 2019, Brighton, UK. BioMed Central, p. 71. EISSN: 1745-6215.

<https://doi.org/10.1186/s13063-019-3688-6>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

MEETING ABSTRACTS

Open Access



Meeting abstracts from the 5th International Clinical Trials Methodology Conference (ICTMC 2019)

Brighton, UK. 06-09 October 2019

Published: 22 October 2019

P-1

Abstract omitted

P-2

Some practical considerations in the design of multi-arm multi-stage designs

Jerome Wulff, Nikolaos Demiris

Cambridge Clinical Trial Unit, Cambridge, United Kingdom

Trials 2019, 20(Suppl 1):P-2

Introduction: In the design of cancer clinical trials, one is often concerned with a number of options in the event that several treatments are of interest.

Methods: We explore in this work the distinct possibilities when four treatments are available, one acting as control and three as potentially efficacious alternatives. This design may be embedded within the context of multi-arm multi-stage (MAMS) trials where one may select a two- or three-stage design.

Potential Results: We explore the application of such designs, including trade-offs between potential gains in the number of patients with additional stages contrasted with patients "lost" due to practical considerations such as patients randomised in dropped arms while waiting for interim analyses and inspection by an Independent Data and Safety Committee. In addition, in cancer studies one may focus on the primary end-point using a time-to-event analysis or a binary outcome by looking at the probability of (potentially progression-free) survival at a specific, clinically meaningful, time point. The effect of such choices is extensively investigated.

Potential Relevance & Impact: We conclude with a discussion of the available software for MAMS designs and their advantages and disadvantages in terms of accuracy.

P-3

The UK plasma based Molecular profiling of Advanced breast cancer to inform Therapeutic CHOices (plasmaMATCH) Trial: A multiple parallel-cohort, phase IIa platform trial aiming to provide proof of principle efficacy for designated targeted therapies in patient subgroups identified through ctDNA screening (CRUK/15/010)

Sarah Kernaghan¹, Laura Moretti¹, Lucy Kilburn¹, Katie Wilkinson¹, Claire

Snowdon¹, James Morden¹, Iain Macpherson², Andrew Wardley³, Rebecca Roylance⁴, Richard Baird⁵, Alistair Ring⁶, Nicholas Turner⁷, Judith M Bliss¹, on behalf of the plasmaMATCH Trial Management Group

¹Clinical Trials and Statistics Unit at The Institute of Cancer Research (ICR-CTS), United Kingdom; ²The Beatson West of Scotland Cancer Centre, Glasgow, United Kingdom; ³The Christie NHS Foundation Trust, Manchester, United Kingdom; ⁴University College London Hospitals NHS Foundation Trust, London, United Kingdom; ⁵Cambridge University Hospitals NHS Foundation Trust, Cambridge, United Kingdom; ⁶The Royal Marsden NHS Foundation Trust, Sutton, United Kingdom; ⁷The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust, London, United Kingdom

Trials 2019, 20(Suppl 1):P-3

Introduction: plasmaMATCH is a novel platform trial which assesses the potential of circulating tumour DNA (ctDNA) screening to direct targeted therapies in advanced breast cancer (ABC) patients. The trial recruited ahead of target and will report initial results within 3years of first patient first visit demonstrating efficiency of this design.

Methods: plasmaMATCH is an open-label, multi-centre phase IIa platform trial, consisting of a ctDNA screening component and five parallel treatment cohorts. Patients with an actionable mutation identified at ctDNA screening are invited to enter Cohorts A-D to receive a targeted treatment matched to the mutation identified (A: ESR1-extended-dose fulvestrant; B: HER2-neratinib+/-fulvestrant; C&D: AKT1 (or PTEN for Cohort D) -AZD5363+/-fulvestrant). Cohort E was added



digital tool, with examples, will inform the NIHR research community about choices and help them identify potential tools to support recruitment and retention.

P-245

Lessons learnt from a multi-centre Type 3 surgical trial

Katie Biggs¹, Daniel Hind¹, Mike Bradburn¹, Lizzie Swaby¹, Steven Brown²

¹University Of Sheffield, Sheffield, UK; ²Sheffield Teaching Hospitals NHS Trust, Sheffield, UK

Trials 2019, 20(Suppl 1):P-245

Background: Pragmatic randomised controlled trials are increasingly being used to evaluate surgical interventions, although they present particular difficulties in regard to recruitment and retention.

Methods

This paper details the procedures and processes related to implementation of a multi-centre pragmatic surgical randomised controlled trial.

Results: Forecasting consent rates based on previous similar trials ensured that the recruitment window was of adequate length. Adequate resource was available for study procedures at multiple clinics in each hospital due to micro-costing of study activities with research partners ensure. A video was produced targeting recruiting staff, which aimed to help recruiters explain the trial, randomisation and equipoise, based on methodological work and experiences from another study. Post-randomisation delays in delivering surgery to one study arm were investigated by assessing the outcomes at the time of randomisation and the day of surgery which provided confidence in the baseline measure. Real-time monitoring of participant drop-out due to delays in surgery meant we were able to extend the recruitment window in a timely fashion. Triangulation of data sources ensured adequate numbers of participants provided primary outcome data.

Discussion: This paper provides a range of evidence- and experience-based approaches which resulted in meeting our study's objectives and these lessons may be transferable.

P-246

Challenges in the design, planning and implementation of trials evaluating group interventions

Katie Biggs¹, Daniel Hind¹, Rebecca Gossage-Worrall¹, Kirsty Sprange², David White¹, Jessica Wright¹, Robin Chatters¹, Katherine Berry³, Diana Papaioannou¹, Mike Bradburn¹, Stephen J Walters¹, Cindy L Cooper¹

¹SCHARR, University of Sheffield, Sheffield, UK; ²NCTU, University of Nottingham, Nottingham, UK; ³School of health Sciences, University of Manchester, Manchester, UK

Trials 2019, 20(Suppl 1):P-246

Background: Evaluating group interventions in randomised controlled trials (RCTs) presents a set of practical problems which may not be immediately obvious and are not present in RCTs of one-to-one interventions.

Methods

Case-based approach summarising Sheffield clinical trial unit's experience in the design and implementation of group interventions across five randomised controlled trials.

Results: Median recruitment across the five trials was 5.8 (range 2.1-16.0) participants per site per month. Group intervention trials can involve a delay in the start of treatment whilst waiting for sufficient numbers to start a group. There was no evidence in our trials that the timing of consent, relative to randomisation, affected post-randomisation attrition, but attrition was a concern for all trial teams. Group facilitator attrition was common in studies where facilitators were employed by the health-system rather than the by the grant holder, and lead to the early closure of one trial. Solutions to this included training 'back-up' and new facilitators. Trials specified that participants had to attend a median of 62.5% (range 16.7% to 80%) of sessions, in order to receive a 'therapeutic dose'; a median of

75.0% (range 34.6% to 97.8%) received a therapeutic dose. Across the five trials, 66.4% of all sessions ran with fewer than the numbers pre-specified as ideal. A variety of methods were used to assess the fidelity of group interventions across the five trials.

Discussion: Investigators should expect delays / difficulties in recruiting groups of the optimal size, plan for both facilitator and participant attrition and consider how group attendance and group size affects treatment fidelity.

P-247

Delivering site set-up training to groups of sites versus individually: a randomised study within a trial

Eleanor Mitchell¹, Alan Montgomery¹, Garry Meakin¹, Rachel Haines¹,

Reuben Ogollah¹, Chris Partlett¹, Kate Walker¹, Jon Dorling², Shalini

Ojha³, on behalf of the FEED1 Collaborative Group

¹Nottingham Clinical Trials Unit, University of Nottingham, Nottingham, United Kingdom; ²Dalhousie University, Halifax, Canada; ³Division of Graduate Entry Medicine & Medical Sciences, University of Nottingham, Nottingham, United Kingdom

Trials 2019, 20(Suppl 1):P-247

Introduction: Site initiation visits (SIVs) are conducted to deliver training to sites before opening them to recruitment, though this can be burdensome during the time-intensive trial set-up period. There is little evidence about the best way to deliver training for sites to perform well. Evaluating methods of training was the highest priority identified at a workshop exploring recruitment and retention of participants to trials. Two systematic reviews investigating training in clinical trials showed a variety of different training methods and more research is needed to determine what kind of training and support can improve recruitment. A small retrospective study showed that, whilst face-to-face training (either at SIV or group training session) was associated with better recruitment than remote training (i.e. telephone or DVD), no difference was seen between the two types of face-to-face training.

Our objective is to compare group-based training during the trial set-up period versus visiting the site to conduct a SIV to investigate the impact of the training method upon key site performance metrics. We will embed a SWAT into the FEED1 trial, funded by the NIHR HTA programme.

Methods: Once selected, sites will be randomised in batches to receiving their site-initiation training during a SIV or group-based training by attending a collaborators' meeting. To allow for non-availability of site staff, two meetings will be held. Outcomes will include recruitment and retention, data quality and protocol compliance (defined as core site performance metrics) and associated costs of each training method.

Timing of potential results: Clinical trial and SWAT results will be available in Q1 of 2023.

Potential relevance and impact: If the intervention is shown to be effective, there could be significant benefits to funders and trial teams, in particular reducing the length of time it takes to set-up a trial and open all sites.

P-248

Making a challenging trial possible: Lessons from the Emergency Department led ECLIPSE trial

Kerry Woolfall¹, Louise Roper¹, Mark D Lyttle⁴, Carrol Gamble⁵, Amy Humphreys⁵, Shrouk Messahel³, Elizabeth Lee³, Joanne Noblet³, Helen Hickey⁵, Naomi E.A Rainford⁵, Anand Iyer², Richard Appleton²

¹University of Liverpool, United Kingdom; ²Neurology Department Alder Hey Children's NHS Foundation Trust, United Kingdom; ³Emergency Department, Alder Hey Children's NHS Foundation Trust, United Kingdom; ⁴Emergency Department, Bristol Royal Hospital for Children, United Kingdom; ⁵University of Liverpool, Clinical Trials Research Centre (CTRC), United Kingdom

Trials 2019, 20(Suppl 1):P-248