

antibiotic resistance, keeping use of key agents below the threshold that may trigger resistance problems.

Transparency declarations

None to declare.

References

- 1 Limmathurotsakul D, Sandoe JAT, Barrett DC et al. 'Antibiotic footprint' as a communication tool to aid reduction of antibiotic consumption. *J Antimicrob Chemother* 2019; **74**: 2122–7.
- 2 López-Lozano JM, Lawes T, Nebot C et al. A nonlinear time-series analysis approach to identify thresholds in associations between population antibiotic use and rates of resistance. *Nat Microbiol* 2019; **4**: 1160–72.
- 3 Lawes T, Lopez-Lozano JM, Nebot CA et al. Effect of a national 4C antibiotic stewardship intervention on the clinical and molecular epidemiology of *Clostridium difficile* infections in a region of Scotland: a non-linear time-series analysis. *Lancet Infect Dis* 2017; **17**: 194–206.
- 4 Lawes T, Lopez-Lozano JM, Nebot CA et al. Effects of national antibiotic stewardship and infection control strategies on hospital-associated and community-associated methicillin-resistant *Staphylococcus aureus* infections across a region of Scotland: a non-linear time-series study. *Lancet Infect Dis* 2015; **15**: 1438–49.
- 5 Lawes T, López-Lozano JM, Nebot C et al. Turning the tide or riding the waves? Impacts of antibiotic stewardship and infection control on MRSA strain dynamics in a Scottish region over 16 years: non-linear time series analysis. *BMJ Open* 2015; **5**: e006596.
- 6 Levy SB. Balancing the drug–resistance equation. *Trends Microbiol* 1994; **2**: 341–2.

J Antimicrob Chemother 2020; **75**: 785–786
doi:10.1093/jac/dkz533
Advance Access publication 22 December 2019

'Antibiotic footprint' as a communication tool to aid reduction of antibiotic consumption—authors' response

Direk Limmathurotsakul ^{1–3**},
Jonathan A. T. Sandoe^{4,5}, David C. Barrett⁶,
Michael Corley⁵, Li Yang Hsu^{7,8}, Marc Mendelson^{9,10},
Peter Collignon^{11,12}, Ramanan Laxminarayan^{13,14},
Sharon J. Peacock ¹⁵ and Philip Howard^{4,5}

¹Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, 10400, Thailand; ²Department of Tropical Hygiene, Faculty of Tropical Medicine, Mahidol University, Bangkok, 10400, Thailand; ³Centre for Tropical Medicine and Global Health, University of Oxford, Oxford OX3 7FZ, UK; ⁴University of Leeds/Leeds Teaching Hospitals NHS Trust, Leeds LS1 3EX, UK; ⁵British Society of Antimicrobial Chemotherapy, Birmingham B1 3NJ, UK; ⁶Bristol Veterinary School, University of Bristol, Bristol BS40 5DU, UK; ⁷Saw Swee Hock School of Public Health, National University of Singapore and National University Health System, 12 Science Drive 2, Singapore 117649, Singapore; ⁸National Centre for Infectious Diseases, Moulmein Road, Singapore 308433, Singapore; ⁹Division of Infectious Diseases & HIV Medicine, Department of Medicine, University of Cape Town, Cape Town, 7925, South Africa; ¹⁰International Society for Infectious Diseases, Brookline, MA 02446, USA; ¹¹Infectious Diseases and Microbiology, Canberra Hospital, Canberra, 2605, Australia; ¹²Medical School, Australian National University, Acton, 2606, Australia; ¹³Center for Disease Dynamics, Economics & Policy, New Delhi, 110024, India; ¹⁴Princeton Environmental Institute, Princeton, NJ 08544, USA; ¹⁵Department of Medicine, University of Cambridge, Cambridge CB2 0QQ, UK

**Corresponding author. E-mail: direk@tropmedres.ac

Sir,

We thank Gould et al.¹ for their comments on our article on the antibiotic footprint, which we proposed as a simple communication to aid public understanding of antibiotic consumption.² We fully agree that the antibiotic footprint could also be used as a tool to support interventions to reduce the overuse of antibiotics.¹ We understand the possibility of presenting the antibiotic footprint together with thresholds of antibiotic consumption in different

hospitals and communities, defined as number of treatment courses, or DDDs per unit of hospital activity or community population, as presented by Gould *et al.*,¹ additionally recognizing that a similar approach could be taken on livestock farms and in other veterinary and one-health contexts. However, complete data on antibiotic usage as proposed¹ are rarely available in low- and middle-income countries (LMICs) and further studies evaluating whether thresholds of antibiotic consumption could be defined for each LMIC are needed. The information on complete antibiotic use and new evidence arising from such studies would certainly allow the communication strategy to be adjusted and improved over time.² Furthermore, we strongly agree with the recent report, 'Reframing Resistance', published by the Wellcome Trust,³ which proposed that communication messages to the public need to be tested and that including the issue of antibiotic overuse in both humans and animals in the right way helps make the issue feel tractable. We foresee that the concept of the antibiotic footprint could be expanded to online individual calculators² and that local thresholds of antibiotic consumption could help in guiding local appropriate antibiotic use in the community.¹

However, further studies to evaluate the use of the antibiotic footprint and antibiotic consumption thresholds to communicate with communities are needed, particularly in different languages and contexts.

Transparency declarations

None to declare.

References

- 1 Gould IM, Nebot C, Aldeyab M *et al.* Comment on: 'Antibiotic footprint' as a communication tool to aid reduction of antibiotic consumption. *J Antimicrob Chemother* 2020; **75**: 784–85.
- 2 Limmathurotsakul D, Sandoe JAT, Barrett DC *et al.* 'Antibiotic footprint' as a communication tool to aid reduction of antibiotic consumption. *J Antimicrob Chemother* 2019; **74**: 2122–7.
- 3 Wellcome Trust. Reframing Resistance. <https://wellcome.ac.uk/sites/default/files/reframing-resistance-report.pdf>.