

This is a repository copy of *Supergravity on a 3-torus : quantum linearization instabilities with a supergroup*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/160490/>

Version: Published Version

Article:

Higuchi, Atsushi orcid.org/0000-0002-3703-7021 and Schmieding, Lasse (2020)
Supergravity on a 3-torus : quantum linearization instabilities with a supergroup. *Classical and Quantum Gravity*. pp. 1-36. ISSN 1361-6382

<https://doi.org/10.1088/1361-6382/ab90a4>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

PAPER • OPEN ACCESS

Supergravity on a three-torus: quantum linearization instabilities with a supergroup

To cite this article: Atsushi Higuchi and Lasse Schmieding 2020 *Class. Quantum Grav.* **37** 165009

View the [article online](#) for updates and enhancements.



AMERICAN
ASTRONOMICAL
SOCIETY

IOP | ebooks™

Your first choice for astronomy, astrophysics,
solar physics, and planetary science ebooks.

Start exploring the collection—download the
first chapter of every title for free.

Supergravity on a three-torus: quantum linearization instabilities with a supergroup

Atsushi Higuchi¹ and Lasse Schmieding¹

Department of Mathematics, University of York, Heslington, York, YO10 5DD, United Kingdom

E-mail: lcs549@york.ac.uk

Received 8 January 2020, revised 1 May 2020

Accepted for publication 6 May 2020

Published 28 July 2020



CrossMark

Abstract

It is well known that linearized gravity in spacetimes with compact Cauchy surfaces and continuous symmetries suffers from linearization instabilities: solutions to classical linearized gravity in such a spacetime must satisfy so-called linearization stability conditions (or constraints) for them to extend to solutions in the full non-linear theory. Moncrief investigated implications of these conditions in linearized quantum gravity in such background spacetimes and found that the quantum linearization stability constraints lead to the requirement that all physical states must be invariant under the symmetries generated by these constraints. He studied these constraints for linearized quantum gravity in flat spacetime with the spatial sections of toroidal topology in detail. Subsequently, his result was reproduced by the method of group-averaging. In this paper the quantum linearization stability conditions are studied for $\mathcal{N} = 1$ simple supergravity in this spacetime. In addition to the linearization stability conditions corresponding to the spacetime symmetries, i.e. spacetime translations, there are also fermionic linearization stability conditions corresponding to the background supersymmetry. We construct all states satisfying these quantum linearization stability conditions, including the fermionic ones, and show that they are obtained by group-averaging over the supergroup of the global supersymmetry of this theory.

Keywords: linearization stability conditions, group averaging, supergravity, quantum gravity

¹ Author to whom any correspondence should be addressed.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

1. Introduction

In physics, the equations of interest are frequently non-linear and difficult to solve. Typically only a small number of solutions are known and these exploit special symmetries. To extract further physics from the known solutions, a common strategy is to perturbatively expand small deviations around the known background solutions. At lowest order in perturbation one obtains linear equations for the perturbations, which are typically easier to analyse. However, it is not guaranteed that all solutions to the linearized equations actually arise as first approximations to solutions of the non-linear equations of the system. Let us illustrate this point in a simple example [1]: for $(x, y) \in \mathbb{R}^2$, consider the algebraic system $x(x^2 + y^2) = 0$, of which the exact solutions are $(0, y)$, where y is any real number. On the other hand, if we consider the linearized equations for perturbations δx and δy around a background (x_0, y_0) , these satisfy $\delta x(x_0^2 + y_0^2) + x_0(2x_0\delta x + 2y_0\delta y) = 0$. If we let the background be $(x_0, y_0) = (0, 0)$, then any pairs $(\delta x, \delta y)$ satisfy the linearized equation of motion, but those with $\delta x \neq 0$ cannot have arisen as linearizations of solutions to the non-linear equation. One characterizes this phenomenon as the equation $x(x^2 + y^2) = 0$ being linearization unstable at $(0, 0)$.

A well-known field-theoretic system with linearization instabilities is electrodynamics in a ‘closed Universe’, i.e. in a spacetime with compact Cauchy surfaces [2, 3]. Consider, for example, the electromagnetic field coupled to a charged matter field in such a spacetime. Note that the conserved total charge of the matter field must vanish in this system. This fact is a simple consequence of Gauss’s law $\vec{\nabla} \cdot \vec{E} \propto \rho$, where \vec{E} and ρ are the electric field and charge density, respectively. The integral over a Cauchy surface of the charge density ρ gives the total charge but the integral of $\vec{\nabla} \cdot \vec{E}$ vanishes because the Cauchy surface is compact. At the level of the linearized theory about vanishing background electromagnetic and matter field the theory is non-interacting. Since the matter field is non-interacting, there is no constraint on the total charge in the linearized theory, but a solution to the linearized (i.e. free) matter field equation does not extend to an exact solution to the full interacting theory unless its total charge Q_e vanishes. The linearization stability condition (LSC) in this case is $Q_e = 0$. If a solution to the linearized equations satisfies this condition, then it extends to an exact solution to the full theory. It is useful to note here that the charge Q_e generates the global gauge symmetry of the free charged matter field.

In gravitational systems, it is known that such linearization instabilities occur for any perturbations around a background which has both Killing symmetries and compact Cauchy surfaces [2, 4–12]. The LSCs which need to be imposed on such a closed Universe are that the generators Q of the background Killing symmetries must vanish when evaluated on any linearized solution. For example, if the spacetime possesses time and space translation symmetries, then the corresponding conserved charges are the energy and momentum, respectively.

In the late seventies Moncrief studied the rôle of these conditions in linearized *quantum* gravity in spacetimes with compact Cauchy surfaces. He proposed that they should be imposed as physical-state conditions as in the Dirac quantization [11, 12], i.e.

$$Q |\text{phys}\rangle = 0. \tag{1.1}$$

Since the conserved charges Q generate the spacetime symmetries (in the component of the identity), he concluded that quantum linearization stability conditions (QLSCs) imply that all physical states must be invariant under the spacetime Killing symmetries. He argued that these constraints can be viewed as a remnant of the diffeomorphism invariance of the non-linear theory.

Imposing the invariance of the physical Hilbert space under the full background symmetries (in the component of the identity) as required by the QLSCs would appear too restrictive. This problem is exemplified by de Sitter space, a spacetime which is physically relevant for inflationary cosmology. This spacetime has Cauchy surfaces with the topology of the three-dimensional sphere, which is compact. Therefore, the QLSCs imply that all physical states of linearized quantum gravity on a de Sitter background ought to be invariant under the full $SO_0(4, 1)$ symmetry group, i.e. the component of the identity of $SO(4, 1)$, of the spacetime. However, this would appear to exclude all states except the vacuum state, which would make the Hilbert space for the theory quite empty [3, 12].

However, there are non-trivial $SO_0(4, 1)$ invariant states that have infinite norm and, hence, are not in the Hilbert space. Moncrief suggested that a Hilbert space consisting of these invariant states could be constructed by dividing the infinite inner product by the infinite volume of the group $SO_0(4, 1)$. This suggestion was taken up in reference [13]. In that work a new inner product for $SO_0(4, 1)$ -invariant states was defined by what would later be termed group-averaging, which is an important ingredient in the refined algebraic quantization [14] and has been well studied in the context of loop quantum gravity. (The group-averaging procedure was also proposed in [15].) In this approach, one defines the invariant states $|\Psi\rangle$ by starting with a non-invariant state $|\psi\rangle$ and averaging against the symmetry group G , assumed here to be an unimodular group such as $SO_0(4, 1)$, to obtain

$$|\Psi\rangle = \int_G dg U(g) |\psi\rangle, \quad (1.2)$$

where U is the unitary operator implementing the symmetry on the states. The state $|\Psi\rangle$ can readily be shown to be invariant, i.e. $\langle\phi|U(g)|\Psi\rangle = \langle\phi|\Psi\rangle$ for any state $|\phi\rangle$ in the Hilbert space by the invariance of the measure dg . If the volume of the symmetry group is finite, the inner product of the invariant states $|\Psi_1\rangle$ and $|\Psi_2\rangle$ obtained from $|\psi_1\rangle$ and $|\psi_2\rangle$ as in (1.2) is

$$\langle\Psi_1|\Psi_2\rangle = V_G \int_G dg \langle\psi_1|U(g)|\psi_2\rangle, \quad (1.3)$$

where V_G is the volume of the group G . If G has infinite volume, e.g. if it is $SO_0(4, 1)$, then one needs to redefine the inner product on the invariant states by removing a factor of the group volume to make them normalizable. Thus, one defines the inner product on the new Hilbert space by

$$\langle\Psi_1|\Psi_2\rangle_{ga} = \int_G dg \langle\psi_1|U(g)|\psi_2\rangle. \quad (1.4)$$

By this group-averaging procedure one obtains an infinite-dimensional Hilbert space of $SO_0(4, 1)$ invariant states for linearized gravity in de Sitter space [3, 13]. The group-averaging procedure was carried out for this spacetime also for other free fields [16, 17] to obtain Hilbert spaces of invariant states. The QLSCs in de Sitter space were also studied in the context of cosmological perturbation [18, 19]. The group-averaging procedure has also been studied extensively in the context of constrained dynamical systems (see e.g. [20–23]). This method was extended to non-unimodular groups in [24].

The group-averaging procedure can also be explicitly carried out for perturbative quantum gravity in static space with topology of $\mathbb{R} \times T^3$, where the spatial Cauchy surfaces are copies of T^3 , the three-dimensional torus, to find the states satisfying the QLSCs. The classical [2] and quantum theory [11, 25] of this model have been studied and the group-averaging procedure can be carried out to obtain a physical Hilbert space of states invariant under the

$\mathbb{R} \times U(1)^3$ symmetry group of the background. Now, if one considers four-dimensional $\mathcal{N} = 1$ simple supergravity [26, 27] on this background spacetime, it is not difficult to see that there are additional fermionic LSCs. The purpose of this paper is to find all states satisfying the bosonic and fermionic QLSCs and show that a Hilbert space of these states can be constructed using the group-averaging procedure over the supergroup of symmetries of the linearized theory.

Let us describe how the LSCs arise for supergravity in this spacetime. Recall that the energy and momentum of a system in general relativity can be expressed as an integral over a two-dimensional surface at infinity of a Cauchy surface (the ADM mass and momentum) in asymptotically-flat spacetime. In classical perturbation theory about Minkowski space, this fact implies that the total energy and momentum of the linearized fields can be expressed as a surface integral at infinity of perturbations of the next order. Then, one expects that in perturbation theory in the flat $\mathbb{R} \times \mathbb{T}^3$ background, the total energy and momentum of the linearized fields vanish because there is no spatial infinity. This is indeed the case, and the vanishing of the total energy and momentum of the linearized field is expressed as the LSCs. Note here that the expression for the total energy, or the Hamiltonian, in this spacetime is not positive definite and, therefore, can vanish for non-trivial field configurations. Now, in supergravity there is a spinor supercharge Q_α , $\alpha = 1, 2, 3, 4$, associated with a global supersymmetry variation, and it is known that this supercharge can be written as an integral over two-dimensional surface at infinity of a Cauchy surface in asymptotically-flat spacetime [28]. This fact again implies that in static three-torus space there are quadratic constraints on the linearized theory corresponding to the vanishing of the supercharge, which are the fermionic LSCs. In this paper we study these fermionic LSCs together with the bosonic ones in linearized quantum supergravity.

The remainder of this paper is organized as follows. In section 2 we present a derivation of the (classical) bosonic and fermionic LSCs for $\mathcal{N} = 1$ simple supergravity in the background of flat $\mathbb{R} \times \mathbb{T}^3$ spacetime. We show that these conditions are of the form that the conserved Noether charges of the linearized theory vanish. In section 3 we discuss linearized supergravity in this spacetime and express the LSCs in terms of the classical analogues of annihilation and creation operators. In section 4 we impose the bosonic QLSCs on the states and recall how this can be understood in the context of group-averaging over the bosonic symmetry group. In section 5 we describe how all physical states satisfying both the bosonic and fermionic QLSCs are found and how a Hilbert space of physical states is constructed. Then we show that the procedure of finding the physical Hilbert space can be interpreted as group-averaging over the supergroup of global supersymmetry. We summarize and discuss our results in section 6. In appendix A we present a gauge transformation of the vierbein field that is proportional to the Lie derivative of a vector field, though it is not necessary for this work. In appendix B we illustrate our derivation of the LSCs in the simple example of electrodynamics in flat $\mathbb{R} \times \mathbb{T}^3$ spacetime. In appendix C we present the proof of some identities used in this paper. In appendix D we discuss some aspects of the zero-momentum sector of the gravitino field. Appendix E presents an example of a two-particle state satisfying all QLSCs. We follow the conventions of reference [29] throughout this paper.

2. The linearization stability conditions

The action for the four-dimensional $\mathcal{N} = 1$ simple supergravity [29, 30] with $8\pi G = 1$ is

$$S = \frac{1}{2} \int d^4x e [R - \bar{\Psi}_\mu \gamma^{\mu\nu\rho} D_\nu \Psi_\rho + X_{(4\Psi)}], \quad (2.1)$$

where $X_{(4\Psi)}$ consists of terms quartic in Ψ_μ (see e.g. [30] for the explicit form of $X_{(4\Psi)}$). Here, e_μ^a are the vierbein fields, $e := \det(e_\mu^a)$, and $\Psi_{\mu\alpha}$, $\alpha = 1, 2, 3, 4$, is the gravitino field, which is a Majorana spinor. The 4×4 gamma matrices γ^a , $a = 0, 1, 2, 3$, satisfy the Clifford relation $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$, where $\eta^{ab} = \text{diag}(-1, 1, 1, 1)$. The indices a, b, c, \dots are raised and lowered by the flat metric η_{ab} whereas the spacetime indices μ, ν, σ, \dots are raised and lowered by the spacetime metric $g_{\mu\nu} = e_\mu^a e_{\nu a}$. The matrices γ^a have the properties $\gamma^{0\dagger} = -\gamma^0$ and $\gamma^{i\dagger} = \gamma^i$, $i = 1, 2, 3$. One defines $\gamma^\mu := e_\mu^a \gamma^a$ and $\gamma^{\mu\nu\rho} := \gamma^{[\mu} \gamma^\nu \gamma^{\rho]}$, where $[\dots]$ indicates total anti-symmetrization. The Riemann tensor is given by

$$R_{\mu\nu}{}^{ab} := \partial_\mu \omega_\nu{}^{ab} - \partial_\nu \omega_\mu{}^{ab} + \omega_\mu{}^{ac} \omega_{\nu c}{}^b - \omega_\nu{}^{ac} \omega_{\mu c}{}^b, \quad (2.2)$$

and the curvature scalar is $R := e_\mu^\alpha e_\nu^\beta R_{\mu\nu}{}^{\alpha\beta}$. The spin connection $\omega_\mu{}^{ab} = \omega_\mu^{[ab]}$ is expressed in terms of e_μ^a as

$$\omega_\mu{}^{ab} = e^{a\nu} \partial_{[\mu} e_{\nu]}{}^b - e^{b\nu} \partial_{[\mu} e_{\nu]}{}^a - \frac{1}{2} (e^{a\nu} e^{b\sigma} - e^{b\nu} e^{a\sigma}) e_\mu^c \partial_\nu e_{c\sigma}. \quad (2.3)$$

One also defines $\bar{\Psi} = \Psi^T C$, where C is a unitary matrix satisfying $C^T = -C$ and $\gamma^{\mu T} = -C \gamma^\mu C^{-1}$. The Majorana condition satisfied by Ψ_μ reads $(\Psi_\mu^\dagger)_\alpha = i(C \gamma^0 \Psi_\mu)_\alpha$ [30]. We later choose a Majorana representation for γ^a in which $C = i\gamma^0$. In this representation we have $(\Psi_\mu^\dagger)_\alpha = (\Psi_\mu)_\alpha$. The gravitino covariant derivative in (2.1) is given by

$$D_\mu \Psi_\nu = \partial_\mu \Psi_\nu + \frac{1}{4} \omega_{\mu ab} \gamma^{ab} \Psi_\nu. \quad (2.4)$$

The action (2.1) is invariant under a local supersymmetry transformation of the form

$$\Delta_\epsilon e_\mu^a = \frac{1}{2} \bar{\epsilon} \gamma^a \Psi_\mu, \quad (2.5)$$

$$\Delta_\epsilon \Psi_\mu = D_\mu \epsilon + Y_{\mu ab} \gamma^{ab} \epsilon, \quad (2.6)$$

where $Y_{\mu ab}$ is quadratic in Ψ_μ and where ϵ is a spacetime dependent Grassmann variable with four components. (See, e.g. [30] for the explicit form of $Y_{\mu ab}$).

We consider this theory on a purely bosonic static background whose spatial sections are flat three-dimensional tori. For the background geometry we therefore take

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu = -dt^2 + dx^2 + dy^2 + dz^2, \quad (2.7)$$

where the spatial coordinates x , y and z are periodic with periods L_1, L_2 and L_3 respectively, and we let $V = L_1 L_2 L_3$ denote the spatial volume.

Let us first discuss the bosonic LSCs for this theory, which are well known as we described in section 1. The diffeomorphism transformation in the direction of a vector ζ^μ of the vierbein field e_μ^a and gravitino field Ψ_μ is²

$$\delta_\zeta e_\mu^a = \zeta^\nu \partial_\nu e_\mu^a + e_\nu^a \partial_\mu \zeta^\nu, \quad (2.8)$$

$$\delta_\zeta \Psi_\mu = \zeta^\nu \partial_\nu \Psi_\mu + \Psi_\nu \partial_\mu \zeta^\nu. \quad (2.9)$$

²The transformation obtained as the commutator of two supersymmetry transformations is different from the diffeomorphism transformation presented here (see, e.g. [29]). The former takes the form $\delta'_\zeta e_\mu^a = \delta_\zeta e_\mu^a - \zeta^\rho \omega_\rho{}^{ab} e_{b\mu}$, $\delta'_\zeta \Psi_\mu = \delta_\zeta \Psi_\mu - \partial_\mu (\zeta^\rho \Psi_\rho)$ to first order in the fields $\bar{\epsilon}_\mu^a$ defined by (2.10) and Ψ_μ . It can be shown that the LSCs corresponding to these two transformations are identical.

Now, we write the vierbein field as

$$e_\mu^a = \delta_\mu^a + \tilde{e}_\mu^a. \quad (2.10)$$

That is, we write e_μ^a as the sum of its background value δ_μ^a and perturbation \tilde{e}_μ^a . Then,

$$\delta_\zeta \tilde{e}_\mu^a = \delta_\nu^a \partial_\mu \zeta^\nu + \zeta^\nu \partial_\nu \tilde{e}_\mu^a + \tilde{e}_\nu^a \partial_\mu \zeta^\nu. \quad (2.11)$$

It is important to note here that the part of $\delta_\zeta \tilde{e}_\mu^a$ that is independent of \tilde{e}_μ^a vanishes if ζ^μ is a constant vector, i.e. a Killing vector of the flat background³.

Since the action (2.1) is invariant under the diffeomorphism transformation given by (2.9) and (2.11), we have

$$\delta_\zeta \tilde{e}_\mu^a \frac{\delta S}{\delta \tilde{e}_\mu^a} + \delta_\zeta \Psi_\mu \frac{\delta S}{\delta \Psi_\mu} = \partial_\mu (\sqrt{-g} J_{(\zeta)}^\mu), \quad (2.12)$$

for some vector field $J_{(\zeta)}^\mu$, which is linear in ζ^μ . Here the variation $\delta S / \delta \Psi_\mu$ is a left-variation, i.e.

$$\delta S = \int d^4x \delta \Psi_\mu \frac{\delta S}{\delta \Psi_\mu}. \quad (2.13)$$

We adopt left-variations for fermionic fields throughout this paper. We then expand $\delta_\zeta \tilde{e}_\mu^a$, $\delta_\zeta \Psi_\mu$, $\delta S / \delta \tilde{e}_\mu^a$, $\delta S / \delta \Psi_\mu$ and $\sqrt{-g} J_{(\zeta)}^\mu$ according to the order in the fields \tilde{e}_μ^a and Ψ_μ , i.e. the number of these fields in the product, as

$$\delta_\zeta \tilde{e}_\mu^a = \delta_\zeta^{(0)} \tilde{e}_\mu^a + \delta_\zeta^{(1)} \tilde{e}_\mu^a, \quad (2.14)$$

$$\delta_\zeta \Psi_\mu = \delta_\zeta^{(1)} \Psi_\mu, \quad (2.15)$$

$$\frac{\delta S}{\delta \tilde{e}_\mu^a} = E_a^{(1)\mu} + E_a^{(2)\mu} + \dots, \quad (2.16)$$

$$\frac{\delta S}{\delta \Psi_\mu} = \mathcal{E}^{(1)\mu} + \mathcal{E}^{(2)\mu} + \dots, \quad (2.17)$$

$$\sqrt{-g} J_{(\zeta)}^\mu = J_{(\zeta)}^{(0)\mu} + J_{(\zeta)}^{(1)\mu} + J_{(\zeta)}^{(2)\mu} + \dots, \quad (2.18)$$

Recall that the background metric is flat. We note that $E_a^{(0)\mu} = 0$ and $\mathcal{E}^{(0)\mu} = 0$ because the flat spacetime with $\Psi_\mu = 0$ satisfies the field equations. That is, $\delta S / \delta \tilde{e}_\mu^a = 0$ and $\delta S / \delta \Psi_\mu = 0$ if $\tilde{e}_\mu^a = 0$ and $\Psi_\mu = 0$.

The identity (2.12) must be satisfied order by order. The first- and second-order equalities read

$$(\delta_\zeta^{(0)} \tilde{e}_\mu^a) E_a^{(1)\mu} = \partial_\mu J_{(\zeta)}^{(1)\mu}, \quad (2.19)$$

$$(\delta_\zeta^{(0)} \tilde{e}_\mu^a) E_a^{(2)\mu} + (\delta_\zeta^{(1)} \tilde{e}_\mu^a) E_a^{(1)\mu} + (\delta_\zeta^{(1)} \Psi_\mu) \mathcal{E}^{(1)\mu} = \partial_\mu J_{(\zeta)}^{(2)\mu}. \quad (2.20)$$

³ If the background has non-zero curvature, $\delta_\xi \tilde{e}_\mu^a$ does not necessarily vanish at lowest order even if ξ is a Killing vector of the background metric. One needs to consider a transformation modified by an infinitesimal local Lorentz transformation in this case to make $\delta_\xi \tilde{e}_\mu^a$ vanish at lowest order. This is done in appendix A.

We emphasize here that these identities hold for any vector ζ^μ and for any field configuration. Now, if the vector field ξ^μ is a constant vector, then $\delta_\xi^{(0)}\tilde{e}_\mu^a = 0$ as can readily be seen from (2.11). Hence, by (2.19), the current $J_{(\xi)}^{(1)\mu}$ is conserved. Then the charge

$$P_{(\xi)}^{(1)} := \int d^3\vec{x} J_{(\xi)}^{(1)0}, \quad (2.21)$$

is conserved for any field configuration. In particular, it is conserved even if the fields are smoothly deformed to 0 in the past or future of the $t = \text{constant}$ Cauchy surface where the integral is evaluated. Since $P_{(\xi)}^{(1)} = 0$ if the fields vanish, the conservation of this charge implies that

$$P_{(\xi)}^{(1)} = 0, \quad (2.22)$$

for any field configuration if ξ^μ is a constant vector. The compactness of the Cauchy surfaces is crucial for this conclusion because this charge is not necessarily conserved if the field configuration is time-dependent at infinity for the case where the Cauchy surfaces are non-compact.

Now, suppose one attempts to solve the field equations for \tilde{e}_μ^a and Ψ_μ order by order. Let $(\tilde{e}_\mu^{(1)a}, \Psi_\mu^{(1)})$ be a solution to the linearized equations and $(\tilde{e}_\mu^{(2)a}, \Psi_\mu^{(2)})$ be the second-order correction. Then,

$$E_a^{(1)\mu}[\tilde{e}^{(1)}] = 0, \quad (2.23)$$

$$\mathcal{E}^{(1)\mu}[\Psi^{(1)}] = 0, \quad (2.24)$$

$$E_a^{(1)\mu}[\tilde{e}^{(2)}] + E_a^{(2)\mu}[\tilde{e}^{(1)}, \Psi^{(1)}] = 0, \quad (2.25)$$

$$\mathcal{E}^{(1)\mu}[\Psi^{(2)}] + \mathcal{E}^{(2)\mu}[\tilde{e}^{(1)}, \Psi^{(1)}] = 0. \quad (2.26)$$

Here $\mathcal{E}^{(2)\mu}[\tilde{e}^{(1)}, \Psi^{(1)}]$ is the vector-spinor $\mathcal{E}^{(2)\mu}$ evaluated with $(\tilde{e}_\mu^a, \Psi_\mu) = (\tilde{e}_\mu^{(1)a}, \Psi_\mu^{(1)})$, and similarly for the others. The identities (2.19) and (2.20) imply

$$\partial_\mu(J_{(\zeta)}^{(1)\mu}[\tilde{e}^{(2)}] + J_{(\zeta)}^{(2)\mu}[\tilde{e}^{(1)}, \Psi^{(1)}]) = 0, \quad (2.27)$$

for any vector field ζ^μ as long as field equations (2.23)–(2.25) are satisfied. The charge corresponding to the conserved current $J_{(\zeta)}^{(1)\mu}[\tilde{e}^{(2)}] + J_{(\zeta)}^{(2)\mu}[\tilde{e}^{(1)}, \Psi^{(1)}]$ must vanish for any ζ^μ because this charge is conserved even if the field ζ^μ is smoothly deformed to zero in the past or future and is evaluated there. Hence, if we define

$$P_{(\zeta)}^{(2)}[\tilde{e}^{(1)}, \Psi^{(1)}] := \int d^3\vec{x} J_{(\zeta)}^{(2)0}[\tilde{e}^{(1)}, \Psi^{(1)}], \quad (2.28)$$

then

$$P_{(\zeta)}^{(2)}[\tilde{e}^{(1)}, \Psi^{(1)}] = -P_{(\zeta)}^{(1)}[\tilde{e}^{(2)}], \quad (2.29)$$

for any vector field ζ^μ as long as the fields $(\tilde{e}_\nu^{(1)b}, \Psi_\nu^{(1)})$ and $(\tilde{e}_\nu^{(2)b}, \Psi_\nu^{(2)})$ are perturbative solutions to the field equations. In particular, if $\zeta^\mu = \xi^\mu$ is a Killing vector, then

$$P_{(\xi)}^{(2)}[\tilde{e}^{(1)}, \Psi^{(1)}] = 0, \quad (2.30)$$

because of (2.22). Equation (2.30) is a consequence of the requirement that the solution $(\tilde{e}_\nu^{(1)b}, \Psi_\nu^{(1)})$ extend to an exact solution and does not follow from the linearized field equations.

This equation is called a linearization stability condition (LSC) and has to be imposed on the solutions to the linearized equations. In appendix B we illustrate the argument leading to (2.30) for electrodynamics. The total charge of the linearized charged field must vanish in this model.

The LSCs from local supersymmetry can be derived in a similar manner. We expand the supersymmetry transformation given by (2.5) and (2.6) according to the number of fields in the product as

$$\Delta_\epsilon \tilde{z}_\mu^a = \Delta_\epsilon^{(1)} \tilde{z}_\mu^a, \quad (2.31)$$

$$\Delta_\epsilon \Psi_\mu = \Delta_\epsilon^{(0)} \Psi_\mu + \Delta_\epsilon^{(1)} \Psi_\mu + \Delta_\epsilon^{(2)} \Psi_\mu + \dots \quad (2.32)$$

It is clear that $\Delta_\epsilon \tilde{z}_\mu^a = \Delta_\epsilon e_\mu^a$ given by (2.5) has no field independent contribution. That is, $\Delta_\epsilon^{(0)} \tilde{z}_\mu^a = 0$. The analogue of the identity (2.12) is

$$\Delta_\epsilon \tilde{z}_\mu^a \frac{\delta S}{\delta \tilde{z}_\mu^a} + \Delta_\epsilon \Psi_\mu \frac{\delta S}{\delta \Psi_\mu} = \partial_\mu (\sqrt{-g} \mathcal{J}_{(\epsilon)}^\mu). \quad (2.33)$$

From this equation one finds

$$(\Delta_\epsilon^{(0)} \Psi_\mu) \mathcal{E}^{(1)\mu} = \partial_\mu \mathcal{J}_{(\epsilon)}^{(1)\mu}, \quad (2.34)$$

$$(\Delta_\epsilon^{(0)} \Psi_\mu) \mathcal{E}^{(2)\mu} + (\Delta_\epsilon^{(1)} \tilde{z}_\mu^a) E_a^{(1)\mu} + (\Delta_\epsilon^{(1)} \Psi_\mu) \mathcal{E}^{(1)\mu} = \partial_\mu \mathcal{J}_{(\epsilon)}^{(2)\mu}, \quad (2.35)$$

for any spinor field ϵ and any field configuration. Since $\Delta_\epsilon^{(0)} \Psi_\mu = \partial_\mu \epsilon = 0$ if ϵ is a constant spinor, the charge defined by

$$Q_{(\epsilon)}^{(1)} := \int d^3 \vec{x} \mathcal{J}_{(\epsilon)}^{(1)0}, \quad (2.36)$$

vanishes for any field configuration if $\epsilon = \varepsilon$ is a constant spinor by the argument which led to (2.22). By the same argument as that led to (2.29), if we define

$$Q_{(\epsilon)}^{(2)}[\tilde{z}^{(1)}, \Psi^{(1)}] := \int d^3 \vec{x} \mathcal{J}_{(\epsilon)}^{(2)0}[\tilde{z}^{(1)}, \Psi^{(1)}], \quad (2.37)$$

then

$$Q_{(\epsilon)}^{(2)}[\tilde{z}^{(1)}, \Psi^{(1)}] = -Q_{(\epsilon)}^{(1)}[\Psi^{(2)}], \quad (2.38)$$

for any spinor field ϵ if $(\tilde{z}_\nu^{(1)b}, \Psi_\nu^{(1)})$ gives a solution to the linearized field equations, $E_a^{(1)\mu}[\tilde{z}^{(1)}] = \mathcal{E}^{(1)\mu}[\Psi^{(1)}] = 0$. In particular, if $\epsilon = \varepsilon$ is a constant spinor, since $Q_{(\varepsilon)}^{(1)}[\Psi^{(2)}] = 0$ in this case, we must have

$$Q_{(\varepsilon)}^{(2)}[\tilde{z}^{(1)}, \Psi^{(1)}] = 0. \quad (2.39)$$

These are the LSCs arising from local supersymmetry on static three-torus space.

Next, we shall derive the conserved currents $J_{(\xi)}^{(2)\mu}[\tilde{z}^{(1)}, \Psi^{(1)}]$ given by (2.20) and $\mathcal{J}_{(\xi)}^{(2)\mu}[\tilde{z}^{(1)}, \Psi^{(1)}]$ given by (2.35) and the corresponding conserved charges $P_{(\xi)}^{(2)}[\tilde{z}^{(1)}, \Psi^{(1)}]$ and $Q_{(\xi)}^{(2)}[\tilde{z}^{(1)}, \Psi^{(1)}]$ for a constant vector ξ^μ and a constant spinor ε . From now on we write $\tilde{z}_\mu^{(1)a} = \tilde{z}_\mu^a$ and $\Psi_\mu^{(1)} = \Psi_\mu$.

For either charge we only need the linearized field equations. To find $E_a^{(1)\mu}$ for \tilde{z}_μ^a , it is useful to note that eR in the Lagrangian density in terms of the vierbein fields equals $\sqrt{-g} R$ given

in terms of the metric tensor $g_{\mu\nu}$. Thus, we may vary $\sqrt{-g}R$ with respect to the metric tensor and then vary the metric tensor with respect to the vierbein fields. Writing $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$, we find the perturbation $h_{\mu\nu}$ in terms of \tilde{e}_μ^a at first order as

$$h_{\mu\nu} = \delta_\mu^a \tilde{e}_{a\nu} + \tilde{e}_{a\mu} \delta_\nu^a. \quad (2.40)$$

Then we find

$$E_a^{(1)\mu} [\tilde{e}] = \frac{1}{2} \delta_a^\nu [-\partial^\mu \partial^\lambda h_{\lambda\nu} - \partial_\nu \partial^\lambda h_\lambda^\mu + \partial^\mu \partial_\nu h + \square h_\nu^\mu + \delta_\nu^\mu \partial^\lambda \partial^\sigma h_{\lambda\sigma} - \delta_\nu^\mu \square h], \quad (2.41)$$

where $h := \eta^{\mu\nu} h_{\mu\nu}$ and $\square := \partial_\lambda \partial^\lambda$, with indices raised and lowered by $\eta_{\mu\nu}$. We readily find

$$\mathcal{E}^{(1)\mu} [\Psi] = -C \gamma^{\mu\nu\rho} \partial_\nu \Psi_\rho. \quad (2.42)$$

To find the bosonic conserved current $J_{(\xi)}^{(2)\mu}$, we use (2.40)–(2.42) together with

$$\delta_{(\xi)}^{(1)} \tilde{e}_\mu^a = \xi^\nu \partial_\nu \tilde{e}_\mu^a, \quad (2.43)$$

$$\delta_{(\xi)}^{(1)} \Psi_\mu = \xi^\nu \partial_\nu \Psi_\mu, \quad (2.44)$$

in (2.20), recalling $\delta_{(\xi)}^{(0)} \tilde{e}_\mu^a = 0$, to find

$$\begin{aligned} \partial_\mu J_{(\xi)}^{(2)\mu} [\tilde{e}, \Psi] &= \frac{1}{4} (\xi^\lambda \partial_\lambda h_{\mu\nu}) [-\partial^\mu \partial^\lambda h_\lambda^\nu - \partial^\nu \partial^\lambda h_\lambda^\mu + \partial^\mu \partial^\nu h + \square h^{\mu\nu} \\ &\quad + \eta^{\mu\nu} \partial^\lambda \partial^\sigma h_{\lambda\sigma} - \eta^{\mu\nu} \square h] - (\xi^\lambda \partial_\lambda \bar{\Psi}_\mu) \gamma^{\mu\nu\rho} \partial_\nu \Psi_\rho. \end{aligned} \quad (2.45)$$

The current $J_{(\xi)}^{(2)\mu} [h, \Psi]$, which depends on \tilde{e}_μ^a only through $h_{\mu\nu}$, is identified with the Noether current for the spacetime translation in the direction of ξ^μ of the decoupled theory consisting of a Fierz–Pauli Lagrangian for the graviton and a Rarita–Schwinger Lagrangian for the Majorana gravitino [30]

$$\mathcal{L} = -\frac{1}{4} \partial^\mu h \partial^\nu h_{\mu\nu} + \frac{1}{8} \partial_\mu h \partial^\mu h + \frac{1}{4} \partial_\sigma h^{\mu\sigma} \partial^\rho h_{\mu\rho} - \frac{1}{8} \partial_\rho h^{\mu\nu} \partial^\rho h_{\mu\nu} - \frac{1}{2} \bar{\Psi}_\mu \gamma^{\mu\nu\rho} \partial_\nu \Psi_\rho. \quad (2.46)$$

It can be given explicitly as

$$\begin{aligned} J_{(\xi)}^{(2)\mu} [h, \Psi] &= -(\xi^\lambda \partial_\lambda h_{\rho\sigma}) \frac{\partial \mathcal{L}}{\partial (\partial_\mu h_{\rho\sigma})} - (\xi^\lambda \partial_\lambda \Psi_\rho) \frac{\partial \mathcal{L}}{\partial (\partial_\mu \Psi_\rho)} + \xi^\mu \mathcal{L} \\ &= \frac{1}{4} [\xi^\lambda \partial_\lambda h \partial_\nu h^{\mu\nu} + \xi^\lambda \partial_\lambda h^{\mu\rho} \partial_\rho h - \xi^\lambda \partial_\lambda h \partial^\mu h \\ &\quad - 2\xi^\lambda \partial_\lambda h^{\mu\rho} \partial^\sigma h_{\rho\sigma} + \xi^\lambda \partial_\lambda h_{\rho\sigma} \partial^\mu h^{\rho\sigma}] \\ &\quad + \frac{1}{4} \xi^\mu \left[-\partial^\rho h \partial^\sigma h_{\rho\sigma} + \frac{1}{2} \partial_\rho h \partial^\rho h + \partial_\sigma h^{\rho\sigma} \partial^\lambda h_{\rho\lambda} - \frac{1}{2} \partial_\lambda h_{\rho\sigma} \partial^\lambda h^{\rho\sigma} \right] \\ &\quad + \frac{1}{2} (\xi^\sigma \bar{\Psi}_\nu \gamma^{\nu\mu\rho} \partial_\sigma \Psi_\rho - \xi^\mu \bar{\Psi}_\nu \gamma^{\nu\sigma\rho} \partial_\sigma \Psi_\rho). \end{aligned} \quad (2.47)$$

Next we find the fermionic conserved current $\mathcal{J}_{(\xi)}^{(2)\mu}$. We note that the first-order part of the global supersymmetry transformation is given by

$$\Delta_{(\varepsilon)}^{(1)} \tilde{z}^a = \frac{1}{2} \bar{\varepsilon} \gamma^a \Psi_\mu, \quad (2.48)$$

$$\Delta_{(\varepsilon)}^{(1)} \Psi_\mu = \frac{1}{4} (\partial_\mu \tilde{e}_{av} \gamma^{\nu a} + \partial_\rho h_{\mu\nu} \gamma^{\nu\rho}) \varepsilon. \quad (2.49)$$

By substituting these formulas and equations (2.41) and (2.42) into (2.35), and using the identity [29]

$$\gamma^{\rho\nu} \gamma^{\sigma\mu\kappa} = 3(\eta^{\nu[\sigma} \gamma^{\mu\kappa]\rho} - \eta^{\rho[\sigma} \gamma^{\mu\kappa]\nu} + \eta^{\nu[\sigma} \gamma^{\mu\kappa]\rho} - \eta^{\rho[\sigma} \gamma^{\mu\kappa]\nu}), \quad (2.50)$$

we find

$$\mathcal{J}_{(\varepsilon)}^{(2)\mu}[h, \Psi] = \frac{1}{4} (\partial_\rho h_{\sigma\nu} \bar{\varepsilon} \gamma^{\rho\nu} \gamma^{\mu\sigma\kappa} \Psi_\kappa - \delta_\rho^a \tilde{z}_{av} \bar{\varepsilon} \gamma^{\rho\nu} \gamma^{\mu\sigma\kappa} \partial_\sigma \Psi_\kappa). \quad (2.51)$$

Note that the second term vanishes if the (linear) local Lorentz invariance is fixed by requiring $\tilde{e}_{av} = \delta_a^\mu \delta_\nu^b \tilde{e}_{b\mu}$. (It vanishes by the linearized field equation for Ψ_μ as well.) With this condition imposed, the term which explicitly depends on $\tilde{e}_{b\nu}$ in the supersymmetry transformation $\Delta_{(\varepsilon)}^{(1)} \Psi_\mu$ given by (2.49) vanishes. With this choice the current $\mathcal{J}_{(\varepsilon)}^{(2)\mu}[h, \Psi]$ and the charge $Q_{(\varepsilon)}^{(2)}[h, \Psi]$ are the Noether current and charge, respectively, for the supersymmetry transformation,

$$\Delta_{(\varepsilon)}^{(1)} h_{\mu\nu} = \frac{1}{2} (\bar{\varepsilon} \gamma_\mu \Psi_\nu + \bar{\varepsilon} \gamma_\nu \Psi_\mu), \quad (2.52)$$

$$\Delta_{(\varepsilon)}^{(1)} \Psi_\mu = \frac{1}{4} \partial_\rho h_{\mu\nu} \gamma^{\nu\rho} \varepsilon, \quad (2.53)$$

of the linearized supergravity Lagrangian given by (2.46).

Thus, the classical LSCs are $P_{(\varepsilon)}^{(2)}[h, \Psi] = 0$ and $Q_{(\varepsilon)}^{(2)}[h, \Psi] = 0$, where $P_{(\varepsilon)}^{(2)}[h, \Psi]$ and $Q_{(\varepsilon)}^{(2)}[h, \Psi]$ are the conserved charges corresponding to the conserved currents $J_{(\varepsilon)}^{(2)\mu}[h, \Psi]$ in (2.47) and $\mathcal{J}_{(\varepsilon)}^{(2)\mu}[h, \Psi]$ in (2.51) (without the second term) respectively (see (2.28) and (2.37)). In the next section we discuss linearized supergravity on static three-torus space at the classical level and express the LSCs in a form suitable for quantization.

3. Linearized supergravity on static three-torus space

The results of the previous section allow us to describe the LSCs entirely in terms of the linearized theory. By linearizing $\mathcal{N} = 1$ simple supergravity in 4 dimensions about static three-torus background spacetime, we have the Lagrangian, which is given here again for convenience:

$$\mathcal{L} = -\frac{1}{4} \partial^\mu h \partial^\nu h_{\mu\nu} + \frac{1}{8} \partial_\mu h \partial^\mu h + \frac{1}{4} \partial_\sigma h^{\mu\sigma} \partial^\rho h_{\mu\rho} - \frac{1}{8} \partial_\rho h^{\mu\nu} \partial^\rho h_{\mu\nu} - \frac{1}{2} \bar{\Psi}_\mu \gamma^{\mu\nu\rho} \partial_\nu \Psi_\rho. \quad (3.1)$$

A suitable real Majorana representation for the γ -matrices is given by

$$\begin{aligned} \gamma^0 &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, & \gamma^1 &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \\ \gamma^2 &= \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}, & \gamma^3 &= \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix}, \end{aligned} \quad (3.2)$$

where σ^1 , σ^2 and σ^3 are the standard Pauli matrices. In this representation the charge conjugation matrix takes the form $C = i\gamma^0$. The action with the Lagrangian density (3.1) is invariant under the following gauge transformations:

$$h_{\mu\nu} \rightarrow h_{\mu\nu} + \partial_\mu \zeta_\nu + \partial_\nu \zeta_\mu, \quad (3.3)$$

$$\Psi_\mu \rightarrow \Psi_\mu + \partial_\mu \epsilon, \quad (3.4)$$

where ζ^μ is an arbitrary vector field and ϵ_α is an arbitrary Majorana spinor field satisfying $\epsilon_\alpha^\dagger = \epsilon_\alpha$. This invariance is a remnant of the diffeomorphism invariance and local supersymmetry of the full theory. We shall fix this gauge freedom completely.

Working on the spatial torus and imposing periodic boundary conditions on the fields allows us to decompose each field as a Fourier series⁴. Due to the periodic boundary conditions, each field has a spatially constant zero-momentum component. The zero-momentum sector of the linearized theory will be seen to contain six bosonic and six fermionic degrees freedom and this sector forms an important ingredient in the QLSCs. Meanwhile, the non-zero momentum sector of the theory contains the usual gravitons and gravitinos, which each have two polarization states. Explicitly we expand the fields as

$$h_{\mu\nu}(t, \vec{x}) = \frac{1}{\sqrt{V}} h_{\mu\nu}^{(0)}(t) + \frac{1}{\sqrt{V}} \sum_{\vec{k} \neq 0} \tilde{h}_{\mu\nu}(t, \vec{k}) e^{i\vec{k} \cdot \vec{x}},$$

$$\Psi_\mu(t, \vec{x}) = \frac{1}{\sqrt{V}} \psi_\mu(t) + \frac{1}{\sqrt{V}} \sum_{\vec{k} \neq 0} \tilde{\Psi}_\mu(t, \vec{k}) e^{i\vec{k} \cdot \vec{x}},$$

where the volume of the torus is $V = L_1 L_2 L_3$ and the L_1 , L_2 and L_3 are the periods of the torus in the x -, y - and z -directions respectively. The periodic boundary condition implies $\vec{k} = \left(\frac{2\pi}{L_1} n_1, \frac{2\pi}{L_2} n_2, \frac{2\pi}{L_3} n_3 \right)$ with n_1 , n_2 and n_3 integers, and reality restricts $h_{\mu\nu}^{(0)}$ to be real, $\tilde{h}_{\mu\nu}(t, -\vec{k})$ to be equal to $\tilde{h}_{\mu\nu}^\dagger(t, \vec{k})$, and similarly for ψ_μ and $\tilde{\Psi}_\mu(t, \vec{k})$.

The Lagrangian for the theory does not provide dynamical coupling between the modes with zero momentum and those with non-zero momentum \vec{k} . Therefore, it is possible to analyse these separately. We begin with the zero-momentum sector of the theory. We write $h_{\mu\nu}^{(0)}(t) = h_{\mu\nu}(t)$, dropping the superscript '(0)', in the rest of this section. The Lagrangian, i.e. the space integral of the Lagrangian density, for this sector reads

$$L_0 = -\frac{1}{8} \partial_0 h^i{}_i \partial_0 h^j{}_j + \frac{1}{8} \partial_0 h^{ij} \partial_0 h_{ij} + \frac{1}{2} \bar{\psi}_i \gamma^0 \gamma^{ij} \partial_0 \psi_j. \quad (3.5)$$

The classical [31] and quantum [25] theory of the graviton contributions have been studied previously using the ADM formalism without fixing the gauge. Here we fix the gauge to extract the physical degrees of freedom in the linearized theory. (This gauge fixing has little to do with the imposition of LSCs discussed later).

The linearized Lagrangian does not provide equations of motion for $h_{\mu 0}$ and $\psi_{0\alpha}$, and these are precisely the components which carry the gauge degrees of freedom. It is therefore possible to gauge-fix these components to vanish by solving $h_{00}(t) = 2\partial_0 \xi_0(t)$, $h_{i0}(t) = \partial_0 \xi_i(t)$,

⁴ It is possible to choose other boundary conditions also compatible with local supersymmetry, such as anti-periodic boundary conditions for both the gravitino Ψ_μ and the local supersymmetry parameter ϵ . However, only the periodic boundary conditions lead to fermionic LSCs.

$i = 1, 2, 3$ and $\psi_0(t) = \partial_0 \epsilon(t)$. The elimination of $h_{0\mu}(t)$ and $\psi_0(t)$ exhausts the gauge freedom in the zero-momentum sector, and all other components are physical. Thus, in the zero-momentum sector the physical degrees of freedom are contained in the symmetric tensor h_{ij} and vector-spinor $\psi_{i\alpha}$, $i = 1, 2, 3$, each of which corresponds to six classical degrees of freedom.

The six bosonic degrees of freedom are contained in the symmetric tensor h_{ij} . To analyse this tensor it is convenient to break it up into the trace and trace-free sectors. Thus, we let

$$h_{ij}(t) = \sqrt{\frac{2}{3}} \delta_{ij} c(t) + 2 \sum_{A=1}^5 T_{ij}^A c_A(t), \quad (3.6)$$

where the T_{ij}^A are a set of five trace-free tensors which satisfy the orthonormality condition $T^{Aij} T_{ij}^B = \delta^{AB}$. We choose them as

$$\begin{aligned} T^1 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad T^2 = \frac{1}{\sqrt{6}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \\ T^3 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad T^4 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\ T^5 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \end{aligned} \quad (3.7)$$

In terms of the six variables c and c^A , the Lagrangian reads

$$L_0 = -\frac{1}{2} (\partial_0 c)^2 + \frac{1}{2} \sum_{A=1}^5 (\partial_0 c_A)^2 + \frac{1}{2} \bar{\psi}_i \gamma^0 \gamma^{ij} \partial_0 \psi_j. \quad (3.8)$$

The momentum conjugate to h_{ij} , i.e. $p^{ij} = \partial L_0 / \partial (\partial_0 h_{ij})$, is

$$\begin{aligned} p^{ij} &= -\frac{1}{4} (\partial_0 h^k{}_k) \delta^{ij} + \frac{1}{4} \partial_0 h^{ij} \\ &= \frac{1}{2} \sqrt{\frac{2}{3}} \delta^{ij} c_P + \frac{1}{2} \sum_A T^{Aij} c_{PA}, \end{aligned} \quad (3.9)$$

where $c_P = -\partial_0 c$ and $c_{PA} = \partial_0 c_A$ are the momenta conjugate to c and c_A , respectively. The time derivative of the zero-momentum field h_{ij} is then

$$\partial_0 h_{ij} = 2 \sum_{A=1}^5 T_{ij}^A c_{PA} - \sqrt{\frac{2}{3}} \delta_{ij} c_P. \quad (3.10)$$

The canonical Poisson bracket relations for c , c_A , c_P and c_{PA} , and equivalently for h_{ij} and p^{kl} , are

$$\{c, c_P\}_P = 1, \quad \{c_A, c_{PB}\}_P = \delta_{AB}, \quad (3.11)$$

$$\{h_{ij}, p^{kl}\}_P = \frac{1}{2} (\delta_i^k \delta_j^l + \delta_i^l \delta_j^k). \quad (3.12)$$

The fermionic part of L_0 is of first order in time derivatives and therefore already defines a constrained dynamical system [32, 33], and we need to use the Dirac bracket as the bracket to be ‘promoted’ to the anti-commutator upon quantization. The momentum variables π_α^i conjugate to $\psi_{i\alpha}$ are

$$\begin{aligned}\pi_\alpha^i &= \frac{\partial L_0}{\partial(\partial_0\psi_{i\alpha})} \\ &= -\frac{1}{2}(\bar{\psi}_j\gamma^0\gamma^{ji})_\alpha,\end{aligned}\quad (3.13)$$

where the derivative of L_0 with respect to $\partial_0\psi_{i\alpha}$ is the left derivative. These momenta are not invertible in terms of the coordinates and velocities, so there are twelve primary constraints,

$$\phi_\alpha^i = \pi_\alpha^i + \frac{1}{2}(\bar{\psi}_j\gamma^0\gamma^{ji})_\alpha \approx 0. \quad (3.14)$$

We note that the Poisson and Dirac brackets for two fermionic fields are symmetric under the exchange of arguments. That is, if the fields A and B are fermionic, then $\{A, B\}_P = \{B, A\}_P$, and similarly for the Dirac bracket.

For the associated primary Hamiltonian H_P we add arbitrary linear combinations of the primary constraints to the canonical Hamiltonian,

$$H_P = \partial_0\psi_i\pi^i - L_0 + \lambda^i\phi_i = \lambda^i\phi_i, \quad (3.15)$$

where the λ_α^i are arbitrary. Notice in particular that the primary Hamiltonian for these modes weakly vanishes, i.e. it vanishes if the constraints are satisfied. The constraints are to be preserved under evolution by the primary Hamiltonian. This requirement leads to consistency conditions

$$\partial_0\phi^i \approx \{\phi^i, H_P\}_P \approx 0. \quad (3.16)$$

To evaluate the Poisson bracket between the constraints, we use the canonical Poisson bracket,

$$\left\{\psi_{i\alpha}, \pi_\beta^j\right\}_P = -\delta_j^i\delta_{\alpha\beta}, \quad (3.17)$$

which allows us to compute the Poisson bracket between the constraints (and therefore the primary Hamiltonian) as

$$\left\{\phi_\alpha^i, \phi_\beta^j\right\}_P = -(C\gamma^0\gamma^{ij})_{\alpha\beta}. \quad (3.18)$$

Hence, we find that the consistency conditions (3.16) imply $\lambda^i = 0$. Thus, the primary Hamiltonian H_P vanishes and, as a result, the fields ψ_i are time independent. This fact can also be deduced from the Euler–Lagrange equation resulting from (3.8). There are no secondary constraints and all the constraints are of second class. To obtain the bracket structure for the theory suitable for subsequent quantization, we compute the Dirac bracket

$$\begin{aligned}\left\{\psi_{i\alpha}, \pi_\beta^j\right\}_D &= \left\{\psi_{i\alpha}, \pi_\beta^j\right\}_P - \left\{\psi_{i\alpha}, \phi_\gamma^k\right\}_P (\{\phi, \phi\}_P)^{-1}_{kl\gamma\delta} \left\{\phi_\delta^l, \pi_\beta^j\right\}_P \\ &= \frac{1}{2}\left\{\psi_{i\alpha}, \pi_\beta^j\right\}_P \\ &= -\frac{1}{2}\delta_i^j\delta_{\alpha\beta}.\end{aligned}\quad (3.19)$$

On the Dirac bracket, we can impose the second-class constraints as strong conditions since the Dirac bracket between second-class constraints vanishes.

Working explicitly in the Majorana representation, where $C = i\gamma^0$, the Dirac bracket for $\psi_{i\alpha}$ evaluates to

$$\{\psi_{i\alpha}, \psi_{j\beta}\}_D = -\frac{i}{2} (\delta_{ij}\delta_{\alpha\beta} - (\gamma_{ij})_{\alpha\beta}). \quad (3.20)$$

To provide some insight into these relations we write

$$\psi_{i\alpha} = \frac{1}{\sqrt{6}}(\gamma_i\eta)_\alpha + \sum_{A=1}^2 T_{ij}^A(\gamma^j\eta^A)_\alpha, \quad (3.21)$$

where η and η_A are Majorana spinors. The matrices T^1 and T^2 are given in (3.7). These symmetric and traceless matrices satisfy

$$T^{Aij}\gamma_j T_{ik}^B\gamma^k = \delta^{AB}. \quad (3.22)$$

One can also show, by a component-by-component calculation,

$$\sum_{A=1}^2 T_{ik}^A\gamma^k T_{j\ell}^A\gamma^\ell = \frac{2}{3}\delta_{ij} - \frac{1}{3}\gamma_{ij}. \quad (3.23)$$

These relations can be used to show that the Dirac bracket relations for $\psi_{i\alpha}$ are equivalent to

$$\{\eta_\alpha, \eta_\beta\}_D = i\delta_{\alpha\beta}, \quad \{\eta_\alpha^A, \eta_\beta^B\}_D = -i\delta^{AB}\delta_{\alpha\beta}, \quad \{\eta_\alpha, \eta_\beta^A\}_D = 0. \quad (3.24)$$

Next, we briefly describe the mode expansion of the non-zero-momentum sector of this theory, which is well known. Letting the non-zero-momentum components of the fields be denoted by $\hat{h}_{\mu\nu}$ and $\hat{\Psi}_\mu$, we can completely fix the gauge freedom in this sector by imposing the following conditions (see, for instance, [29, 30]):

$$\hat{h}_{\mu 0} = \hat{h} = \partial^i \hat{h}_{ij} = 0, \quad (3.25)$$

$$\gamma^i \hat{\Psi}_i = \hat{\Psi}_0 = \partial^i \hat{\Psi}_i = 0. \quad (3.26)$$

Then we can write the expansion of the graviton as follows:

$$\hat{h}_{ij}(\vec{x}, t) = \sqrt{\frac{2}{V}} \sum_{\vec{k} \neq 0} \sum_{\lambda=\pm} \frac{1}{\sqrt{k}} \left[H_{ij}^\lambda(\vec{k}) a_\lambda(\vec{k}) e^{ik \cdot x} + H_{ij}^{\lambda*}(\vec{k}) a_\lambda^\dagger(\vec{k}) e^{-ik \cdot x} \right], \quad (3.27)$$

where $k := |\vec{k}|$ and $k^\mu = (k, \vec{k})$. We have let the complex conjugate of $a_\lambda(\vec{k})$ be denoted by $a_\lambda^\dagger(\vec{k})$, anticipating quantization. The symmetric and traceless polarization tensors are given by

$$H_{ij}^\lambda(\vec{k}) = \epsilon_i^\lambda(\vec{k}) \epsilon_j^\lambda(\vec{k}), \quad (3.28)$$

where the polarization vectors $\epsilon_i^\lambda(\vec{k})$ are given by

$$\epsilon_i^\pm(\vec{k}) = \frac{1}{\sqrt{2}}(\hat{e}_i^{(1)}(\vec{k}) \pm i\hat{e}_i^{(2)}(\vec{k})). \quad (3.29)$$

The unit spatial vectors $\hat{e}_i^{(1)}(\vec{k})$, $\hat{e}_i^{(2)}(\vec{k})$ and $\hat{e}_i^{(3)}(\vec{k}) = \vec{k}/k$ form a right-handed orthonormal system in this order. The polarization vectors $\epsilon_i^\pm(\vec{k})$ satisfy $\epsilon^{\lambda*}(\vec{k}) \cdot \epsilon^\lambda(\vec{k}) = \delta^{\lambda\lambda'}$ and $k^i \epsilon_i^\lambda(\vec{k}) = 0$.

As a result, $H_{ij}^\lambda(\vec{k})$ satisfy $k^i H_{ij}^\lambda(\vec{k}) = 0$ and $H_{ij}^{\lambda*}(\vec{k}) H^{\lambda'ij}(\vec{k}) = \delta^{\lambda\lambda'}$. The Lagrangian density for the non-zero-momentum sector of the graviton field in this gauge can be found from (3.1) as

$$\mathcal{L}_{TT} = -\frac{1}{8} \partial_\rho \hat{h}^{ij} \partial^\rho \hat{h}_{ij}. \quad (3.30)$$

The standard procedure to find the Poisson bracket relations for the coefficients $a_\lambda(\vec{k})$ and $a_\lambda^\dagger(\vec{k})$ leads to

$$\left\{ a_\lambda(\vec{k}), a_{\lambda'}^\dagger(\vec{k}') \right\}_P = -i \delta_{\vec{k}, \vec{k}'} \delta_{\lambda\lambda'}, \quad (3.31)$$

with all other brackets among $a_\lambda(\vec{k})$ and $a_\lambda^\dagger(\vec{k})$ vanishing.

The non-zero-momentum sector of the gravitino field can similarly be expanded into modes as

$$\hat{\Psi}_{i\alpha} = \frac{1}{\sqrt{V}} \sum_{\vec{k} \neq 0} \sum_{\lambda=\pm} \frac{1}{\sqrt{2k}} \left[\epsilon_i^\lambda(\vec{k}) u_\alpha^\lambda(\vec{k}) b_\lambda(\vec{k}) e^{ik \cdot x} + \epsilon_i^{\lambda*}(\vec{k}) u_\alpha^{\lambda*}(\vec{k}) b_\lambda^\dagger(\vec{k}) e^{-ik \cdot x} \right], \quad (3.32)$$

where $u^\pm(\vec{k})$ are eigenspinors of $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$ and $\gamma^0 \hat{k} \cdot \vec{\gamma}$, where $\hat{k} := \vec{k}/k$, with eigenvalues ± 1 and -1 , respectively. We normalize them by requiring $u^{\lambda\dagger}(\vec{k}) u^{\lambda'}(\vec{k}') = 2k \delta^{\lambda\lambda'}$. The fermionic coefficients $b_\lambda(\vec{k})$ and $b_\lambda^\dagger(\vec{k})$ satisfy the classical analogues of the anti-commutation relations for the annihilation and creation operators:

$$\left\{ b_\lambda(\vec{k}), b_{\lambda'}^\dagger(\vec{k}') \right\}_D = -i \delta_{\vec{k}, \vec{k}'} \delta_{\lambda\lambda'}, \quad (3.33)$$

with all other brackets among $b_\lambda(\vec{k})$ and $b_\lambda^\dagger(\vec{k})$ vanishing.

4. Imposing the bosonic linearization stability conditions

We quantize the linearized theory by promoting the physical degrees of freedom to operators acting on some Hilbert space, and we impose

$$[(\text{anti}) - \text{commutator}]_\pm = i\hbar \{ \text{Poisson/Dirac Bracket} \}, \quad (4.1)$$

as the algebraic relations between the operators, where $[A, B]_\pm := AB \pm BA$. We work in units with $\hbar = 1$. Thus, equations (3.11) and (3.31) become

$$[c, c_P]_- = i, \quad [c_A, c_{PB}]_- = i \delta_{AB}, \quad (4.2)$$

$$\left[a_\lambda(\vec{k}), a_{\lambda'}^\dagger(\vec{k}') \right]_- = \delta_{\lambda\lambda'} \delta_{\vec{k}, \vec{k}'}, \quad (4.3)$$

respectively. On the other hand, the relations (3.24) and (3.33) become

$$[\eta_\alpha, \eta_\beta]_+ = -\delta_{\alpha\beta}, \quad [\eta_\alpha^A, \eta_\beta^B]_+ = \delta^{AB} \delta_{\alpha\beta}, \quad [\eta_\alpha, \eta_\beta^A]_+ = 0, \quad (4.4)$$

$$\left[b_\lambda(\vec{k}), b_{\lambda'}^\dagger(\vec{k}') \right]_+ = \delta_{\lambda\lambda'} \delta_{\vec{k}, \vec{k}'}, \quad (4.5)$$

respectively.

After imposing our gauge conditions and using the field equations, the time component of the conserved Noether current $J_{(\xi)}^{(2)\mu}[h, \Psi]$ for the spacetime translation symmetry becomes

$$J_{(\xi)}^{(2)0} = \frac{\sqrt{V}}{8} \xi_0 [\partial_0 h_{ij} \partial_0 h^{ij} - \partial_0 h^j{}_i \partial_0 h^i{}_j] - \frac{1}{4} \xi_\nu \partial^\nu \hat{h}^{ij} \partial_0 \hat{h}_{ij} + \frac{1}{8} \xi_0 \partial_\nu \hat{h}_{ij} \partial^\nu \hat{h}^{ij} - \frac{i}{2} \xi_\nu \hat{\Psi}_i^T \partial^\nu \hat{\Psi}^i, \quad (4.6)$$

where we have dropped the cross terms between the zero-momentum and non-zero-momentum sectors because they do not contribute to the space integral of $J_{(\xi)}^{(2)0}$, which gives the conserved charge. Let us write

$$\xi_0 H + \xi_i P^i = \int d^3 \vec{x} J_{(\xi)}^{(2)0}. \quad (4.7)$$

By substituting (3.9), (3.27) and (3.32) into (4.6) and integrating the result over space, one finds

$$H = -\frac{1}{2} c_P^2 + \sum_{A=1}^5 \frac{1}{2} c_{PA}^2 + \sum_{\vec{k} \neq 0} \sum_{\lambda=\pm} |\vec{k}| \left(a_\lambda^\dagger(\vec{k}) a_\lambda(\vec{k}) + b_\lambda^\dagger(\vec{k}) b_\lambda(\vec{k}) \right), \quad (4.8)$$

$$\vec{P} = \sum_{\vec{k} \neq 0} \sum_{\lambda=\pm} \vec{k} \left(a_\lambda^\dagger(\vec{k}) a_\lambda(\vec{k}) + b_\lambda^\dagger(\vec{k}) b_\lambda(\vec{k}) \right), \quad (4.9)$$

the bosonic sector of which agrees with the conserved charges given in [25]. Note that the zero-point energy which had to be renormalized away in the pure-gravity case is absent here because of supersymmetry. Note also that the Hamiltonian H is not positive definite.

The Hilbert space for the theory can be constructed as the tensor product of a non-zero momentum sector and a sector with zero momentum. In the sector with non-zero momentum, we have the usual Fock spaces for the gravitons and gravitinos with the number operators for the gravitons and gravitinos (with a given momentum and a polarization) being $a_\lambda^\dagger(\vec{k}) a_\lambda(\vec{k})$ and $b_\lambda^\dagger(\vec{k}) b_\lambda(\vec{k})$, respectively. A suitable basis of states is given by states with definite number of particles in each momentum and polarization. Thus, the normalized state with $n_{B\lambda}(\vec{k})$ gravitons and $n_{F\lambda}(\vec{k})$ gravitinos with momentum \vec{k} and polarization λ can be given as

$$|\{n_B\}\rangle \otimes |\{n_F\}\rangle = \prod_{\vec{k}, \lambda=\pm} \left[\frac{1}{\sqrt{n_{B\lambda}(\vec{k})!}} (a_\lambda^\dagger(\vec{k}))^{n_{B\lambda}(\vec{k})} (b_\lambda^\dagger(\vec{k}))^{n_{F\lambda}(\vec{k})} |0\rangle \right], \quad (4.10)$$

where the vacuum state $|0\rangle$ satisfies $a_\lambda(\vec{k})|0\rangle = b_\lambda(\vec{k})|0\rangle = 0$ for all \vec{k} and λ . (For the gravitino $n_{F\lambda}(\vec{k}) = 0$ or 1, of course).

For the graviton zero-modes, we represent the commutation rules $[c, c_P]_- = i$ and $[c_A, c_{PB}]_- = i\delta_{AB}$ on wave functions which are functions of the variables c and c_A by

$$c \mapsto \text{multiply by } c, \quad c_P \mapsto -i \frac{\partial}{\partial c}, \quad (4.11)$$

$$c_A \mapsto \text{multiply by } c_A, \quad c_{PA} \mapsto -i \frac{\partial}{\partial c_A}. \quad (4.12)$$

Therefore, if we take some state which is proportional to a single eigenstate of the number operators,

$$|\text{state}\rangle = \Psi \otimes |\{n_B\}\rangle \otimes |\{n_F\}\rangle, \quad (4.13)$$

where Ψ is some wave function of c and c_A , then the Hamiltonian and momentum operators, (4.8) and (4.9), acting on such a state take the form

$$H |\text{state}\rangle = \frac{1}{2} \left(+ \frac{\partial^2}{\partial c^2} - \sum_{A=1}^5 \frac{\partial^2}{\partial c_A^2} + M^2 \right) |\text{state}\rangle, \quad (4.14)$$

$$\vec{P} |\text{state}\rangle = \sum_{\vec{k} \neq 0} \sum_{\lambda=\pm} \vec{k} \left(n_{B\lambda}(\vec{k}) + n_{F\lambda}(\vec{k}) \right) |\text{state}\rangle, \quad (4.15)$$

where we have defined a ‘squared mass’ for these states by

$$M^2 = 2 \sum_{\vec{k} \neq 0} \sum_{\lambda=\pm} |\vec{k}| \left(n_{B\lambda}(\vec{k}) + n_{F\lambda}(\vec{k}) \right). \quad (4.16)$$

The operators H and \vec{P} do not contain the zero-momentum sector of the gravitino field. They appear in the global supercharges as we shall see in the next section.

As discussed in section 2, the conserved charges H and \vec{P} must vanish classically for the classical solutions to the linearized equations if they were to be extendible to exact solutions. Moncrief proposed that in quantum theory these linearization stability conditions (LSCs) should be imposed as constraints on the physical states. Thus, the constraints on the physical states $|\text{phys}\rangle$ are

$$H |\text{phys}\rangle = 0, \quad \vec{P} |\text{phys}\rangle = 0. \quad (4.17)$$

In [25] the group-averaging procedure was used to find all states satisfying these constraints and define an inner product among these states for pure gravity on static three-torus space. We apply this procedure to $\mathcal{N} = 1$ simple supergravity in this section.

We start with the Hilbert space \mathcal{H}_0 of superpositions of the states defined by (4.13). Take two states in this Hilbert space of the form $|\varphi_1\rangle = \Psi_1(c, c_A) \otimes |\Phi_1\rangle$ and $|\varphi_2\rangle = \Psi_2(c, c_A) \otimes |\Phi_2\rangle$, where $|\Phi_1\rangle$ and $|\Phi_2\rangle$ are states in the Fock space \mathcal{F} of non-zero-momentum gravitons and gravitinos. The inner product between these states is

$$\langle \varphi_1 | \varphi_2 \rangle_{\mathcal{H}_0} = \langle \Phi_1 | \Phi_2 \rangle_{\mathcal{F}} \int dc d^5 \vec{c} \Psi_1^* \Psi_2, \quad (4.18)$$

where \vec{c} is the vector with components c_A , $A = 1, 2, 3, 4, 5$. Let us choose both $|\Phi_1\rangle$ and $|\Phi_2\rangle$ to have a definite value of M^2 defined by (4.16). (If these states have different values of M^2 , then $\langle \Phi_1 | \Phi_2 \rangle_{\mathcal{F}} = 0$.) Then, by expressing $\Psi_I(c, \vec{c})$, $I = 1, 2$, as a Fourier integral in the six-dimensional space with coordinates (c, \vec{c}) , these states can be given as

$$|\varphi_I\rangle = \int \frac{dp^0}{2\pi} \frac{d^5 \vec{p}}{(2\pi)^5} F_I(p^0, \vec{p}) e^{-ip^0 c + i\vec{p} \cdot \vec{c}} \otimes |\Phi_I\rangle, \quad (4.19)$$

where \vec{p} is also a five-dimensional vector and $\vec{p} \cdot \vec{c} = p^A c_A$. The inner product for these states is

$$\langle \varphi_1 | \varphi_2 \rangle_{\mathcal{H}_0} = \langle \Phi_1 | \Phi_2 \rangle_{\mathcal{F}} \int \frac{dp^0}{2\pi} \frac{d^5 \vec{p}}{(2\pi)^5} F_1^*(p^0, \vec{p}) F_2(p^0, \vec{p}). \quad (4.20)$$

Since the operators H and \vec{P} are the generators of spacetime translations, we can construct states satisfying (4.17) by averaging the states $|\varphi_I\rangle$ over this translation group as follows:

$$|\varphi_I^{(B)}\rangle = \frac{1}{2V} \int_{-\infty}^{\infty} d\alpha^0 \left(\prod_{i=1}^3 \int_0^{L_i} d\alpha_i \right) \exp(i\alpha^0 H - i\vec{\alpha} \cdot \vec{P}) |\varphi_I\rangle. \quad (4.21)$$

The integral over space acts as the projector onto the sector of the Fock space with zero total momentum:

$$\frac{1}{V} \left(\prod_{i=1}^3 \int_0^{L_i} d\alpha_i \right) \exp(-i\vec{\alpha} \cdot \vec{P}) |\Phi_I\rangle = \mathcal{P}_{\vec{p}=0} |\Phi_I\rangle. \quad (4.22)$$

The integral over α^0 can readily be evaluated using the representation (4.19) since

$$H e^{-ip^0 c + i\vec{p} \cdot \vec{c}} \otimes |\Phi_I\rangle = \frac{1}{2} [-(p^0)^2 + \vec{p}^2 + M^2] e^{-ip^0 c + i\vec{p} \cdot \vec{c}} \otimes |\Phi_I\rangle. \quad (4.23)$$

Thus we find, with the notation $|\Phi_I^{(\vec{p}=0)}\rangle = \mathcal{P}_{\vec{p}=0} |\Phi_I\rangle$,

$$\begin{aligned} |\varphi_I^{(B)}\rangle &= \int \frac{d^5 p}{(2\pi)^5} \frac{d^5 \vec{p}}{(2\pi)^5} 2\pi \delta((p^0)^2 - \vec{p}^2 - M^2) F_I(p^0, \vec{p}) e^{-ip^0 c + i\vec{p} \cdot \vec{c}} \otimes |\Phi_I^{(\vec{p}=0)}\rangle \\ &= \int \frac{d^5 \vec{p}}{(2\pi)^5} \left[f_I^{(+)}(\vec{p}) e^{-iE(\vec{p})c + i\vec{p} \cdot \vec{c}} + f_I^{(-)}(\vec{p}) e^{iE(\vec{p})c + i\vec{p} \cdot \vec{c}} \right] \otimes |\Phi_I^{(\vec{p}=0)}\rangle, \end{aligned} \quad (4.24)$$

where $E(\vec{p}) = \sqrt{\vec{p}^2 + M^2}$ and where

$$f_I^{(\pm)}(\vec{p}) = \frac{F_I(\pm E(\vec{p}), \vec{p})}{2E(\vec{p})}. \quad (4.25)$$

Although we obtained the invariant state $|\varphi_I^{(B)}\rangle$ by averaging $|\varphi_I\rangle$ over the spacetime translation group, it is easy to show that any state with definite value of M^2 satisfying the constraints (4.17) is of this form. (The zero-momentum sector is a solution to the six-dimensional Klein–Gordon equation with mass M).

The states $|\varphi_I^{(B)}\rangle$ are indeed invariant, i.e. they satisfy the QLSCs given by (4.17). However, they have infinite norm and, hence, are not in the Hilbert space \mathcal{H}_0 : the inner product $\langle \varphi_I^{(B)} | \varphi_I^{(B)} \rangle_{\mathcal{H}_0}$ computed using (4.20) is infinite because of the δ -function in (4.24). The group-averaging inner product for the Hilbert space \mathcal{H}_B of invariant states is defined by

$$\begin{aligned} \langle \varphi_1^{(B)} | \varphi_2^{(B)} \rangle_{\mathcal{H}_B} &= \frac{1}{2V} \int_{-\infty}^{\infty} d\alpha^0 \left(\prod_{i=1}^3 \int_0^{L_i} d\alpha_i \right) \langle \varphi_1 | \exp(i\alpha^0 H - i\vec{\alpha} \cdot \vec{P}) | \varphi_2 \rangle_{\mathcal{H}_0} \\ &= \langle \varphi_1 | \varphi_2^{(B)} \rangle_{\mathcal{H}_0} \\ &= 2 \int \frac{d^5 \vec{p}}{(2\pi)^5} E(\vec{p}) \left[f_1^{(+)*}(\vec{p}) f_2^{(+)}(\vec{p}) + f_1^{(-)*}(\vec{p}) f_2^{(-)}(\vec{p}) \right] \langle \Phi_1^{(\vec{p}=0)} | \Phi_2^{(\vec{p}=0)} \rangle_{\mathcal{F}}. \end{aligned} \quad (4.26)$$

Notice that, although this inner product is defined in terms of the ‘seed states’ $|\varphi_I\rangle$, it depends only on the invariant states $|\varphi_I^{(B)}\rangle$. This inner product is equivalent to that proposed in [34] in the context of quantum cosmology.

Since the constraint $H|\text{phys}\rangle = 0$ is a Klein–Gordon equation, it is tempting to use the Klein–Gordon inner product for the Hilbert space \mathcal{H}_B :

$$\langle \varphi_1^{(B)} | \varphi_2^{(B)} \rangle_{\text{KG}} = i \int d^5 c \left(\Psi_1^* \frac{\partial \Psi_2}{\partial c} - \frac{\partial \Psi_1^*}{\partial c} \Psi_2 \right) \langle \Phi_1^{(\vec{P}=0)} | \Phi_2^{(\vec{P}=0)} \rangle_{\mathcal{F}}. \quad (4.27)$$

This inner product can readily be evaluated as

$$\langle \varphi_1^{(B)} | \varphi_2^{(B)} \rangle_{\text{KG}} = 2 \int \frac{d^5 \vec{p}}{(2\pi)^5} E(\vec{p}) \left[f_1^{(+)*}(\vec{p}) f_2^{(+)}(\vec{p}) - f_1^{(-)*}(\vec{p}) f_2^{(-)}(\vec{p}) \right] \langle \Phi_1^{(\vec{P}=0)} | \Phi_2^{(\vec{P}=0)} \rangle_{\mathcal{F}}, \quad (4.28)$$

which is identical with the group-averaging inner product, $\langle \varphi_1^{(B)} | \varphi_2^{(B)} \rangle_{\mathcal{H}_B}$, given by (4.26) except for the minus sign in the second term. Although the Klein–Gordon inner product is more widely used in quantum cosmology, it can be used only if the space of geometries considered has the structure of spacetime [35]. (In our example, it is the six-dimensional Minkowski space.) The group-averaging inner product, which we use in this paper, has wider applicability. For example, it can be used in recollapsing quantum cosmology as shown in [34].

In the next section we impose the fermionic QLSCs derived in the previous section. We shall find that the states satisfying these constraints and the inner product among them can be found by group-averaging over the relevant supergroup.

5. Imposing the fermionic linearization stability conditions

The time component of the conserved fermionic current $\mathcal{J}_{(\varepsilon)}^{(2)\mu}[h, \Psi]$ is

$$\mathcal{J}_{(\varepsilon)}^{(2)0}[h, \Psi] = \frac{1}{4} \left[\partial_0 h \bar{\varepsilon} \gamma^i \psi_i - \partial_0 h_{ij} \bar{\varepsilon} \gamma^i \psi^j - \partial_0 \hat{h}_{ij} \bar{\varepsilon} \gamma^j \hat{\Psi}^i + \partial_\ell \hat{h}_{ij} \bar{\varepsilon} \gamma^0 \gamma^\ell \gamma^j \hat{\Psi}^i \right], \quad (5.1)$$

where we dropped the cross terms between the zero-momentum and non-zero-momentum sectors since they do not contribute to the space integral. By letting

$$\bar{\varepsilon} Q = - \int d^3 \vec{x} \mathcal{J}_{(\varepsilon)}^{(2)0}, \quad (5.2)$$

and substituting (3.10) and (3.21) into (5.1) and integrating over space, we find

$$Q = Q^{(0)} + \hat{Q}, \quad (5.3)$$

with the zero- and non-zero-momentum contributions respectively given by

$$Q^{(0)} = \frac{1}{2} \left(c_P \eta + \sum_{A=1}^5 \sum_{B=1}^2 T^{Ai}{}_j T^{Bj}{}_{ik} \gamma^j \gamma^k c_{PA} \eta^B \right), \quad (5.4)$$

$$\hat{Q} = \frac{1}{4} \int d^3 \vec{x} \left(\partial_0 \hat{h}^{ij} \gamma_j \hat{\Psi}_i - \hat{h}^{ij} \gamma_j \partial_0 \hat{\Psi}_i \right), \quad (5.5)$$

where we have integrated by parts in the second term of \hat{Q} and used $\gamma^j \partial_j \hat{\Psi}_i = -\gamma^0 \partial_0 \hat{\Psi}_i$.

One is readily able to find the mode expansion for \hat{Q} , which then yields

$$Q = \frac{1}{2} \left(c_P \eta + \sum_{A=1}^5 \sum_{B=1}^2 T^{Ai} T_{ik}^B \gamma^j \gamma^k c_{PA} \eta^B \right) - \frac{i}{2} \sum_{\vec{k} \neq 0} \sum_{\lambda=\pm} \left[\epsilon^\lambda(\vec{k}) \cdot \vec{\gamma} u^{\lambda*}(\vec{k}) a_\lambda(\vec{k}) b_\lambda^\dagger(\vec{k}) - \epsilon^{\lambda*}(\vec{k}) \cdot \vec{\gamma} u^\lambda(\vec{k}) a_\lambda^\dagger(\vec{k}) b_\lambda(\vec{k}) \right]. \quad (5.6)$$

This charge and the bosonic charges, $P^\mu = (H, \vec{P})$, satisfy the supersymmetry algebra. That is, $[P^\mu, Q]_- = 0$, $[P^\mu, P^\nu]_- = 0$ and

$$[Q_\alpha, Q_\beta]_+ = \frac{1}{2} (\gamma_\mu \gamma^0)_{\alpha\beta} P^\mu. \quad (5.7)$$

Since the contribution to Q with different \vec{k} anti-commute, equation (5.7) holds for each \vec{k} including $\vec{k} = 0$. In deriving (5.7) we have used the following identities for $\vec{k} = 0$ and $\vec{k} \neq 0$ proved in appendix C:

$$\sum_{B=1}^2 \sum_{A=1}^5 \sum_{A'=1}^5 T^{Ai} T_{ik}^B T_{i'j'}^{A'} T^{Bi'} T_{k'l'} \gamma^j \gamma^k \gamma^{k'} \gamma^{j'} c_{PA} c_{PA'} = \sum_{A=1}^5 (c_{PA})^2, \quad (5.8)$$

$$\left[\epsilon^{\pm*}(\vec{k}) \cdot \vec{\gamma} u^\pm(\vec{k}) \right]_\alpha \left[\epsilon^\pm(\vec{k}) \cdot \vec{\gamma} u^{\pm*}(\vec{k}) \right]_\beta + (\alpha \leftrightarrow \beta) = 2(\gamma \cdot k \gamma^0)_{\alpha\beta}. \quad (5.9)$$

We also recall that γ^μ are real, γ_5 is purely imaginary and that $\epsilon^{\pm*}(\vec{k}) = \epsilon^\mp(\vec{k})$ and $u^{\pm*}(\vec{k}) = u^\mp(\vec{k})$.

One can represent the anti-commutation relations (4.4) satisfied by the twelve fermionic operators η_α and η_α^A in the zero-momentum sector of the gravitino field by a 64-dimensional indefinite-metric Hilbert space (or Krein space) \mathcal{H}_{0F} as shown in appendix D. The basis vectors of this space can be chosen such that 32 of them are normalized with positive norm and the other 32 are normalized with negative norm. [We say here that $|v\rangle$ is normalized with positive (negative) norm if $\langle v|v\rangle = 1$ ($\langle v|v\rangle = -1$).] An important property of \mathcal{H}_{0F} used later is that the 16-dimensional subspace of \mathcal{H}_{0F} of states annihilated by the operators d_1 and d_2 defined by

$$d_1 := (\eta_1 + i\eta_2)/\sqrt{2}, \quad d_2 := (\eta_3 + i\eta_4)/\sqrt{2}, \quad (5.10)$$

is a positive-norm subspace because this subspace is spanned by the states of the form (D.5) with $n_1 = n_2 = 0$ (with $M = 2$) [see (D.6)]. The Hilbert space of states satisfying the bosonic QLSCs is in fact the tensor product $\mathcal{H}_B \otimes \mathcal{H}_{0F}$. We call this tensor product space also \mathcal{H}_B from now on in order not to complicate the notation.

Now we are in a position to find the physical states $|\text{phys}\rangle$ satisfying the fermionic QLSCs, $Q_\alpha |\text{phys}\rangle = 0$, as well as the bosonic ones, $P^\mu |\text{phys}\rangle = 0$. Since the operators P^μ commute with Q_α , we specialize to the Hilbert space \mathcal{H}_B of states satisfying the bosonic QLSCs and identify P^μ with the null operator. Thus, we have $[Q_\alpha, Q_\beta]_+ = 0$ in place of (5.7). Splitting the supercharge into zero- and non-zero-momentum contributions, these anti-commutation relations of the supercharge can be rewritten as

$$[Q_\alpha^{(0)}, Q_\beta^{(0)}]_+ = -\frac{1}{4} M^2 \delta_{\alpha\beta} = -[\widehat{Q}_\alpha, \widehat{Q}_\beta]_+, \quad (5.11)$$

where M^2 is defined by (4.16). We specialize to an eigenspace of the operator M^2 without loss of generality. Thus, we treat M as if it were a non-negative number.

Assume $M > 0$. Then, in each common eigenspace of the operators c_P and c_{PA} with eigenvalues p_0 and p_A , respectively, satisfying $p_0^2 - \vec{p}^2 = M^2$, the operators $Q_\alpha^{(0)}$ given by (5.4) is of the form

$$Q_\alpha^{(0)} = \frac{M}{2} \left(\pm \eta_\alpha \cosh \theta + \sum_{B=1}^2 \sum_{\beta=1}^4 C_{\alpha B}^\beta \eta_\beta^B \sinh \theta \right), \quad (5.12)$$

where $C_{\alpha B}^\beta$ can be regarded as a 4×8 matrix with the rows and columns labelled by α and (B, β) , respectively. Here, $\cosh \theta = |c_P|/M$ and $\sinh \theta = |\vec{c}_P|/M$, where $\vec{c}_P = (c_{P1}, c_{P2}, c_{P3}, c_{P4}, c_{P5})$. The four eight-dimensional column vectors of the matrix $C_{\alpha B}^\beta$ are orthonormal, i.e.

$$\sum_{B=1}^2 \sum_{\beta=1}^4 C_{\alpha_1 B}^\beta C_{\alpha_2 B}^\beta = \delta_{\alpha_1 \alpha_2}. \quad (5.13)$$

Then by appendix D there is a unitary operator U , i.e. an operator preserving the inner product, on \mathcal{H}_{0F} such that

$$Q_\alpha^{(0)} = (M/2)U\eta_\alpha U^\dagger. \quad (5.14)$$

Instead of directly working with $Q_\alpha^{(0)}$ and \widehat{Q}_α , it is more convenient to combine them into annihilation- and creation-type operators [33] as

$$a_1 = \frac{\sqrt{2}}{M}(Q_1^{(0)} + iQ_2^{(0)}), \quad a_2 = \frac{\sqrt{2}}{M}(Q_3^{(0)} + iQ_4^{(0)}), \quad (5.15)$$

$$b_1 = \frac{\sqrt{2}}{M}(\widehat{Q}_1 + i\widehat{Q}_2), \quad b_2 = \frac{\sqrt{2}}{M}(\widehat{Q}_3 + i\widehat{Q}_4), \quad (5.16)$$

provided that $M > 0$.⁵ These new operators satisfy the anti-commutation relations of Fermi oscillators (up to a sign), i.e. $a_i^2 = b_i^2 = a_i^{\dagger 2} = b_i^{\dagger 2} = 0$ and

$$[a_i, a_j^\dagger]_+ = -\delta_{ij}, \quad [b_i, b_j^\dagger]_+ = \delta_{ij}. \quad (5.17)$$

Now we construct all states $|\varphi^{(BF)}\rangle \in \mathcal{H}_B$ satisfying $Q_\alpha |\varphi^{(BF)}\rangle = 0$, $\alpha = 1, 2, 3, 4$. These constraints can be organized as

$$(a_i + b_i) |\varphi^{(BF)}\rangle = (a_i^\dagger + b_i^\dagger) |\varphi^{(BF)}\rangle = 0, \quad i = 1, 2. \quad (5.18)$$

Note that all states are linear combinations of the states each annihilated by either b_1 or b_1^\dagger since $|\varphi^{(B)}\rangle = b_1^\dagger b_1 |\varphi^{(B)}\rangle + b_1 b_1^\dagger |\varphi^{(B)}\rangle$ for any state $|\varphi^{(B)}\rangle \in \mathcal{H}_B$. The same is true for each pair of operators, (b_2, b_2^\dagger) , (a_1, a_1^\dagger) and (a_2, a_2^\dagger) . This means that a general state is a superposition of states, each belonging to a 16-dimensional Fock space built on a state $|\chi^{(B)}\rangle$ satisfying $a_i |\chi^{(B)}\rangle = b_i |\chi^{(B)}\rangle = 0$, $i = 1, 2$, by applying the creation-type operators a_i^\dagger and b_i^\dagger . Notice that equations (5.14) and (5.15) imply $a_1 = Ud_1 U^\dagger$ and $a_2 = Ud_2 U^\dagger$, where U is a unitary

⁵For $M = 0$, formally one can proceed in a similar manner by considering (δ -function normalisable) plane wave states $\Psi(c, \vec{c}) = e^{\pm i p c + i \vec{p} \vec{c}}$. On such states the supercharge takes the form $Q = \frac{p}{2}(\pm \eta + R)$, with $[\eta_\alpha, \eta_\beta]_+ = -\delta_{\alpha\beta}$ and $[R_\alpha, R_\beta]_+ = \delta_{\alpha\beta}$. Then, we define ladder-type operators $a_1 = (\eta_1 + i\eta_2)/\sqrt{2}$, $a_2 = (\eta_3 + i\eta_4)/\sqrt{2}$, $b_1 = (R_1 + iR_2)/\sqrt{2}$ and $b_2 = (R_3 + iR_4)/\sqrt{2}$ and proceed in a manner similar to the case with $M > 0$.

operator on \mathcal{H}_{0F} and where d_1 and d_2 are defined by (5.10). Since the states annihilated by d_1 and d_2 have positive norm as we stated before, we have $\langle \chi^{(B)} | \chi^{(B)} \rangle_{\mathcal{H}_B} > 0$.

We label the 16 possible states in such a Fock space as follows:

$$(a_1^\dagger)^{m_1} (a_2^\dagger)^{m_2} (b_1^\dagger)^{n_1} (b_2^\dagger)^{n_2} |\chi^{(B)}\rangle = |m_1 m_2 n_1 n_2\rangle, \quad (5.19)$$

with each m_1, m_2, n_1, n_2 being either 0 or 1. For example, we define

$$|\chi^{(B)}\rangle = |0000\rangle, \quad a_1^\dagger |\chi^{(B)}\rangle = |1000\rangle, \quad a_1^\dagger a_2^\dagger b_1^\dagger b_2^\dagger |\chi^{(B)}\rangle = |1111\rangle. \quad (5.20)$$

Thus, we look for states satisfying the fermionic QLSCs in the form

$$|\varphi^{(BF)}\rangle = \sum_{m_1=0}^1 \sum_{m_2=0}^1 \sum_{n_1=0}^1 \sum_{n_2=0}^1 c_{m_1 m_2 n_1 n_2} |m_1 m_2 n_1 n_2\rangle. \quad (5.21)$$

We find

$$(a_1 + b_1) |\varphi^{(BF)}\rangle = \sum_{m_1=0}^1 \sum_{m_2=0}^1 \sum_{n_1=0}^1 \sum_{n_2=0}^1 c_{m_1 m_2 n_1 n_2} [-m_1 |0m_2 n_1 n_2\rangle + (-1)^{m_1+m_2} n_1 |m_1 m_2 0 n_2\rangle]. \quad (5.22)$$

The coefficient of the term $|0m_2 1n_2\rangle$ in this equation is $-c_{1m_2 1n_2}$. Hence $c_{1m_1 n} = 0$ for all m and n . We can conclude similarly that $c_{0m0n} = c_{m0n0} = c_{m1n1} = 0$ by using the other constraints. Thus, $c_{m_1 m_2 n_1 n_2} = 0$ if $m_1 = n_1$ or $m_2 = n_2$. Hence the invariant state $|\varphi^{(BF)}\rangle \in \mathcal{H}_B$, i.e. the state satisfying the fermionic QLSCs, must be of the following form:

$$|\varphi^{(BF)}\rangle = A |1100\rangle + B |0110\rangle + C |1001\rangle + D |0011\rangle, \quad (5.23)$$

where A, B, C and D are constants. Note that the four states $|1100\rangle, |0110\rangle, |1001\rangle$ and $|0011\rangle$ are mutually orthogonal and satisfy

$$\begin{aligned} \langle 1100 | 1100 \rangle &= \langle 0011 | 0011 \rangle = \langle \chi^{(B)} | \chi^{(B)} \rangle_{\mathcal{H}_B}, \\ \langle 0110 | 0110 \rangle &= \langle 1001 | 1001 \rangle = -\langle \chi^{(B)} | \chi^{(B)} \rangle_{\mathcal{H}_B}. \end{aligned} \quad (5.24)$$

In particular, the two states $|0110\rangle$ and $|1001\rangle$ have negative norm.

By applying the constraints (5.18) we find the following unique solution up to an overall normalization:

$$|\varphi^{(BF)}\rangle \propto |\varphi_P^{(BF)}\rangle := |1100\rangle - |0110\rangle + |1001\rangle + |0011\rangle. \quad (5.25)$$

Thus, there is precisely one possible combination which satisfies the fermionic QLSCs in each Fock space spanned by $(a_1^\dagger)^{n_1} (a_2^\dagger)^{n_2} (b_1^\dagger)^{n_1} (b_2^\dagger)^{n_2} |\chi^{(B)}\rangle$, where $|\chi^{(B)}\rangle$ is any state with fixed positive M^2 , satisfying the bosonic QLSCs and the conditions

$$(Q_1^{(0)} + iQ_2^{(0)}) |\chi^{(B)}\rangle = (Q_3^{(0)} + iQ_4^{(0)}) |\chi^{(B)}\rangle = 0, \quad (5.26)$$

$$(\widehat{Q}_1 + i\widehat{Q}_2) |\chi^{(B)}\rangle = (\widehat{Q}_1 + i\widehat{Q}_2) |\chi^{(B)}\rangle = 0. \quad (5.27)$$

Now, in section 4 it was shown that all states satisfying the bosonic QLSCs are obtained by group-averaging. Here we show that the same is true for the fermionic QLSCs. That is, the state $|\varphi^{(BF)}\rangle$ in (5.23) is obtained as

$$|\varphi^{(BF)}\rangle = - \int d^4\theta e^{-\bar{\theta}Q} |\varphi^{(B)}\rangle, \tag{5.28}$$

for some linear combination $|\varphi^{(B)}\rangle$ of the states $|m_1 m_2 n_1 n_2\rangle$, where $\theta = (\theta_1, \theta_2, \theta_3, \theta_4)$ are Grassmann numbers and where $\bar{\theta} = i\theta^T \gamma^0$ and $d^4\theta = d\theta_4 d\theta_3 d\theta_2 d\theta_1$. (The minus sign in (5.28) has been introduced for convenience.) By the usual rules, $\int d\theta_\alpha = 0$ and $\int d\theta_\alpha \theta_\alpha = 1$, $\alpha = 1, 2, 3, 4$, this equation becomes

$$\begin{aligned} |\varphi^{(BF)}\rangle &= -Q_1 Q_2 Q_3 Q_4 |\varphi^{(B)}\rangle \\ &= \frac{1}{4} (Q_1 - iQ_2)(Q_1 + iQ_2)(Q_3 - iQ_4)(Q_3 + iQ_4) |\varphi^{(B)}\rangle \\ &= \frac{M^4}{16} (a_1^\dagger + b_1^\dagger)(a_1 + b_1)(a_2^\dagger + b_2^\dagger)(a_2 + b_2) |\varphi^{(B)}\rangle. \end{aligned} \tag{5.29}$$

One readily finds that $Q_1 Q_2 Q_3 Q_4 |m_1 m_2 n_1 n_2\rangle = 0$ unless $|m_1 m_2 n_1 n_2\rangle = |1100\rangle, |0110\rangle, |1001\rangle$ or $|0011\rangle$ and that

$$(a_1^\dagger + b_1^\dagger)(a_1 + b_1)(a_2^\dagger + b_2^\dagger)(a_2 + b_2) |1100\rangle = |\varphi_P^{(BF)}\rangle, \tag{5.30}$$

$$(a_1^\dagger + b_1^\dagger)(a_1 + b_1)(a_2^\dagger + b_2^\dagger)(a_2 + b_2) |0110\rangle = |\varphi_P^{(BF)}\rangle, \tag{5.31}$$

$$(a_1^\dagger + b_1^\dagger)(a_1 + b_1)(a_2^\dagger + b_2^\dagger)(a_2 + b_2) |1001\rangle = -|\varphi_P^{(BF)}\rangle, \tag{5.32}$$

$$(a_1^\dagger + b_1^\dagger)(a_1 + b_1)(a_2^\dagger + b_2^\dagger)(a_2 + b_2) |0011\rangle = |\varphi_P^{(BF)}\rangle, \tag{5.33}$$

where the state $|\varphi_P^{(BF)}\rangle$ is defined by (5.25). Thus, starting from any linear combination $|\varphi^{(B)}\rangle$ of $|m_1 m_2 n_1 n_2\rangle$, we find

$$- \int d^4\theta e^{-\bar{\theta}Q} |\varphi^{(B)}\rangle = -Q_1 Q_2 Q_3 Q_4 |\varphi^{(B)}\rangle = \kappa |\varphi_P^{(BF)}\rangle, \tag{5.34}$$

with $\kappa \in \mathbb{C}$.

Now, an explicit computation shows that $\langle \varphi_P^{(BF)} | \varphi_P^{(BF)} \rangle_{\mathcal{H}_B} = 0$. In fact this can be deduced immediately from (5.34) because $Q_\alpha^2 = 0$ on the space \mathcal{H}_B of states satisfying the bosonic QLSCs. However, the group-averaging formula (5.34) suggests that one can proceed in analogy with the bosonic case to define a new inner product. Thus, for two states $|\varphi_1^{(B)}\rangle, |\varphi_2^{(B)}\rangle \in \mathcal{H}_B$ we obtain two states satisfying the fermionic QLSCs as

$$|\varphi_I^{(BF)}\rangle = -Q_1 Q_2 Q_3 Q_4 |\varphi_I^{(B)}\rangle, \tag{5.35}$$

and define the new inner product by

$$\begin{aligned} \langle \varphi_1^{(BF)} | \varphi_2^{(BF)} \rangle_{\mathcal{H}_{BF}} &= -\langle \varphi_1^{(B)} | Q_1 Q_2 Q_3 Q_4 | \varphi_2^{(B)} \rangle_{\mathcal{H}_B} \\ &= \langle \varphi_1^{(B)} | \varphi_2^{(BF)} \rangle_{\mathcal{H}_B}. \end{aligned} \tag{5.36}$$

Equations (5.30)–(5.33) and (5.25) imply that, if

$$|\varphi_I^{(B)}\rangle = \kappa_I^{(1)} |1100\rangle + \kappa_I^{(2)} |0110\rangle + \kappa_I^{(3)} |1001\rangle + \kappa_I^{(4)} |0011\rangle, \tag{5.37}$$

then

$$|\varphi_I^{(BF)}\rangle = \frac{M^4}{16} \lambda_I |\varphi_P^{(B)}\rangle, \tag{5.38}$$

where

$$\lambda_I = \kappa_I^{(1)} + \kappa_I^{(2)} - \kappa_I^{(3)} + \kappa_I^{(4)}, \tag{5.39}$$

and

$$\langle \varphi_1^{(BF)} | \varphi_2^{(BF)} \rangle_{\mathcal{H}_{BF}} = \frac{M^4}{16} \lambda_1^* \lambda_2 \langle \chi^{(B)} | \chi^{(B)} \rangle. \tag{5.40}$$

Thus, our new inner product $\langle \cdot | \cdot \rangle_{\mathcal{H}_{BF}}$ is positive definite⁶. Although it is defined in terms of the ‘seed states’ $|\varphi_I^{(B)}\rangle$, which do not satisfy the fermionic QLSCs, the new inner product depends only on the invariant states $|\varphi_I^{(BF)}\rangle$.

We constructed the states satisfying the bosonic and fermionic QLSCs step by step. We first found the states satisfying the bosonic QLSCs and an inner product among them in the previous section, following [25]. Then, we found the states satisfying the fermionic QLSCs as well among these states and defined an inner product for these invariant states. Our construction can in fact be understood as group-averaging over the supergroup generated by P^μ and Q_α . Starting from a state $|\varphi_I\rangle$, $I = 1, 2$, in the original Hilbert space \mathcal{H}_0 , we define a state satisfying all constraints by integrating over the supergroup [37, 38]:

$$|\varphi_I^{(BF)}\rangle = -\frac{1}{2V} \int d^4\alpha, d^4\theta \exp(-i\alpha \cdot P - \bar{\theta}Q) |\varphi_I\rangle, \tag{5.41}$$

where $\theta = (\theta_1, \theta_2, \theta_3, \theta_4)$ are Grassmann numbers and $\alpha^\mu = (\alpha^0, \vec{\alpha})$ are commuting numbers. As P^μ and Q_α commute, we can write this integral as

$$\begin{aligned} |\varphi_I^{(BF)}\rangle &= -\frac{1}{2V} \int d^4\theta \exp(-\bar{\theta}Q) \int d^4\alpha \exp(-i\alpha \cdot P) |\varphi_I\rangle \\ &= -Q_1 Q_2 Q_3 Q_4 |\varphi_I^{(B)}\rangle, \end{aligned} \tag{5.42}$$

where $|\varphi_I^{(B)}\rangle$ satisfies only the bosonic QLSCs:

$$|\varphi_I^{(B)}\rangle = \frac{1}{2V} \int d^4\alpha \exp(-i\alpha \cdot P) |\varphi_I\rangle. \tag{5.43}$$

We have shown that all states satisfying both the bosonic and fermionic QLSCs can be obtained in this manner. The inner product $\langle \cdot | \cdot \rangle_{\mathcal{H}_{BF}}$ we have defined is

$$\begin{aligned} \langle \varphi_1^{(BF)} | \varphi_2^{(BF)} \rangle_{\mathcal{H}_{BF}} &= -\frac{1}{2V} \int d^4\alpha d^4\theta \langle \varphi_1 | \exp(-i\alpha \cdot P - \bar{\theta}Q) | \varphi_2 \rangle_{\mathcal{H}_0} \\ &= -\int d^4\theta \langle \varphi_1^{(B)} | \exp(-\bar{\theta}Q) | \varphi_2^{(B)} \rangle_{\mathcal{H}_B}. \end{aligned} \tag{5.44}$$

In appendix E we present an example of a state with two particles with non-zero momenta which satisfies all QLSCs.

⁶The definition of a positive-definite inner product used here is similar to that in appendix D of reference [36].

6. Summary and discussion

In this paper we pointed out that there are fermionic linearization stability conditions as well as bosonic ones in four-dimensional $\mathcal{N} = 1$ simple supergravity in the background of static three-torus space. Then we showed that states satisfying both fermionic and bosonic quantum linearization stability conditions (QLSCs) can be constructed by group-averaging over the supergroup of global supersymmetry and spacetime translation symmetry.

States satisfying the bosonic QLSCs have infinite norm in the original Hilbert space. This infinity results from the infinite volume of the symmetry group generated by the LSCs. Roughly speaking, this infinite volume is factored out in the group-averaging inner product. It is interesting that the inner product of states satisfying all QLSCs have zero norm in the Hilbert space of states satisfying only the bosonic ones. The finite group-averaging inner product is obtained by factoring out zero in this case.

As the bosonic QLSCs can be interpreted as a remnant of diffeomorphism invariance of the full generally covariant theory, one should be able to interpret the fermionic QLSCs as a remnant of full local supersymmetry in the context of canonical quantization [39, 40]. The bosonic QLSCs, $H|\text{phys}\rangle = 0$ and $\vec{P}|\text{phys}\rangle = 0$, have a natural physical picture. These conditions imply that all physical states are invariant under spacetime translations. This means that there is no meaning in the position and time coordinates of an event relative to the background spacetime of static three-torus. However, the physical states still encode relative positions and relative time differences between two or more events. Thus, the bosonic QLSCs can be seen as a manifestation of Mach's principle in quantum general relativity. (See e.g. [41] for a discussion of Mach's principle in general relativity.) On the other hand, it is not clear if there is a simple interpretation of the fermionic QLSCs. It would be interesting to find one.

It would also be interesting to investigate whether there are analogues of LSCs in string theory. A preliminary investigation in this direction [31] did not find such analogues in Bosonic string theory, but since string theory contains general relativity, we believe there should be analogues of LSCs in (super) string theory on any background spacetime with compact Cauchy surfaces.

The supergroup relevant to this work was a simple one with an abelian bosonic subgroup. It would be interesting to investigate the group-averaging procedure for general supergroups and establish general properties in analogy with the bosonic case studied in [24].

Acknowledgments

LS was supported by a PhD studentship from the Engineering and Physical Sciences Research Council (EPSRC) (EP/N509802/1).

Appendix A. Modified transformation of the vierbein in general spacetime

In this appendix we show that one can modify the infinitesimal diffeomorphism transformation on e_μ^a by an infinitesimal local Lorentz transformation so that the transform is proportional to $\nabla_\mu \zeta_\nu + \nabla_\nu \zeta_\mu$. If we transform e_μ^a by diffeomorphism in the direction of ζ^μ and by a local Lorentz transformation, we have

$$\delta^{(m)} e_\mu^a = \zeta^\rho \nabla_\rho e_\mu^a + e_\rho^a \nabla_\mu \zeta^\rho - s^{ab} e_{b\mu}, \quad (\text{A.1})$$

where s^{ab} is anti-symmetric and where $\nabla_\rho e_\mu^a = \partial_\rho e_\mu^a - \Gamma_{\rho\mu}^\sigma e_\sigma^a$ with the Levi-Civita connection $\Gamma_{\rho\mu}^\sigma$. Our task is to choose s^{ab} so that $\delta^{(m)} e_\mu^a$ is proportional to $\nabla_\mu \zeta_\nu + \nabla_\nu \zeta_\mu$. To this end, we require

$$e_{a[\mu} \delta^{(m)} e_{\nu]}^a = (\zeta^\rho \nabla_\rho e_{c[\nu]} e_{\mu]}^c + e_{c\rho} e_{[\mu}^c \nabla_{\nu]} \zeta^\rho - s_{ab} e_{[\mu}^a e_{\nu]}^b) = 0. \quad (\text{A.2})$$

Since

$$D_\rho e_\nu^c = \nabla_\rho e_\nu^c + \omega_\mu^{cb} e_{b\nu} = 0, \quad (\text{A.3})$$

we find

$$s_{ab} e_{[\mu}^a e_{\nu]}^b = -\zeta^\rho \omega_{\rho cb} e_{[\mu}^c e_{\nu]}^b + \nabla_{[\nu} \zeta_{\mu]}. \quad (\text{A.4})$$

Thus,

$$s^{ab} = -\zeta^\rho \omega_\rho^{ab} + \nabla_{[\nu} \zeta_{\mu]} e^{a\mu} e^{b\nu}. \quad (\text{A.5})$$

Then

$$\begin{aligned} \delta^{(m)} e_\mu^a &= \zeta^\rho \nabla_\rho e_\mu^a + e_\rho^a \nabla_\mu \zeta^\rho + \zeta^\rho \omega_\rho^{ab} + \frac{1}{2} (\nabla_\lambda \zeta_\nu - \nabla_\nu \zeta_\lambda) e^{a\lambda} e^{b\nu} e_{b\mu} \\ &= \zeta^\rho D_\rho e_\mu^a + e_\lambda^a \nabla_\mu \zeta^\lambda + \frac{1}{2} (\nabla_\lambda \zeta_\mu - \nabla_\mu \zeta_\lambda) e^{a\lambda} \\ &= \frac{1}{2} (\nabla_\lambda \zeta_\mu + \nabla_\mu \zeta_\lambda) e^{a\lambda}. \end{aligned} \quad (\text{A.6})$$

Appendix B. Linearization stability condition for electrodynamics

In this appendix we discuss the linearization stability condition for quantum electrodynamics in a static three-torus space. The Lagrangian density is

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \bar{\psi} \gamma^\mu (\partial_\mu - ieA_\mu) \psi - m \bar{\psi} \psi, \quad (\text{B.1})$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and $\bar{\psi} = i\psi^\dagger \gamma^0$. The field equations are

$$\frac{\delta S}{\delta A_\mu} = \partial_\nu F^{\nu\mu} - ie \bar{\psi} \gamma^\mu \psi = 0, \quad (\text{B.2})$$

$$\frac{\delta S}{\delta \psi^\dagger} = i\gamma^0 [\gamma^\mu (\partial_\mu - ieA_\mu) \psi - m\psi] = 0, \quad (\text{B.3})$$

$$\frac{\delta S}{\delta \psi} = (\partial_\mu + ieA_\mu) \bar{\psi} \gamma^\mu + m \bar{\psi} = 0, \quad (\text{B.4})$$

where S is the action obtained by integrating \mathcal{L} over spacetime. We have defined the left-hand side of (B.3) and (B.4) as the left functional derivative of the action S .

The field equations (B.3) and (B.4) imply that the current $J_{(E)}^\mu = ie \bar{\psi} \gamma^\mu \psi$ is conserved. As a result, the total charge, which is the integral of $J_{(E)}^0$ over the space,

$$Q_{(E)} = e \int d^3 \vec{x} \psi^\dagger \psi, \quad (\text{B.5})$$

is time independent. This charge must vanish because the zeroth component of (B.2) implies $\partial_i F^{i0} = e\psi^\dagger\psi$ and hence

$$Q_{(E)} = \int d^3\vec{x} \partial_i F^{i0}, \tag{B.6}$$

which must vanish by Gauss’s divergence theorem. (Recall our space is compact.) At linearized level, the fields A_μ and ψ are decoupled, and the condition $Q_{(E)} = 0$ does not automatically arise in the linearized theory. Hence this condition has to be imposed as a linearization stability condition on the linearized solution for it to extend to an exact solution. The aim of this appendix is to illustrate how the argument in section 3 works in this simple model.

The action is invariant under the gauge transformation $\delta A_\mu = \partial_\mu \Lambda$, $\delta \psi = ie\Lambda\psi$ for any function Λ of the spacetime point. The argument in section 3 leads to the conclusion that

$$\begin{aligned} \partial_\mu \Lambda (\partial_\nu F^{\nu\mu} - ie\bar{\psi}\gamma^\mu\psi) - ie\Lambda\bar{\psi} [\gamma^\mu(\partial_\mu - ieA_\mu)\psi - m\psi] - ie\Lambda \{ [(\partial_\mu + ieA_\mu)\bar{\psi}] \gamma^\mu + m\bar{\psi} \} \psi \\ = \partial_\mu J_{(\Lambda)}^\mu [A, \psi], \end{aligned} \tag{B.7}$$

for some current $J_{(\Lambda)}^\mu [A, \psi]$ for any Λ . We indeed find that this equation is satisfied with

$$J_{(\Lambda)}^\mu [A, \psi] = \Lambda (\partial_\nu F^{\nu\mu} - ie\bar{\psi}\gamma^\mu\psi). \tag{B.8}$$

The identity (B.7) is satisfied order by order in the fields A_μ and ψ , i.e.

$$\partial_\mu \Lambda \partial_\nu F^{\nu\mu} = \partial_\mu J_{(\Lambda)}^{(1)\mu} [A], \tag{B.9}$$

$$- ie\partial_\mu \Lambda \bar{\psi}\gamma^\mu\psi - ie\Lambda\bar{\psi}\gamma^\mu\partial_\mu\psi - ie\Lambda(\partial_\mu\bar{\psi})\gamma^\mu\psi = \partial_\mu J_{(\Lambda)}^{(2)\mu} [A, \psi], \tag{B.10}$$

where

$$J_{(\Lambda)}^{(1)\mu} [A] = \Lambda \partial_\nu F^{\nu\mu}, \tag{B.11}$$

$$J_{(\Lambda)}^{(2)\mu} [A, \psi] = -ie\Lambda\bar{\psi}\gamma^\mu\psi. \tag{B.12}$$

Now, if $\Lambda(x) = \lambda$ is constant, then the left-hand side of (B.9) vanishes. Hence the current $J_{(\lambda)}^{(1)\mu} [A] = \lambda\partial_\nu F^{\nu\mu}$ is conserved for any field configuration A_μ . In particular, we can smoothly deform A_μ to zero in the far future or past of any given Cauchy surface while keeping the value of the conserved charge

$$Q_{(E)}^{(1)} [A] = \int d^3\vec{x} \partial_i F^{i0}, \tag{B.13}$$

unchanged. It follows that $Q_{(E)}^{(1)} [A] = 0$ for any field A_μ . Of course, one can readily verify this fact by Gauss’s divergence theorem, as we observed before.

Now, consider solving the field equations order by order by letting

$$A_\mu = A_\mu^{(1)} + A_\mu^{(2)} + \dots, \tag{B.14}$$

$$\psi = \psi^{(1)} + \psi^{(2)} + \dots. \tag{B.15}$$

Then, with the definition $F_{\mu\nu}^{(I)} := \partial_\mu A_\nu^{(I)} - \partial_\nu A_\mu^{(I)}$, $I = 1, 2$, the field equations (B.2)–(B.4) become

$$\partial_\nu F^{(1)\nu\mu} = 0, \tag{B.16}$$

$$i\gamma^0 [\gamma^\mu \partial_\mu \psi^{(1)} - m\psi^{(1)}] = 0, \tag{B.17}$$

$$\partial_\mu \overline{\psi^{(1)}} \gamma^\mu + m\overline{\psi^{(1)}} = 0, \tag{B.18}$$

$$\partial_\nu F^{(2)\nu\mu} - ie\overline{\psi^{(1)}} \gamma^\mu \psi^{(1)} = 0. \tag{B.19}$$

Then, equations (B.9) and (B.10) imply that the current

$$J_{(\Lambda)}^{(1)\mu}[A^{(2)}] + J_{(\Lambda)}^{(2)\mu}[A^{(1)}, \psi^{(1)}] = \Lambda(\partial_\nu F^{(2)\nu\mu} - ie\overline{\psi^{(1)}} \gamma^\mu \psi^{(1)}), \tag{B.20}$$

is conserved for any $\Lambda(x)$. This implies that the corresponding charge must vanish, which is rather obvious in this example because, in fact, $J_{(\Lambda)}^{(1)\mu}[A^{(2)}] + J_{(\Lambda)}^{(2)\mu}[A^{(1)}, \psi^{(1)}] = 0$ by (B.19). In particular, if we define

$$Q_{(E)}^{(2)}[\psi^{(1)}] = -e \int d^3\vec{x} \psi^{(1)\dagger} \psi^{(1)}, \tag{B.21}$$

then, since $e\lambda\psi^{(1)\dagger}\psi^{(1)} = ie\lambda\overline{\psi^{(1)}}\gamma^0\psi^{(1)} = -J_{(\lambda)}^{(2)0}[A^{(1)}, \psi^{(1)}]$, we find

$$Q_{(E)}^{(2)}[\psi^{(1)}] = -Q_{(E)}^{(1)}[A^{(2)}] = 0, \tag{B.22}$$

which is the linearization stability condition for electrodynamics in static torus space. In this example, it is the linearization of the condition (B.6) in the exact theory.

Appendix C. Proof of identities (5.8) and (5.9)

We first note that the matrices $T_{ij}^B, B = 1, 2$, are diagonal with $T_{kk}^{Bi} = \delta_k^i q_k^B$, where $q_1^1 = -q_2^1 = 1/\sqrt{2}, q_3^1 = 0, q_1^2 = q_2^2 = 1/\sqrt{6}$ and $q_3^2 = -2/\sqrt{6}$. Hence

$$\begin{aligned} X &:= \sum_{B=1}^2 \sum_{A=1}^5 \sum_{A'=1}^5 T_{ij}^A T^{Bi}{}_k T_{i'j'}^{A'} T^{Bj'}{}_{k'} \gamma^j \gamma^k \gamma^{k'} \gamma^{j'} c_{PA} c_{PA'} \\ &= \sum_{k=1}^3 \sum_{k'=1}^3 \sum_{A=1}^5 \sum_{A'=1}^5 T_{kj}^A T_{k'j'}^{A'} \sum_{B=1}^2 q_k^B q_{k'}^B \gamma^j \gamma^k \gamma^{k'} \gamma^{j'} c_{PA} c_{PA'}. \end{aligned} \tag{C.1}$$

By a component-by-component calculation we find

$$\sum_{B=1}^2 q_k^B q_{k'}^B = \delta_{kk'} - \frac{1}{3}. \tag{C.2}$$

By substituting this formula into (C.1) we obtain

$$X = \sum_{A=1}^5 \sum_{A'=1}^5 T^{Ak}{}_j T_{k'j'}^{A'} \gamma^j \gamma^k \gamma^{k'} \gamma^{j'} c_{PA} c_{PA'} - \frac{1}{3} \sum_{A=1}^5 \sum_{A'=1}^5 T_{kj}^A T_{k'j'}^{A'} \gamma^j \gamma^k \gamma^{k'} \gamma^{j'} c_{PA} c_{PA'}. \tag{C.3}$$

The second sum vanishes because the matrices T_{ij}^A are symmetric and traceless so that $T_{kj}^A \gamma^j \gamma^k = T_{jk}^A (\delta^{jk} + \gamma^{jk}) = 0$. Since the first sum is of the form $S_{jj'} \gamma^j \gamma^{j'}$ where $S_{jj'}$ is symmetric, we may replace $\gamma^j \gamma^{j'}$ by $\delta^{jj'}$. Hence

$$X = \sum_{A=1}^5 \sum_{A'=1}^5 T^{Ak} T_{kj}^A \delta^{j'l} c_{PA} c_{PA'} = \sum_{A=1}^5 \delta^{AA'} c_{PA} c_{PA'} = \sum_{A=1}^5 (c_{PA})^2, \quad (\text{C.4})$$

which is equation (5.8).

To show (5.9) we note that, since $u^\pm(\vec{k})$ is a simultaneous eigenspinor of $\gamma^0 \hat{k} \cdot \vec{\gamma}$ and γ_5 with eigenvalues -1 and ± 1 , respectively, with the normalization $u^{\pm\dagger}(\vec{k})u^\pm(\vec{k}) = 2k$, the matrix $M_{\alpha\beta} = u^\pm(\vec{k})_\alpha u^{\pm*}(\vec{k})_\beta$ is the product of the projection operator onto the eigenspace of $\gamma^0 \hat{k} \cdot \vec{\gamma}$ with eigenvalue -1 and that of γ_5 with eigenvalue ± 1 (multiplied by $2k$) since these projection operators commute. Hence, we have

$$\begin{aligned} u^\pm(\vec{k})_\alpha u^{\pm*}(\vec{k})_\beta &= \frac{k}{2} \left[(1 - \gamma^0 \hat{k} \cdot \vec{\gamma})(1 \pm \gamma_5) \right]_{\alpha\beta} \\ &= \frac{1}{2} \left[(1 \pm \gamma_5) k \cdot \gamma \gamma^0 \right]_{\alpha\beta}, \end{aligned} \quad (\text{C.5})$$

where we have used $k \cdot \gamma = k(-\gamma^0 + \hat{k} \cdot \vec{\gamma})$ so that $k(1 - \gamma^0 \hat{k} \cdot \vec{\gamma}) = k \cdot \gamma \gamma^0$. Hence

$$\left[\epsilon^{\pm*}(\vec{k}) \cdot \vec{\gamma} u^\pm(\vec{k}) \right]_\alpha \left[\epsilon^\pm(\vec{k}) \cdot \vec{\gamma} u^{\pm*}(\vec{k}) \right]_\beta = \frac{1}{2} \left[(1 \mp \gamma_5) k \cdot \gamma (\epsilon^{\pm*}(\vec{k}) \cdot \vec{\gamma})(\epsilon^\pm(\vec{k}) \cdot \vec{\gamma}) \gamma^0 \right]_{\alpha\beta}, \quad (\text{C.6})$$

where we have used the fact that $k \cdot \gamma$, γ^0 and γ_5 all anti-commute with $\epsilon^\pm(\vec{k}) \cdot \vec{\gamma}$ and $\epsilon^{\pm*}(\vec{k}) \cdot \vec{\gamma}$. Now the identities $(\epsilon^{\pm*}(\vec{k}) \cdot \vec{\gamma})(\epsilon^\pm(\vec{k}) \cdot \vec{\gamma}) = 1 \pm \gamma^0 \hat{k} \cdot \vec{\gamma} \gamma_5$ and $k \cdot \gamma \gamma^0 \hat{k} \cdot \vec{\gamma} = k \cdot \gamma$ imply $k \cdot \gamma (\epsilon^{\pm*}(\vec{k}) \cdot \vec{\gamma})(\epsilon^\pm(\vec{k}) \cdot \vec{\gamma}) = k \cdot \gamma (1 \pm \gamma_5)$. By substituting this identity into (C.6) we find

$$\left[\epsilon^{\pm*}(\vec{k}) \cdot \vec{\gamma} u^\pm(\vec{k}) \right]_\alpha \left[\epsilon^\pm(\vec{k}) \cdot \vec{\gamma} u^{\pm*}(\vec{k}) \right]_\beta = \left[(1 \mp \gamma_5) k \cdot \gamma \gamma^0 \right]_{\alpha\beta}. \quad (\text{C.7})$$

Finally, by noting that $[(1 \mp \gamma_5) k \cdot \gamma \gamma^0]_{\beta\alpha} = [(1 \pm \gamma_5) k \cdot \gamma \gamma^0]_{\alpha\beta}$ we find (5.9).

Appendix D. Zero-momentum sector of the gravitino field

In this appendix we provide some details of the (indefinite-metric) Hilbert space describing the zero-momentum sector of the gravitino field. Some of the material presented here can be found in [33]. We make the discussion general and treat the fermionic algebra where the self-adjoint operators η_α , $\alpha = 1, 2, \dots, 2M$ and $\eta_\alpha^{(+)}$, $\alpha = 1, 2, \dots, 2N$ satisfying

$$[\eta_\alpha, \eta_\beta]_+ = -\delta_{\alpha\beta}, \quad [\eta_\alpha^{(+)}, \eta_\beta^{(+)}]_+ = \delta_{\alpha\beta}, \quad [\eta_\alpha, \eta_\beta^{(+)}]_+ = 0. \quad (\text{D.1})$$

The zero-momentum sector of the gravitino field corresponds to $M = 2$ and $N = 4$.

We represent this algebra as follows. We define annihilation-type operators as

$$d_a := \frac{1}{\sqrt{2}}(\eta_{2a-1} + i\eta_{2a}), \quad a = 1, 2, \dots, M, \quad (\text{D.2})$$

$$d_a^{(+)} := \frac{1}{\sqrt{2}}(\eta_{2a-1}^{(+)} + i\eta_{2a}^{(+)}), \quad a = 1, 2, \dots, N. \quad (\text{D.3})$$

These operators and their adjoint, the creation-type operators, have the following non-zero anti-commutators:

$$[d_a, d_b^\dagger]_+ = -\delta_{ab}, \quad [d_a^{(+)}, d_b^{(+)\dagger}]_+ = \delta_{ab}. \tag{D.4}$$

We define the state $|0_{0F}\rangle$ by requiring $d_a|0_{0F}\rangle = d_a^{(+)}|0_{0F}\rangle = 0$ for all a and $\langle 0_{0F}|0_{0F}\rangle = 1$. Then the 2^{M+N} -dimensional (indefinite-metric) Hilbert space representing the algebra (D.1) is spanned by the following orthogonal states:

$$|\{n\}, \{n^{(+)}\}\rangle = \prod_{a=1}^M (d_a^\dagger)^{n_a} \prod_{b=1}^N (d_b^{(+)\dagger})^{n_b} |0_{0F}\rangle, \tag{D.5}$$

where n_a and $n_b^{(+)}$ are either 0 or 1 and where $\{n\} = \{n_1, n_2, \dots, n_M\}$ and $\{n^{(+)}\} = \{n_1^{(+)}, n_2^{(+)}, \dots, n_N^{(+)}\}$. One readily finds

$$\langle \{n\}, \{n^{(+)}\} | \{n\}, \{n^{(+)}\} \rangle = \begin{cases} 1 & \text{if } n_1 + n_2 + \dots + n_M \text{ is even,} \\ -1 & \text{if } n_1 + n_2 + \dots + n_M \text{ is odd.} \end{cases} \tag{D.6}$$

If Ω is an anti-self-adjoint operator satisfying $\Omega^\dagger = -\Omega$, then the operator $\exp \Omega$, which can be defined as a power series, is unitary in the sense that it preserves the inner product. Of particular interest are the following unitary operators:

$$U_{\alpha\beta}(\theta) := \exp\left(\frac{\theta}{2}\eta_\alpha\eta_\beta\right) = \cos\frac{\theta}{2} + \eta_\alpha\eta_\beta \sin\frac{\theta}{2}, \quad \alpha \neq \beta, \tag{D.7}$$

$$V_{\alpha\beta}(\theta) := \exp\left(\frac{\theta}{2}\eta_\alpha\eta_\beta^{(+)}\right) = \cosh\frac{\theta}{2} + \eta_\alpha\eta_\beta^{(+)} \sinh\frac{\theta}{2}, \tag{D.8}$$

$$W_{\alpha\beta}(\theta) := \exp\left(\frac{\theta}{2}\eta_\alpha^{(+)}\eta_\beta^{(+)}\right) = \cos\frac{\theta}{2} + \eta_\alpha^{(+)}\eta_\beta^{(+)} \sin\frac{\theta}{2}, \quad \alpha \neq \beta. \tag{D.9}$$

These unitary operators act on η_α and $\eta_\alpha^{(+)}$ as follows:

$$U_{\alpha\beta}(\theta)\eta_\alpha U_{\alpha\beta}(\theta)^\dagger = \eta_\alpha \cos\theta + \eta_\beta \sin\theta, \tag{D.10}$$

$$W_{\alpha\beta}(\theta)\eta_\alpha^{(+)} W_{\alpha\beta}(\theta)^\dagger = \eta_\alpha^{(+)} \cos\theta - \eta_\beta^{(+)} \sin\theta, \tag{D.11}$$

$$V_{\alpha\beta}(\theta)\eta_\alpha V_{\alpha\beta}(\theta)^\dagger = \eta_\alpha \cosh\theta + \eta_\beta^{(+)} \sinh\theta, \tag{D.12}$$

$$V_{\alpha\beta}(\theta)\eta_\beta^{(+)} V_{\alpha\beta}(\theta)^\dagger = \eta_\beta^{(+)} \cosh\theta + \eta_\alpha \sinh\theta. \tag{D.13}$$

Let us define $Y := \prod_{\alpha=1}^{2M} \eta_\alpha \prod_{\beta=1}^{2N} \eta_\beta^{(+)}$. Then, Y is unitary, i.e. $Y^\dagger = Y^{-1}$. The operator $W_\alpha := Y\eta_\alpha^{(+)}$, $\alpha = 1, 2, \dots, 2N$, is also unitary. We find $W_\alpha\eta_\alpha^{(+)}W_\alpha^\dagger = -\eta_\alpha^{(+)}$ whereas $W_\alpha\eta_\beta^{(+)}W_\alpha^\dagger = \eta_\beta^{(+)}$ if $\beta \neq \alpha$ and $W_\alpha\eta_\beta W_\alpha^\dagger = \eta_\beta$ for all β .

Since any product of $U_{\alpha\beta}(\theta)$, $W_{\alpha\beta}(\theta)$, $V_\alpha(\theta)$ and W_α is unitary, we can conclude the following. Suppose that $M \leq N$ and let

$$\tilde{X}_\alpha = A_\alpha^\beta \eta_\beta \cosh\theta_\alpha - C_\alpha^\beta \eta_\beta^{(+)} \sinh\theta_\alpha, \tag{D.14}$$

where the index $\alpha = 1, 2, \dots, 2M$, is not summed over, and where (A_α^β) and (C_α^β) are a $2M \times 2M$ matrix and a $2M \times 2N$ matrix, respectively, satisfying $A_{\alpha_1}^{\beta_1} A_{\alpha_2}^{\beta_2} \delta_{\beta_1\beta_2} = C_{\alpha_1}^{\beta_1} C_{\alpha_2}^{\beta_2} \delta_{\beta_1\beta_2} = \delta_{\alpha_1\alpha_2}$ and $\det(A^\alpha_\beta) = 1$.⁷ Then there is a unitary operator U such that $\tilde{X}_\alpha = U\eta_\alpha U^\dagger$.

⁷ The unitarity of the operators W_α allows us to have $\det(C^\alpha_\beta) = -1$.

Appendix E. Example of a state satisfying all quantum linearization stability conditions

In this appendix we present a concrete example of a state satisfying all fermionic as well as bosonic QLSCs. We consider an example where there are two particles, one going in the positive x -direction and the other in the negative x -direction with momentum $k > 0$ and both with positive helicity. In this sector of the theory $M^2 = 4k$.

To find the non-zero-momentum contribution to the supercharge in this sector we need to find the spinors $u^+(\vec{k})$ with $\vec{k} = \pm k\hat{e}_1$, where \hat{e}_1 is the unit vector in the x -direction. It is an eigenspinor of $\gamma^0\gamma^1$ and $i\gamma^2\gamma^3$ with eigenvalue -1 for both and, hence, an eigenspinor of $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$ with eigenvalue 1 . It is normalized as $u^{+\dagger}(\pm k\hat{e}_1)u^+(\pm k\hat{e}_1) = 2k$. We find from (3.2)

$$\gamma^0\gamma^1 = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \quad i\gamma^2\gamma^3 = \begin{pmatrix} \sigma_2 & 0 \\ 0 & \sigma_2 \end{pmatrix}. \quad (\text{E.1})$$

Hence we can take

$$u^+(k\hat{e}_1) = \sqrt{\frac{k}{2}} \begin{pmatrix} 1 \\ -i \\ 1 \\ -i \end{pmatrix}, \quad u^{+*}(k\hat{e}_1) = \sqrt{\frac{k}{2}} \begin{pmatrix} 1 \\ i \\ 1 \\ i \end{pmatrix}. \quad (\text{E.2})$$

Now, $\{\hat{e}_2, \hat{e}_3, \hat{e}_1\}$ form a right handed basis so that we can choose

$$\epsilon^+(k\hat{e}_1) \cdot \vec{\gamma} = \frac{1}{\sqrt{2}}(\gamma^2 + i\gamma^3) = \frac{1}{\sqrt{2}}\gamma^2(1 + i\gamma^2\gamma^3). \quad (\text{E.3})$$

Since $u^{+*}(k\hat{e}_1)$ is an eigenspinor of $i\gamma^2\gamma^3$ with eigenvalue $+1$, we have

$$\epsilon^+(k\hat{e}_1) \cdot \vec{\gamma} u^{+*}(k\hat{e}_1) = \sqrt{k}\gamma^2 \begin{pmatrix} 1 \\ i \\ 1 \\ i \end{pmatrix} = \sqrt{k} \begin{pmatrix} i \\ 1 \\ i \\ 1 \end{pmatrix}, \quad (\text{E.4})$$

and

$$\epsilon^{+*}(k\hat{e}_1) \cdot \vec{\gamma} u^+(k\hat{e}_1) = \sqrt{k}\gamma^2 \begin{pmatrix} 1 \\ -i \\ 1 \\ -i \end{pmatrix} = \sqrt{k} \begin{pmatrix} -i \\ 1 \\ -i \\ 1 \end{pmatrix}. \quad (\text{E.5})$$

The spinor $u^+(-k\hat{e}_1)$ is an eigenspinor of $\gamma^0\gamma^1$ and $i\gamma^2\gamma^3$ with eigenvalue $+1$ for both. We can choose

$$\epsilon^+(-k\hat{e}_1) \cdot \vec{\gamma} = \frac{1}{\sqrt{2}}\gamma^2(1 - i\gamma^2\gamma^3). \quad (\text{E.6})$$

Proceeding similarly as above, we find

$$\epsilon^+(-k\hat{e}_1) \cdot \vec{\gamma} u^+(-k\hat{e}_1) = \sqrt{k}\gamma^2 \begin{pmatrix} 1 \\ i \\ -1 \\ -i \end{pmatrix} = \sqrt{k} \begin{pmatrix} -i \\ -1 \\ i \\ 1 \end{pmatrix}, \quad (\text{E.7})$$

and

$$\epsilon^{+*}(-k\hat{e}_1) \cdot \vec{\gamma} u^+(-k\hat{e}_1) = \sqrt{k} \gamma^2 \begin{pmatrix} 1 \\ -i \\ -1 \\ i \end{pmatrix} = \sqrt{k} \begin{pmatrix} i \\ -1 \\ -i \\ 1 \end{pmatrix}, \quad (\text{E.8})$$

Thus, the relevant part of the $\vec{k} \neq 0$ contribution to the supercharge

$$\hat{Q} = -\frac{i}{2} \sum_{\vec{k} \neq 0} \sum_{\lambda = \pm} \left[\epsilon^\lambda(\vec{k}) \cdot \gamma u^{\lambda*}(\vec{k}) a_\lambda(\vec{k}) b_\lambda^\dagger(\vec{k}) - \epsilon^{\lambda*}(\vec{k}) \cdot \gamma u^\lambda(\vec{k}) a_\lambda^\dagger(\vec{k}) b_\lambda(\vec{k}) \right], \quad (\text{E.9})$$

is, with the notation $a_{(\pm)} = a_+(\pm k\hat{e}_1)$ and $b_{(\pm)} = b_+(\pm k\hat{e}_1)$,

$$\hat{Q} = \frac{\sqrt{k}}{2} \begin{pmatrix} [a_{(+)} b_{(+)}^\dagger + a_{(+)}^\dagger b_{(+)}] - [a_{(-)} b_{(-)}^\dagger + a_{(-)}^\dagger b_{(-)}] \\ -i [a_{(+)} b_{(+)}^\dagger - a_{(+)}^\dagger b_{(+)}] + i [a_{(-)} b_{(-)}^\dagger - a_{(-)}^\dagger b_{(-)}] \\ [a_{(+)} b_{(+)}^\dagger + a_{(+)}^\dagger b_{(+)}] + [a_{(-)} b_{(-)}^\dagger + a_{(-)}^\dagger b_{(-)}] \\ -i [a_{(+)} b_{(+)}^\dagger - a_{(+)}^\dagger b_{(+)}] - i [a_{(-)} b_{(-)}^\dagger - a_{(-)}^\dagger b_{(-)}] \end{pmatrix}. \quad (\text{E.10})$$

Proceeding as in section 5 to construct a state satisfying the fermionic QLSCs we define

$$a_1 = \frac{\sqrt{2}}{M} (Q_1^{(0)} + iQ_2^{(0)}), \quad a_2 = \frac{\sqrt{2}}{M} (Q_3^{(0)} + iQ_4^{(0)}), \quad (\text{E.11})$$

as before and also

$$b_1 = \frac{1}{\sqrt{2k}} (\hat{Q}_1 + i\hat{Q}_2) = \frac{1}{\sqrt{2}} (a_{(+)} b_{(+)}^\dagger - a_{(-)} b_{(-)}^\dagger), \quad (\text{E.12})$$

$$b_2 = \frac{1}{\sqrt{2k}} (\hat{Q}_3 + i\hat{Q}_4) = \frac{1}{\sqrt{2}} (a_{(+)} b_{(+)}^\dagger + a_{(-)} b_{(-)}^\dagger), \quad (\text{E.13})$$

in the relevant subspace. Then in the two-particle sector, $M^2 = 4k$, it is easy to see that

$$[a_i, a_j^\dagger]_+ = -\delta_{ij}, \quad [b_i, b_j^\dagger]_+ = \delta_{ij}. \quad (\text{E.14})$$

We define a state $|\varphi^{(B)}\rangle$ obeying the bosonic QLSCs as follows. We take the bosonic zero-momentum sector wave function to be given by the Gaussian

$$\Psi(c, \vec{c}) = N e^{-c^2} e^{-\vec{c}^2}, \quad (\text{E.15})$$

where N is a constant, and the non-zero-momentum part of the state contains two positive helicity gravitons with momentum $\pm k\hat{e}_1$. Then the state obeying the bosonic QLSCs by group-averaging is given by (4.24), which reads

$$|\varphi^{(B)}\rangle = \int \frac{d^5 \vec{p}}{(2\pi)^5} [f^{(+)}(\vec{p}) e^{-iE(\vec{p})c + i\vec{p}\cdot\vec{c}} + f^{(-)}(\vec{p}) e^{iE(\vec{p})c + i\vec{p}\cdot\vec{c}}] \otimes (a_{(+)}^\dagger a_{(-)}^\dagger |0\rangle), \quad (\text{E.16})$$

where the state $|0\rangle$ is the Fock vacuum in the non-zero-momentum sector, $E(\vec{p}) = \sqrt{\vec{p}^2 + M^2}$ and

$$f^{(\pm)}(\vec{p}) = \frac{\pi^3 N e^{-M^2/4} e^{-\vec{p}^2/2}}{2\sqrt{\vec{p}^2 + M^2}}. \quad (\text{E.17})$$

The norm of this state in the Hilbert space \mathcal{H}_B is then given by (4.26), which reads

$$\langle \varphi^{(B)} | \varphi^{(B)} \rangle_{\mathcal{H}_B} = \frac{\pi^3}{12} |N|^2 e^{-M^2/2} \int_0^\infty dp \frac{p^4 e^{-p^2}}{\sqrt{p^2 + M^2}} < \infty. \quad (\text{E.18})$$

We choose the normalisation constant N so that $\langle \varphi^{(B)} | \varphi^{(B)} \rangle_{\mathcal{H}_B} = 1$.

Now we need to consider the fermionic zero-momentum sector. As we stated in section 5, we need to consider the tensor product of \mathcal{H}_B and the 64-dimensional Hilbert space \mathcal{H}_{0F} describing the fermionic zero-momentum sector. We call this tensor product \mathcal{H}_B as in section 5. As in appendix D let $|0_{0F}\rangle$ be the normalized state annihilated by $d_1 = (\eta_1 + i\eta_2)/\sqrt{2}$, $d_2 = (\eta_3 + i\eta_4)/\sqrt{2}$, $d_1^B = (\eta_1^B + i\eta_2^B)/\sqrt{2}$ and $d_2^B = (\eta_3^B + i\eta_4^B)/\sqrt{2}$, $B = 1, 2$. Let \bar{a}_1 and \bar{a}_2 be the operators a_1 and a_2 , respectively, restricted to the eigenspace of c_P and c_{PA} with eigenvalues p_0 and p_A , respectively. Define $|p_0, \vec{p}\rangle \in \mathcal{H}_{0F}$ by $|p_0, \vec{p}\rangle := C_{p_0, \vec{p}} \bar{a}_1 \bar{a}_2 d_2^\dagger d_1^\dagger |0_{0F}\rangle$, where $C_{p_0, \vec{p}}$ is a normalization constant such that $\langle p_0, \vec{p} | p_0, \vec{p} \rangle_{\mathcal{H}_{0F}} = 1$. Then, $\bar{a}_1 |p_0, \vec{p}\rangle = \bar{a}_2 |p_0, \vec{p}\rangle = 0$ because $\bar{a}_1^2 = \bar{a}_2^2 = 0$ and $\bar{a}_1 \bar{a}_2 = -\bar{a}_2 \bar{a}_1$. Now, we let

$$\begin{aligned} |\varphi^{(B)}\rangle &= \int \frac{d^5 \vec{p}}{(2\pi)^5} [f^{(+)}(\vec{p}) e^{-iE(\vec{p})c + i\vec{p}\cdot\vec{c}} \otimes |E(\vec{p}), \vec{p}\rangle + f^{(-)}(\vec{p}) e^{iE(\vec{p})c + i\vec{p}\cdot\vec{c}} \otimes |-E(\vec{p}), \vec{p}\rangle] \\ &\quad \otimes (a_{(+)}^\dagger a_{(-)}^\dagger |0\rangle). \end{aligned} \quad (\text{E.19})$$

Then, $a_1 |\varphi^{(B)}\rangle = a_2 |\varphi^{(B)}\rangle = 0$ and $\langle \varphi^{(B)} | \varphi^{(B)} \rangle_{\mathcal{H}_B} = 1$. Notice also that because $|\varphi^{(B)}\rangle$ contains only gravitons, it obeys $b_1^\dagger |\varphi^{(B)}\rangle = b_2^\dagger |\varphi^{(B)}\rangle = 0$.

In the notation of section 5 we can thus take $|\varphi^{(B)}\rangle \propto |0011\rangle$. Indeed, by the group-averaging procedure described in section 5, we obtain a state $|\varphi^{(BF)}\rangle \in \mathcal{H}_{BF}$ obeying the bosonic and fermionic QLSCs by applying $-Q_1 Q_2 Q_3 Q_4$. That is,

$$|\varphi^{(BF)}\rangle = -Q_1 Q_2 Q_3 Q_4 |\varphi^{(B)}\rangle \quad (\text{E.20})$$

$$= \frac{1}{4} (Q_1 - iQ_2)(Q_1 + iQ_2)(Q_3 - iQ_4)(Q_3 + iQ_4) |\varphi^{(B)}\rangle \quad (\text{E.21})$$

$$= \frac{M^4}{16} (a_1^\dagger + b_1^\dagger)(a_1 + b_1)(a_2^\dagger + b_2^\dagger)(a_2 + b_2) |\varphi^{(B)}\rangle. \quad (\text{E.22})$$

By writing $|\varphi^{(B)}\rangle = a_{(+)}^\dagger a_{(-)}^\dagger |\Psi^{(B)}\rangle$ this can be evaluated as

$$\begin{aligned} |\varphi^{(BF)}\rangle &= \frac{M^4}{16} \left[a_{(+)}^\dagger a_{(-)}^\dagger |\Psi^{(B)}\rangle - \frac{1}{\sqrt{2}} (a_1^\dagger - a_2^\dagger) a_{(+)}^\dagger b_{(-)}^\dagger |\Psi^{(B)}\rangle \right. \\ &\quad \left. + \frac{1}{\sqrt{2}} (a_1^\dagger + a_2^\dagger) b_{(+)}^\dagger a_{(-)}^\dagger |\Psi^{(B)}\rangle - a_1^\dagger a_2^\dagger b_{(+)}^\dagger b_{(-)}^\dagger |\Psi^{(B)}\rangle \right]. \end{aligned} \quad (\text{E.23})$$

In particular, by the group-averaging procedure, the norm of this state is given by

$$\langle \varphi^{(BF)} | \varphi^{(BF)} \rangle_{\mathcal{H}_{BF}} = \langle \varphi^{(B)} | \varphi^{(B)} \rangle_{\mathcal{H}_B} = \frac{M^4}{16}, \quad (\text{E.24})$$

as would be expected by the procedure presented in section 5.

ORCID iDs

Atsushi Higuchi  <https://orcid.org/0000-0002-3703-7021>

Lasse Schmieding  <https://orcid.org/0000-0002-5198-1342>

References

- [1] Altas E 2018 Linearization instability in gravity theories (arXiv:1808.04722)
- [2] Brill D R and Deser S 1973 Instability of closed spaces in general relativity *Commun. Math. Phys.* **32** 291–304
- [3] Higuchi A 1991 Quantum linearization instabilities of de sitter spacetime I *Class. Quantum Grav.* **8** 1961
- [4] Brill D 1972 Isolated solutions in general relativity *Gravitation: Problems and Prospects* (Kiev: Naukova Dumka) p 17
- [5] Fischer A E and Marsden J E 1973 Linearization stability of the Einstein equations *Bull. Am. Math. Soc.* **79** 997–1003
- [6] Moncrief V 1975 Spacetime symmetries and linearization stability of the Einstein equations I *J. Math. Phys.* **16** 493–8
- [7] Moncrief V 1976 Space–time symmetries and linearization stability of the Einstein equations II *J. Math. Phys.* **17** 1893–902
- [8] Fischer A E, Marsden J E and Moncrief V 1980 The structure of the space of solutions of einstein’s equations I. One Killing field *Ann. Inst. Henri Poincaré. A* **33** 147–94
- [9] Arms J M, Marsden J E and Moncrief V 1981 Symmetry and bifurcations of momentum mappings *Commun. Math. Phys.* **78** 455–78
- [10] Arms J M, Marsden J E and Moncrief V 1982 The structure of the space of solutions of Einstein’s equations II. Several Killing fields and the Einstein–Yang–Mills equations *Ann. Phys.* **144** 81–106
- [11] Moncrief V 1978 Invariant states and quantized gravitational perturbations *Phys. Rev. D* **18** 983
- [12] Moncrief V 1979 Quantum linearization instabilities *Gen. Relativ. Gravit.* **10** 93–7
- [13] Higuchi A 1991 Quantum linearization instabilities of de Sitter spacetime II *Class. Quantum Grav.* **8** 1983
- [14] Ashtekar A, Lewandowski J, Marolf D, Mourao J and Thiemann T 1995 Quantization of diffeomorphism invariant theories of connections with local degrees of freedom *J. Math. Phys.* **36** 6456–93
- [15] Landsman N P 1995 Rieffel induction as generalized quantum Marsden–Weinstein reduction *J. Geom. Phys.* **15** 285–319
- [16] Marolf D and Morrison I A 2009 Group averaging for de sitter free fields *Class. Quantum Grav.* **26** 235003
- [17] Marolf D and Morrison I A 2009 Group averaging of massless scalar fields in $1+1$ de Sitter *Class. Quantum Grav.* **26** 035001
- [18] Losic B and Unruh W G 2006 On leading order gravitational backreactions in de Sitter spacetime *Phys. Rev. D* **74** 023511
- [19] Losic B 2007 Dynamics of backreactions around pure de Sitter spacetime *J. Phys. A* **40** 6951–6
- [20] Louko J and Rovelli C 2000 Refined algebraic quantization in the oscillator representation of $SL(2, \mathbb{R})$ *J. Math. Phys.* **41** 132–55
- [21] Louko J and Molgado A 2005 Superselection sectors in the Ashtekar–Horowitz–Boulware model *Class. Quantum Grav.* **22** 4007–20
- [22] Louko J 2006 Group averaging, positive definiteness and superselection sectors *J. Phys.: Conf. Ser.* **33** 142–9
- [23] Gomberoff A and Marolf D 1999 On group averaging for $SO(n, 1)$ *Int. J. Mod. Phys. D* **8** 519–35
- [24] Giulini D and Marolf D 1999 A uniqueness theorem for constraint quantization *Class. Quantum Grav.* **16** 2489–505
- [25] Higuchi A 1991 Linearized quantum gravity in flat space with toroidal topology *Class. Quantum Grav.* **8** 2023
- [26] Deser S and Zumino B 1976 Consistent supergravity *Phys. Lett. B* **62** 335

- [27] Freedman D Z, van Nieuwenhuizen P and Ferrara S 1976 Progress toward a theory of supergravity *Phys. Rev. D* **13** 3214–8
- [28] Hull C M 1983 The positivity of gravitational energy and global supersymmetry *Commun. Math. Phys.* **90** 545–61
- [29] Freedman D Z and Van Proeyen A 2012 *Supergravity* (Cambridge: Cambridge University Press)
- [30] Van Nieuwenhuizen P 1981 Supergravity *Phys. Rep.* **68** 189–398
- [31] Higuchi A 2001 Possible constraints on string theory in closed space with symmetries *Noncommutative Structures in Mathematics and Physics* NATO Science Series (*Series II: Mathematics, Physics and Chemistry*) vol 22, ed S Duplij, J Wess (Berlin: Springer) pp 465–73
- [32] Dirac P A M 1964 *Lectures on Quantum Mechanics* (New York: Academic)
- [33] Henneaux M and Teitelboim C 1994 *Quantization of Gauge Systems* (Princeton, NJ: Princeton University Press)
- [34] Marolf D 1995 Quantum observables and recollapsing dynamics *Class. Quantum Grav.* **12** 1199–220
- [35] Kuchar K V 2011 Time and interpretations of quantum gravity *Int. J. Mod. Phys.D* **20** 3–86
- [36] Gibbons J and Higuchi A 2015 Removing the Faddeev–Popov zero modes from Yang–Mills theory in spacetimes with compact spatial sections *Phys. Rev. D* **91** 024006
- [37] Fayet P and Ferrara S 1977 Supersymmetry. *Phys. Rep.* **32** 249–334
- [38] A R 2007 *Supermanifolds: Theory and Applications* (Singapore: World Scientific)
- [39] D’Eath P D 1984 The canonical quantization of supergravity *Phys. Rev. D* **29** 2199
D’Eath P D 1985 *Phys. Rev. D* **32** 1593 erratum
- [40] Carroll S M, Freedman D Z, Ortiz M E and Page D N 1994 Physical states in canonically quantized supergravity *Nucl. Phys. B* **423** 661–87
- [41] Barbour J B and Bertotti B 1982 Mach’s principle and the structure of dynamical theories *Proc. R. Soc. A* **382** 295–306