

This is a repository copy of *JNK signalling regulates antioxidant responses in neurons*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/164368/>

Version: Published Version

Article:

Ugbode, Christopher orcid.org/0000-0002-6023-8294, Garnham, Nathan Arron, Fort Aznar, Laura et al. (3 more authors) (2020) JNK signalling regulates antioxidant responses in neurons. *Redox Biology*. 101712. ISSN 2213-2317

<https://doi.org/10.1016/j.redox.2020.101712>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Research Paper

JNK signalling regulates antioxidant responses in neurons

Chris Ugboade^{a,b}, Nathan Garnham^{a,b}, Laura Fort-Aznar^{a,b}, Gareth J.O. Evans^{a,b},
Sangeeta Chawla^{a,b,1,**}, Sean T. Sweeney^{a,b,1,*}

^a Department of Biology, University of York, York, YO10 5DD, UK

^b York Biomedical Research Institute, University of York, York, YO10 5DD, UK

ARTICLE INFO

Keywords:

Drosophila
Glutathione
Hydrogen Peroxide
ROS

ABSTRACT

Reactive oxygen species (ROS) are generated during physiological bouts of synaptic activity and as a consequence of pathological conditions in the central nervous system. How neurons respond to and distinguish between ROS in these different contexts is currently unknown. In *Drosophila* mutants with enhanced JNK activity, lower levels of ROS are observed and these animals are resistant to both changes in ROS and changes in synapse morphology induced by oxidative stress. In wild type flies, disrupting JNK-AP-1 signalling perturbs redox homeostasis suggesting JNK activity positively regulates neuronal antioxidant defense. We validated this hypothesis in mammalian neurons, finding that JNK activity regulates the expression of the antioxidant gene *Srxn-1*, in a c-Jun dependent manner. We describe a conserved 'adaptive' role for neuronal JNK in the maintenance of redox homeostasis that is relevant to several neurodegenerative diseases.

1. Introduction

Active neurons generate reactive oxygen species (ROS) predominantly as a by-product of mitochondrial respiration. ROS levels are neutralized by constitutive and adaptive reductive mechanisms operating in neurons and glia, including the glutathione system [1–3]. In this manner, the amplitude and temporal dynamics of the ROS signal are controlled, damage is limited and transient ROS signals can be interpreted in part to support the growth and plasticity of neurons [4,5]. A physiological level of ROS has been demonstrated to regulate a range of nervous system processes, including neuronal development, synaptic plasticity, and neural circuit tuning [5–7].

In many neurodegenerative disorders the reductive capacity of neurons is overwhelmed, contributing to disease progression [4,8]. These damaging levels of ROS, termed oxidative stress, overwhelm neuronal antioxidant defenses. A central component of the neuronal response to ROS is the activation of c-Jun N-terminal kinase, JNK. ROS have been shown to activate JNK-AP-1 signalling [9] which regulates neuronal growth and plasticity [10–13]. In *Drosophila* models, excessive ROS driving increases in synaptic growth are seen in both activity generated excitotoxicity and lysosomal storage disease [5,12,14]. Similar changes in synaptic structure can be induced directly by

application of oxidants such as paraquat [12] or diethylmaleate (DEM) [5], or through genetic activation of JNK via manipulation of the JNKK *hemipterous* [15] (*hep*) or the JNKKK *Wallenda* [11] (*wnd*), upstream activators of JNK.

Whether JNK activation by ROS in neurons is protective or degenerative is unknown. Recent evidence indicates JNK-AP-1 activity can prevent degeneration of injured axons [16] while JNK inhibition prevents neuronal loss after injury by promoting AKT signalling [17].

We recently found that knockdown of enzymes involved in antioxidant defense, superoxide dismutase and catalase, reshapes neuronal morphology at the *Drosophila* larval neuromuscular junction (NMJ) [5]. Similarly, mis-expressing antioxidant enzymes to manipulate levels of individual ROS species has profound effects on synaptic and dendritic growth and arborisation [5,12]. This suggests that plasticity at the synapse is finely tuned to the level of ROS. Given that JNK is known to regulate synaptic plasticity in response to ROS and that knockdown of antioxidant genes can change synapse structure, we hypothesized that JNK-induced structural changes in neurons are mediated in part by an antioxidant response.


To understand if JNK activity regulates antioxidant responses in neurons, we monitored ROS levels in *Drosophila* with mutations in *highwire* (*hiw*) and *puckered* (*puc*^{E69}). In *hiw* mutants, unrestrained

* Corresponding author. Department of Biology, University of York, York, YO10 5DD, UK.

** Corresponding author. Department of Biology, University of York, York, YO10 5DD, UK.

E-mail addresses: sangeeta.chawla@york.ac.uk (S. Chawla), sean.sweeney@york.ac.uk (S.T. Sweeney).

¹ These authors contributed equally to this work.



(caption on next page)

Fig. 1. JNK activity regulates neuronal antioxidant responses in *Drosophila*. A DEM induces oxidative stress in wild type flies but not *puckered* (*puc*^{E69/+}) or *highwire* (*hiw*) mutants. Quantification of hydrogen peroxide levels (Amplex red fluorescence) in wild type, *puc*^{E69/+} or *hiw* mutant flies administered vehicle (0.16% ethanol) or DEM (5 mM). B Quantification of hydrogen peroxide levels in *hiw* mutants and *hiw* mutants pan-neuronally expressing dominant negative *jnk* (*hiw*; *nSyb* > *bsk*^{R53K}), *wnd* (*hiw*; *nSyb* > *wnd*^{DN}), *ask1* (*hiw*; *nSyb* > *ask1*^{DN}), *jun* (*hiw*; *nSyb* > *jun*^{DN}) and *fos* (*hiw*; *nSyb* > *fos*^{DN}). C Quantification of hydrogen peroxide levels in wild type flies, pan-neuronally expressing dominant negative *jnk* (*nSyb* > *bsk*^{R53K}), *wnd* (*nSyb* > *wnd*^{DN}) and *ask1* (*nSyb* > *ask1*^{DN}). A-C Graphs show Amplex red fluorescence values normalized to the average signal of wild type flies. Data are plotted as mean ± SEM (**p < 0.01; ***p < 0.001; one-way ANOVA; minimum 15 flies per genotype). D Representative micrographs showing synaptic overgrowth at the *Drosophila* third instar larval NMJ (Muscle 6/7, hemi-segment A3) in wild type, *puc*^{E69/+} mutants and wild type flies expressing dominant negative *fos* (*fos*^{DN}), *jun* (*jun*^{DN}) and *ask1* (*ask1*^{DN}), reared on food containing ethanol or DEM. Scale bar = 30 μm. E Quantification of mean normalized bouton number from (D) genotypes. Data are plotted as mean ± SEM (***p < 0.001; one-way ANOVA; minimum 15 NMJ's analysed per genotype). F Schematic showing the contribution of JNK to neuronal antioxidant stress responses in *Drosophila*. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

wallenda (*wnd*) activity activates JNK and downstream AP-1 signalling to generate exuberant synaptic overgrowth [11,18]. *Puc* encodes a dual specificity phosphatase, transcriptionally activated by AP-1 that counteracts JNK activity [19]. In both the *hiw* and *puc*^{E69/+} mutant backgrounds we identified enhanced antioxidant defenses and low levels of ROS. In *hiw*, this enhanced antioxidant defense can be reversed in neurons by down-regulating JNK-AP-1 signalling. Conversely, both *puc*^{E69/+} and *hiw* flies are resistant to chemically induced oxidative stress, while inhibition of JNK-AP-1 signalling increases ROS levels in wild type neurons. We show that in mammalian neurons exposed to oxidative stress, the JNK-AP-1 pathway coordinates the transcription and translation of the antioxidant gene *Sulfiredoxin-1*, which was localised predominantly in synaptic compartments. Our data indicates that activation of JNK in neurons drives antioxidant responses to shape neuronal morphology.

2. Results

2.1. JNK activity drives antioxidant responses in *Drosophila*

To quantify levels of ROS in *Drosophila*, we used the Amplex red assay for detection of hydrogen peroxide [20]. We have previously found that food containing 5 mM and 10 mM DEM which depletes glutathione levels, is sufficient to induce an overgrowth at the larval NMJ [5], with 10 mM concentrations being detrimental to survival (Fig. S1, 6% of flies pupate). To understand whether DEM influences ROS levels in flies, we raised flies on food containing ethanol (vehicle) or 5 mM DEM. Using Amplex red we found that wild type flies have an increased ROS burden when raised on food containing 5 mM DEM (Fig. 1A). To investigate how JNK activity influences ROS levels under these conditions, we reared *puc*^{E69/+} and *hiw* mutants on the same food and found that both genotypes have significantly lower levels of ROS than wild type flies in vehicle-containing food. Moreover, both genotypes failed to show an increase in ROS when reared on DEM-containing food (Fig. 1A) suggesting that they are resistant to the oxidative stress generated by DEM.

Given that both *puc*^{E69/+} and *hiw* mutants have lower levels of ROS than wild type flies and are resistant to chemically-induced oxidative stress, we hypothesized that if JNK activity could be reduced in neurons of these flies, then ROS levels would increase back to wild type levels. To test this, we used *hiw* mutants and expressed dominant negative transgenes of *jnk*, *wnd*, *ask1*, *jun* and *fos* pan-neuronally using *nSyb*-Gal4 (Fig. 1B). We found that pan-neuronal expression of these dominant negative transgenes increased ROS levels of *hiw* mutants back to wild type controls, except for *ask1*^{DN}. This data indicates that manipulation of JNK, its upstream regulators and downstream targets, regulates ROS levels in neurons.

To further evaluate the relationship between JNK activity and ROS levels, we investigated whether expression of dominant negative *jnk*, *ask1* and *wnd* affects ROS levels in wild type flies. We found that expression of all dominant negative transgenes, pan-neuronally, significantly increases the ROS levels of wild type flies (Fig. 1C). Indeed, the levels observed were similar to those observed when wild type flies are reared on food containing DEM.

The elaborate synaptic morphology of *hiw* mutant flies has been extensively described [11] and is a consequence of activation of the JNK-AP-1 pathway. We therefore chose to evaluate the synaptic morphology of *puc*^{E69/+} neurons by analyzing the NMJ at Muscle 6/7, hemi-segment A3. We first monitored the effect of DEM in wild type larvae, finding that wild type larvae fed with DEM develop an overgrown NMJ, compared to larvae reared on normal food (Fig. 1D). Similar to *hiw* mutants, we found that *puc*^{E69/+} larvae have overgrown NMJ synapses (Fig. 1D). Feeding of DEM has no effect on the NMJ of *puc*^{E69/+} larvae, indicating that JNK activity alone is sufficient to change synaptic morphology, which correlates with an antioxidant response (decrease in ROS levels from Fig. 1A). Furthermore, the DEM-induced changes in synaptic morphology in wild type flies, can be rescued when dominant negative transgenes of *jun*, *fos* and *ask1* are expressed pre- and post-synaptically using *spin*GAL4 (Fig. 1D, E). Our observations in *Drosophila* show that JNK activity regulates synaptic morphology but also antioxidant responses. In *puc*^{E69/+} and *hiw* flies, JNK-AP-1 signalling generates an antioxidant response which lowers the total levels of ROS in flies and protects them from induced oxidative stress while also causing an increase in NMJ synapse size. In wild type flies, inducing oxidative stress changes synaptic morphology, generates a stress response and these responses can be regulated by JNK (Fig. 1F).

2.2. DEM induces oxidative stress in primary mammalian neurons

JNK activity in mammalian neurons is often associated with apoptosis and activated JNK is commonly used as a hallmark of cell death in neurodegenerative disease. Given that JNK-AP-1 signalling in the fly generates an antioxidant response and decreasing JNK activity in fly neurons increases ROS levels, we hypothesized that antioxidant responses in mammalian neurons would also be regulated by JNK-AP-1. Having previously validated the effects of DEM in the fly, we established a mammalian model of oxidative stress, using DEM to deplete cellular glutathione. We used this model to assay antioxidant responses after DEM treatment. We first characterized the effect of DEM on cellular GSH levels and ROS production in mature primary rat neurons, using a glutathione assay measuring total glutathione (oxidized and reduced forms). Fig. S2A shows that application of DEM to primary cortical neuronal cultures caused a concentration and time-dependent decrease in cellular glutathione. A 1 h and 4 h treatment of DEM at concentrations of 10 μM and 100 μM caused a rapid decrease in GSH (Fig. S2A). Lower concentrations of 1 μM DEM reduced cellular GSH content with less severity. Given the extent of GSH depletion by DEM, we investigated whether DEM induces mitochondrial toxicity by monitoring WST-1 absorbance (Water Soluble Tetrazolium Salts) as a measure of mitochondrial stress. Compared to vehicle treated neurons, DEM induced a small but significant decrease in mitochondrial capacity, however this was much less than treatment with equivalent concentrations of paraquat (PQ) and rotenone (RT) (Fig. S2B). To determine if DEM induces a ROS burden as it does in our fly model we measured H₂O₂ levels in neurons treated with DEM using Amplex red reagent. The Amplex red assay revealed that DEM (100 μM) significantly increased in H₂O₂ levels (Fig. S2C), after 24 h compared to ethanol treated controls (Fig. S2C). DEM-induced oxidative stress is significantly attenuated when neurons




Fig. 2. ROS induced JNK phosphorylation triggers neuronal antioxidant responses. **A** Representative Western blot of neurons treated with Bicuculline (50 μ M, Bic) and 4-Aminopyridine (500 μ M, 4AP) to increase synaptic activity (Bic/4AP) or DEM (100 μ M) for 1 h, to induce oxidative stress. JNK activity was inhibited by a 1 h pre-treatment with SU-3327 (SU, 700 nM). **B** Quantification of total phospho-JNK/pan-JNK ratios (* $p < 0.05$; ** $p < 0.01$; one-way ANOVA; $n = 3$ biological replicates). **C, D** The mRNA levels of *Srxn-1* and *c-Fos* were determined using quantitative PCR (qPCR) and were normalized to *Gapdh* mRNA levels. mRNA levels are expressed relative to untreated controls and shown as means \pm SEM (* $p < 0.05$; ** $p < 0.01$; unpaired *t*-test; $n = 4$ biological replicates) **C** *Srxn-1* mRNA is induced by oxidative stress in a JNK dependent manner. **D** JNK signalling downstream of oxidative stress does not require *c-Fos*. **E** JNK regulates SRXN-1 expression under oxidative stress but not during synaptic activity. Representative Western blot probed for SRXN-1, Pan-JNK, phospho-JNK, Pan-ERK, phospho-ERK and GAPDH antibodies. Neurons were treated with Bic/4AP (50/500 μ M) or DEM (100 μ M) in the presence or absence of either the JNK inhibitor SU 3327 (700 nM) or the NMDA receptor blocker MK801 (10 μ M) for 30 min (lanes 2/4/6) or 4 h (lanes 3/5/7/8/9).

are pre-incubated with catalase. The duration and concentrations of DEM that we employed deplete glutathione and induce an increase in ROS with minimal toxicity to mitochondrial function.

To model chronic oxidative stress, we treated neurons for longer time points (48 h) with DEM (100 μ M) and found that DEM causes a retraction of the dendritic arbor. This effect is rescued when neurons are pre-incubated with Catalase (100 nM, Fig. S2D) and when the components that produce glutathione, the catalytic and modifying subunits of glutamate cysteine ligase (GCL), are transduced into neurons (Fig. S3).

2.3. Oxidative stress activates JNK-dependent SRXN-1 expression, in primary mammalian neurons

JNK enzymes are key mediators of cellular responses to stress in neurons [21], including oxidative stress [22,23], and alterations in JNK activity have previously been linked to changes in dendritic complexity [24]. Phosphorylation of threonine 183 and tyrosine 185 increase JNK activity [21]. To characterize the effectiveness of both DEM and the JNK inhibitor SU 3327 on regulating JNK phosphorylation and activation, we treated primary neurons with DEM (100 μ M, 1 h) alone or in the presence of SU 3327 (700 nM, 1 h pre-treatment) and assayed JNK phosphorylation (Fig. 2A). We found a significant increase in phospho-JNK/total JNK ratios when neurons were treated with DEM, which was significantly attenuated when JNK activity was blocked with SU 3327 (Fig. 2 A,B).

Bicuculline, a competitive antagonist of GABA_A (γ -aminobutyric acid) receptors, and 4-aminopyridine, a voltage activated potassium

channel blocker, are commonly used pharmacological tools that increase neuronal activity in cultures of cortical neurons [25]. These tools have shown that increased neuronal activity can upregulate neuronal antioxidant genes, in an ERK-AP-1 dependent manner. A putative antioxidant target gene of this signalling pathway, is *Sulfiredoxin-1* (*Srxn-1*) [26]. To identify whether JNK-AP-1 signalling could regulate neuronal antioxidant genes, we compared our experiments with neurons treated with Bicuculline (Bic, 50 μ M) and 4-Aminopyridine (4-AP, 500 μ M) as a positive control (Bic/4AP).

Using DEM, we found that JNK activity regulates *Srxn-1* transcription. DEM (10 μ M) induced a 1.73-fold induction in *Srxn-1* mRNA expression compared to the ethanol treated controls (Fig. 2C). Similar levels of *Srxn-1* mRNA were observed at higher concentrations of DEM (Fig. S4C). When neurons were pre-treated with the JNK inhibitor SU 3327 for 24 h, DEM failed to induce *Srxn-1* mRNA. Similar to DEM treatment, increasing neuronal activity with Bic/4AP increased *Srxn-1* mRNA (1.32-fold), however this induction was unaffected by JNK inhibition. Synaptic activity is known to induce expression of *c-Fos*, an AP-1 component. We determined whether oxidative stress also induced *c-Fos* transcription. Fig. 2D shows that there was no change in *c-Fos* mRNA levels with DEM treatment whereas Bic/4AP treatment robustly induced *c-Fos* transcription. Similar to activity-induced *Srxn-1* induction, *c-Fos* induction by synaptic activity was unaffected by inhibition of JNK (Fig. 2D) suggesting that synaptic activity and oxidative stress recruit signalling pathways differentially to activate potentially divergent gene transcription responses.

We next evaluated whether JNK activity regulates SRXN-1 protein




Fig. 3. JNK Recruits c-Jun for SRXN-1 expression.

A, B Oxidative stress reduces c-Fos and increases c-Jun. Representative images showing c-Fos (top panels) and c-Jun (bottom panels) immunoreactivity in neurons left untreated (Control) or treated for 1 h with 100 μ M DEM. Neurons were co-stained for the neuronal marker NeuN and nuclei were labelled with DAPI. **B** Quantification of immunoreactivity in **A**, of neurons stimulated for 1 h or 4 h with 100 μ M DEM. Data are plotted as mean \pm SEM. Analysis represents NeuN positive neurons. c-Fos levels were assessed in 118 neurons and c-Jun immunoreactivity was quantified in 81 neurons across 3 biological replicates (** $p < 0.01$; *** $p < 0.001$ one-way ANOVA followed by a Dunnett's post-hoc test). Scale bar = 25 μ m. **C** Representative Western blot of neuronal lysates immunoprecipitated with a c-Jun antibody in conditions of synaptic activity (Bic/4AP) or oxidative stress (DEM 100 μ M) in the presence or absence of the JNK inhibitor SU 3327 (700 nM, 1 h). **D** Representative images of c-Jun immunofluorescence in HEK293T cells transduced with lentivirus encoding empty vector, or pLenti c-Jun. Scale bar = 25 μ m. **E** Western blot showing c-Jun, GAPDH and SRXN-1 levels in neurons transduced with lentivirus containing empty vector (1) or c-Jun (3) overnight, or treated with the Nrf2 agonist, tert-Butylhydroquinone (TBHQ, 10 μ M, 4) overnight. HEK293T cells transduced with pLenti c-Jun (2) were run as a positive control. **F** A schematic demonstrating JNK and c-Jun dependent regulation of SRXN-1 during oxidative stress.

expression. **Fig. 2E** shows that treatment of neurons with DEM (100 μ M) rapidly increased SRXN-1 protein expression within 30 min, with high levels detected at 4 h as assessed by Western blot using an antibody that we first validated by overexpressing human SRXN-1 in HEK293T cells (**Fig S4A**). DEM-induced SRXN-1 expression was attenuated when neurons were pre-treated with SU 3327 (700 nM, 1 h pre-incubation). As expected, and in agreement with previous work [26], synaptic activity also induced SRXN-1 expression after a 4 h treatment with Bic/4AP, which was attenuated when neurons were pre-treated for 1 h with the non-competitive NMDA receptor antagonist MK801 (10 μ M) but was unaffected by JNK inhibition. Moreover, synaptic activity but not DEM, induced ERK activation (**Fig. 2E**). Taken together, these data indicate that oxidative stress activates JNK to coordinate an antioxidant response in neurons, which is distinct from antioxidant responses coordinated by synaptic activity.

2.4. JNK-c-Jun signalling regulates antioxidant responses in mammalian neurons

To identify the cellular mediators that co-ordinate SRXN-1 expression under conditions of synaptic activity or oxidative stress, we assessed the expression and interaction of putative AP-1 components under the two conditions. Cortical neurons were treated with Bic/4AP or DEM for 1 or 4 h and c-Fos or c-Jun immunofluorescence was quantified in NeuN positive neurons. In neurons treated with DEM (**Fig. 3A** and **B**) c-Fos

immunofluorescence was found to decrease. In contrast, DEM induced a significant increase in c-Jun immunofluorescence. As expected, in neurons stimulated with Bic/4AP for 1 or 4 h, both c-Fos and c-Jun immunofluorescence was found to increase (**Figs. S4, B,D**). These data suggest that oxidative stress relies on Jun proteins to mount an antioxidant response in the absence of c-Fos. To further investigate the differential involvement of specific AP-1 components we immunoprecipitated c-Jun from cortical neurons treated for 1 h with either DEM or Bic/4AP and assessed its phosphorylation and association with c-Fos protein. While both DEM and Bic/4AP induce c-Jun expression and phosphorylation, c-Jun co-immunoprecipitated with c-Fos following Bic/4AP-induced synaptic activity but not during oxidative stress mediated by DEM (**Fig. 3C**). In mammals, c-Jun can form heterodimers with other members of the Jun family to form the dimeric AP-1 transcription factor. To identify if c-Jun was associated with another Jun family member, we probed membranes for JunB and JunD and found no association between either protein with c-Jun in DEM-treated neurons (**Fig. S4E**). To determine whether c-Jun alone is sufficient to induce SRXN-1 expression, lentiviral particles encoding human c-Jun (**Fig. 3D**) were transduced into cortical neurons. Lentiviral c-Jun expression was sufficient to induce SRXN-1 expression in neurons (**Fig. 3E**), however activation of Nuclear Factor Erythroid 2-related factor 2 (Nrf2) using the well characterized agonist TBHQ (10 μ M, 16 h) had no effect on neuronal expression of SRXN-1. Taken together, these data suggest that oxidative stress activates JNK, which selectively recruits c-Jun to induce




Fig. 4. SRXN-1 localises to synaptic terminals in rodent neurons and rescues DEM induced retraction

A SRXN-1 overexpression rescues DEM-induced dendritic retraction. Representative micrographs of mature neurons transfected with PSD95-GFP constructs alone (left panels) or in combination with Flag-tagged human SRXN-1 (right panels) \pm 100 μ M DEM (48hr). Cells stained with anti-GFP (green), anti SRXN-1 (Flag antibody, red) and nuclear staining with DAPI (blue). Scale bar = 100 μ m. **B** Overexpressed human SRXN-1 co-localises with PSD95-GFP in dendritic spines. Representative images show an overlay of GFP and Flag immunofluorescence. Magnified images of dendritic spines outlined in white boxes (PSD95-GFP + SRXN-1, scale bar = 5 μ m). **C** Endogenous SRXN-1 protein is present at synapses. Representative Western blot of synaptosome fractions prepared from mouse brain, probed with antibodies for the nuclear protein FOXO3a, the pre-synaptic marker synapsin 1 (Syn1), the mitochondrial protein ATP5A1, JNK and SRXN-1 as indicated. **D** Schematic representation of likely signalling pathways mediating *Srxn-1* transcriptional induction in response to synaptic activity and oxidative stress. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

SRXN-1 expression in neurons (Fig. 3F).

2.5. SRXN-1 is localised at synaptic terminals and prevents dendrite loss

To understand how SRXN-1 contributes to neuronal antioxidant defenses we investigated SRXN-1 localization in neurons. Having previously identified that chronic treatment of cortical neurons with DEM causes a loss of dendrites, we transfected neurons with PSD95-GFP alone or in combination with a plasmid encoding N-terminal FLAG-tagged human SRXN-1 and treated cells with DEM (48 h). We found that overexpression of SRXN-1 prevented dendrite loss caused by long-term treatment of DEM (Fig. 4A). Interestingly, we also observed overexpressed SRXN-1 immunoreactivity in distal dendrites and on closer examination found that overexpressed SRXN-1 was localised to dendrites and dendritic spines (Fig. 4B). SRXN-1 co-localised with PSD-95-GFP, as

indicated by white arrowheads in Fig. 4B.

To further investigate the synaptic localization of SRXN-1, we assessed its distribution in subcellular fractions of mouse brain using the antibody validated in heterologous cells (Fig. S4A). Although characterized as a cytoplasmic enzyme [27], SRXN-1 was not observed in the synaptic supernatant fractions, but in the LP1 (synaptosomal) and LP2 (synaptic vesicle) membrane fractions. The efficacy of the fractionation was confirmed by enrichment of the transcription factor FOXO3a in supernatant fractions over the pellet fractions. The P2 fraction represents a crude synaptosomal fraction comprising microsomes and isolated pre- and post-synaptic nerve terminals. SRXN-1 was most enriched in the LP1 fraction, indicative of synaptic membranes and small organelles, such as mitochondria. Immunoblotting for the alpha subunit of the mitochondrial F1 ATP synthase (ATP5A1) confirmed the presence of mitochondrial proteins in the LP1 fraction. SRXN-1 immunoreactivity

was also present in the LP2 fraction, which is enriched in synaptic vesicles (SV), as shown by an enrichment of the SV protein synapsin I (Fig. 4C). JNK immunoreactivity was observed in all fractions, but was enriched in P2, LP1 and LP2, in-line with the distribution of SRXN-1. These data suggest that SRXN-1 associates with synaptic membranes and is consistent with the co-localization with PSD95 we observed by fluorescence microscopy. Therefore, in neurons, antioxidant proteins such as SRXN-1 and sensors of oxidative stress such as JNK are located at synapses, placing them at functionally important sites to orchestrate the adaptive antioxidant response (Fig. 4D).

3. Discussion

Here we have uncovered a role for JNK signalling in the regulation of neuronal antioxidant capacity using *Drosophila* and mammalian neurons in culture. We and others have shown previously that JNK regulates changes in synaptic morphology independently of any stimuli [11] or in response to ROS [5,12,14]. We now show that JNK activation drives an antioxidant response, which shapes cellular reductive capacity and synaptic morphology.

We show that *Drosophila* mutants that have enhanced JNK activity have low levels of ROS and furthermore, are resistant to chemically induced oxidative stress. These reduced ROS levels in mutant flies can be reinstated back to levels observed in wild type flies by inhibiting the JNK-AP-1 pathway. In addition to changes in ROS levels, we show that activation of the JNK-AP-1 pathway leads to overgrown synapses at the *Drosophila* NMJ suggesting that the status of JNK signalling has implications for structural synaptic plasticity. In mammalian cortical neurons we identified an 'adaptive' JNK-dependent antioxidant response when neurons are challenged with oxidative stress. This adaptive response relies on JNK's downstream effector c-Jun, and results in increased abundance of the antioxidant protein SRXN-1. In contrast, JNK-independent SRXN-1 expression is triggered with increased synaptic activity of cortical neurons. We therefore implicate JNK signalling in the regulation of neuronal antioxidant defenses and identify distinct differences in neuronal responses to ROS generated in conditions of synaptic activity and oxidative stress.

3.1. Constitutive JNK activation drives an antioxidant response in *Drosophila*

Genetic and pharmacological activation of JNK changes neuronal morphology and this structural change is dependent on AP-1 [15,28]. JNK-AP-1 signalling has long been known to positively regulate growth and strength at the larval NMJ [15]. JNK has also been shown to mediate the effect of ROS generated during oxidative stress associated with excitotoxicity [14] and lysosomal storage disease [12]. Combined with reports that activation of the JNK pathway after neuronal injury coordinates a regenerative response [16] and that AP-1 signalling regulates dendrite growth during both development and during conditions of synaptic activity; JNK-AP-1 signalling has been demonstrated as a crucial regulator of neuronal homeostasis. Both *puc*^{E69}/+ heterozygotes and *hiw* mutants, which have increased JNK activity [11,19], showed reduced total ROS levels and rendered flies resistant to chemically induced oxidative stress. The reduced ROS levels in *puc*^{E69}/+ heterozygotes and *hiw* mutants flies could be reversed by inhibition of neuronal JNK-AP-1, suggesting that the altered ROS originates from neurons. The reduced total ROS levels, arising from protective JNK signalling, in *hiw* mutants might explain the recently reported resistance of neurons in *highwire* null flies to the damaging effects of physical blows, which mimic traumatic brain injury [29].

A key question raised by our work is whether the JNK-dependent changes in synaptic morphology are driven by the increased antioxidant defenses. We have previously identified DEM as a regulator of neuronal morphology at the NMJ [5] and here we show that this DEM-induced change in synaptic morphology is regulated by JNK.

Furthermore, we show that *puc*^{E69}/+ larvae have overgrown NMJ's which are not influenced by oxidative stress induced by DEM. Given that *puc*^{E69}/+ and *hiw* larvae have elevated JNK activity, reduced ROS, elaborate NMJ's and are resistant to chemically induced oxidative stress, our data suggests that changes in neuronal plasticity in these models is at least, in part, driven by an antioxidant response.

3.2. JNK coordinates antioxidant responses in mammalian neurons

It is known that synaptic activity can regulate the expression of neuronal antioxidant genes [26]. Furthermore, we have recently shown that ROS generated from synaptic activity acts as a signalling molecule to regulate neuronal plasticity, in a PI3K/DJ-1 β dependent manner [5] implicating synaptic activity and ROS in the regulation of synaptic plasticity. What is currently unknown is how neurons recognize and respond to differences in physiological and pathological ROS. Our data shows that neurons recruit different signalling pathways to mediate ROS generated by synaptic activity or oxidative stress, but converge to regulate the same antioxidant gene, *Srxn-1*.

Increasing synaptic activity activates ERK1/2 [30,31] that can in turn phosphorylate AP-1 components [32,33]. Consistent with this, we found that increasing neuronal activity using Bicuculline and 4-AP regulates *Srxn-1* mRNA and protein expression [26]. Conversely, inducing oxidative stress in neurons with DEM did not affect ERK phosphorylation of AP-1 but did activate JNK and SRXN-1 expression. JNK inhibition abolished increased SRXN-1 in conditions of oxidative stress but not during synaptic activity. This suggests that neurons use distinct kinases to upregulate SRXN-1 expression in response to synaptic activity or oxidative stress.

Synaptic activity has been demonstrated to boost glutathione synthesis, providing a positive feedback loop, where accumulating levels of ROS from increased synaptic activity are neutralized by increases in cellular glutathione [26]. JNK activity is also regulated by cellular glutathione levels. In non-stressed cells, JNK is tethered to monomeric glutathione S-transferase Pi (GST-Pi) and increased H₂O₂ can trigger the detachment and oligomerization of GST-Pi, conjugating ROS to glutathione [34]. This suggests that JNK activity is tuned to the redox thresholds of the cell [35]. This is important given that glutathione homeostasis is altered in many neurodegenerative diseases [60,61] and decreased glutathione levels have been observed in mouse models of Alzheimer's Disease [62] and in AD patients [63].

Both ERK1/2 and JNK kinases regulate AP-1 [21,32], raising the possibility that the mechanisms converge at the level of this dimeric transcription factor. The AP-1 heterodimer of c-Fos and c-Jun [36] is well characterized but the conditions driving differential heterodimer composition are not well understood. In our work, a notable difference between synaptic activity and oxidative stress conditions was the lack of c-Fos transcriptional/translational induction by DEM and reduced dimerization with c-Jun. This indicates that DEM-induced AP-1 activity and *Srxn1* transcription is c-Fos independent (Fig. 4C and D). The mammalian AP-1 complex can contain Fos, Jun, Fra, and ATF components and multiple hetero- and homo-dimeric partners are capable of binding to AP-1 DNA binding sites [37,38]. Our immunoprecipitation of c-Jun showed a clear association with c-Fos during conditions of increased neuronal activity, but this association was absent during conditions of oxidative stress. Moreover, total c-Jun increased during oxidative stress and we did not detect JunB or JunD in immunoprecipitates suggesting that c-Jun may homodimerise to generate an antioxidant response in neurons. Our data identify a clear divergence between physiological and stress-dependent signalling in neurons, where activation of JNK and lack of c-Fos defines the stress response.

3.3. SRXN-1 is present at the synapse

SRXN-1 is an oxidoreductase that reduces sulfenylated proteins such as 2-cys peroxiredoxins (PRDX), small redox sensitive proteins widely

expressed in mammalian neurons [39]. Both cysteine residues in PRDX can undergo reversible hyper-oxidation by peroxides to form sulfinic acid (Cys-SO₂H) [40]. Hyper-oxidation of PRDX proteins inactivates their peroxidase activity. This inactivation is known to occur in neurons where it can be seen to follow a circadian time course [41], and occur during glutamate induced excitotoxicity [42] in addition to oxidative stress conditions [43]. SRXN-1 catalyses the reduction of Cys-SO₂H in an ATP dependent manner, restoring PRDX activity [44]. Both PRDX and SRXN-1 are regulated by synaptic activity [26]. Our data indicate that the cellular levels of SRXN-1 can also be tuned by the intracellular concentration of GSH and H₂O₂ levels.

We additionally show that SRXN-1 can be found at cellular locations that are particularly vulnerable to ROS - synaptic terminals and dendrites [45]. Overexpressed human SRXN-1 localised to dendrites and dendritic spines and endogenous SRXN-1 protein was particularly enriched in synaptosomal membrane fractions. We also found that JNK kinases were present in synaptosomes. The localization of SRXN-1, a cytosolic protein, in a synaptic membrane fraction is intriguing. However, in non-neuronal cells, SRXN-1 has been reported to translocate from the cytosol to a 'heavy membrane' fraction enriched in mitochondria in response to oxidative stress [46]. Mitochondrial SRXN-1 has a role in re-activation of oxidized peroxiredoxins [40,47] and our data indicate that SRXN-1 might reside constitutively with neuronal mitochondria, consistent with a greater basal ROS burden in neurons.

Taken together, our data from both fly and mammalian models identify a fundamental conserved role of JNK signalling in regulating neuronal redox homeostasis through an adaptive antioxidant response. JNK signalling is activated specifically by oxidative stress and activates c-Jun containing AP-1 transcription factors independently of c-Fos. In this manner, JNK imparts mechanistic specificity to activate AP-1 under conditions that have relevance to a number of neurodegenerative diseases.

4. Materials and methods

4.1. Culture of neurons and cell lines

Timed-mated female Wistar rats (Charles River UK) (RRID: RGD_737929) were maintained in accordance with the UK Animals (Scientific Procedures) Act (1986). Cortices were dissected from post-natal day 1 (P1) mixed sex rat pups. Animals were euthanised using pentobarbital injection followed by cervical dislocation, according to Home Office guidelines. Cortical cell suspensions were obtained as previously described and cytosine arabinoside (AraC, 2.4 μM final concentration) was added to the growth medium at 1 day *in vitro* (DIV) [48].

Neurons were transfected at 12 DIV with PSD95-GFP (a kind gift from David Bredt [49]) and FLAG-tagged Human SRXN-1 (purchased from Origene: RC207654) for 5 h using Lipofectamine 2000 (11668019, Thermo Scientific). Experiments were performed in a defined culture medium (Transfection medium (TM) contains: 90 ml SGG; 114 mM NaCl, 26.1 mM NaHCO₃, 5.3 mM KCl, 1 mM MgCl₂, 2 mM CaCl₂, 10 mM HEPES, 1 mM glycine, 30 mM glucose, 0.5 mM C₃H₃NaO₃ and 10 ml MEM (51200046, Thermo Scientific) supplemented with 100x ITS (1x final concentration; insulin/transferrin/selenium, 41400045, Thermo Scientific), penicillin (50 U/ml) and streptomycin (50 μg/ml)). After 24 h incubation in TM, cells were treated with DEM, bicuculline, catalase, paraquat, rotenone, 4-Aminopyridine or SU 3327 at concentrations and durations stated in the figure legends.

HEK293-FT (RRID:CVCL_6911, Thermo Scientific) cells were grown in DMEM containing 10% FBS. Cells were passaged and at 60% confluency, transfected with 10 μg of either pCMV entry vector (empty) or pCMV Human SRXN-1. After 3 days, cells were lysed as described below.

4.2. Lentiviral preparation

Lentiviruses were prepared using the 2nd generation system. Packaging plasmids were produced by the Trono lab (psPAX2 and pMD2.G were gifts from Didier Trono (Addgene plasmid #12260; <http://n2t.net/addgene:12260>; RRID:Addgene_12260 and Addgene plasmid #12259; <http://n2t.net/addgene:12259>; RRID:Addgene_12259). pLenti c-Jun was created by cloning human c-Jun from PMIEG3-c-Jun (a gift from Alexander Dent (Addgene plasmid #40348) [50] using BAMHI and XHOI into pLenti-puro (a gift from Ie-Ming Shih (Addgene plasmid #39481; <http://n2t.net/addgene:39481>; RRID:Addgene_39481)) [51]. Viral particles were generated and transduced as described previously [52].

4.3. Hydrogen peroxide assay

Experiments were performed in phenol red free TM. TM containing 50 μM Amplex Red reagent and 0.1 U/ml HRP was added to neurons in 12 well dishes (500 μl per well) for 15 min after which cells were treated with different concentrations of DEM, or catalase. Fluorescence at 590 nm was measured periodically and compared against a H₂O₂ standard curve for quantification. For fly experiments, Amplex red reagent was made up in Hemolymph solution (HL3) containing NaCl (128 mM), KCl (2 mM), CaCl₂·2H₂O (1.8 mM) MgCl₂·6H₂O (4 mM), HEPES (5 mM) and sucrose (35.5 mM) at pH 7.2. Each fly was crushed in 250 μl of Amplex red reagent using a pestle, vortexed and left at 25 °C, in the dark for 90 min. After 90 min, all samples were spun down for 1 min and each biological replicate was read in duplicate. Only males were used for amplex red assays to avoid issues with dosage compensation.

4.4. WST-1 assay

After treatment, primary rat neurons were washed with TM and then incubated with phenol red free TM containing WST-1 reagent (Sigma, Cat# 5015944001, 20 μl/ml) for 4 h at 37 °C. 200 μl was transferred to a 96 well plate and absorbance measured using a plate reader (BMG Fluostar λ = 440 nm).

4.5. Immunocytochemistry

Cells were washed with phosphate buffered saline (PBS) and fixed for 30 min at room temperature with 4% paraformaldehyde (containing 4% sucrose; Sigma). Cells were permeabilized in 0.5% NP40 in PBS for 5 min at room temperature. All primary and secondary antibodies, dilutions and suppliers can be found in Table 1. Primary antibodies were incubated overnight at 4 °C. Corresponding Alexafluor secondary antibodies (1:500, Thermo Scientific) were incubated for 1 h at room temperature before mounting with Fluoromount (Sigma).

4.6. Microscopy and image analysis

Images were collected on an inverted Zeiss microscope (880) with 20x or 63x Plan Apochromat objectives using Zeiss filter sets for DAPI and Alexa 488, 546 or 633. Images were taken at a resolution of 2048 × 2048 pixels.

4.7. Synaptosome preparation

The subcellular fractionation of mouse brain was performed as previously described [53]. Briefly, seven mixed sex mouse forebrains were dissected and transferred into cold homogenization buffer (320 mM sucrose, 1 mM EDTA, 5 mM Tris, pH 7.4, 4 °C), homogenized in a glass Teflon homogenizer, and centrifuged in an SS-34 rotor (1000g, 10 min, 4 °C). The supernatant (S1) was collected and the pellet re-suspended in 15 ml of homogenization buffer and centrifuged again (1000g for 10 min). S1 supernatants were combined, and pellets (P1) collected. S1

lysates were further centrifuged at 13000g for 20 min at 4 °C. Supernatant (S2) was collected and the pellet (P2) re-suspended in 21 ml homogenization buffer. The P2 pellet was further centrifuged at 13000g for 20 min at 4 °C. P2 was re-suspended in 320 mM sucrose (300 µl/forebrain), transferred into a glass-Teflon homogenizer with 9 vol of ice-cold distilled water and immediately homogenized for 3 up/down strokes at 2000 rpm. Homogenates were incubated on ice for 30 min with 0.1 volume of 1 M HEPES-NaOH (pH 7.4) and centrifuged at 25000g for 20 min at 4 °C. Pellets (LP1, synaptosomal membranes) were collected and the supernatant (LS1) further centrifuged in a 70 Ti rotor at 165,000g for 2 h at 4 °C. The resulting supernatant (LS2) was collected and the pellet (LP2, synaptic vesicles) re-suspended in 40 mM sucrose (2 ml). Protein concentration was assayed using Bradford reagent and 20 µg of each fraction was prepared in 1x Laemmli buffer for western blotting.

4.8. Western blotting

Cells were lysed in RIPA containing phosSTOP phosphatase inhibitors (4906845001, Roche) and complete EDTA-free protease inhibitors (04693132001, Roche) as previously described [54]. Lysates were run on Novex pre-cast mini gels (NuPAGE 4–12% Bis-Tris Gels, NP0322BOX, Thermo Scientific) in either 1 x MES or 1 x MOPS buffer. Antibodies used for immunoblotting are detailed in Table 1.

4.9. Co-immunoprecipitation

Neurons in 10 cm dishes were lysed as described. Lysates were incubated with primary antibodies for c-Jun at a 1:50 dilution and rotated overnight at 4 °C. Lysates were incubated with 15 µl (30 µl bead slurry) Protein G, Sepharose (GE Healthcare, 17-0618-01) for 2 h at 4 °C rotating. Samples were spun down, transferred to spin-x columns (Sigma, CLS8163-100 EA), washed 3 times with lysis buffer and the beads finally incubated with 4 x Laemmli (containing mercaptoethanol) for 10 min, then eluted through the spin columns. Samples were further heated at 95 °C for 5 min before western blotting. Proteins were detected using conformation specific antibodies (Cell Signalling, L27A9, mAb #3678) which prevent detection of heavy and light antibody chains.

4.10. Glutathione assay

Total levels of glutathione were assayed using a colorimetric Glutathione Assay Kit (CS0260, Sigma). Briefly, at 14 DIV, primary neurons in 35 mm dishes were treated with DEM in transfection medium for 1, 4 and 24 h. Cells were then washed with PBS (4 °C), lysed with 5% sulfosalicylic acid and snap frozen in liquid nitrogen. Lysates were then defrosted at 37 °C, centrifuged at 10,000g for 10 min and 10 µl samples were used for the glutathione assay in accordance with the manufacturer's instructions.

4.11. Quantitative PCR

Real-time quantitative PCR on the resulting cDNA was performed using a Fast SYBR Green Master mix (Applied Biosystems, 4385612) and gene specific primers. Relative expression of genes was determined by the 2- $\Delta\Delta$ CT method and Gapdh for normalization. Primers used were: Gapdh forward: 5'-AAACCCATCACCATCTTCCA-3' and Gapdh reverse: 5'-GTGGTTCACACCCATCACAA-3'; c-Fos forward: 5'-AGAATCCGAAGGAAAGGAA-3' and c-Fos reverse: 5'-ATTGAGAA-GAGGCAGGGTGA-3'; Srxn1 forward: 5'-GACGTCCTCTGGATCAAAG-3' and Srxn1 reverse: 5'-GCAGGAATGGTCTCTCTG-3'.

4.12. Drosophila stocks and husbandry

Drosophila were raised on 4–24® instant *Drosophila* medium (Carolina Biological Supply Company, USA) supplemented with a yeast

Table 1
Antibodies used in this study.

Name	Supplier	Dilution	Species	RRID
Anti- GFP	Synaptic Systems	1:1000	Guinea Pig	RRID: AB_11042617
Anti- Synapsin 1	MERCK	1:1000	Rabbit	RRID: AB_2200400
Anti- GCLC	SCBT	1:500	Mouse	RRID: AB_2736837
Anti- GCLM	SCBT	1:500	Mouse	RRID: AB_831789
Anti- Phospho JNK	Cell Signalling Technology	1:1000 (WB)	Mouse	RRID: AB_2307321
Anti SAPK/JNK Antibody	Cell Signalling Technology	1:1000 (WB)	Rabbit	RRID: AB_2250373
Anti FOXO3a	Cell Signalling Technology	1:1000 (WB)	Rabbit	RRID: AB_2636990
Anti- FLAG	Sigma	1:1000	Mouse	RRID: AB_259529
Anti- FLAG	Cell Signalling Technology	1:1000	Rabbit	RRID: AB_2217020
Anti- c-Jun	Cell Signalling Technology	1:1000 (WB)	Rabbit	RRID: AB_2130165
Anti- Phospho c-Jun	Cell Signalling Technology	1:1000 (WB)	Rabbit	RRID: AB_2130162
Anti- NeuN	Millipore	1:2000	Guinea Pig	RRID: AB_11205592
Anti- c-Fos	Cell Signalling Technology	1:2000 (WB)	Rabbit	RRID: AB_2106617
Anti- GAPDH	Millipore	1:10,000 (WB)	Mouse	RRID: AB_2107445
Anti- JunB	SCBT	1:1000 (WB)	Mouse	RRID: AB_2130023
Anti- JunD	SCBT	1:1000 (WB)	Mouse	RRID: AB_10650101
Anti- SRXN-1	SCBT	1:250 (WB)	Mouse	RRID: AB_2286615
Anti- ATP5A1	Proteintech	1:1000 (WB)	Rabbit	RRID: AB_2061761
Anti- p44/42 MAPK (Erk1/2)	Cell Signalling Technology	1:1000 (WB)	Rabbit	RRID: AB_330744
Anti- Phospho-p44/42 MAPK (Erk1/2)	Cell Signalling Technology	1:1000 (WB)	Mouse	RRID: AB_331768
Drosophila NMJ Antibodies				
Anti- Horseradish peroxidase- Cy3 (HRP-Cy3)	Jackson ImmunoResearch Labs	1:200	Goat	RRID: AB_2307391
Anti- synaptotagmin (Anti-SYT91)	Sweeney Lab	1:2000	Rabbit	RRID: AB_2713991
Secondary Ab's				
Gt anti Ms Alexa 488	Life Technologies	1:500	Goat	RRID: AB_2576217
Gt anti Rb Alexa 546	Life Technologies	1:500	Goat	RRID: AB_10563566
Gt anti Ms Alexa 546	Life Technologies	1:500	Goat	RRID: AB_2534071
Gt anti Rb Alexa 488	Life Technologies	1:500	Goat	RRID: AB_2576217
Peroxidase-AffiniPure Goat Anti-Guinea Pig IgG (H + L) antibody	Jackson ImmunoResearch Labs	1:10,000 (WB)	Goat	RRID: AB_2337402
Peroxidase-AffiniPure Goat Anti-Rabbit IgG	Jackson ImmunoResearch Labs	1:10,000 (WB)	Goat	RRID: AB_2307391
Peroxidase-AffiniPure Goat Anti-Mouse IgG (H + L) antibody	Jackson ImmunoResearch Labs	1:10,000 (WB)	Goat	RRID: AB_10015289
Fluorescein (FITC) AffiniPure Goat	Jackson ImmunoResearch Labs	1:200	Goat	RRID: AB_2337972

(continued on next page)

Table 1 (continued)

Name	Supplier	Dilution	Species	RRID
Anti-Rabbit IgG (H + L)				
IP Ab's				
Mouse Anti-rabbit IgG (Conformation Specific) (L27A9) mAb #3678	Cell Signalling Technology	1:5000 (WB)	Mouse	RRID: AB_1549606
Anti-mouse IgG, HRP-linked Antibody #7076	Cell Signalling Technology	1:10,000 (WB)	Horse	RRID: AB_330924

sucrose solution (5% w/v inactivated yeast, 10% w/v sucrose in ddH₂O, 100 g/500 ml) and maintained at 25 °C on a 12 h light:dark cycle. Prior to mixing with 4–24® instant media, vehicle (Ethanol) and DEM were added at the desired concentration (Ethanol 0.16%, DEM 0–10 mM). The following stocks were obtained from Bloomington *Drosophila* stock center: Canton-S (CS), *w¹¹¹⁸*, *UAS-bsk^{K53R}* (#9311), *UAS-fos^{DN}* (#7214), *UAS-jun^{DN}* (#7217), *Act5C-Gal4* (Actin-Gal4), *nSyb-Gal4*. *UAS-ask1^{K618M}* (*ask1^{DN}*) was a kind gift from Masayuki Miura [55]. *UAS-wnd^{DN}* (*wnd^{KD}*) and *hiw^{ND9}* were kind gifts from Aaron DiAntonio [11,56]. *puc^{E69}/TM6b* flies were obtained from Alfonso Martinez-Arias [57]. *SpinGal4/TM6b* was obtained from Daisuke Yamamoto [58]. The JNK dominant negative (*bsk^{K53R}*) [59] and the *highwire* mutant (*hiw^{ND9}*) have been described previously [18]. All analyses were performed in males. *hiw^{ND9}* females were crosses to Canton-S males to produce male *hiw^{ND9}* animals in an outcrossed background. For *hiw^{ND9}* crosses with dominant negative transgenes, *hiw^{ND9}*; *nsyb-Gal4* females were crossed with males carrying the UAS transgenes. All wild types were an outcross of Canton S to *w¹¹¹⁸*.

4.13. Survival analysis

Hatched 1st instar larvae were collected and raised in standard food with different DEM concentrations (0 mM, 1 mM, 5 mM and 10 mM). For each survival experiment, at least 2 vials, each containing 50 larvae, were maintained until fly eclosion. The number of eclosed flies was recorded.

4.14. Immunohistochemistry and NMJ analysis

Third instar wandering larvae were dissected, fixed, antibody stained, imaged and analysed as described previously [13]. All NMJ analysis was performed double-blind. Primary antibodies detailed in Table 1. Confocal microscopy was performed using a Zeiss LSM 880 on an Axio Observer.Z1 invert confocal microscope (Zeiss). Z-stacked projections of NMJ's and VNCs were obtained using a Plan Neofluar 40x/0.75 NA oil objective. NMJ lengths were measured from stacked NMJ images using the NeuronJ plugin for ImageJ (National Institutes of Health) as described previously [13,54].

4.15. Antibodies

Author contributions

C.U., S.C., and S.T.S. designed research; C.U., N.G., S.C., and L.F. performed research; G.J.O.E. contributed reagents/analytic tools; C.U., S.C., and L.F. analysed data; and C.U., S.C., and S.T.S. wrote the paper.

Declaration of competing interest

The authors declare no competing interests.

Acknowledgments

This work was supported by BBSRC Project Grants (BB/I012273/1) awarded to STS and (BB/M002322/1) awarded to STS and SC, and was part-funded by The Wellcome Trust [204829] awarded to CU, through the Centre for Future Health (CFH) at the University of York. We thank Addgene (Cambridge, MA, USA.) for providing plasmid stocks. We thank Lucy Rudd for technical assistance and the Bioscience Technology Facility at the University of York for providing access to confocal microscopes.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.redox.2020.101712>.

References

- [1] R. Dringen, Metabolism and functions of glutathione in brain, *Prog. Neurobiol.* 62 (2000) 649–671.
- [2] V.J. Milton, S.T. Sweeney, Oxidative stress in synapse development and function, *Develop. Neurobiol.* 72 (2012) 100–110.
- [3] P.S. Baxter, G.E. Hardingham, Adaptive regulation of the brain's antioxidant defences by neurons and astrocytes, *Free Radic. Biol. Med.* 100 (2016) 147–152.
- [4] K.J. Barnham, C.L. Masters, A.I. Bush, Neurodegenerative diseases and oxidative stress, *Nat. Rev. Drug Discov.* 3 (2004) 205.
- [5] M.C. Oswald, P.S. Brooks, M.F. Zwart, A. Mukherjee, R.J. West, C.N. Giachello, K. Morarach, R.A. Baines, S.T. Sweeney, M. Landgraf, Reactive oxygen species regulate activity-dependent neuronal plasticity in *Drosophila*, *eLife* 7 (2018), e39393.
- [6] G. Li, J. Gong, H. Lei, J. Liu, X.Z.S. Xu, Promotion of behavior and neuronal function by reactive oxygen species in *C. elegans*, *Nat. Commun.* 7 (2016), 13234.
- [7] M.C.W. Oswald, N. Garnham, S.T. Sweeney, M. Landgraf, Regulation of neuronal development and function by ROS, *FEBS Lett.* 592 (2018) 679–691.
- [8] M.T. Lin, M.F. Beal, Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases, *Nature* 443 (2006) 787.
- [9] J. Zhang, X. Wang, V. Vikash, Q. Ye, D. Wu, Y. Liu, W. Dong, ROS and ROS-mediated cellular signaling, *Oxid. Med. Cell. Longev.* 2016 (2016), 4350965.
- [10] W. Shen, B. Ganetzky, Autophagy promotes synapse development in *Drosophila*, *J. Cell Biol.* 187 (2009) 71–79.
- [11] C.A. Collins, Y.P. Wairkar, S.L. Johnson, A. DiAntonio, Highwire restrains synaptic growth by attenuating a MAP kinase signal, *Neuron* 51 (2006) 57–69.
- [12] V.J. Milton, H.E. Jarrett, K. Gowers, S. Chalak, L. Briggs, I.M. Robinson, S. T. Sweeney, Oxidative stress induces overgrowth of the *Drosophila* neuromuscular junction, *Proc. Natl. Acad. Sci. U. S. A.* 108 (2011) 17521–17526.
- [13] R.J. West, Y. Lu, B. Marie, F.B. Gao, S.T. Sweeney, Rab8, POSH, and TAK1 regulate synaptic growth in a *Drosophila* model of frontotemporal dementia, *J. Cell Biol.* 208 (2015) 931–947.
- [14] J.J. Peng, S.H. Lin, Y.T. Liu, H.C. Lin, T.N. Li, C.K. Yao, A circuit-dependent ROS feedback loop mediates glutamate excitotoxicity to sculpt the *Drosophila* motor system, *Elife* 8:e47372 (2019).
- [15] P.D. Etter, R. Narayanan, Z. Navratilova, C. Patel, D. Bohmann, H. Jasper, M. Ramaswami, Synaptic and genomic responses to JNK and AP-1 signaling in *Drosophila* neurons, *BMC Neurosci.* 6 (2005) 39.
- [16] Y. Hao, T.J. Waller, D.M. Nye, J. Li, Y. Zhang, R.I. Hume, M.M. Rolls, C.A. Collins, Degeneration of injured axons and dendrites requires restraint of a protective JNK signaling pathway by the transmembrane protein raw, *J. Neurosci.* 39 (2019) 8457–8470.
- [17] J. Yang, Z. Wu, N. Renier, D.J. Simon, K. Uryu, D.S. Park, P.A. Greer, C. Tournier, R.J. Davis, M. Tessier-Lavigne, Pathological axonal death through a MAPK cascade that triggers a local energy deficit, *Cell* 160 (2015) 161–176.
- [18] H.I. Wan, A. DiAntonio, R.D. Fetter, K. Bergstrom, R. Strauss, C.S. Goodman, Highwire regulates synaptic growth in *Drosophila*, *Neuron* 26 (2000) 313–329.
- [19] E. Martín-Blanco, A. Gampel, J. Ring, K. Virdee, N. Kirov, A.M. Tolkovsky, A. Martinez-Arias, Puckered encodes a phosphatase that mediates a feedback loop regulating JNK activity during dorsal closure in *Drosophila*, *Genes Dev.* 12 (1998) 557–570.
- [20] Y.W. Fridell, A. Sanchez-Blanco, B.A. Silvia, S.L. Helfand, Targeted expression of the human uncoupling protein 2 (hUCP2) to adult neurons extends life span in the fly, *Cell Metabol.* 1 (2005) 145–152.
- [21] E.T. Coffey, Nuclear and cytosolic JNK signalling in neurons, *Nat. Rev. Neurosci.* 15 (2014) 285.
- [22] W.S. Choi, G. Abel, H. Klintworth, R.A. Flavell, Z. Xia, JNK3 mediates paraquat- and rotenone-induced dopaminergic neuron death, *J. Neuropathol. Exp. Neurol.* 69 (2010) 511–520.

- [23] Y. Luo, H. Umegaki, X. Wang, R. Abe, G.S. Roth, Dopamine induces apoptosis through an oxidation-involved SAPK/JNK activation pathway, *J. Biol. Chem.* 273 (1998) 3756–3764.
- [24] E. Komulainen, J. Zdrojewska, E. Freemantle, H. Mohammad, N. Kuleskaya, P. Deshpande, F. Marchisella, R. Mysore, P. Hollos, K.A. Michelsen, et al., JNK1 controls dendritic field size in L2/3 and L5 of the motor cortex, constrains soma size, and influences fine motor coordination, *Front. Cell. Neurosci.* 8 (2014) 272.
- [25] G.E. Hardingham, F.J.L. Arnold, H. Bading, Nuclear calcium signaling controls CREB-mediated gene expression triggered by synaptic activity, *Nat. Neurosci.* 4 (2001) 261–267.
- [26] S. Papadia, F.X. Soriano, F. Leveille, M.A. Martel, K.A. Dakin, H.H. Hansen, A. Kaindl, M. Sifringer, J. Fowler, V. Stefovskaya, et al., Synaptic NMDA receptor activity boosts intrinsic antioxidant defenses, *Nat. Neurosci.* 11 (2008) 476–487.
- [27] T.S. Chang, W. Jeong, H.A. Woo, S.M. Lee, S. Park, S.G. Rhee, Characterization of mammalian sulfiredoxin and its reactivation of hyperoxidized peroxiredoxin through reduction of cysteine sulfenic acid in the active site to cysteine, *J. Biol. Chem.* 279 (2004) 50994–51001.
- [28] S. Sanyal, D.J. Sandstrom, C.A. Hoeffer, M. Ramaswami, AP-1 functions upstream of CREB to control synaptic plasticity in *Drosophila*, *Nature* 416 (2002) 870–874.
- [29] C.S. Hill, J. Sreedharan, A. Loreto, D.K. Menon, M.P. Coleman, Loss of highwire protects against the deleterious effects of traumatic brain injury in *Drosophila melanogaster*, *Front. Neurol.* 11 (2020), 401–401.
- [30] S.E. Hoey, R.J. Williams, M.S. Perkinson, Synaptic NMDA receptor activation stimulates α -secretase amyloid precursor protein processing and inhibits amyloid- β production, *J. Neurosci.* 29 (2009) 4442–4460.
- [31] G.E. Hardingham, F.J.L. Arnold, H. Bading, A calcium microdomain near NMDA receptors: on switch for ERK-dependent synapse-to-nucleus communication, *Nat. Neurosci.* 4 (2001) 565.
- [32] M. Dai, M. Feng, Y. Ye, X. Wu, D. Liu, M. Liao, W. Cao, Exogenous avian leukosis virus-induced activation of the ERK/AP1 pathway is required for virus replication and correlates with virus-induced tumorigenesis, *Sci. Rep.* 6 (2016) 19226.
- [33] M. Karin, The regulation of AP-1 activity by mitogen-activated protein kinases, *J. Biol. Chem.* 270 (1995) 16483–16486.
- [34] V. Adler, Z. Yin, S.Y. Fuchs, M. Benezra, L. Rosario, K.D. Tew, M.R. Pincus, M. Sardana, C.J. Henderson, C.R. Wolf, et al., Regulation of JNK signaling by GSTp, *EMBO J.* 18 (1999) 1321–1334.
- [35] M. Saitoh, H. Nishitoh, M. Fujii, K. Takeda, K. Tobiume, Y. Sawada, M. Kawabata, K. Miyazono, H. Ichijo, Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1, *EMBO J.* 17 (1998) 2596–2606.
- [36] K.F. Bell, B. Al-Mubarak, M.A. Martel, S. McKay, N. Wheelan, P. Hasel, N. Markus, P. Baxter, R.F. Deighton, A. Serio, et al., Neuronal development is promoted by weakened intrinsic antioxidant defences due to epigenetic repression of Nrf2, *Nat. Commun.* 6 (2015) 7066.
- [37] F. Mechta-Grigoriou, D. Gerald, M. Yaniv, The mammalian Jun proteins: redundancy and specificity, *Oncogene* 20 (2001) 2378–2389.
- [38] M. Karin, Z-g Liu, E. Zandi, AP-1 function and regulation, *Curr. Opin. Cell Biol.* 9 (1997) 240–246.
- [39] J. Goemaere, B. Knoops, Peroxiredoxin distribution in the mouse brain with emphasis on neuronal populations affected in neurodegenerative disorders, *J. Comp. Neurol.* 520 (2012) 258–280.
- [40] H.A. Woo, W. Jeong, T.S. Chang, K.J. Park, S.J. Park, J.S. Yang, S.G. Rhee, Reduction of cysteine sulfenic acid by sulfiredoxin is specific to 2-cys peroxiredoxins, *J. Biol. Chem.* 280 (2005) 3125–3128.
- [41] R.S. Edgar, E.W. Green, Y. Zhao, G. van Ooijen, M. Olmedo, X. Qin, Y. Xu, M. Pan, U.K. Valekunja, K.A. Feeney, et al., Peroxiredoxins are conserved markers of circadian rhythms, *Nature* 485 (2012) 459–464.
- [42] F. Léveillé, F.X. Soriano, S. Papadia, G.E. Hardingham, Excitotoxic insults lead to peroxiredoxin hyperoxidation, *Oxid. Med. Cell. Longev.* 2 (2009) 110–113.
- [43] P.S. Baxter, K.F. Bell, P. Hasel, A.M. Kaindl, M. Fricker, D. Thomson, S.P. Cregan, T. H. Gillingwater, G.E. Hardingham, Synaptic NMDA receptor activity is coupled to the transcriptional control of the glutathione system, *Nat. Commun.* 6 (2015) 6761.
- [44] S.G. Rhee, W. Jeong, T.S. Chang, H.A. Woo, Sulfiredoxin, the cysteine sulfenic acid reductase specific to 2-Cys peroxiredoxin: its discovery, mechanism of action, and biological significance, *Kidney Int.* 72 (2007) S3–S8.
- [45] P. Hasel, S. McKay, J. Qiu, G.E. Hardingham, Selective dendritic susceptibility to bioenergetic, excitotoxic and redox perturbations in cortical neurons, *Biochim. Biophys. Acta* 1853 (2015) 2066–2076.
- [46] Y.H. Noh, J.Y. Baek, W. Jeong, S.G. Rhee, T.S. Chang, Sulfiredoxin translocation into mitochondria plays a crucial role in reducing hyperoxidized peroxiredoxin III, *J. Biol. Chem.* 284 (2009) 8470–8477.
- [47] C.R. Sunico, A. Sultan, T. Nakamura, N. Dolatabadi, J. Parker, B. Shan, X. Han, J. R. Yates 3rd, E. Masliah, R. Ambasadhan, et al., Role of sulfiredoxin as a peroxiredoxin-2 denitrosylase in human iPSC-derived dopaminergic neurons, *Proc. Natl. Acad. Sci. U. S. A.* 113 (2016) E7564–E7571.
- [48] R. Suman, G. Smith, K.E. Hazel, R. Kasprowitz, M. Coles, P. O'Toole, S. Chawla, Label-free imaging to study phenotypic behavioural traits of cells in complex co-cultures, *Sci. Rep.* 6 (2016) 22032.
- [49] S.E. Craven, A.E. El-Husseini, D.S. Bredt, Synaptic targeting of the postsynaptic density protein PSD-95 mediated by lipid and protein motifs, *Neuron* 22 (1999) 497–509.
- [50] Z.Y. Wang, H. Sato, S. Kusam, S. Sehra, L.M. Toney, A.L. Dent, Regulation of IL-10 gene expression in Th2 cells by Jun proteins, *J. Immunol.* 174 (2005) 2098–2105.
- [51] B. Guan, T.L. Wang, J. M. Shih, ARID1A, a factor that promotes formation of SWI/SNF-mediated chromatin remodeling, is a tumor suppressor in gynecologic cancers, *Canc. Res.* 71 (2011) 6718–6727.
- [52] Barde I, Salmon P, Trono D *Production And Titration of Lentiviral Vectors*. In Gerfen CR, Holmes A, Sibley D, Skolnick P, Wray S (eds) pp Unit 4.21-Unit 4.21. John Wiley & Sons.
- [53] G.J.O. Evans, Subcellular fractionation of the brain: preparation of synaptosomes and synaptic vesicles, *Cold Spring Harb. Protoc.* 5 (2015) 462–466.
- [54] R.J.H. West, C. Ugbo, F.B. Gao, S.T. Sweeney, The pro-apoptotic JNK scaffold POSH/SH3RF1 mediates CHMP2B/intron5-associated toxicity in animal models of frontotemporal dementia, *Hum. Mol. Genet.* 27 (2018) 1382–1395.
- [55] E. Kuranaga, H. Kanuka, T. Igaki, K. Sawamoto, H. Ichijo, H. Okano, M. Miura, Reaper-mediated inhibition of DIAP1-induced DTRAF1 degradation results in activation of JNK in *Drosophila*, *Nat. Cell Biol.* 4 (2002) 705–710.
- [56] C. Wu, R.W. Daniels, A. DiAntonio, DfSn collaborates with Highwire to down-regulate the Wallenda/DLK kinase and restrain synaptic terminal growth, *Neural Dev.* 2 (2007) 16.
- [57] J.M. Ring, A. Martinez Arias, puckered, a gene involved in position-specific cell differentiation in the dorsal epidermis of the *Drosophila* larva, *Dev. Suppl.* (1993) 251–259.
- [58] Y. Nakano, K. Fujitani, J. Kurihara, J. Ragan, K. Usui-Aoki, L. Shimoda, T. Lukacsovich, K. Suzuki, M. Sezaki, Y. Sano, et al., Mutations in the novel membrane protein spinster interfere with programmed cell death and cause neural degeneration in *Drosophila melanogaster*, *Mol. Cell Biol.* 21 (2001) 3775–3788.
- [59] U. Weber, N. Paricio, M. Mlodzik, Jun mediates Frizzled-induced R3/R4 cell fate distinction and planar polarity determination in the *Drosophila* eye, *Development* 127 (2000) 3619–3629.
- [60] K. Aoyama, T. Nakaki, Impaired glutathione synthesis in neurodegeneration, *Int. J. Mol. Sci.* 14 (2013) 21021–21044.
- [61] F. Gu, V. Chauhan, A. Chauhan, Glutathione redox imbalance in brain disorders, *Curr. Opin. Clin. Nutr. Metab. Care* 18 (2015) 89–95.
- [62] R. Resende, P.I. Moreira, T. Proenca, A. Deshpande, J. Busciglio, C. Pereira, C. R. Oliveira, Brain oxidative stress in a triple-transgenic mouse model of Alzheimer disease, *Free Radic. Biol. Med.* 44 (2008) 2051–2057.
- [63] P.K. Mandal, S. Saharan, M. Tripathi, G. Murari, Brain glutathione levels – A novel biomarker for mild cognitive impairment and Alzheimer's disease, *Biol. Psychiatr.* 78 (2015) 702–710.