

## PRIMER NOTE

# Isolation of microsatellite loci in the Capricorn silvereye, *Zosterops lateralis chlorocephalus* (Aves: Zosteropidae)

F. D. FRENTIU,\*† C. L. LANGE,‡ T. BURKE† and I. P. F. OWENS§

\*Department of Zoology and Entomology, University of Queensland, St. Lucia 4072, Australia, †Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, UK, ‡Centre for Identification and Diagnostics, University of Queensland, St. Lucia 4072, Australia, §Department of Biology, Imperial College at Silwood Park, Berkshire, SL5 7PY, UK

## Abstract

The Capricorn silvereye (*Zosterops lateralis chlorocephalus*) is ideally suited to investigating the genetic basis of body size evolution. We have isolated and characterized a set of microsatellite markers for this species. Seven out of 11 loci were polymorphic. The number of alleles detected ranged from two to five and observed heterozygosities between 0.12 and 0.67. One locus, ZL49, was found to be sex-linked. This moderate level of diversity is consistent with that expected in an isolated, island population.

*Keywords:* microsatellite, passerine, silvereye, white-eye, Zosteropidae, *Zosterops*

Received 24 March 2003; revision received 14 May 2003; accepted 22 May 2003

The Capricorn silvereye (*Zosterops lateralis chlorocephalus*) has evolved to be up to 40% larger than its ancestor, the continental race *Z. l. familiaris* (Degnan 1993). This species inhabits the islands of the southern Great Barrier Reef, Australia and, because its biology is well known, it provides an ideal population in which to investigate the evolution of body size shifts. The genetic basis of such morphological change can be investigated through new, molecular-marker assisted methods for the estimation of inheritance. Although a set of four polymorphic microsatellites already exists for this species (Degnan *et al.* 1999), additional markers are necessary for the resolution of relatedness amongst individuals required for the application of molecular-marker methods.

To construct a microsatellite library, DNA was extracted from the liver tissue of a single individual using a cetyltrimethyl ammonium bromide protocol (Graham *et al.* 1994). A genomic library was constructed and screened using a protocol modified from the method of Fisher & Bachmann (1998). Library construction involved the digestion of genomic DNA with *RsaI* restriction enzyme (Roche), the ligation of 21-mer (5'-CTCTTGCTTACGCGTGGACTA-3') and 25-mer (5' phosphate-TAGTCCACGCGTAAGCAAGAGCAC-3') *MluI* adaptors to the digested genomic DNA and amplification by polymerase chain reaction (PCR). The

amplification products were denatured and hybridized to 19 single-stranded biotin-labelled (5' end) oligonucleotides, which were then captured by Streptavidin-coated magnetic beads (Roche). The oligonucleotides had the following motifs: (CT)<sub>15'</sub>, (CA)<sub>20'</sub>, (CAT)<sub>14'</sub>, (ACA)<sub>14'</sub>, (CTG)<sub>10'</sub>, (ACT)<sub>14'</sub>, (AGA)<sub>14'</sub>, (CTA)<sub>14'</sub>, (CTT)<sub>14'</sub>, (GAC)<sub>10'</sub>, (CAG)<sub>10'</sub>, (AGC)<sub>8'</sub>, (ACC)<sub>8'</sub>, (AAC)<sub>8'</sub>, (TAT)<sub>8'</sub>, (CCCT)<sub>6'</sub>, (CTAG)<sub>6'</sub>, (GATA)<sub>6'</sub>, (GACA)<sub>6'</sub>. The enriched genomic products were eluted from the magnetic beads and amplified again using the *MluI* adaptors. The amplified products were restricted with *MluI* enzyme (Roche) to remove the *MluI* adaptors. Prepared products were then cloned into pUC19 vector (Stratagene) and transformed into Max Efficiency DH5 $\alpha$ <sup>TM</sup> competent bacterial cells (GibcoBRL). Two hundred clones were amplified with the pUC<sup>-</sup> and pUC<sup>+</sup> primers (5'-CAG-GAAACAGCTATGACC-3' and 5'-GTTTTCCCAGTCAC-GACG-3', respectively). Amplified products were sequenced with pUC<sup>-</sup> primer to detect microsatellite presence in clones. Clones containing microsatellite repeats and flanking regions suitable for primer design were sequenced in the reverse direction with the pUC<sup>+</sup> primer. Thirteen consensus sequences were submitted to the European Bioinformatics Institute's EMBL database (Accession nos AJ517993–AJ518005). The sequences were confirmed unique in the EMBL and GenBank databases using BLASTN 2.2.4 software (Altschul *et al.* 1997) and primer pairs were designed by eye. To reduce nonspecific amplification (Brownstein *et al.* 1996), a GTTT 'pigtail' was added to the

Correspondence: Francesca Frentiu. Fax: + 617 3365 1655; E-mail: ffrentiu@zen.uq.edu.au

**Table 1** Characterization of seven polymorphic microsatellite loci in the Capricorn silvereeye, *Zosterops lateralis chlorocephalus*

Locus	EMBL Accession no.	Repeat motif (5'-3')*	Primer sequence (5'-3') and 5' primer modification (underlined)	$T_a$ (°C)	MgCl <sub>2</sub> (mM)	$n$	No. of alleles	Obs. allele size range (bp)	$H_O$	$H_E$
ZL41	AJ517994	(GACA) <sub>7</sub>	F: <u>NED</u> -TATGAGAAAGTGTAAAGAAAGG R: <u>GTTT</u> -CGACCAAAGTGCAGTTATC	53.0	1.5	45	5	82–92	0.60†	0.69
ZL44	AJ517996	(CT) <sub>9</sub>	F: <u>HEX</u> -CTGTCCCTGCCTCTCATC R: ACCATGGCAGAGGCCACCAA	53.0	1.0	39	3	220–229	0.39	0.39
ZL45	AJ517997	(GACA) <sub>5</sub>	F: <u>6-FAM</u> -CCGGAGCACCACGCACAGC R: <u>GTTT</u> -GGGTCCAAGCGCCTCGAG	57.5	1.0	49	2	108–116	0.12	0.19
ZL46	AJ517998	(GTT) <sub>7</sub>	F: GTCAGTGCCTGTGCTTTGAT R: <u>HEX</u> -AACCTGAAATTACACTTCT	54.2	1.5	52	4	127–150	0.42	0.46
ZL49	AJ518000	(GTCT) <sub>7</sub>	F: TGTCTCGTGCAAGATGG R: <u>NED-C</u> -TTGCAAGCTGATGTCTTAT	59.3	1.5	52	4	154–170	0.39‡	0.40
ZL50	AJ518001	(GCT) <sub>7</sub>	F: <u>NED-T</u> -AAGGTGCCGAGGTCCTGT R: <u>G</u> -TTTGCATGAGTGCATGCTGG	59.3	1.5	46	3	126–132	0.59	0.49
ZL54	AJ518005	(CA) <sub>13</sub>	F: <u>NED</u> -CACGACTTCTCAAGCAGAC R: GAGCCTTGCACAAACCGAC	57.5	1.5	51	4	118–126	0.67	0.62

\*From sequenced clone.

†Loci deviating significantly ( $P < 0.007$ ) from Hardy–Weinberg equilibrium. ‡Observed and expected heterozygosities calculated using males only.

$n$ , number of Heron Island individuals tested;  $T_a$ , annealing temperature;  $H_O$ , observed heterozygosity;  $H_E$ , expected heterozygosity.

5' end of ZL41R and ZL45R primers (Table 1). An extra base (C, T and G, respectively) was added to the 5' end of ZL49R, ZL50F and ZL50R (Table 1) to prevent these primers annealing to themselves.

Silvereeye DNA for PCR was extracted from blood using a salt precipitation method (Nicholls *et al.* 2000). PCR reactions were performed in 10- $\mu$ L volumes and contained 15 ng of DNA, 1  $\mu$ M of each primer, 0.2 mM of each dNTP, 0.25 U of DNA polymerase (Thermoprime<sup>Plus</sup>, ABgene), in the manufacturer's buffer [final concentrations 20 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 75 mM Tris-HCl pH 9.0, 0.01% (w/v) Tween], including 1.0 or 1.5 mM MgCl<sub>2</sub> (Table 1). PCR cycling was performed in a Hybaid Touchdown Thermocycler (Thermo Hybaid) and followed a pattern of initial denaturation for 3 min at 94 °C, followed by 33 cycles of 45 s at 94 °C, 45 s at annealing temperature ( $T_a$ , Table 1) and a final 20 s at 72 °C. PCR reaction conditions were optimized for all loci using both a thermal gradient of annealing temperatures (50–60 °C) and a magnesium chloride (MgCl<sub>2</sub>) concentration gradient (0.625 mM–2.5 mM per reaction). Eleven out of 13 primer sets successfully amplified products of the expected size. One primer in each pair was labelled with one of three fluorescent dyes: HEX<sup>TM</sup>, 6-FAM<sup>TM</sup> and NED<sup>TM</sup> (Applied Biosystems) (Table 1). PCR products were analysed using an ABI 377 DNA Sequencer (Applied Biosystems).

Products were initially tested for polymorphism on 20 putatively unrelated individuals from Heron Island, the largest population of Capricorn silvereeyes. Seven loci were polymorphic (Table 1) and four were monomorphic (further details are available on the Molecular Ecology Notes database: <http://snook.bio.indiana.edu/MENotes/home>).

Polymorphic loci were then characterized using larger sample sizes (39–52 individuals), also obtained from the Heron Island population. Observed and expected heterozygosities were calculated using the program CERVUS version 2.0 (Marshall *et al.* 1998). Departures from Hardy–Weinberg equilibrium and linkage among loci were tested using GENEPOP (<http://wbiomed.curtin.edu.au/genepop>; Raymond & Rousset 1995). Significant deviations from Hardy–Weinberg equilibrium were found at two loci, ZL41 and ZL49 ( $P$ -values  $\leq 0.000$ , Bonferroni  $\alpha = 0.007$ ,  $k = 7$ ), possibly indicating the presence of null alleles at ZL41. No loci were found to be in linkage disequilibrium, but an indication of sex linkage was found for ZL49 ( $P = 0.0019$ , Bonferroni  $\alpha = 0.0018$ ,  $k = 28$ ). In a sample of 52 individuals genotyped at this locus, and which had had their sex determined molecularly, all females (ZW) were hemizygous ( $n = 26$ ), with 62% of males (ZZ) being homozygous and 38% heterozygous ( $n = 16$  and  $n = 10$ , respectively). These results suggest that ZL49 is situated on the Z chromosome. Once females were removed from the sample, the locus was observed to be in Hardy–Weinberg equilibrium (Table 1).

Moderate to low levels of polymorphism were observed in this set of loci (2–5 alleles in 39–52 individuals). This low level of polymorphism is consistent with that expected from a small, isolated, island population. A similar pattern has been found previously, with this subspecies showing lower levels of genetic diversity than others in the *Zosterops lateralis* complex (Degnan 1993; Degnan *et al.* 1999). Although we have not tested the utility of these primers across many taxa, ZL44 and ZL45 amplify polymorphic loci in the house

sparrow, *Passer domesticus*, and the magpie, *Pica pica*, respectively (N. Ockendon and D. Galvez, personal communication, see the Sheffield Molecular Genetics Facility's primer database for details: <http://www.shef.ac.uk/misc/groups/molecol/birdmarkers.html>; Dawson *et al.* 2003). The primers we have presented here may prove to have higher levels of diversity in other passerine birds.

### Acknowledgements

The authors are grateful for funding assistance from the Australian Research Council, Natural Environment Research Council (UK) and a Walter and Eliza Hall Travelling Scholarship awarded by the University of Queensland to FDF. Andy Krupa and John Chittock (University of Sheffield) and Leon Scott (Centre for Identification and Diagnostics) provided technical assistance. This work was assisted by the NERC-funded Sheffield Molecular Genetics Facility, UK. Deborah A. Dawson, Lisa Pope, Fiona Manson and Michael Phillips provided comments on the manuscript.

### References

- Altschul SF, Madden TL, Schäffer AA, *et al.* (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, **25**, 3389–3402.
- Brownstein MJ, Carpten JD, Smith JR (1996) Modulation of non-templated nucleotide addition by Taq DNA Polymerase: Primer modifications that facilitate genotyping. *Biotechniques*, **20**, 1004–1010.
- Dawson D, Greig C, Burket (2003) Bird microsatellite primers cross-utility online database: <http://www.shef.ac.uk/misc/groups/molecol/birdmarkers.html>. University of Sheffield, UK.
- Degnan SM (1993) Genetic variability and population differentiation inferred from DNA fingerprinting in silvereyes (Aves: Zosteropidae). *Evolution*, **47**, 1105–1117.
- Degnan SM, Robertson BC, Clegg SM, Moritz CC (1999) Microsatellite primers for studies of gene flow and mating systems in white-eyes (Zosterops). *Molecular Ecology*, **8**, 157–168.
- Fisher D, Bachmann K (1998) Microsatellite enrichment in organisms with large genomes (*Allium cepa* L.). *Biotechniques*, **24**, 796–802.
- Graham GC, Mayers P, Henry RJ (1994) A simple and rapid method for the preparation of fungal genomic DNA for PCR and RAPD analysis. *Biotechniques*, **16**, 48–50.
- Marshall TC, Slate J, Kruuk L, Pemberton JM (1998) Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology*, **7**, 639–655.
- Nicholls JA, Double MC, Rouell DM, Magrath RD (2000) The evolution of cooperative and pair breeding in thornbills *Acanthiza* (Pardalotidae). *Journal of Avian Biology*, **31**, 165–176.
- Raymond M, Rousset F (1995) GENEPOP (Version 1.2): Population genetics software for exact tests and ecumenicism. *Journal of Heredity*, **86**, 248–249.